

$\Sigma(3170)$ Bumps

$I(J^P) = 1(?^?)$ Status: *

OMITTED FROM SUMMARY TABLE

Seen by AMIRZADEH 79 as a narrow 6.5-standard-deviation enhancement in the reaction $K^- p \rightarrow Y^{*+} \pi^-$ using data from independent high statistics bubble chamber experiments at 8.25 and 6.5 GeV/c. The dominant decay modes are multibody, multistrange final states and the production is via isospin-3/2 baryon exchange. Isospin 1 is favored.

Not seen in a $K^- p$ experiment in LASS at 11 GeV/c (ASTON 85B).

$\Sigma(3170)$ MASS (PRODUCTION EXPERIMENTS)

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
≈ 3170 OUR ESTIMATE				
3170 ± 5	35	AMIRZADEH 79	HBC	$K^- p \rightarrow Y^{*+} \pi^-$

$\Sigma(3170)$ WIDTH (PRODUCTION EXPERIMENTS)

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 20	35	¹ AMIRZADEH 79	HBC	$K^- p \rightarrow Y^{*+} \pi^-$

$\Sigma(3170)$ DECAY MODES (PRODUCTION EXPERIMENTS)

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \Lambda K \bar{K} \pi$'s	seen
$\Gamma_2 \quad \Sigma K \bar{K} \pi$'s	seen
$\Gamma_3 \quad \Xi K \pi$'s	seen

$\Sigma(3170)$ BRANCHING RATIOS (PRODUCTION EXPERIMENTS)

$\Gamma(\Lambda K \bar{K} \pi$'s)/$\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	AMIRZADEH 79 HBC $K^- p \rightarrow Y^{*+} \pi^-$
$\Gamma(\Sigma K \bar{K} \pi$'s)/$\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	AMIRZADEH 79 HBC $K^- p \rightarrow Y^{*+} \pi^-$

$\Gamma(\Xi K \pi's)/\Gamma_{total}$

Γ_3/Γ

VALUE

DOCUMENT ID

TECN

COMMENT

seen

AMIRZADEH 79 HBC $K^- p \rightarrow \Upsilon^{*+} \pi^-$

**$\Sigma(3170)$ FOOTNOTES
(PRODUCTION EXPERIMENTS)**

¹ Observed width consistent with experimental resolution.

**$\Sigma(3170)$ REFERENCES
(PRODUCTION EXPERIMENTS)**

ASTON	85B	PR D32 2270	+Carnegie+	(SLAC, CARL, CNRC, CINC)
AMIRZADEH	79	PL 89B 125	+	(BIRM, CERN, GLAS, MSU, CURIN, CAVE+) I
Also	80	Toronto Conf. 263	Kinson+	(BIRM, CERN, GLAS, MSU, CURIN) I