

# Quark and Lepton Compositeness, Searches for

A REVIEW GOES HERE – Check our WWW List of Reviews

## SCALE LIMITS for Contact Interactions: $\Lambda(eeee)$

Limits are for  $\Lambda_{LL}^\pm$  only. For other cases, see each reference.

$\Lambda_{LL}^+(TeV)$	$\Lambda_{LL}^-(TeV)$	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;3.1</b>	<b>&gt;3.8</b>	95	ABBIENDI	99	OPAL $E_{cm} = 130-136, 161-172,$ $183 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>2.7	>2.4	95	ACCIARRI	98J L3	$E_{cm} = 130-172 \text{ GeV}$
>3.0	>2.5	95	ACKERSTAFF	98v OPAL	$E_{cm} = 130-172 \text{ GeV}$
>2.4	>2.2	95	ACKERSTAFF	97C OPAL	$E_{cm} = 130-136, 161 \text{ GeV}$
>1.7	>2.3	95	<sup>1</sup> ARIMA	97 VNS	$E_{cm} = 57.77 \text{ GeV}$
>1.6	>2.0	95	<sup>2</sup> BUSKULIC	93Q ALEP	$E_{cm} = 88.25-94.25 \text{ GeV}$
>1.6		95	<sup>2,3</sup> BUSKULIC	93Q RVUE	
	>2.2	95	BUSKULIC	93Q RVUE	
	>3.6	95	<sup>4</sup> KROHA	92 RVUE	
>1.3		95	<sup>4</sup> KROHA	92 RVUE	
>0.7	>2.8	95	BEHREND	91C CELL	$E_{cm} = 35 \text{ GeV}$
>1.3	>1.3	95	KIM	89 AMY	$E_{cm} = 50-57 \text{ GeV}$
>1.4	>3.3	95	<sup>5</sup> BRAUNSCH...	88 TASS	$E_{cm} = 12-46.8 \text{ GeV}$
>1.0	>0.7	95	<sup>6</sup> FERNANDEZ	87B MAC	$E_{cm} = 29 \text{ GeV}$
>1.1	>1.4	95	<sup>7</sup> BARTEL	86C JADE	$E_{cm} = 12-46.8 \text{ GeV}$
>1.17	>0.87	95	<sup>8</sup> DERRICK	86 HRS	$E_{cm} = 29 \text{ GeV}$
>1.1	>0.76	95	<sup>9</sup> BERGER	85B PLUT	$E_{cm} = 34.7 \text{ GeV}$

<sup>1</sup>  $Z-Z'$  mixing is assumed to be zero.

<sup>2</sup> BUSKULIC 93Q uses the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

<sup>3</sup> This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

<sup>4</sup> KROHA 92 limit is from fit to BERGER 85B, BARTEL 86c, DERRICK 86B, FERNANDEZ 87B, BRAUNSCHWEIG 88, BEHREND 91B, and BEHREND 91C. The fit gives  $\eta/\Lambda_{LL}^2 = +0.230 \pm 0.206 \text{ TeV}^{-2}$ .

<sup>5</sup> BRAUNSCHWEIG 88 assumed  $m_Z = 92 \text{ GeV}$  and  $\sin^2\theta_W = 0.23$ .

<sup>6</sup> FERNANDEZ 87B assumed  $\sin^2\theta_W = 0.22$ .

<sup>7</sup> BARTEL 86C assumed  $m_Z = 93 \text{ GeV}$  and  $\sin^2\theta_W = 0.217$ .

<sup>8</sup> DERRICK 86 assumed  $m_Z = 93 \text{ GeV}$  and  $g_V^2 = (-1/2 + 2\sin^2\theta_W)^2 = 0.004$ .

<sup>9</sup> BERGER 85B assumed  $m_Z = 93 \text{ GeV}$  and  $\sin^2\theta_W = 0.217$ .

## SCALE LIMITS for Contact Interactions: $\Lambda(ee\mu\mu)$

Limits are for  $\Lambda_{LL}^\pm$  only. For other cases, see each reference.

$\Lambda_{LL}^+(TeV)$	$\Lambda_{LL}^-(TeV)$	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;4.5</b>	<b>&gt;4.3</b>	95	ABBIENDI	99	OPAL $E_{cm} = 130-136, 161-172,$ $183 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>3.6	>2.4	95	ACCIARRI	98J L3	$E_{cm} = 130\text{--}172 \text{ GeV}$
>2.9	>3.4	95	ACKERSTAFF	98V OPAL	$E_{cm} = 130\text{--}172 \text{ GeV}$
>3.1	>2.0	95	MIURA	98 VNS	$E_{cm} = 57.77 \text{ GeV}$
>2.4	>2.9	95	ACKERSTAFF	97C OPAL	$E_{cm} = 130\text{--}136, 161 \text{ GeV}$
>1.7	>2.2	95	<sup>10</sup> VELISSARIS	94 AMY	$E_{cm} = 57.8 \text{ GeV}$
>1.3	>1.5	95	<sup>10</sup> BUSKULIC	93Q ALEP	$E_{cm} = 88.25\text{--}94.25 \text{ GeV}$
>2.6	>1.9	95	<sup>10,11</sup> BUSKULIC	93Q RVUE	
>2.3	>2.0	95	HOWELL	92 TOPZ	$E_{cm} = 52\text{--}61.4 \text{ GeV}$
	>1.7	95	<sup>12</sup> KROHA	92 RVUE	
>2.5	>1.5	95	BEHREND	91C CELL	$E_{cm} = 35\text{--}43 \text{ GeV}$
>1.6	>2.0	95	<sup>13</sup> ABE	90I VNS	$E_{cm} = 50\text{--}60.8 \text{ GeV}$
>1.9	>1.0	95	KIM	89 AMY	$E_{cm} = 50\text{--}57 \text{ GeV}$
>2.3	>1.3	95	BRAUNSCH...	88D TASS	$E_{cm} = 30\text{--}46.8 \text{ GeV}$
>4.4	>2.1	95	<sup>14</sup> BARTEL	86C JADE	$E_{cm} = 12\text{--}46.8 \text{ GeV}$
>2.9	>0.86	95	<sup>15</sup> BERGER	85 PLUT	$E_{cm} = 34.7 \text{ GeV}$

<sup>10</sup> BUSKULIC 93Q and VELISSARIS 94 use the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

<sup>11</sup> This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

<sup>12</sup> KROHA 92 limit is from fit to BARTEL 86C, BEHREND 87C, BRAUNSCHWEIG 88D, BRAUNSCHWEIG 89C, ABE 90I, and BEHREND 91C. The fit gives  $\eta/\Lambda_{LL}^2 = -0.155 \pm 0.095 \text{ TeV}^{-2}$ .

<sup>13</sup> ABE 90I assumed  $m_Z = 91.163 \text{ GeV}$  and  $\sin^2\theta_W = 0.231$ .

<sup>14</sup> BARTEL 86C assumed  $m_Z = 93 \text{ GeV}$  and  $\sin^2\theta_W = 0.217$ .

<sup>15</sup> BERGER 85 assumed  $m_Z = 93 \text{ GeV}$  and  $\sin^2\theta_W = 0.217$ .

## SCALE LIMITS for Contact Interactions: $\Lambda(ee\tau\tau)$

Limits are for  $\Lambda_{LL}^\pm$  only. For other cases, see each reference.

$\Lambda_{LL}^+(TeV)$	$\Lambda_{LL}^-(TeV)$	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;3.8</b>	<b>&gt;4.0</b>	95	ABBIENDI	99	OPAL $E_{cm} = 130\text{--}136, 161\text{--}172,$ 183 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

>2.4	>2.8	95	ACCIARRI	98J L3	$E_{cm} = 130\text{--}172 \text{ GeV}$
>2.3	>3.7	95	ACKERSTAFF	98V OPAL	$E_{cm} = 130\text{--}172 \text{ GeV}$
>1.9	>3.0	95	ACKERSTAFF	97C OPAL	$E_{cm} = 130\text{--}136, 161 \text{ GeV}$
>1.4	>2.0	95	<sup>16</sup> VELISSARIS	94 AMY	$E_{cm} = 57.8 \text{ GeV}$
>1.0	>1.5	95	<sup>16</sup> BUSKULIC	93Q ALEP	$E_{cm} = 88.25\text{--}94.25 \text{ GeV}$
>1.8	>2.3	95	<sup>16,17</sup> BUSKULIC	93Q RVUE	
>1.9	>1.7	95	HOWELL	92 TOPZ	$E_{cm} = 52\text{--}61.4 \text{ GeV}$
>1.9	>2.9	95	<sup>18</sup> KROHA	92 RVUE	
>1.6	>2.3	95	BEHREND	91C CELL	$E_{cm} = 35\text{--}43 \text{ GeV}$
>1.8	>1.3	95	<sup>19</sup> ABE	90I VNS	$E_{cm} = 50\text{--}60.8 \text{ GeV}$
>2.2	>3.2	95	<sup>20</sup> BARTEL	86 JADE	$E_{cm} = 12\text{--}46.8 \text{ GeV}$

<sup>16</sup> BUSKULIC 93Q and VELISSARIS 94 use the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

- <sup>17</sup> This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.
- <sup>18</sup> KROHA 92 limit is from fit to BARTEL 86C BEHREND 89B, BRAUNSCHWEIG 89C, ABE 90I, and BEHREND 91C. The fit gives  $\eta/\Lambda_{LL}^2 = +0.095 \pm 0.120 \text{ TeV}^{-2}$ .
- <sup>19</sup> ABE 90I assumed  $m_Z = 91.163 \text{ GeV}$  and  $\sin^2\theta_W = 0.231$ .
- <sup>20</sup> BARTEL 86 assumed  $m_Z = 93 \text{ GeV}$  and  $\sin^2\theta_W = 0.217$ .

## SCALE LIMITS for Contact Interactions: $\Lambda(\ell\ell\ell\ell)$

Lepton universality assumed. Limits are for  $\Lambda_{LL}^\pm$  only. For other cases, see each reference.

$\Lambda_{LL}^+(\text{TeV})$	$\Lambda_{LL}^-(\text{TeV})$	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;5.2</b>	<b>&gt;5.3</b>	95	ABBIENDI	99	OPAL $E_{cm} = 130-136, 161-172,$ 183 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

- >4.0 >3.1 95 21 ACCIARRI 98J L3  $E_{cm} = 130-172 \text{ GeV}$   
 >3.4 >4.4 95 ACKERSTAFF 98V OPAL  $E_{cm} = 130-172 \text{ GeV}$   
 >2.7 >3.8 95 ACKERSTAFF 97C OPAL  $E_{cm} = 130-136, 161 \text{ GeV}$   
 >3.0 >2.3 95 21,22 BUSKULIC 93Q ALEP  $E_{cm} = 88.25-94.25 \text{ GeV}$   
 >3.5 >2.8 95 22,23 BUSKULIC 93Q RVUE  
 >2.5 >2.2 95 24 HOWELL 92 TOPZ  $E_{cm} = 52-61.4 \text{ GeV}$   
 >3.4 >2.7 95 25 KROHA 92 RVUE

21 From  $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$ , and  $\tau^+ \tau^-$ .

22 BUSKULIC 93Q uses the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

23 This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

24 HOWELL 92 limit is from  $e^+ e^- \rightarrow \mu^+ \mu^-$  and  $\tau^+ \tau^-$ .

25 KROHA 92 limit is from fit to most PEP/PETRA/TRISTAN data. The fit gives  $\eta/\Lambda_{LL}^2 = -0.0200 \pm 0.0666 \text{ TeV}^{-2}$ .

## SCALE LIMITS for Contact Interactions: $\Lambda(eeqq)$

Limits are for  $\Lambda_{LL}^\pm$  only. For other cases, see each reference.

$\Lambda_{LL}^+(\text{TeV})$	$\Lambda_{LL}^-(\text{TeV})$	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;4.4</b>	<b>&gt;2.8</b>	95	26 ABBIENDI	99	OPAL ( $eeqq$ )
<b>&gt;4.0</b>	<b>&gt;4.8</b>	95	27 ABBIENDI	99	OPAL ( $eebb$ )

• • • We do not use the following data for averages, fits, limits, etc. • • •

- >3.0 >2.1 95 28 ACCIARRI 98J L3 ( $eeqq$ )  
 >3.4 >2.2 95 29 ACKERSTAFF 98V OPAL ( $eeqq$ )

>4.0	>2.8	95	30	ACKERSTAFF	98V OPAL	( $e e b b$ )
>2.5	>3.7	95	31	ABE	97T CDF	( $e e q q$ ) (isosinglet)
>2.5	>2.1	95	32	ACKERSTAFF	97C OPAL	( $e e q q$ )
>3.1	>2.9	95	33	ACKERSTAFF	97C OPAL	( $e e b b$ )
>7.4	>11.7	95	34	DEANDREA	97 RVUE	$e e u u$ , atomic parity violation
>2.3	>1.0	95	35	AID	95 H1	( $e e q q$ ) ( $u, d$ quarks)
1.7	>2.2	95	36	ABE	91D CDF	( $e e q q$ ) ( $u, d$ quarks)
>1.2		95	37	ADACHI	91 TOPZ	( $e e q q$ ) (flavor-universal)
	>1.6	95	37	ADACHI	91 TOPZ	( $e e q q$ ) (flavor-universal)
>0.6	>1.7	95	38	BEHREND	91C CELL	( $e e c c$ )
>1.1	>1.0	95	38	BEHREND	91C CELL	( $e e b b$ )
>0.9		95	39	ABE	89L VNS	( $e e q q$ ) (flavor-universal)
	>1.7	95	39	ABE	89L VNS	( $e e q q$ ) (flavor-universal)
>1.05	>1.61	95	40	HAGIWARA	89 RVUE	( $e e c c$ )
>1.21	>0.53	95	41	HAGIWARA	89 RVUE	( $e e b b$ )

26 ABBIENDI 99 limits are from  $e^+ e^- \rightarrow q\bar{q}$  cross section at 130–136, 161–172, 183 GeV.

27 ABBIENDI 99 limits are from  $R_b$  at 130–136, 161–172, 183 GeV.

28 ACCIARRI 98J limits are from  $e^+ e^- \rightarrow q\bar{q}$  cross section at  $E_{cm} = 130$ –172 GeV.

29 ACKERSTAFF 98V limits are from  $e^+ e^- \rightarrow q\bar{q}$  at  $E_{cm} = 130$ –172 GeV.

30 ACKERSTAFF 98V limits are from  $R_b$  measurements at  $E_{cm} = 130$ –172 GeV.

31 ABE 97T limits are from  $e^+ e^-$  mass distribution in  $\bar{p}p \rightarrow e^+ e^- X$  at  $E_{cm} = 1.8$  TeV.

32 ACKERSTAFF 97C limits are from  $e^+ e^- \rightarrow q\bar{q}$  cross section at  $E_{cm} = 130$ –136 GeV and 161 GeV.

33 ACKERSTAFF 97C limits are  $R_b$  measurements at  $E_{cm} = 133$  GeV and 161 GeV.

34 DEANDREA 97 limit is from atomic parity violation of cesium. The limit is eluded if the contact interactions are parity conserving.

35 AID 95 limits are from the  $Q^2$  spectrum measurement of  $ep \rightarrow eX$ .

36 ABE 91D limits are from  $e^+ e^-$  mass distribution in  $p\bar{p} \rightarrow e^+ e^- X$  at  $E_{cm} = 1.8$  TeV.

37 ADACHI 91 limits are from differential jet cross section. Universality of  $\Lambda(e e q q)$  for five flavors is assumed.

38 BEHREND 91C is from data at  $E_{cm} = 35$ –43 GeV.

39 ABE 89L limits are from jet charge asymmetry. Universality of  $\Lambda(e e q q)$  for five flavors is assumed.

40 The HAGIWARA 89 limit is derived from forward-backward asymmetry measurements of  $D/D^*$  mesons by ALTHOFF 83C, BARTEL 84E, and BARINGER 88.

41 The HAGIWARA 89 limit is derived from forward-backward asymmetry measurement of  $b$  hadrons by BARTEL 84D.

### SCALE LIMITS for Contact Interactions: $\Lambda(\mu\mu qq)$

$\Lambda_{LL}^+(TeV)$	$\Lambda_{LL}^-(TeV)$	CL%	DOCUMENT ID	TECN	COMMENT
>2.9	>4.2	95	42	ABE	97T CDF ( $\mu\mu qq$ ) (isosinglet)
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
>1.4	>1.6	95	ABE	92B CDF	( $\mu\mu qq$ ) (isosinglet)

<sup>42</sup> ABE 97T limits are from  $\mu^+ \mu^-$  mass distribution in  $\bar{p}p \rightarrow \mu^+ \mu^- X$  at  $E_{cm}=1.8$  TeV.

---

### SCALE LIMITS for Contact Interactions: $\Lambda(\ell\nu\ell\nu)$

VALUE (TeV)	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;3.10</b>	90	43 JODIDIO	86 SPEC	$\Lambda_{LR}^\pm(\nu_\mu \nu_e \mu e)$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
>3.8	44 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^+(\tau \nu_\tau e \nu_e)$	
>8.1	44 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^-(\tau \nu_\tau e \nu_e)$	
>4.1	45 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^+(\tau \nu_\tau \mu \nu_\mu)$	
>6.5	45 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^-(\tau \nu_\tau \mu \nu_\mu)$	
43 JODIDIO 86 limit is from $\mu^+ \rightarrow \bar{\nu}_\mu e^+ \nu_e$ . Chirality invariant interactions $L = (g^2/\Lambda^2)$ $[\eta_{LL} (\bar{\nu}_\mu L \gamma^\alpha \mu_L) (\bar{e}_L \gamma_\alpha \nu_e L) + \eta_{LR} (\bar{\nu}_\mu L \gamma^\alpha \nu_e L) (\bar{e}_R \gamma_\alpha \mu_R)]$ with $g^2/4\pi = 1$ and $(\eta_{LL}, \eta_{LR}) = (0, \pm 1)$ are taken. No limits are given for $\Lambda_{LL}^\pm$ with $(\eta_{LL}, \eta_{LR}) = (\pm 1, 0)$ . For more general constraints with right-handed neutrinos and chirality nonconserving contact interactions, see their text.				
44 DIAZCRUZ 94 limits are from $(\tau \rightarrow e \nu \nu)$ and assume flavor-dependent contact interactions with $\Lambda(\tau \nu_\tau e \nu_e) \ll \Lambda(\mu \nu_\mu e \nu_e)$ .				
45 DIAZCRUZ 94 limits are from $(\tau \rightarrow \mu \nu \nu)$ and assume flavor-dependent contact interactions with $\Lambda(\tau \nu_\tau \mu \nu_\mu) \ll \Lambda(\mu \nu_\mu e \nu_e)$ .				

---

### SCALE LIMITS for Contact Interactions: $\Lambda(qqqq)$

Limits are for  $\Lambda_{LL}^\pm$  with color-singlet isoscalar exchanges among  $u_L$ 's and  $d_L$ 's only.

See EICHEN 84 for details.

VALUE (TeV)	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;1.9</b>	95	46 ABBOTT	98G D0	$p\bar{p} \rightarrow$ dijet angl. $\Lambda_{LL}^+$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
>1.6	95	47 ABE	96 CDF	$p\bar{p} \rightarrow$ jets inclusive
>1.3	95	48 ABE	96S CDF	$p\bar{p} \rightarrow$ dijet angl.; $\Lambda_{LL}^+$
>1.4	95	49 ABE	93G CDF	$p\bar{p} \rightarrow$ dijet mass
>1.0	99	50 ABE	92D CDF	$p\bar{p} \rightarrow$ jets inclusive
>0.825	95	51 ABE	92M CDF	$p\bar{p} \rightarrow$ dijet angl.
>0.700	95	52 ALITTI	91B UA2	$p\bar{p} \rightarrow$ jets inclusive
>0.330	95	50 ABE	89 CDF	$p\bar{p} \rightarrow$ jets inclusive
>0.400	95	53 ABE	89H CDF	$p\bar{p} \rightarrow$ dijet angl.
>0.415	95	54 ARNISON	86C UA1	$p\bar{p} \rightarrow$ jets inclusive
>0.370	95	55 ARNISON	86D UA1	$p\bar{p} \rightarrow$ dijet angl.
>0.275	95	56 APPEL	85 UA2	$p\bar{p} \rightarrow$ jets inclusive
		57 BAGNAIA	84C UA2	Repl. by APPEL 85

46 ABBOTT 98G limit is from dijet angular distribution in  $p\bar{p}$  collisions at  $E_{cm}=1.8$  TeV.

47 ABE 96 finds that the inclusive jet cross section for  $E_T > 200$  GeV is significantly higher than the  $\mathcal{O}(\alpha_s^3)$  perturbative QCD prediction. This could be interpreted as the effect of a contact interaction with  $\Lambda_{LL} \sim 1.6$  TeV. However, ABE 96 state that uncertainty in the parton distribution functions, higher-order QCD corrections, and the detector calibration may possibly account for the effect.

- 48 ABE 96S limit is from dijet angular distribution in  $p\bar{p}$  collisions at  $E_{cm} = 1.8$  TeV. The limit for  $\Lambda_{LL}^-$  is  $> 1.4$  TeV. ABE 96S also obtain limits for flavor symmetric contact interactions among all quark flavors:  $\Lambda_{LL}^+ > 1.8$  TeV and  $\Lambda_{LL}^- > 1.6$  TeV.
- 49 ABE 93G limit is from dijet mass distribution in  $p\bar{p}$  collisions at  $E_{cm} = 1.8$  TeV. The limit is the weakest from several choices of structure functions and renormalization scale.
- 50 Limit is from inclusive jet cross-section data in  $p\bar{p}$  collisions at  $E_{cm} = 1.8$  TeV. The limit takes into account uncertainties in choice of structure functions and in choice of process scale.
- 51 ABE 92M limit is from dijet angular distribution for  $m_{dijet} > 550$  GeV in  $p\bar{p}$  collisions at  $E_{cm} = 1.8$  TeV.
- 52 ALITTI 91B limit is from inclusive jet cross section in  $p\bar{p}$  collisions at  $E_{cm} = 630$  GeV. The limit takes into account uncertainties in choice of structure functions and in choice of process scale.
- 53 ABE 89H limit is from dijet angular distribution for  $m_{dijet} > 200$  GeV at the Fermilab Tevatron Collider with  $E_{cm} = 1.8$  TeV. The QCD prediction is quite insensitive to choice of structure functions and choice of process scale.
- 54 ARNISON 86C limit is from the study of inclusive high- $p_T$  jet distributions at the CERN  $\bar{p}p$  collider ( $E_{cm} = 546$  and 630 GeV). The QCD prediction renormalized to the low- $p_T$  region gives a good fit to the data.
- 55 ARNISON 86D limit is from the study of dijet angular distribution in the range  $240 < m(\text{dijet}) < 300$  GeV at the CERN  $\bar{p}p$  collider ( $E_{cm} = 630$  GeV). QCD prediction using EHLQ structure function (EICHTEN 84) with  $\Lambda_{QCD} = 0.2$  GeV for the choice of  $Q^2 = p_T^2$  gives the best fit to the data.
- 56 APPEL 85 limit is from the study of inclusive high- $p_T$  jet distributions at the CERN  $\bar{p}p$  collider ( $E_{cm} = 630$  GeV). The QCD prediction renormalized to the low- $p_T$  region gives a good description of the data.
- 57 BAGNAIA 84C limit is from the study of jet  $p_T$  and dijet mass distributions at the CERN  $\bar{p}p$  collider ( $E_{cm} = 540$  GeV). The limit suffers from the uncertainties in comparing the data with the QCD prediction.

### SCALE LIMITS for Contact Interactions: $\Lambda(\nu\nu qq)$

Limits are for  $\Lambda_{LL}^\pm$  only. For other cases, see each reference.

$\Lambda_{LL}^+(TeV)$	$\Lambda_{LL}^-(TeV)$	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;5.0</b>	<b>&gt;5.4</b>	95	58 MCFARLAND 98	CCFR	$\nu N$ scattering

<sup>58</sup> MCFARLAND 98 assumed a flavor universal interaction. Neutrinos were mostly of muon type.

## MASS LIMITS for Excited $e$ ( $e^*$ )

Most  $e^+ e^-$  experiments assume one-photon or  $Z$  exchange. The limits from some  $e^+ e^-$  experiments which depend on  $\lambda$  have assumed transition couplings which are chirality violating ( $\eta_L = \eta_R$ ). However they can be interpreted as limits for chirality-conserving interactions after multiplying the coupling value  $\lambda$  by  $\sqrt{2}$ ; see Note.

Excited leptons have the same quantum numbers as other ortholeptons. See also the searches for ortholeptons in the "Searches for Heavy Leptons" section.

### Limits for Excited $e$ ( $e^*$ ) from Pair Production

These limits are obtained from  $e^+ e^- \rightarrow e^+ e^-$  and thus rely only on the (electroweak) charge of  $e^*$ . Form factor effects are ignored unless noted. For the case of limits from  $Z$  decay, the  $e^*$  coupling is assumed to be of sequential type. Possible  $t$  channel contribution from transition magnetic coupling is neglected. All limits assume  $e^* \rightarrow e\gamma$  decay except the limits from , ( $Z$ ).

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;85.0</b>	95	59 ACKERSTAFF 98C	OPAL	$e^+ e^- \rightarrow e^* e^*$ Homodoublet type
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
60 BARATE	98U	ALEP	$Z \rightarrow e^* e^*$	
>79.6	95	61,62 ABREU	97B DLPH	$e^+ e^- \rightarrow e^* e^*$ Homodoublet type
>77.9	95	61,63 ABREU	97B DLPH	$e^+ e^- \rightarrow e^* e^*$ Sequential type
>79.7	95	61 ACCIARRI	97G L3	$e^+ e^- \rightarrow e^* e^*$ Sequential type
>79.9	95	61,64 ACKERSTAFF	97 OPAL	$e^+ e^- \rightarrow e^* e^*$ Homodoublet type
>62.5	95	65 ABREU	96K DLPH	$e^+ e^- \rightarrow e^* e^*$ Homodoublet type
>64.7	95	66 ACCIARRI	96D L3	$e^+ e^- \rightarrow e^* e^*$ Sequential type
>66.5	95	66 ALEXANDER	96Q OPAL	$e^+ e^- \rightarrow e^* e^*$ Homodoublet type
>65.2	95	66 BUSKULIC	96W ALEP	$e^+ e^- \rightarrow e^* e^*$ Sequential type
>45.6	95	ADRIANI	93M L3	$Z \rightarrow e^* e^*$
>45.6	95	ABREU	92C DLPH	$Z \rightarrow e^* e^*$
>29.8	95	67 BARDADIN-...	92 RVUE	, ( $Z$ )
>26.1	95	68 DECOMP	92 ALEP	$Z \rightarrow e^* e^*; , (Z)$
>46.1	95	DECAMP	92 ALEP	$Z \rightarrow e^* e^*$
>33	95	68 ABREU	91F DLPH	$Z \rightarrow e^* e^*; , (Z)$
>45.0	95	69 ADEVA	90F L3	$Z \rightarrow e^* e^*$
>44.9	95	AKRAWY	90I OPAL	$Z \rightarrow e^* e^*$
>44.6	95	70 DECOMP	90G ALEP	$e^+ e^- \rightarrow e^* e^*$
>30.2	95	ADACHI	89B TOPZ	$e^+ e^- \rightarrow e^* e^*$
>28.3	95	KIM	89 AMY	$e^+ e^- \rightarrow e^* e^*$
>27.9	95	71 ABE	88B VNS	$e^+ e^- \rightarrow e^* e^*$

<sup>59</sup> From  $e^+ e^-$  collisions at  $\sqrt{s}=170-172$  GeV. ACKERSTAFF 98C also obtain limit from  $e^* \rightarrow \nu W$  decay mode:  $m_{e^*} > 81.3$  GeV.

<sup>60</sup> BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane.

<sup>61</sup> From  $e^+ e^-$  collisions at  $\sqrt{s}= 161$  GeV.

<sup>62</sup> ABREU 97B also obtain limit from charged current decay mode  $e^* \rightarrow \nu W$ ,  $m_{e^*} > 70.9$  GeV.

<sup>63</sup> ABREU 97B also obtain limit from charged current decay mode  $e^* \rightarrow \nu W$ ,  $m_{e^*} > 44.6$  GeV.

<sup>64</sup> ACKERSTAFF 97 also obtain limit from charged current decay mode  $e^* \rightarrow \nu W$ ,  $m_{\nu_e^*} > 77.1$  GeV.

<sup>65</sup> From  $e^+ e^-$  collisions at  $\sqrt{s} = 130$ –136 GeV.

<sup>66</sup> From  $e^+ e^-$  collisions at  $\sqrt{s} = 130$ –140 GeV.

<sup>67</sup> BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on  $\Delta, (Z) < 36$  MeV.

<sup>68</sup> Limit is independent of  $e^*$  decay mode.

<sup>69</sup> ADEVA 90F is superseded by ADRIANI 93M.

<sup>70</sup> Superseded by DECOMP 92.

<sup>71</sup> ABE 88B limits assume  $e^+ e^- \rightarrow e^+ e^-$  with one photon exchange only and  $e^* \rightarrow e\gamma$  giving  $ee\gamma\gamma$ .

## Limits for Excited $e$ ( $e^*$ ) from Single Production

These limits are from  $e^+ e^- \rightarrow e^* e$ ,  $W \rightarrow e^* \nu$ , or  $e p \rightarrow e^* X$  and depend on transition magnetic coupling between  $e$  and  $e^*$ . All limits assume  $e^* \rightarrow e\gamma$  decay except as noted. Limits from LEP, UA2, and H1 are for chiral coupling, whereas all other limits are for nonchiral coupling,  $\eta_L = \eta_R = 1$ . In most papers, the limit is expressed in the form of an excluded region in the  $\lambda - m_{e^*}$  plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT	
none	20–170	95	72 ACCIARRI	98T L3	$e\gamma \rightarrow e^* \rightarrow ee\gamma$
none	30–200	95	73 BREITWEG	97C ZEUS	$e p \rightarrow e^* X$
>89	95	ADRIANI	93M L3	$Z \rightarrow ee^*, \lambda_Z > 0.5$	
>88	95	ABREU	92C DLPH	$Z \rightarrow ee^*, \lambda_Z > 0.5$	
<b>&gt;91</b>	95	DECAMP	92 ALEP	$Z \rightarrow ee^*, \lambda_Z > 1$	
>87	95	AKRAWY	90I OPAL	$Z \rightarrow ee^*, \lambda_Z > 0.5$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

74	ACKERSTAFF	98C OPAL	$e^+ e^- \rightarrow ee^*$	
75	BARATE	98U ALEP	$e^+ e^- \rightarrow ee^*$	
76,77	ABREU	97B DLPH	$e^+ e^- \rightarrow ee^*$	
76,78	ACCIARRI	97G L3	$e^+ e^- \rightarrow ee^*$	
79	ACKERSTAFF	97 OPAL	$e^+ e^- \rightarrow ee^*$	
80	ADLOFF	97 H1	Lepton-flavor violation	
81	ABREU	96K DLPH	$e^+ e^- \rightarrow ee^*$	
82	ACCIARRI	96D L3	$e^+ e^- \rightarrow ee^*$	
83	ALEXANDER	96Q OPAL	$e^+ e^- \rightarrow ee^*$	
84	BUSKULIC	96W ALEP	$e^+ e^- \rightarrow ee^*$	
85	DERRICK	95B ZEUS	$e p \rightarrow e^* X$	
86	ABT	93 H1	$e p \rightarrow e^* X$	

>86	95	ADRIANI	93M L3	$\lambda_\gamma > 0.04$
	87	DERRICK	93B ZEUS	Superseded by DERRICK 95B
>86	95	ABREU	92C DLPH	$e^+ e^- \rightarrow ee^*, \lambda_\gamma > 0.1$
>88	95	88 ADEVA	90F L3	$Z \rightarrow ee^*, \lambda_Z > 0.5$
>86	95	88 ADEVA	90F L3	$Z \rightarrow ee^*, \lambda_Z > 0.04$
>81	95	89 DECAMP	90G ALEP	$Z \rightarrow ee^*, \lambda_Z > 1$
>50	95	ADACHI	89B TOPZ	$e^+ e^- \rightarrow ee^*, \lambda_\gamma > 0.04$
>56	95	KIM	89 AMY	$e^+ e^- \rightarrow ee^*, \lambda_\gamma > 0.03$
none 23–54	95	90 ABE	88B VNS	$e^+ e^- \rightarrow ee^*, \lambda_\gamma > 0.04$
>75	95	91 ANSARI	87D UA2	$W \rightarrow e^* \nu; \lambda_W > 0.7$
>63	95	91 ANSARI	87D UA2	$W \rightarrow e^* \nu; \lambda_W > 0.2$
>40	95	91 ANSARI	87D UA2	$W \rightarrow e^* \nu; \lambda_W > 0.09$

72 ACCIARRI 98T search for single  $e^*$  production in quasi-real Compton scattering. The limit is for  $|\lambda| > 1.0 \times 10^{-1}$  and non-chiral coupling of  $e^*$ . See their Fig. 7 for the exclusion plot in the mass-coupling plane.

73 BREITWEG 97C search for single  $e^*$  production in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ .  $f = -f' = 2\Lambda/m_{e^*}$  is assumed for the  $e^*$  coupling. See their Fig. 9 for the exclusion plot in the mass-coupling plane.

74 ACKERSTAFF 98C from  $e^+ e^-$  collisions at  $\sqrt{s} = 170$ –172 GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane.

75 BARATE 98U is from  $e^+ e^-$  collision at  $\sqrt{s} = M_Z$ . See their Fig. 12 for limits in mass-coupling plane.

76 From  $e^+ e^-$  collisions at  $\sqrt{s} = 161$  GeV.

77 See Fig. 4a and Fig. 5a of ABREU 97B for the exclusion limit in the mass-coupling plane.

78 See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane.

79 ACKERSTAFF 97 result is from  $e^+ e^-$  collisions at  $\sqrt{s} = 161$  GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.

80 ADLOFF 97 search for single  $e^*$  production in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ . See their Fig. 4 for the rejection limits on the product of the production cross section and the branching ratio into a specific decay channel.

81 ABREU 96K result is from  $e^+ e^-$  collisions at  $\sqrt{s} = 130$ –136 GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane.

82 ACCIARRI 96D result is from  $e^+ e^-$  collisions at  $\sqrt{s} = 130$ –140 GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane.

83 ALEXANDER 96Q result is from  $e^+ e^-$  collisions at  $\sqrt{s} = 130$ –140 GeV. See their Fig. 3a for the exclusion limit in the mass-coupling plane.

84 BUSKULIC 96W result is from  $e^+ e^-$  collisions at  $\sqrt{s} = 130$ –140 GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.

85 DERRICK 95B search for single  $e^*$  production via  $e^* e\gamma$  coupling in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ . See their Fig. 13 for the exclusion plot in the  $m_{e^*}$ – $\lambda_\gamma$  plane.

86 ABT 93 search for single  $e^*$  production via  $e^* e\gamma$  coupling in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ . See their Fig. 4 for exclusion plot in the  $m_{e^*}$ – $\lambda_\gamma$  plane.

87 DERRICK 93B search for single  $e^*$  production via  $e^* e\gamma$  coupling in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma, eZ, \nu W$ . See their Fig. 3 for exclusion plot in the  $m_{e^*}$ – $\lambda_\gamma$  plane.

88 Superseded by ADRIANI 93M.

89 Superseded by DECAMP 92.

90 ABE 88B limits use  $e^+ e^- \rightarrow ee^*$  where t-channel photon exchange dominates giving  $e\gamma(e)$  (quasi-real compton scattering).

91 ANSARI 87D is at  $E_{cm} = 546$ –630 GeV.

## Limits for Excited e ( $e^*$ ) from $e^+ e^- \rightarrow \gamma\gamma$

These limits are derived from indirect effects due to  $e^*$  exchange in the  $t$  channel and depend on transition magnetic coupling between  $e$  and  $e^*$ . All limits are for  $\lambda_\gamma = 1$ . All limits except ABE 89J are for nonchiral coupling with  $\eta_L = \eta_R = 1$ .

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

<i>VALUE</i> (GeV)	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<b>&gt;250</b>	95	BARATE	98J ALEP	$\sqrt{s} = 183$ GeV
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
>231	95	ABREU	98J DLPH	$\sqrt{s} = 130\text{--}183$ GeV
>194	95	ACKERSTAFF	98 OPAL	$\sqrt{s} = 130\text{--}172$ GeV
>227	95	ACKER..,K...	98B OPAL	$\sqrt{s} = 183$ GeV
>160	95	92 BARATE	98U ALEP	
>210	95	93 ACCIARRI	97W L3	$\sqrt{s} = 161, 172$ GeV
>129	95	ACCIARRI	96L L3	$\sqrt{s} = 133$ GeV
>147	95	ALEXANDER	96K OPAL	
>136	95	BUSKULIC	96Z ALEP	$\sqrt{s} = 130, 136$ GeV
>146	95	ACCIARRI	95G L3	
	94	BUSKULIC	93Q ALEP	
>127	95	ADRIANI	92B L3	
>114	95	96 BARDADIN-...	92 RVUE	
> 99	95	DECAMP	92 ALEP	
	97	SHIMOZAWA	92 TOPZ	
>100	95	ABREU	91E DLPH	
>116	95	AKRAWY	91F OPAL	
> 83	95	ADEVA	90K L3	
> 82	95	AKRAWY	90F OPAL	
> 68	95	98 ABE	89J VNS	$\eta_L = 1, \eta_R = 0$
> 90.2	95	ADACHI	89B TOPZ	
> 65	95	KIM	89 AMY	

92 BARATE 98U is from  $e^+ e^-$  collision at  $\sqrt{s} = M_Z$ . See their Fig. 5 for limits in mass-coupling plane

93 ACCIARRI 97W also obtain a limit on  $e^*$  with chiral coupling,  $m_{e^*} > 157$  GeV (95%CL).

94 BUSKULIC 93Q obtain  $\Lambda^+ > 121$  GeV (95%CL) from ALEPH experiment and  $\Lambda^+ > 135$  GeV from combined TRISTAN and ALEPH data. These limits roughly correspond to limits on  $m_{e^*}$ .

95 ADRIANI 92B superseded by ACCIARRI 95G.

96 BARDADIN-OTWINOWSKA 92 limit from fit to the combined data of DECAMP 92, ABREU 91E, ADEVA 90K, AKRAWY 91F.

97 SHIMOZAWA 92 fit the data to the limiting form of the cross section with  $m_{e^*} \gg E_{cm}$  and obtain  $m_{e^*} > 168$  GeV at 95%CL. Use of the full form would reduce this limit by a few GeV. The statistically unexpected large value is due to fluctuation in the data.

98 The ABE 89J limit assumes chiral coupling. This corresponds to  $\lambda_\gamma = 0.7$  for nonchiral coupling.

## Indirect Limits for Excited $e$ ( $e^*$ )

These limits make use of loop effects involving  $e^*$  and are therefore subject to theoretical uncertainty.

<u>VALUE (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
99 DORENBOSCH 89	89 CHRM $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$ and $\nu_\mu e \rightarrow \nu_\mu e$		
100 GRIFOLS 86	86 THEO $\nu_\mu e \rightarrow \nu_\mu e$		
101 RENARD 82	82 THEO $g-2$ of electron		
99 DORENBOSCH 89	obtain the limit $\lambda_\gamma^2 \Lambda_{\text{cut}}^2 / m_{e^*}^2 < 2.6$ (95% CL), where $\Lambda_{\text{cut}}$ is the cutoff scale, based on the one-loop calculation by GRIFOLS 86. If one assumes that $\Lambda_{\text{cut}} = 1$ TeV and $\lambda_\gamma = 1$ , one obtains $m_{e^*} > 620$ GeV. However, one generally expects $\lambda_\gamma \approx m_{e^*}/\Lambda_{\text{cut}}$ in composite models.		
100 GRIFOLS 86	uses $\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$ data from CHARM Collaboration to derive mass limits which depend on the scale of compositeness.		
101 RENARD 82	derived from $g-2$ data limits on mass and couplings of $e^*$ and $\mu^*$ . See figures 2 and 3 of the paper.		

## MASS LIMITS for Excited $\mu$ ( $\mu^*$ )

### Limits for Excited $\mu$ ( $\mu^*$ ) from Pair Production

These limits are obtained from  $e^+ e^- \rightarrow \mu^+ \mu^-$  and thus rely only on the (electroweak) charge of  $\mu^*$ . Form factor effects are ignored unless noted. For the case of limits from  $Z$  decay, the  $\mu^*$  coupling is assumed to be of sequential type. All limits assume  $\mu^* \rightarrow \mu \gamma$  decay except for the limits from , (Z).

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

<u>VALUE (GeV)</u>	<u>CL %</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
>85.3	95	102 ACKERSTAFF 98C OPAL	$e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type	
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
>79.6	95	104,105 ABREU	97B DLPH $e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type	
>78.4	95	104,106 ABREU	97B DLPH $e^+ e^- \rightarrow \mu^* \mu^*$ Sequential type	
>79.9	95	104 ACCIARRI	97G L3 $e^+ e^- \rightarrow \mu^* \mu^*$ Sequential type	
>80.0	95	104,107 ACKERSTAFF	97 OPAL $e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type	
>62.6	95	108 ABREU	96K DLPH $e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type	
>64.9	95	109 ACCIARRI	96D L3 $e^+ e^- \rightarrow \mu^* \mu^*$ Sequential type	
>66.8	95	109 ALEXANDER	96Q OPAL $e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type	
>65.4	95	109 BUSKULIC	96W ALEP $e^+ e^- \rightarrow \mu^* \mu^*$ Sequential type	
>45.6	95	ADRIANI	93M L3 $Z \rightarrow \mu^* \mu^*$	

>45.6	95	ABREU	92C	DLPH	$Z \rightarrow \mu^* \mu^*$
>29.8	95	110 BARDADIN...	92	RVUE	, (Z)
>26.1	95	111 DECOMP	92	ALEP	$Z \rightarrow \mu^* \mu^* ; , (Z)$
>46.1	95	DECOMP	92	ALEP	$Z \rightarrow \mu^* \mu^*$
>33	95	111 ABREU	91F	DLPH	$Z \rightarrow \mu^* \mu^* ; , (Z)$
>45.3	95	112 ADEVA	90F	L3	$Z \rightarrow \mu^* \mu^*$
>44.9	95	AKRAWY	90I	OPAL	$Z \rightarrow \mu^* \mu^*$
>44.6	95	113 DECOMP	90G	ALEP	$e^+ e^- \rightarrow \mu^* \mu^*$
>29.9	95	ADACHI	89B	TOPZ	$e^+ e^- \rightarrow \mu^* \mu^*$
>28.3	95	KIM	89	AMY	$e^+ e^- \rightarrow \mu^* \mu^*$

102 From  $e^+ e^-$  collisions at  $\sqrt{s}=170\text{--}172$  GeV. ACKERSTAFF 98C also obtain limit from  $\mu^* \rightarrow \nu W$  decay mode:  $m_{\mu^*} > 81.3$  GeV.

103 BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane. ■

104 From  $e^+ e^-$  collisions at  $\sqrt{s}=161$  GeV.

105 ABREU 97B also obtain limit from charged current decay mode  $\mu^* \rightarrow \nu W$ ,  $m_{\mu^*} > 70.9$  GeV.

106 ABREU 97B also obtain limit from charged current decay mode  $\mu^* \rightarrow \nu W$ ,  $m_{\mu^*} > 44.6$  GeV.

107 ACKERSTAFF 97 also obtain limit from charged current decay mode  $\mu^* \rightarrow \nu W$ ,  $m_{\nu_\mu^*} > 77.1$  GeV.

108 From  $e^+ e^-$  collisions at  $\sqrt{s}=130\text{--}136$  GeV.

109 From  $e^+ e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV.

110 BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on  $\Delta, (Z) < 36$  MeV.

111 Limit is independent of  $\mu^*$  decay mode.

112 Superseded by ADRIANI 93M.

113 Superseded by DECOMP 92.

### Limits for Excited $\mu$ ( $\mu^*$ ) from Single Production

These limits are from  $e^+ e^- \rightarrow \mu^* \mu$  and depend on transition magnetic coupling between  $\mu$  and  $\mu^*$ . All limits assume  $\mu^* \rightarrow \mu \gamma$  decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling,  $\eta_L = \eta_R = 1$ . In most papers, the limit is expressed in the form of an excluded region in the  $\lambda - m_{\mu^*}$  plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
>89	95	ADRIANI	93M L3	$Z \rightarrow \mu \mu^*, \lambda_Z > 0.5$
>88	95	ABREU	92C DLPH	$Z \rightarrow \mu \mu^*, \lambda_Z > 0.5$
<b>&gt;91</b>	95	DECAMP	92 ALEP	$Z \rightarrow \mu \mu^*, \lambda_Z > 1$
>87	95	AKRAWY	90I OPAL	$Z \rightarrow \mu \mu^*, \lambda_Z > 1$

• • • We do not use the following data for averages, fits, limits, etc. • • •

114	ACKERSTAFF	98C OPAL	$e^+ e^- \rightarrow \mu\mu^*$	
115	BARATE	98U ALEP	$Z \rightarrow \mu\mu^*$	
116,117	ABREU	97B DLPH	$e^+ e^- \rightarrow \mu\mu^*$	
116,118	ACCIARRI	97G L3	$e^+ e^- \rightarrow \mu\mu^*$	
119	ACKERSTAFF	97 OPAL	$e^+ e^- \rightarrow \mu\mu^*$	
120	ABREU	96K DLPH	$e^+ e^- \rightarrow \mu\mu^*$	
121	ACCIARRI	96D L3	$e^+ e^- \rightarrow \mu\mu^*$	
122	ALEXANDER	96Q OPAL	$e^+ e^- \rightarrow \mu\mu^*$	
123	BUSKULIC	96W ALEP	$e^+ e^- \rightarrow \mu\mu^*$	
>85	95	124 ADEVA	90F L3	$Z \rightarrow \mu\mu^*, \lambda_Z > 1$
>75	95	124 ADEVA	90F L3	$Z \rightarrow \mu\mu^*, \lambda_Z > 0.1$
>80	95	125 DECOMP	90G ALEP	$e^+ e^- \rightarrow \mu\mu^*, \lambda_Z = 1$
>50	95	ADACHI	89B TOPZ	$e^+ e^- \rightarrow \mu\mu^*, \lambda_\gamma = 0.7$
>46	95	KIM	89 AMY	$e^+ e^- \rightarrow \mu\mu^*, \lambda_\gamma = 0.2$

114 ACKERSTAFF 98C from  $e^+ e^-$  collisions at  $\sqrt{s}=170-172$  GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane.

115 BARATE 98U obtain limits on the  $Z\mu\mu^*$  coupling. See their Fig. 12 for limits in mass-coupling plane

116 From  $e^+ e^-$  collisions at  $\sqrt{s}= 161$  GeV.

117 See Fig. 4a and Fig. 5a of ABREU 97B for the exclusion limit in the mass-coupling plane.

118 See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane.

119 ACKERSTAFF 97 result is from  $e^+ e^-$  collisions at  $\sqrt{s}= 161$  GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.

120 ABREU 96K result is from  $e^+ e^-$  collisions at  $\sqrt{s}= 130-136$  GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane.

121 ACCIARRI 96D result is from  $e^+ e^-$  collisions at  $\sqrt{s}= 130-140$  GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane.

122 ALEXANDER 96Q result is from  $e^+ e^-$  collisions at  $\sqrt{s}= 130-140$  GeV. See their Fig. 3a for the exclusion limit in the mass-coupling plane.

123 BUSKULIC 96W result is from  $e^+ e^-$  collisions at  $\sqrt{s}= 130-140$  GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.

124 Superseded by ADRIANI 93M.

125 Superseded by DECOMP 92.

### Indirect Limits for Excited $\mu$ ( $\mu^*$ )

These limits make use of loop effects involving  $\mu^*$  and are therefore subject to theoretical uncertainty.

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

126	RENARD	82 THEO	$g-2$ of muon
-----	--------	---------	---------------

126 RENARD 82 derived from  $g-2$  data limits on mass and couplings of  $e^*$  and  $\mu^*$ . See figures 2 and 3 of the paper.

## MASS LIMITS for Excited $\tau$ ( $\tau^*$ )

### Limits for Excited $\tau$ ( $\tau^*$ ) from Pair Production

These limits are obtained from  $e^+ e^- \rightarrow \tau^+ \tau^-$  and thus rely only on the (electroweak) charge of  $\tau^*$ . Form factor effects are ignored unless noted. For the case of limits from  $Z$  decay, the  $\tau^*$  coupling is assumed to be of sequential type. All limits assume  $\tau^* \rightarrow \tau \gamma$  decay except for the limits from , (Z).

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
>84.6	95	127 ACKERSTAFF 98C	OPAL	$e^+ e^- \rightarrow \tau^* \tau^*$ Homodoublet type

• • • We do not use the following data for averages, fits, limits, etc. • • •

>84.6	95	127 ACKERSTAFF 98C	OPAL	$e^+ e^- \rightarrow \tau^* \tau^*$ Homodoublet type
>79.4	95	129,130 ABREU	97B DLPH	$e^+ e^- \rightarrow \tau^* \tau^*$ Homodoublet type
>77.4	95	129,131 ABREU	97B DLPH	$e^+ e^- \rightarrow \tau^* \tau^*$ Sequential type
>79.3	95	129 ACCIARRI	97G L3	$e^+ e^- \rightarrow \tau^* \tau^*$ Sequential type
>79.1	95	129,132 ACKERSTAFF	97 OPAL	$e^+ e^- \rightarrow \tau^* \tau^*$ Homodoublet type
>62.2	95	133 ABREU	96K DLPH	$e^+ e^- \rightarrow \tau^* \tau^*$ Homodoublet type
>64.2	95	134 ACCIARRI	96D L3	$e^+ e^- \rightarrow \tau^* \tau^*$ Sequential type
>65.3	95	134 ALEXANDER	96Q OPAL	$e^+ e^- \rightarrow \tau^* \tau^*$ Homodoublet type
>64.8	95	134 BUSKULIC	96W ALEP	$e^+ e^- \rightarrow \tau^* \tau^*$ Sequential type
>45.6	95	ADRIANI	93M L3	$Z \rightarrow \tau^* \tau^*$
>45.3	95	ABREU	92C DLPH	$Z \rightarrow \tau^* \tau^*$
>29.8	95	135 BARDADIN-...	92 RVUE	, (Z)
>26.1	95	136 DECAMP	92 ALEP	$Z \rightarrow \tau^* \tau^*; , (Z)$
>46.0	95	DECAMP	92 ALEP	$Z \rightarrow \tau^* \tau^*$
>33	95	136 ABREU	91F DLPH	$Z \rightarrow \tau^* \tau^*; , (Z)$
>45.5	95	137 ADEVA	90L L3	$Z \rightarrow \tau^* \tau^*$
>44.9	95	AKRAWY	90I OPAL	$Z \rightarrow \tau^* \tau^*$
>41.2	95	138 DECAMP	90G ALEP	$e^+ e^- \rightarrow \tau^* \tau^*$
>29.0	95	ADACHI	89B TOPZ	$e^+ e^- \rightarrow \tau^* \tau^*$

127 From  $e^+ e^-$  collisions at  $\sqrt{s}=170\text{--}172$  GeV. ACKERSTAFF 98C also obtain limit from  $\tau^* \rightarrow \nu W$  decay mode:  $m_{\tau^*} > 81.3$  GeV.

128 BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane.

129 From  $e^+ e^-$  collisions at  $\sqrt{s}=161$  GeV.

130 ABREU 97B also obtain limit from charged current decay mode  $\tau^* \rightarrow \nu W$ ,  $m_{\tau^*} > 70.9$  GeV.

131 ABREU 97B also obtain limit from charged current decay mode  $\tau^* \rightarrow \nu W$ ,  $m_{\tau^*} > 44.6$  GeV.

132 ACKERSTAFF 97 also obtain limit from charged current decay mode  $\tau^* \rightarrow \nu W$ ,  $m_{\nu_\tau^*} > 77.1$  GeV.

133 From  $e^+ e^-$  collisions at  $\sqrt{s}=130\text{--}136$  GeV.

134 From  $e^+ e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV.

135 BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on  $\Delta, (Z) < 36$  MeV.

136 Limit is independent of  $\tau^*$  decay mode.

137 Superseded by ADRIANI 93M.

138 Superseded by DECAMP 92.

## Limits for Excited $\tau$ ( $\tau^*$ ) from Single Production

These limits are from  $e^+ e^- \rightarrow \tau^* \tau$  and depend on transition magnetic coupling between  $\tau$  and  $\tau^*$ . All limits assume  $\tau^* \rightarrow \tau \gamma$  decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling,  $\eta_L = \eta_R = 1$ . In most papers, the limit is expressed in the form of an excluded region in the  $\lambda - m_{\tau^*}$  plane. See the original papers.

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
>88	95	ADRIANI	93M L3	$Z \rightarrow \tau \tau^*, \lambda_Z > 0.5$
>87	95	ABREU	92C DLPH	$Z \rightarrow \tau \tau^*, \lambda_Z > 0.5$
<b>&gt;90</b>	95	DECAMP	92 ALEP	$Z \rightarrow \tau \tau^*, \lambda_Z > 0.18$
>86.5	95	AKRAWY	90I OPAL	$Z \rightarrow \tau \tau^*, \lambda_Z > 1$

• • • We do not use the following data for averages, fits, limits, etc. • • •

139	ACKERSTAFF	98C OPAL	$e^+ e^- \rightarrow \tau \tau^*$	
140	BARATE	98U ALEP	$Z \rightarrow \tau \tau^*$	
141,142	ABREU	97B DLPH	$e^+ e^- \rightarrow \tau \tau^*$	
141,143	ACCIARRI	97G L3	$e^+ e^- \rightarrow \tau \tau^*$	
144	ACKERSTAFF	97 OPAL	$e^+ e^- \rightarrow \tau \tau^*$	
145	ABREU	96K DLPH	$e^+ e^- \rightarrow \tau \tau^*$	
146	ACCIARRI	96D L3	$e^+ e^- \rightarrow \tau \tau^*$	
147	ALEXANDER	96Q OPAL	$e^+ e^- \rightarrow \tau \tau^*$	
148	BUSKULIC	96W ALEP	$e^+ e^- \rightarrow \tau \tau^*$	
>88	95	149 ADEVA	90L L3	$Z \rightarrow \tau \tau^*, \lambda_Z > 1$
>59	95	150 DECAMP	90G ALEP	$Z \rightarrow \tau \tau^*, \lambda_Z = 1$
>40	95	151 BARTEL	86 JADE	$e^+ e^- \rightarrow \tau \tau^*, \lambda_\gamma = 1$
>41.4	95	152 BEHREND	86 CELL	$e^+ e^- \rightarrow \tau \tau^*, \lambda_\gamma = 1$
>40.8	95	152 BEHREND	86 CELL	$e^+ e^- \rightarrow \tau \tau^*, \lambda_\gamma = 0.7$

139 ACKERSTAFF 98C from  $e^+ e^-$  collisions at  $\sqrt{s}=170-172$  GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane.

140 BARATE 98U obtain limits on the  $Z \tau \tau^*$  coupling. See their Fig. 12 for limits in mass-coupling plane

141 From  $e^+ e^-$  collisions at  $\sqrt{s}=161$  GeV.

142 See Fig. 4a and Fig. 5a of ABREU 97B for the exclusion limit in the mass-coupling plane.

143 See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane.

144 ACKERSTAFF 97 result is from  $e^+ e^-$  collisions at  $\sqrt{s}=161$  GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.

145 ABREU 96K result is from  $e^+ e^-$  collisions at  $\sqrt{s}=130-136$  GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane.

146 ACCIARRI 96D result is from  $e^+ e^-$  collisions at  $\sqrt{s}=130-140$  GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane.

147 ALEXANDER 96Q result is from  $e^+ e^-$  collisions at  $\sqrt{s}=130-140$  GeV. See their Fig. 3a for the exclusion limit in the mass-coupling plane.

148 BUSKULIC 96W result is from  $e^+ e^-$  collisions at  $\sqrt{s}=130-140$  GeV. See their Fig. 3 for the exclusion limit in the mass-coupling plane.

149 Superseded by ADRIANI 93M.

150 Superseded by DECAMP 92.

151 BARTEL 86 is at  $E_{cm} = 30-46.78$  GeV.

152 BEHREND 86 limit is at  $E_{cm} = 33-46.8$  GeV.

## MASS LIMITS for Excited Neutrino ( $\nu^*$ )

### Limits for Excited $\nu$ ( $\nu^*$ ) from Pair Production

These limits are obtained from  $e^+ e^- \rightarrow \nu^* \nu^*$  and thus rely only on the (electroweak) charge of  $\nu^*$ . Form factor effects are ignored unless noted. The  $\nu^*$  coupling is assumed to be of sequential type unless otherwise noted. Limits assume  $\nu^* \rightarrow \nu \gamma$  decay except for the  $,(Z)$  measurement which makes no assumption about decay mode.

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;84.9</b>	95	153 ACKERSTAFF 98C	OPAL	$e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type

• • • We do not use the following data for averages, fits, limits, etc. • • •

	154	BARATE	98U ALEP	$Z \rightarrow \nu^* \nu^*$
>77.6	95	155,156 ABREU	97B DLPH	$e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type
>64.4	95	155,157 ABREU	97B DLPH	$e^+ e^- \rightarrow \nu^* \nu^*$ Sequential type
>71.2	95	155,158 ACCIARRI	97G L3	$e^+ e^- \rightarrow \nu^* \nu^*$ Sequential type
>77.8	95	155,159 ACKERSTAFF	97 OPAL	$e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type
>61.4	95	160,161 ACCIARRI	96D L3	$e^+ e^- \rightarrow \nu^* \nu^*$ Sequential type
>65.0	95	162,163 ALEXANDER	96Q OPAL	$e^+ e^- \rightarrow \nu^* \nu^*$ Homodoublet type
>63.6	95	160 BUSKULIC	96W ALEP	$e^+ e^- \rightarrow \nu^* \nu^*$ Sequential type
>43.7	95	164 BARDADIN-...	92 RVUE	$,(Z)$
>47	95	165 DECAMP	92 ALEP	
>42.6	95	166 DECAMP	92 ALEP	$,(Z)$
>35.4	95	167,168 DECAMP	900 ALEP	$,(Z)$
>46	95	168,169 DECAMP	900 ALEP	

153 From  $e^+ e^-$  collisions at  $\sqrt{s}=170\text{--}172$  GeV. ACKERSTAFF 98C also obtain limit from charged decay modes:  $m_{\nu_e^*} > 84.1$  GeV,  $m_{\nu_\mu^*} > 83.9$  GeV, and  $m_{\nu_\tau^*} > 79.4$  GeV.

154 BARATE 98U obtain limits on the form factor. See their Fig. 14 for limits in mass-form factor plane.

155 From  $e^+ e^-$  collisions at  $\sqrt{s}=161$  GeV.

156 ABREU 97B also obtain limits from charged current decay modes,  $m_{\nu^*} > 56.4$  GeV.

157 ABREU 97B also obtain limits from charged current decay modes,  $m_{\nu^*} > 44.9$  GeV.

158 ACCIARRI 97G also obtain limits from charged current decay mode  $\nu_e^* \rightarrow e W$ ,  $m_{\nu^*} > 64.5$  GeV.

159 ACKERSTAFF 97 also obtain limits from charged current decay modes  $m_{\nu_e^*} > 78.3$  GeV,  $m_{\nu_\mu^*} > 78.9$  GeV,  $m_{\nu_\tau^*} > 76.2$  GeV.

160 From  $e^+ e^-$  collisions at  $\sqrt{s}=130\text{--}140$  GeV.

161 ACCIARRI 96D also obtain limit from  $\nu^* \rightarrow e W$  decay mode:  $m_{\nu^*} > 57.3$  GeV.

162 From  $e^+ e^-$  collisions at  $\sqrt{s}=130\text{--}136$  GeV.

163 ALEXANDER 96Q also obtain limits from charged current decay modes:  $m_{\nu_e^*} > 66.2$  GeV,  $m_{\nu_\mu^*} > 66.5$  GeV,  $m_{\nu_\tau^*} > 64.7$  GeV.

164 BARDADIN-OTWINOWSKA 92 limit is for Dirac  $\nu^*$ . Based on  $\Delta, (Z) < 36$  MeV. The limit is 36.4 GeV for Majorana  $\nu^*$ , 45.4 GeV for homodoublet  $\nu^*$ .

165 Limit is based on  $B(Z \rightarrow \nu^* \bar{\nu}^*) \times B(\nu^* \rightarrow \nu \gamma)^2 < 5 \times 10^{-5}$  (95%CL) assuming Dirac  $\nu^*$ ,  $B(\nu^* \rightarrow \nu \gamma) = 1$ .

166 Limit is for Dirac  $\nu^*$ . The limit is 34.6 GeV for Majorana  $\nu^*$ , 45.4 GeV for homodoublet  $\nu^*$ .

- 167 DECAMP 900 limit is from excess  $\Delta, (Z) < 89$  MeV. The above value is for Dirac  $\nu^*$ ; 26.6 GeV for Majorana  $\nu^*$ ; 44.8 GeV for homodoublet  $\nu^*$ .  
 168 Superseded by DECAMP 92.  
 169 DECAMP 900 limit based on  $B(Z \rightarrow \nu^* \nu^*) \cdot B(\nu^* \rightarrow \nu \gamma)^2 < 7 \times 10^{-5}$  (95%CL), assuming Dirac  $\nu^*$ ,  $B(\nu^* \rightarrow \nu \gamma) = 1$ .

## Limits for Excited $\nu$ ( $\nu^*$ ) from Single Production

These limits are from  $Z \rightarrow \nu \nu^*$  or  $e p \rightarrow \nu^* X$  and depend on transition magnetic coupling between  $\nu/e$  and  $\nu^*$ . Assumptions about  $\nu^*$  decay mode are given in footnotes.

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
none 40–96	95	170 BREITWEG	97C ZEUS	$e p \rightarrow \nu^* X$
<b>&gt;91</b>	95	ADRIANI	93M L3	$\lambda_Z > 1, \nu^* \rightarrow \nu \gamma$
>89	95	ADRIANI	93M L3	$\lambda_Z > 1, \nu_e^* \rightarrow e W$
<b>&gt;91</b>	95	171 DECAMP	92 ALEP	$\lambda_Z > 1$

• • • We do not use the following data for averages, fits, limits, etc. • • •

172	ACKERSTAFF	98C OPAL	$e^+ e^- \rightarrow \nu^* \nu^*$
173	BARATE	98U ALEP	$Z \rightarrow \nu \nu^*$
174,175	ABREU	97B DLPH	$e^+ e^- \rightarrow \nu \nu^*$
176	ABREU	97I DLPH	$\nu^* \rightarrow \ell W, \nu Z$
177	ABREU	97J DLPH	$\nu^* \rightarrow \nu \gamma$
174,178	ACCIARRI	97G L3	$e^+ e^- \rightarrow \nu \nu^*$
179	ACKERSTAFF	97 OPAL	$e^+ e^- \rightarrow \nu \nu^*$
180	ADLOFF	97 H1	Lepton-flavor violation
181	ACCIARRI	96D L3	$e^+ e^- \rightarrow \nu \nu^*$
182	ALEXANDER	96Q OPAL	$e^+ e^- \rightarrow \nu \nu^*$
183	BUSKULIC	96W ALEP	$e^+ e^- \rightarrow \nu \nu^*$
184	DERRICK	95B ZEUS	$e p \rightarrow \nu^* X$
185	ABT	93 H1	$e p \rightarrow \nu^* X$
>87	95	ADRIANI	$\lambda_Z > 0.1, \nu^* \rightarrow \nu \gamma$
>74	95	ADRIANI	$\lambda_Z > 0.1, \nu_e^* \rightarrow e W$
	186	BARDADIN-...	92 RVUE
>74	95	171 DECAMP	$\lambda_Z > 0.034$
>91	95	187,188 ADEVA	$\lambda_Z > 1$
>83	95	188 ADEVA	$\lambda_Z > 0.1, \nu^* \rightarrow \nu \gamma$
>74	95	188 ADEVA	$\lambda_Z > 0.1, \nu_e^* \rightarrow e W$
>90	95	189,190 DECAMP	$\lambda_Z > 1$
>74.7	95	189,190 DECAMP	$\lambda_Z > 0.06$

170 BREITWEG 97C search for single  $\nu^*$  production in  $e p$  collisions with the decay  $\nu^* \rightarrow \nu \gamma$ .  $f = -f' = 2\Lambda/m_{\nu^*}$  is assumed for the  $\nu^*$  coupling. See their Fig. 10 for the exclusion plot in the mass-coupling plane.

171 DECAMP 92 limit is based on  $B(Z \rightarrow \nu^* \bar{\nu}) \times B(\nu^* \rightarrow \nu \gamma) < 2.7 \times 10^{-5}$  (95%CL) assuming Dirac  $\nu^*$ ,  $B(\nu^* \rightarrow \nu \gamma) = 1$ .

172 ACKERSTAFF 98C from  $e^+ e^-$  collisions at  $\sqrt{s} = 170-172$  GeV. See their Fig. 11 for the exclusion limit in the mass-coupling plane.

173 BARATE 98U obtain limits on the  $Z \nu \nu^*$  coupling. See their Fig. 13 for limits in mass-coupling plane.

174 From  $e^+ e^-$  collisions at  $\sqrt{s} = 161$  GeV.

175 See Fig. 4b and Fig. 5b of ABREU 97B for the exclusion limit in the mass-coupling plane.

- 176 ABREU 97I limit is from  $Z \rightarrow \nu\nu^*$ . See their Fig. 12 for the exclusion limit in the mass-coupling plane.
- 177 ABREU 97J limit is from  $Z \rightarrow \nu\nu^*$ . See their Fig. 5 for the exclusion limit in the mass-coupling plane.
- 178 See Fig. 2 and Fig. 3 of ACCIARRI 97G for the exclusion limit in the mass-coupling plane.
- 179 ACKERSTAFF 97 result is from  $e^+ e^-$  collisions at  $\sqrt{s} = 161$  GeV, for homodoublet  $\nu^*$ . See their Fig. 3 for the exclusion limit in the mass-coupling plane.
- 180 ADLOFF 97 search for single  $e^*$  production in  $e p$  collisions with the decays  $e^* \rightarrow e\gamma$ ,  $eZ$ ,  $\nu W$ . See their Fig. 4 for the rejection limits on the product of the production cross section and the branching ratio.
- 181 ACCIARRI 96D result is from  $e^+ e^-$  collisions at  $\sqrt{s} = 130-140$  GeV. See their Fig. 2 for the exclusion limit in the mass-coupling plane.
- 182 ALEXANDER 96Q result is from  $e^+ e^-$  collisions at  $\sqrt{s} = 130-140$  GeV for homodoublet  $\nu^*$ . See their Fig. 3b and Fig. 3c for the exclusion limit in the mass-coupling plane.
- 183 BUSKULIC 96W result is from  $e^+ e^-$  collisions at  $\sqrt{s} = 130-140$  GeV. See their Fig. 4 for the exclusion limit in the mass-coupling plane.
- 184 DERRICK 95B search for single  $\nu^*$  production via  $\nu^* e W$  coupling in  $e p$  collisions with the decays  $\nu^* \rightarrow \nu\gamma$ ,  $\nu Z$ ,  $e W$ . See their Fig. 14 for the exclusion plot in the  $m_{\nu^*}-\lambda_W$  plane.
- 185 ABT 93 search for single  $\nu^*$  production via  $\nu^* e W$  coupling in  $e p$  collisions with the decays  $\nu^* \rightarrow \nu\gamma$ ,  $\nu Z$ ,  $e W$ . See their Fig. 4 for exclusion plot in the  $m_{\nu^*}-\lambda_W$  plane.
- 186 See Fig. 5 of BARDADIN-OTWINOWSKA 92 for combined limit of ADEVA 900, DECAMP 900, and DECOMP 92.
- 187 Limit is either for  $\nu^* \rightarrow \nu\gamma$  or  $\nu^* \rightarrow e W$ .
- 188 Superseded by ADRIANI 93M.
- 189 DECOMP 900 limit based on  $B(Z \rightarrow \nu\nu^*) \cdot B(\nu^* \rightarrow \nu\gamma) < 6 \times 10^{-5}$  (95%CL), assuming  $B(\nu^* \rightarrow \nu\gamma) = 1$ .
- 190 Superseded by DECOMP 92.

## MASS LIMITS for Excited $q$ ( $q^*$ )

### Limits for Excited $q$ ( $q^*$ ) from Pair Production

These limits are obtained from  $e^+ e^- \rightarrow q^* \bar{q}^*$  and thus rely only on the (electroweak) charge of the  $q^*$ . Form factor effects are ignored unless noted. Assumptions about the  $q^*$  decay are given in the comments and footnotes.

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;45.6</b>	95	191	ADRIANI	93M L3 $u$ or $d$ type, $Z \rightarrow q^* q^*$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
		192	BARATE	98U ALEP $Z \rightarrow q^* q^*$
		193	ADRIANI	92F L3 $Z \rightarrow q^* q^*$
>41.7	95	194	BARDADIN-...	92 RVUE $u$ -type, , ( $Z$ )
>44.7	95	194	BARDADIN-...	92 RVUE $d$ -type, , ( $Z$ )
>40.6	95	195	DECAMP	92 ALEP $u$ -type, , ( $Z$ )
>44.2	95	195	DECAMP	92 ALEP $d$ -type, , ( $Z$ )
>45	95	196	DECAMP	92 ALEP $u$ or $d$ type, $Z \rightarrow q^* q^*$

>45	95	195 ABREU	91F DLPH	$u$ -type, , $(Z)$
>45	95	195 ABREU	91F DLPH	$d$ -type, , $(Z)$
>21.1	95	197 BEHREND	86C CELL	$e(q^*) = -1/3, q^* \rightarrow qg$
>22.3	95	197 BEHREND	86C CELL	$e(q^*) = 2/3, q^* \rightarrow qg$
>22.5	95	197 BEHREND	86C CELL	$e(q^*) = -1/3, q^* \rightarrow q\gamma$
>23.2	95	197 BEHREND	86C CELL	$e(q^*) = 2/3, q^* \rightarrow q\gamma$

191 ADRIANI 93M limit is valid for  $B(q^* \rightarrow qg) > 0.25$  (0.17) for up (down) type.

192 BARATE 98U obtain limits on the form factor. See their Fig. 16 for limits in mass-form factor plane. ■

193 ADRIANI 92F search for  $Z \rightarrow q^* \bar{q}^*$  followed with  $q^* \rightarrow q\gamma$  decays and give the limit  $\sigma_Z \cdot B(Z \rightarrow q^* \bar{q}^*) \cdot B^2(q^* \rightarrow q\gamma) < 2$  pb at 95%CL. Assuming five flavors of degenerate  $q^*$  of homodoublet type,  $B(q^* \rightarrow q\gamma) < 4\%$  is obtained for  $m_{q^*} < 45$  GeV.

194 BARDADIN-OTWINOWSKA 92 limit based on  $\Delta, (Z) < 36$  MeV.

195 These limits are independent of decay modes.

196 Limit is for  $B(q^* \rightarrow qg) + B(q^* \rightarrow q\gamma) = 1$ .

197 BEHREND 86C search for  $e^+ e^- \rightarrow q^* \bar{q}^*$  for  $m_{q^*} > 5$  GeV. But  $m < 5$  GeV excluded by total hadronic cross section. The limits are for point-like photon couplings of excited quarks.

### Limits for Excited $q$ ( $q^*$ ) from Single Production

These limits are from  $e^+ e^- \rightarrow q^* \bar{q}$  or  $p\bar{p} \rightarrow q^* X$  and depend on transition magnetic couplings between  $q$  and  $q^*$ . Assumptions about  $q^*$  decay mode are given in the footnotes and comments.

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
<b>&gt;570 (CL = 95%) OUR EVALUATION</b>				
none 200–520 and 580–760	95	198 ABE	97G CDF	$p\bar{p} \rightarrow q^* X, q^* \rightarrow 2$ jets
none 40–169	95	199 BREITWEG	97C ZEUS	$e p \rightarrow q^* X$
<b>none 80–570</b>	95	200 ABE	95N CDF	$p\bar{p} \rightarrow q^* X, q^* \rightarrow qg$ $q\gamma, qW$
>288	90	201 ALITTI	93 UA2	$p\bar{p} \rightarrow q^* X, q^* \rightarrow qg$
<b>&gt; 88</b>	95	202 DECAMP	92 ALEP	$Z \rightarrow q\bar{q}^*, \lambda_Z > 1$
> 86	95	202 AKRAWY	90J OPAL	$Z \rightarrow q\bar{q}^*, \lambda_Z > 1.2$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
		203 BARATE	98U ALEP	$Z \rightarrow q\bar{q}^*$
		204 ADLOFF	97 H1	Lepton-flavor violation
		205 DERRICK	95B ZEUS	$e p \rightarrow q^* X$
none 80–540	95	206 ABE	94 CDF	$p\bar{p} \rightarrow q^* X, q^* \rightarrow q\gamma,$ $qW$
> 79	95	207 ADRIANI	93M L3	$\lambda_Z(L3) > 0.06$
		208 ABREU	92D DLPH	$Z \rightarrow q\bar{q}^*$
		209 ADRIANI	92F L3	$Z \rightarrow q\bar{q}^*$
> 75	95	207 DECAMP	92 ALEP	$Z \rightarrow q\bar{q}^*, \lambda_Z > 1$
		210 ALBAJAR	89 UA1	$p\bar{p} \rightarrow q^* X,$ $q^* \rightarrow qW$
> 39	95	211 BEHREND	86C CELL	$e^+ e^- \rightarrow q^* \bar{q} (q^* \rightarrow qg, q\gamma), \lambda_\gamma = 1$

198 ABE 97G search for new particle decaying to dijets.

- 199 BREITWEG 97C search for single  $q^*$  production in  $e p$  collisions with the decays  $q^* \rightarrow q\gamma$ ,  $qW$ .  $f_s=0$ , and  $f=f'=2\Lambda/m_{q^*}$  is assumed for the  $q^*$  coupling. See their Fig. 11 for the exclusion plot in the mass-coupling plane.
- 200 ABE 95N assume a degenerate  $u^*$  and  $d^*$  with  $f_s=f=f'=\Lambda/m_{q^*}$ . See their Fig. 4 for the excluded region in  $m_{q^*} - f$  plane.
- 201 ALITTI 93 search for resonances in the two-jet invariant mass. The limit is for  $f_s = f = f' = \Lambda/m_{q^*}$ .  $u^*$  and  $d^*$  are assumed to be degenerate. If not, the limit for  $u^*$  ( $d^*$ ) is 277 (247) GeV if  $m_{d^*} \gg m_{u^*}$  ( $m_{u^*} \gg m_{d^*}$ ).
- 202 Assumes  $B(q^* \rightarrow q\gamma) = 0.1$ .
- 203 BARATE 98U obtain limits on the  $Z q q^*$  coupling. See their Fig. 16 for limits in mass-coupling plane
- 204 ADLOFF 97 search for single  $q^*$  production in  $e p$  collisions with the decay  $q^* \rightarrow q\gamma$ . See their Fig. 6 for the rejection limits on the product of the production cross section and the branching ratio.
- 205 DERRICK 95B search for single  $q^*$  production via  $q^* q\gamma$  coupling in  $e p$  collisions with the decays  $q^* \rightarrow qW$ ,  $qZ$ ,  $qg$ ,  $q\gamma$ . See their Fig. 15 for the exclusion plot in the  $m_{q^*} - \lambda\gamma$  plane.
- 206 ABE 94 search for resonances in jet- $\gamma$  and jet- $W$  invariant mass in  $p\bar{p}$  collisions at  $E_{cm} = 1.8$  TeV. The limit is for  $f_s = f = f' = \Lambda/m_{q^*}$  and  $u^*$  and  $d^*$  are assumed to be degenerate. See their Fig. 4 for the excluded region in  $m_{q^*} - f$  plane.
- 207 Assumes  $B(q^* \rightarrow qg) = 1$ .
- 208 ABREU 92D give  $\sigma(e^+ e^- \rightarrow Z \rightarrow q^* \bar{q} \text{ or } q\bar{q}^*) \times B(q^* \rightarrow q\gamma) < 15 \text{ pb}$  (95% CL) for  $m_{q^*} < 80$  GeV.
- 209 ADRIANI 92F search for  $Z \rightarrow q q^*$  with  $q^* \rightarrow q\gamma$  and give the limit  $\sigma_Z \cdot B(Z \rightarrow q q^*) \cdot B(q^* \rightarrow q\gamma) < (2-10) \text{ pb}$  (95% CL) for  $m_{q^*} = (46-82)$  GeV.
- 210 ALBAJAR 89 give  $\sigma(q^* \rightarrow W + \text{jet})/\sigma(W) < 0.019$  (90% CL) for  $m_{q^*} > 220$  GeV.
- 211 BEHREND 86C has  $E_{cm} = 42.5-46.8$  GeV. See their Fig. 3 for excluded region in the  $m_{q^*} - (\lambda_\gamma/m_{q^*})^2$  plane. The limit is for  $\lambda_\gamma = 1$  with  $\eta_L = \eta_R = 1$ .

### MASS LIMITS for Color Sextet Quarks ( $q_6$ )

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
>84	95	212 ABE	89D CDF	$p\bar{p} \rightarrow q_6 \bar{q}_6$

212 ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color sextet quark is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. A limit of 121 GeV is obtained for a color decuplet.

### MASS LIMITS for Color Octet Charged Leptons ( $\ell_8$ )

$$\lambda \equiv m_{\ell_8}/\Lambda$$

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
>86	95	213 ABE	89D CDF	Stable $\ell_8$ : $p\bar{p} \rightarrow \ell_8 \bar{\ell}_8$

• • • We do not use the following data for averages, fits, limits, etc. • • •

	214 ABT	93 H1	$\nu_8: e p \rightarrow \nu_8 X$
none 3.0–30.3	95 215 KIM	90 AMY	$\nu_8: e^+ e^- \rightarrow e e +$ jets
none 3.5–30.3	95 215 KIM	90 AMY	$\mu_8: e^+ e^- \rightarrow \mu\mu +$ jets
	216 KIM	90 AMY	$\nu_8: e^+ e^- \rightarrow gg; R$
>19.8	95 217 BARTEL	87B JADE	$\nu_8, \mu_8, \tau_8: e^+ e^-; R$
none 5–23.2	95 217 BARTEL	87B JADE	$\mu_8: e^+ e^- \rightarrow \mu\mu +$ jets
	218 BARTEL	85K JADE	$\nu_8: e^+ e^- \rightarrow gg; R$

213 ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color octet lepton is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. The limit improves to 99 GeV if it always fragments into a unit-charged hadron.

214 ABT 93 search for  $\nu_8$  production via  $e$ -gluon fusion in  $e p$  collisions with  $\nu_8 \rightarrow e g$ . See their Fig. 3 for exclusion plot in the  $m_{\nu_8}$ –Λ plane for  $m_{\nu_8} = 35$ –220 GeV.

215 KIM 90 is at  $E_{cm} = 50$ –60.8 GeV. The same assumptions as in BARTEL 87B are used.

216 KIM 90 result  $(m_{\nu_8} \Lambda_M)^{1/2} > 178.4$  GeV (95%CL,  $\alpha_s = 0.16$  used) is subject to the same restriction as for BARTEL 85K.

217 BARTEL 87B is at  $E_{cm} = 46.3$ –46.78 GeV. The limits assume  $\nu_8$  pair production cross sections to be eight times larger than those of the corresponding heavy lepton pair production.

218 In BARTEL 85K,  $R$  can be affected by  $e^+ e^- \rightarrow gg$  via  $e_q$  exchange. Their limit  $m_{\nu_8} > 173$  GeV (CL=95%) at  $\lambda = m_{\nu_8}/\Lambda_M = 1$  ( $\eta_L = \eta_R = 1$ ) is not listed above because the cross section is sensitive to the product  $\eta_L \eta_R$ , which should be absent in ordinary theory with electronic chiral invariance.

## MASS LIMITS for Color Octet Neutrinos ( $\nu_8$ )

$$\lambda \equiv m_{\nu_8}/\Lambda$$

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
>110	90	219 BARGER	89 RVUE	$\nu_8: p\bar{p} \rightarrow \nu_8 \bar{\nu}_8$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
none 3.8–29.8	95	220 KIM	90 AMY	$\nu_8: e^+ e^- \rightarrow$ acoplanar jets
none 9–21.9	95	221 BARTEL	87B JADE	$\nu_8: e^+ e^- \rightarrow$ acoplanar jets

219 BARGER 89 used ABE 89B limit for events with large missing transverse momentum. Two-body decay  $\nu_8 \rightarrow \nu g$  is assumed.

220 KIM 90 is at  $E_{cm} = 50$ –60.8 GeV. The same assumptions as in BARTEL 87B are used.

221 BARTEL 87B is at  $E_{cm} = 46.3$ –46.78 GeV. The limit assumes the  $\nu_8$  pair production cross section to be eight times larger than that of the corresponding heavy neutrino pair production. This assumption is not valid in general for the weak couplings, and the limit can be sensitive to its  $SU(2)_L \times U(1)_Y$  quantum numbers.

## MASS LIMITS for $W_8$ (Color Octet $W$ Boson)

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
222	ALBAJAR	89 UA1	$p\bar{p} \rightarrow W_8 X,$ $W_8 \rightarrow W g$
222 ALBAJAR 89 give $\sigma(W_8 \rightarrow W + \text{jet})/\sigma(W) < 0.019$ (90% CL) for $m_{W_8} > 220$ GeV.			

## Limits on $Z Z \gamma$ Coupling

Limits are for the electric dipole transition form factor for  $Z \rightarrow \gamma Z^*$  parametrized as  $f(s') = \beta(s'/m_Z^2 - 1)$ , where  $s'$  is the virtual  $Z$  mass. In the Standard Model  $\beta \sim 10^{-5}$ .

VALUE	CL %	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
<0.80	95	ADRIANI	92J L3	$Z \rightarrow \gamma \nu \bar{\nu}$

## REFERENCES FOR Searches for Quark and Lepton Compositeness

ABBIENDI	99	EPJ C6 1	G. Abbiendi+	(OPAL Collab.)
ABBOTT	98G	PRL 80 666	B. Abbott+	(D0 Collab.)
ABREU	98J	PL B433 429	P. Abreu+	(DELPHI Collab.)
ACCIARRI	98J	PL B433 163	M. Acciarri+	(L3 Collab.)
ACCIARRI	98T	PL B439 183	M. Acciarri+	(L3 Collab.)
ACKERSTAFF	98	EPJ C1 21	K. Ackerstaff+	(OPAL Collab.)
ACKERSTAFF	98C	EPJ C1 45	K. Ackerstaff+	(OPAL Collab.)
ACKERSTAFF	98V	EPJ C2 441	K. Ackerstaff+	(OPAL Collab.)
ACKER...K...	98B	PL B438 379	K. Ackerstaff+	(OPAL Collab.)
BARATE	98J	PL B429 201	R. Barate+	(ALEPH Collab.)
BARATE	98U	EPJ C4 571	R. Barate+	(ALEPH Collab.)
MCFARLAND	98	EPJ C1 509	K.S. McFarland+	(CCFR/NuTeV Collab.)
MIURA	98	PR D57 5345	M. Miura+	(VENUS Collab.)
ABE	97G	PR D55 R5263	+Akimoto, Akopian, Albrow, Amendolia+	(CDF Collab.)
ABE	97T	PRL 79 2198	+Akimoto, Akopian, Albrow, Amendolia+	(CDF Collab.)
ABREU	97B	PL B393 245	+Adam, Adye, Ajinenko, Alekseev+	(DELPHI Collab.)
ABREU	97I	ZPHY C74 57	+Adam, Adye, Ajinenko, Alekseev+	(DELPHI Collab.)
Also	97L	ZPHY C75 580 erratum	Abreu, Adam, Adye, Ajinenko+	(DELPHI Collab.)
ABREU	97J	ZPHY C74 577	P. Abreu+	(DELPHI Collab.)
ACCIARRI	97G	PL B401 139	+Adriani, Aguilar-Benitez, Ahlen, Alpat+	(L3 Collab.)
ACCIARRI	97W	PL B413 159	M. Acciarri+	(L3 Collab.)
ACKERSTAFF	97	PL B391 197	+Alexander, Allison, Altekamp, Ametewee+	(OPAL Collab.)
ACKERSTAFF	97C	PL B391 221	+Alexander, Allison, Altekamp, Ametewee+	(OPAL Collab.)
ADLOFF	97	NP B483 44	+Aid, Anderson, Andreev, Andrieu, Arndt+	(H1 Collab.)
ARIMA	97	PR D55 19	+Odaka, Ogawa, Shirai, Tsuboyama+	(VENUS Collab.)
BREITWEG	97C	ZPHY C76 631	+Derrick, Krakauer, Magill+	(ZEUS Collab.)
DEANDREA	97	PL B409 277		(MARS)
ABE	96	PRL 77 438	+Akimoto, Akopian, Albrow+	(CDF Collab.)
ABE	96S	PRL 77 5336	+Akimoto, Akopian, Albrow, Amendolia+	(CDF Collab.)
ABREU	96K	PL B380 480	+Adam, Adye, Agasi, Ajinenko+	(DELPHI Collab.)
ACCIARRI	96D	PL B370 211	+Adam, Adriani, Aguilar-Benitez, Ahlen+	(L3 Collab.)
ACCIARRI	96L	PL B384 323	+Adam, Adriani, Aguilar-Benitez+	(L3 Collab.)
ALEXANDER	96K	PL B377 222	+Allison, Altekamp, Ametewee+	(OPAL Collab.)
ALEXANDER	96Q	PL B386 463	+De Bonis, Decamp, Ghez, Goy, Lees+	(ALEPH Collab.)
BUSKULIC	96W	PL B385 445	+De Bonis, Decamp, Ghez+	(ALEPH Collab.)
BUSKULIC	96Z	PL B384 333	+Albrow, Amendolia, Amidei, Antos+	(CDF Collab.)
ABE	95N	PRL 74 3538	+Adam, Adriani, Aguilar-Benitez, Ahlen+	(L3 Collab.)
ACCIARRI	95G	PL B353 136	+Andreev, Andrieu, Appuhn, Arpagaus+	(H1 Collab.)
AID	95	PL B353 578	+Krakauer, Magill, Musgrave, Repond+	(ZEUS Collab.)
DERRICK	95B	ZPHY C65 627		

ABE	94	PRL 72 3004	+Albrow, Amidei, Anway-Wiese, Apolinari+	(CDF Collab.)
DIAZCRUZ	94	PR D49 R2149	Diaz Cruz, Sampayo	(CINV)
VELISSARIS	94	PL B331 227	+Lusin, Chung, Park, Cho, Bodek, Kim+	(AMY Collab.)
ABE	93G	PRL 71 2542	+Albrow, Akimoto, Amidei, Anway-Wiese+	(CDF Collab.)
ABT	93	NP B396 3	+Andreev, Andrieu, Appuhn, Arpagaus+	(H1 Collab.)
ADRIANI	93M	PRPL 236 1	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+	(L3 Collab.)
ALITTI	93	NP B400 3	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
BUSKULIC	93Q	ZPHY C59 215	+Decamp, Goy, Lees, Minard, Mours+	(ALEPH Collab.)
DERRICK	93B	PL B316 207	+Krakauer, Magill, Musgrave, Repond+	(ZEUS Collab.)
ABE	92B	PRL 68 1463	+Amidei, Apolinari, Atac, Auchincloss+	(CDF Collab.)
ABE	92D	PRL 68 1104	+Amidei, Apolinari, Atac, Auchincloss+	(CDF Collab.)
ABE	92M	PL 69 2896	+Amidei, Anway-Wiese, Apolinari, Atac+	(CDF Collab.)
ABREU	92C	ZPHY C53 41	+Adam, Adami, Adye, Akesson+	(DELPHI Collab.)
ABREU	92D	ZPHY C53 555	+Adam, Adami, Adye, Akesson, Alekseev+	(DELPHI Collab.)
ADRIANI	92B	PL B288 404	+Aguilar-Benitez, Ahlen, Akbari, Alcaraz+	(L3 Collab.)
ADRIANI	92F	PL B292 472	+Aguilar-Benitez, Ahlen, Akbari, Alcarez+	(L3 Collab.)
ADRIANI	92J	PL B297 469	+Aguilar-Benitez, Ahlen, Alcarez, Aloisio+	(L3 Collab.)
BARDADIN...	92	ZPHY C55 163	Bardadin-Otwinowska	(CLER)
DECAMP	92	PRPL 216 253	+Deschizeaux, Goy, Lees, Minard+	(ALEPH Collab.)
HOWELL	92	PL B291 206	+Koltick, Tauchi, Miyamoto, Kichimi+	(TOPAZ Collab.)
KROHA	92	PR D46 58		(ROCH)
PDG	92	PR D45, 1 June, Part II	Hikasa, Barnett, Stone+	(KEK, LBL, BOST+)
SHIMOZAWA	92	PL B284 144	+Fujimoto, Abe, Adachi, Doser+	(TOPAZ Collab.)
ABE	91D	PRL 67 2418	+Amidei, Apolinari, Atac, Auchincloss+	(CDF Collab.)
ABREU	91E	PL B268 296	+Adam, Adami, Adye, Akesson+	(DELPHI Collab.)
ABREU	91F	NP B367 511	+Adam, Adami, Adye, Akesson+	(DELPHI Collab.)
ADACHI	91	PL B255 613	+Anazawa, Doser, Enomoto+	(TOPAZ Collab.)
AKRAWY	91F	PL B257 531	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ALITTI	91B	PL B257 232	+Ansari, Autiero, Bareyre, Blaylock+	(UA2 Collab.)
BEHREND	91B	ZPHY C51 143	+Criegee, Field, Franke, Jung+	(CELLO Collab.)
BEHREND	91C	ZPHY C51 149	+Criegee, Field, Franke, Jung, Meyer+	(CELLO Collab.)
Also	91B	ZPHY C51 143	Behrend, Criegee, Field, Franke, Jung+	(CELLO Collab.)
ABE	90I	ZPHY C48 13	+Amako, Arai, Asano, Chiba+	(VENUS Collab.)
ADEVA	90F	PL B247 177	+Adriani, Aguilar-Benitez, Akbari, Alcarazz+	(L3 Collab.)
ADEVA	90K	PL B250 199	+Adriani, Aguilar-Benitez, Akbari, Alcarez+	(L3 Collab.)
ADEVA	90L	PL B250 205	+Adriani, Aguilar-Benitez, Akbari, Alcarazz+	(L3 Collab.)
ADEVA	90O	PL B252 525	+Adriani, Aguilar-Benitez, Akbari, Alcarazz+	(L3 Collab.)
AKRAWY	90F	PL B241 133	+Alexander, Allison, Allport+	(OPAL Collab.)
AKRAWY	90I	PL B244 135	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
AKRAWY	90J	PL B246 285	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
DECAMP	90G	PL B236 501	+Deschizeaux, Lees, Minard+	(ALEPH Collab.)
DECAMP	90O	PL B250 172	+Deschizeaux, Goy, Lees+	(ALEPH Collab.)
KIM	90	PL B240 243	+Breedon, Ko, Lander, Maeshima, Malchow+	(AMY Collab.)
ABE	89	PRL 62 613	+Amidei, Apolinari, Ascoli, Atac+	(CDF Collab.)
ABE	89B	PRL 62 1825	+Amidei, Apolinari, Ascoli, Atac+	(CDF Collab.)
ABE	89D	PRL 63 1447	+Amidei, Apolinari, Ascoli, Atac+	(CDF Collab.)
ABE	89H	PRL 62 3020	+Amidei, Apolinari, Ascoli, Atac+	(CDF Collab.)
ABE	89J	ZPHY C45 175	+Amako, Arai, Fukawa+	(VENUS Collab.)
ABE	89L	PL B232 425	+Amako, Arai, Asano, Chiba+	(VENUS Collab.)
ADACHI	89B	PL B228 553	+Aihara, Doser, Enomoto, Fujii+	(TOPAZ Collab.)
ALBAJAR	89	ZPHY C44 15	+Albrow, Allkofer, Arnison, Astbury+	(UA1 Collab.)
BARGER	89	PL B220 464	+Hagiwara, Han, Zeppenfeld	(WISC, KEK)
BEHREND	89B	PL B222 163	+Criegee, Dainton, Field, Franke+	(CELLO Collab.)
BRAUNSCH...	89C	ZPHY C43 549	Braunschweig, Gerhards, Kirschfink+	(TASSO Collab.)
DORENBOS...	89	ZPHY C41 567	Dorenbosch, Udo, Allaby, Amaldi+	(CHARM Collab.)
HAGIWARA	89	PL B219 369	+Sakuda, Terunuma	(KEK, DURH, HIRO)
KIM	89	PL B223 476	+Kim, Kang, Lee, Myung, Bacala	(AMY Collab.)
ABE	88B	PL B213 400	+Amako, Arai, Asano, Chiba, Chiba+	(VENUS Collab.)
BARINGER	88	PL B206 551	+Bylsma, De Bonte, Koltick, Low+	(HRS Collab.)
BRAUNSCH...	88	ZPHY C37 171	Braunschweig, Gerhards+	(TASSO Collab.)
BRAUNSCH...	88D	ZPHY C40 163	Braunschweig, Gerhards, Kirschfink+	(TASSO Collab.)
ANSARI	87D	PL B195 613	+Bagnaia, Banner+	(UA2 Collab.)
BARTEL	87B	ZPHY C36 15	+Becker, Felst+	(JADE Collab.)
BEHREND	87C	PL B191 209	+Buerger, Criegee, Dainton+	(CELLO Collab.)
FERNANDEZ	87B	PR D35 10	+Ford, Qi, Read, Smith, Camporesi+	(MAC Collab.)
ARNISON	86C	PL B172 461	+Albrow, Allkofer+	(UA1 Collab.)
ARNISON	86D	PL B177 244	+Albajar, Albrow+	(UA1 Collab.)
BARTEL	86	ZPHY C31 359	+Becker, Felst, Haidt+	(JADE Collab.)

BARTEL	86C	ZPHY C30 371	+Becker, Cords, Felst, Haidt+	(JADE Collab.)
BEHREND	86	PL 168B 420	+Buerger, Criegee, Fenner+	(CELLO Collab.)
BEHREND	86C	PL B181 178	+Buerger, Criegee, Dainton+	(CELLO Collab.)
DERRICK	86	PL 166B 463	+Gan, Kooijman, Loos+	(HRS Collab.)
Also	86B	PR D34 3286	Derrick, Gan, Kooijman, Loos, Musgrave+	(HRS Collab.)
DERRICK	86B	PR D34 3286	+Gan, Kooijman, Loos, Musgrave+	(HRS Collab.)
GRIFOLS	86	PL 168B 264	+Peris	(BARC)
JODIDIO	86	PR D34 1967	+Balke, Carr, Gidal, Shinsky+	(LBL, NWES, TRIU)
Also	88	PR D37 237 erratum	Jodidio, Balke, Carr+	(LBL, NWES, TRIU)
APPEL	85	PL 160B 349	+Bagnaia, Banner+	(UA2 Collab.)
BARTEL	85K	PL 160B 337	+Becker, Cords, Eichler+	(JADE Collab.)
BERGER	85	ZPHY C28 1	+Genzel, Lackas, Pielorz+	(PLUTO Collab.)
BERGER	85B	ZPHY C27 341	+Deuter, Genzel, Lackas, Pielorz+	(PLUTO Collab.)
BAGNAIA	84C	PL 138B 430	+Banner, Battiston+	(UA2 Collab.)
BARTEL	84D	PL 146B 437	+Becker, Bowdery, Cords+	(JADE Collab.)
BARTEL	84E	PL 146B 121	+Becker, Bowdery, Cords, Felst+	(JADE Collab.)
EICHEN	84	RMP 56 579	+Hinchliffe, Lane, Quigg	(FNAL, LBL, OSU)
ALTHOFF	83C	PL 126B 493	+Fischer, Burkhardt+	(TASSO Collab.)
RENARD	82	PL 116B 264		(CERN)