

# b-baryon ADMIXTURE ( $\Lambda_b, \Xi_b, \Sigma_b, \Omega_b$ )

## b-baryon ADMIXTURE MEAN LIFE

Each measurement of the  $b$ -baryon mean life is an average over an admixture of various  $b$  baryons which decay weakly. Different techniques emphasize different admixtures of produced particles, which could result in a different  $b$ -baryon mean life.

“OUR EVALUATION” is an average of the data listed below performed by the LEP  $B$  Lifetimes Working Group as described in our review “Production and Decay of  $b$ -flavored Hadrons” in the  $B^\pm$  Section of these Listings. The averaging procedure takes into account correlations between the measurements and asymmetric lifetime errors.

<u>VALUE (<math>10^{-12}</math> s)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.20 ± 0.07 OUR EVALUATION</b>				
1.20 ± 0.08 ± 0.06		<sup>1</sup> BARATE	98D ALEP	$e^+e^- \rightarrow Z$
1.46 <sup>+0.22+0.07</sup> <sub>-0.21-0.09</sub>		ABREU	96D DLPH	Excess $\Lambda\ell^-\pi^+$ , decay lengths
1.10 <sup>+0.19</sup> <sub>-0.17</sub> ± 0.09		ABREU	96D DLPH	Excess $\Lambda\mu^-$ impact parameters
1.16 ± 0.11 ± 0.06		AKERS	96 OPAL	Excess $\Lambda\ell^-$ , decay lengths and impact parameters
1.27 <sup>+0.35</sup> <sub>-0.29</sub> ± 0.09		ABREU	95S DLPH	Excess $p\mu^-$ , decay lengths
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.25 ± 0.11 ± 0.05		<sup>2</sup> ABREU	96D DLPH	Combined result
1.05 <sup>+0.12</sup> <sub>-0.11</sub> ± 0.09	290	BUSKULIC	95L ALEP	Repl. by BARATE 98D
1.04 <sup>+0.48</sup> <sub>-0.38</sub> ± 0.10	11	<sup>3</sup> ABREU	93F DLPH	Excess $\Lambda\mu^-$ , decay lengths
1.05 <sup>+0.23</sup> <sub>-0.20</sub> ± 0.08	157	<sup>4</sup> AKERS	93 OPAL	Excess $\Lambda\ell^-$ , decay lengths
1.12 <sup>+0.32</sup> <sub>-0.29</sub> ± 0.16	101	<sup>5</sup> BUSKULIC	92I ALEP	Excess $\Lambda\ell^-$ , impact parameters

<sup>1</sup> Measured using the excess of  $\Lambda\ell^-$ , lepton impact parameter.

<sup>2</sup> Combined result of the three ABREU 96D methods and ABREU 95S.

<sup>3</sup> ABREU 93F superseded by ABREU 96D.

<sup>4</sup> AKERS 93 superseded by AKERS 96.

<sup>5</sup> BUSKULIC 92I superseded by BUSKULIC 95L.

## **$b$ -baryon ADMIXTURE ( $\Lambda_b, \Xi_b, \Sigma_b, \Omega_b$ )**

These branching fractions are actually an average over weakly decaying  $b$ -baryons weighted by their production rates in  $Z$  decay (or high-energy  $p\bar{p}$ ), branching ratios, and detection efficiencies. They scale with the LEP  $b$ -baryon production fraction  $B(b \rightarrow b\text{-baryon})$  and are evaluated for our value  $B(b \rightarrow b\text{-baryon}) = (10.1^{+3.9}_{-3.1})\%$ .

The branching fractions  $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{ anything})$  and  $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})$  are not pure measurements because the underlying measured products of these with  $B(b \rightarrow b\text{-baryon})$  were used to determine  $B(b \rightarrow b\text{-baryon})$ , as described in the note "Production and Decay of  $b$ -Flavored Hadrons."

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $p\mu^- \bar{\nu}$ anything	$(4.9 \pm 2.4)\%$
$\Gamma_2$ $p\ell \bar{\nu}_\ell$ anything	$(4.7^{+1.6}_{-2.0})\%$
$\Gamma_3$ $p$ anything	$(58^{+24}_{-28})\%$
$\Gamma_4$ $\Lambda \ell^- \bar{\nu}_\ell$ anything	$(3.1^{+1.0}_{-1.2})\%$
$\Gamma_5$ $\Lambda \ell^+ \nu_\ell$ anything	
$\Gamma_6$ $\Lambda$ anything	
$\Gamma_7$ $\Lambda_c^+ \ell^- \bar{\nu}_\ell$ anything	
$\Gamma_8$ $\Lambda/\bar{\Lambda}$ anything	$(35^{+12}_{-14})\%$
$\Gamma_9$ $\Xi^- \ell^- \bar{\nu}_\ell$ anything	$(5.5^{+2.0}_{-2.4}) \times 10^{-3}$

## **$b$ -baryon ADMIXTURE ( $\Lambda_b, \Xi_b, \Sigma_b, \Omega_b$ ) BRANCHING RATIOS**

$\Gamma(p\mu^- \bar{\nu} \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.049^{+0.018+0.015}_{-0.015-0.019}$	125	<sup>6</sup> ABREU	95s DLPH	$e^+e^- \rightarrow Z$	

<sup>6</sup> ABREU 95s reports  $[B(b\text{-baryon} \rightarrow p\mu^- \bar{\nu} \text{ anything}) \times B(\bar{b} \rightarrow b\text{-baryon})] = 0.0049 \pm 0.0011^{+0.0015}_{-0.0011}$ . We divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (10.1^{+3.9}_{-3.1}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\ell \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_2/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT		
$0.047 \pm 0.008^{+0.014}_{-0.018}$	<sup>7</sup> BARATE	98v ALEP	$e^+e^- \rightarrow Z$		

<sup>7</sup> BARATE 98v reports  $[B(b\text{-baryon} \rightarrow p\ell \bar{\nu}_\ell \text{ anything}) \times B(\bar{b} \rightarrow b\text{-baryon})] = (4.72 \pm 0.66 \pm 0.44) \times 10^{-3}$ . We divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (10.1^{+3.9}_{-3.1}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\rho\ell\bar{\nu}_\ell \text{ anything})/\Gamma(\rho \text{ anything})$

$\Gamma_2/\Gamma_3$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.080 \pm 0.012 \pm 0.014$	BARATE	98V ALEP	$e^+e^- \rightarrow Z$

### $\Gamma(\Lambda\ell^-\bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$

$\Gamma_4/\Gamma$

The values and averages in this section serve only to show what values result if one assumes our  $B(b \rightarrow b\text{-baryon})$ . They cannot be thought of as measurements since the underlying product branching fractions were also used to determine  $B(b \rightarrow b\text{-baryon})$  as described in the note on "Production and Decay of  $b$ -Flavored Hadrons."

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.031^{+0.010}_{-0.012}$				<b>OUR AVERAGE</b>

$0.032 \pm 0.004^{+0.010}_{-0.012}$		<sup>8</sup> BARATE	98D ALEP	$e^+e^- \rightarrow Z$
$0.029 \pm 0.003^{+0.009}_{-0.011}$		<sup>9</sup> AKERS	96 OPAL	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
$0.030 \pm 0.007^{+0.009}_{-0.011}$	262	<sup>10</sup> ABREU	95S DLPH	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
$0.060 \pm 0.012^{+0.019}_{-0.023}$	290	<sup>11</sup> BUSKULIC	95L ALEP	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	157	<sup>12</sup> AKERS	93 OPAL	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
$0.069 \pm 0.020^{+0.021}_{-0.027}$	101	<sup>13</sup> BUSKULIC	92I ALEP	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$

<sup>8</sup> BARATE 98D reports  $[B(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{ anything}) \times B(\bar{b} \rightarrow b\text{-baryon})] = 0.00326 \pm 0.00016 \pm 0.00039$ . We divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (10.1^{+3.9}_{-3.1}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Measured using the excess of  $\Lambda\ell^-$ , lepton impact parameter.

<sup>9</sup> AKERS 96 reports  $[B(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{ anything}) \times B(\bar{b} \rightarrow b\text{-baryon})] = 0.00291 \pm 0.00023 \pm 0.00025$ . We divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (10.1^{+3.9}_{-3.1}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>10</sup> ABREU 95S reports  $[B(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{ anything}) \times B(\bar{b} \rightarrow b\text{-baryon})] = 0.0030 \pm 0.0006 \pm 0.0004$ . We divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (10.1^{+3.9}_{-3.1}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>11</sup> BUSKULIC 95L reports  $[B(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{ anything}) \times B(\bar{b} \rightarrow b\text{-baryon})] = 0.0061 \pm 0.0006 \pm 0.0010$ . We divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (10.1^{+3.9}_{-3.1}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>12</sup> AKERS 93 superseded by AKERS 96.

<sup>13</sup> BUSKULIC 92I reports  $[B(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{ anything}) \times B(\bar{b} \rightarrow b\text{-baryon})] = 0.0070 \pm 0.0010 \pm 0.0018$ . We divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (10.1^{+3.9}_{-3.1}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by BUSKULIC 95L.

### $\Gamma(\Lambda\ell^+\nu_\ell \text{ anything})/\Gamma(\Lambda \text{ anything})$

$\Gamma_5/\Gamma_6$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.070 \pm 0.012 \pm 0.007$	ACKERSTAFF	97N OPAL	$e^+e^- \rightarrow Z$

$\Gamma(\Lambda/\bar{\Lambda}\text{anything})/\Gamma_{\text{total}}$

$\Gamma_8/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

**0.35<sup>+0.12</sup><sub>-0.14</sub> OUR AVERAGE**

0.39 ± 0.06<sup>+0.12</sup><sub>-0.15</sub> 14 ACKERSTAFF 97N OPAL e<sup>+</sup>e<sup>-</sup> → Z

0.22<sup>+0.12+0.07</sup><sub>-0.08-0.09</sub> 15 ABREU 95C DLPH e<sup>+</sup>e<sup>-</sup> → Z

14 ACKERSTAFF 97N reports [B(*b*-baryon →  $\Lambda/\bar{\Lambda}$ anything) × B( $\bar{b}$  → *b*-baryon)] = 0.0393 ± 0.0046 ± 0.0037. We divide by our best value B( $\bar{b}$  → *b*-baryon) = (10.1<sup>+3.9</sup><sub>-3.1</sub>) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

15 ABREU 95C reports 0.28<sup>+0.17</sup><sub>-0.12</sub> for B( $\bar{b}$  → *b*-baryon) = 0.08 ± 0.02. We rescale to our best value B( $\bar{b}$  → *b*-baryon) = (10.1<sup>+3.9</sup><sub>-3.1</sub>) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Xi^- \ell^- \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}$

$\Gamma_9/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

**0.0055<sup>+0.0020</sup><sub>-0.0024</sub> OUR AVERAGE**

0.0053 ± 0.0013<sup>+0.0016</sup><sub>-0.0021</sub> 16 BUSKULIC 96T ALEP Excess  $\Xi^- \ell^-$  over  $\Xi^- \ell^+$

0.0058 ± 0.0023<sup>+0.0018</sup><sub>-0.0023</sub> 17 ABREU 95V DLPH Excess  $\Xi^- \ell^-$  over  $\Xi^- \ell^+$

16 BUSKULIC 96T reports [B(*b*-baryon →  $\Xi^- \ell^- \bar{\nu}_\ell$ anything) × B( $\bar{b}$  → *b*-baryon)] = 0.00054 ± 0.00011 ± 0.00008. We divide by our best value B( $\bar{b}$  → *b*-baryon) = (10.1<sup>+3.9</sup><sub>-3.1</sub>) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

17 ABREU 95V reports [B(*b*-baryon →  $\Xi^- \ell^- \bar{\nu}_\ell$ anything) × B( $\bar{b}$  → *b*-baryon)] = 0.00059 ± 0.00021 ± 0.0001. We divide by our best value B( $\bar{b}$  → *b*-baryon) = (10.1<sup>+3.9</sup><sub>-3.1</sub>) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

***b*-baryon ADMIXTURE ( $\Lambda_b$ ,  $\Xi_b$ ,  $\Sigma_b$ ,  $\Omega_b$ ) REFERENCES**

BARATE	98D	EPJ C2 197	R. Barate+	(ALEPH Collab.)
BARATE	98V	EPJ C5 205	R. Barate+	(ALEPH Collab.)
ACKERSTAFF	97N	ZPHY C74 423	K. Ackerstaff+	(OPAL Collab.)
ABREU	96D	ZPHY C71 199	+Adam, Adye, Agasi+	(DELPHI Collab.)
AKERS	96	ZPHY C69 195	+Alexander, Allison, Altekamp+	(OPAL Collab.)
BUSKULIC	96T	PL B384 449	+De Bonis, Decamp, Ghez+	(ALEPH Collab.)
ABREU	95C	PL B347 447	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95S	ZPHY C68 375	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU	95V	ZPHY C68 541	+Adam, Adye, Agasi+	(DELPHI Collab.)
BUSKULIC	95L	PL B357 685	+Casper, De Bonis, Decamp+	(ALEPH Collab.)
ABREU	93F	PL B311 379	+Adam, Adye, Agasi+	(DELPHI Collab.)
AKERS	93	PL B316 435	+Alexander, Allison, Anderson+	(OPAL Collab.)
BUSKULIC	92I	PL B297 449	+Decamp, Goy, Lees+	(ALEPH Collab.)