



$$I(J^P) = 0(0^-)$$

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

B_s^0 MASS

The fit uses m_{B^+} , $(m_{B^0} - m_{B^+})$, $m_{B_s^0}$, and $(m_{B_s^0} - (m_{B^+} + m_{B^0})/2)$ to determine m_{B^+} , m_{B^0} , $m_{B_s^0}$, and the mass differences.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5369.3± 2.0 OUR FIT				
5369.6± 2.4 OUR AVERAGE				
5369.9± 2.3±1.3	32	¹ ABE	96B CDF	$p\bar{p}$ at 1.8 TeV
5374 ±16 ±2	3	ABREU	94D DLPH	$e^+e^- \rightarrow Z$
5359 ±19 ±7	1	¹ AKERS	94J OPAL	$e^+e^- \rightarrow Z$
5368.6± 5.6±1.5	2	BUSKULIC	93G ALEP	$e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5370 ±40	6	² AKERS	94J OPAL	$e^+e^- \rightarrow Z$
5383.3± 4.5±5.0	14	ABE	93F CDF	Repl by ABE 96B
¹ From the decay $B_s \rightarrow J/\psi(1S)\phi$.				
² From the decay $B_s \rightarrow D_s^- \pi^+$.				

$m_{B_s^0} - m_B$

m_B is the average of our B masses $(m_{B^\pm} + m_{B^0})/2$. The fits uses m_{B^+} , $(m_{B^0} - m_{B^+})$, $m_{B_s^0}$, and $m_{B_s^0} - m_B$ to determine m_{B^+} , m_{B^0} , $m_{B_s^0}$, and the mass differences.

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
90.2±2.2 OUR FIT				
89.7±2.7±1.2		ABE	96B CDF	$p\bar{p}$ at 1.8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
80 to 130	68	LEE-FRANZINI 90	CSB2	$e^+e^- \rightarrow \gamma(5S)$

$m_{B_{sH}^0} - m_{B_{sL}^0}$

See the B_s^0 - \overline{B}_s^0 MIXING section near the end of these B_s^0 Listings.

B_s^0 MEAN LIFE

"OUR EVALUATION" is an average of the data listed below performed by the LEP B Lifetimes Working Group as described in our review "Production and Decay of b -flavored Hadrons" in the B^\pm Section of the Listings. The averaging procedure takes into account correlations between the measurements and asymmetric lifetime errors.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.54±0.07 OUR EVALUATION				
1.34 $^{+0.23}_{-0.19}$ ± 0.05		3 ABE	98B CDF	$p\bar{p}$ at 1.8 TeV
1.72 $^{+0.20}_{-0.19}$ ± 0.18		4 ACKERSTAFF	98F OPAL	$e^+ e^- \rightarrow Z$
1.50 $^{+0.16}_{-0.15}$ ± 0.04		5 ACKERSTAFF	98G OPAL	$e^+ e^- \rightarrow Z$
1.47 ± 0.14 ± 0.08		6 BARATE	98C ALEP	$e^+ e^- \rightarrow Z$
1.56 $^{+0.29}_{-0.26}$ ± 0.08		5 ABREU	96F DLPH	$e^+ e^- \rightarrow Z$
1.65 $^{+0.34}_{-0.31}$ ± 0.12		6 ABREU	96F DLPH	$e^+ e^- \rightarrow Z$
1.76 ± 0.20 ± 0.15		7 ABREU	96F DLPH	$e^+ e^- \rightarrow Z$
1.60 ± 0.26 ± 0.13		8 ABREU	96F DLPH	$e^+ e^- \rightarrow Z$
1.54 $^{+0.14}_{-0.13}$ ± 0.04		5 BUSKULIC	96M ALEP	$e^+ e^- \rightarrow Z$
1.42 $^{+0.27}_{-0.23}$ ± 0.11	76	5 ABE	95R CDF	$p\bar{p}$ at 1.8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.51 ± 0.11		9 BARATE	98C ALEP	$e^+ e^- \rightarrow Z$
1.34 $^{+0.23}_{-0.19}$ ± 0.05		10 ABE	96N CDF	Repl. by ABE 98B
1.67 ± 0.14		11 ABREU	96F DLPH	$e^+ e^- \rightarrow Z$
1.61 $^{+0.30}_{-0.29}$ ± 0.18	90	6 BUSKULIC	96E ALEP	Repl. by BARATE 98C
1.74 $^{+1.08}_{-0.69}$ ± 0.07	8	12 ABE	95R CDF	Sup. by ABE 96N
1.54 $^{+0.25}_{-0.21}$ ± 0.06	79	5 AKERS	95G OPAL	Repl. by ACKER-STAFF 98G
1.59 $^{+0.17}_{-0.15}$ ± 0.03	134	5 BUSKULIC	95O ALEP	Sup. by BUSKULIC 96M
0.96 ± 0.37	41	13 ABREU	94E DLPH	Sup. by ABREU 96F
1.92 $^{+0.45}_{-0.35}$ ± 0.04	31	5 BUSKULIC	94C ALEP	Sup. by BUSKULIC 95O
1.13 $^{+0.35}_{-0.26}$ ± 0.09	22	5 ACTON	93H OPAL	Sup. by AKERS 95G

³ Measured using fully reconstructed $B_s \rightarrow J/\psi(1S)\phi$ decay.

⁴ ACKERSTAFF 98F use fully reconstructed $D_s^- \rightarrow \phi\pi^-$ and $D_s^- \rightarrow K^{*0}K^-$ in the inclusive B_s^0 decay.

⁵ Measured using $D_s^- \ell^+$ vertices.

⁶ Measured using D_s hadron vertices.

⁷ Measured using $\phi\ell$ vertices.

⁸ Measured using inclusive D_s vertices.

⁹ Combined results from $D_s^- \ell^+$ and D_s hadron.

¹⁰ ABE 96N uses 58 ± 12 exclusive $B_s \rightarrow J/\psi(1S)\phi$ events.

¹¹ Combined result for the four ABREU 96F methods.¹² Exclusive reconstruction of $B_s \rightarrow \psi\phi$.¹³ ABREU 94E uses the flight-distance distribution of D_s vertices, ϕ -lepton vertices, and $D_s\mu$ vertices.

$$|\Delta\Gamma_{B_s^0}|/\Gamma_{B_s^0}$$

$\Gamma_{B_s^0}$ and $|\Delta\Gamma_{B_s^0}|$ are the decay rate average and difference between two B_s^0 CP eigenstates.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.67	95	14 ACCIARRI	98S L3	$e^+e^- \rightarrow Z$

¹⁴ ACCIARRI 98S assumes a $\tau_{B_s^0} = 1.49 \pm 0.06$ ps and PDG 98 values of b production fraction.

B_s^0 DECAY MODES

These branching fractions all scale with $B(\bar{b} \rightarrow B_s^0)$, the LEP B_s^0 production fraction. The first four were evaluated using $B(\bar{b} \rightarrow B_s^0) = (10.5^{+1.8}_{-1.7})\%$ and the rest assume $B(\bar{b} \rightarrow B_s^0) = 12\%$.

The branching fraction $B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{anything})$ is not a pure measurement since the measured product branching fraction $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{anything})$ was used to determine $B(\bar{b} \rightarrow B_s^0)$, as described in the note on "Production and Decay of b -Flavored Hadrons."

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 D_s^- \text{anything}$	(92 ± 33) %	
$\Gamma_2 D_s^- \ell^+ \nu_\ell \text{anything}$	[a] (8.1 \pm 2.5) %	
$\Gamma_3 D_s^- \pi^+$	< 13 %	
$\Gamma_4 D_s^+ (*) + D_s^- (*) -$	< 21.8 %	90%
$\Gamma_5 J/\psi(1S)\phi$	(9.3 \pm 3.3) $\times 10^{-4}$	
$\Gamma_6 J/\psi(1S)\pi^0$	< 1.2 $\times 10^{-3}$	90%
$\Gamma_7 J/\psi(1S)\eta$	< 3.8 $\times 10^{-3}$	90%
$\Gamma_8 \psi(2S)\phi$	seen	
$\Gamma_9 \pi^+ \pi^-$	< 1.7 $\times 10^{-4}$	90%
$\Gamma_{10} \pi^0 \pi^0$	< 2.1 $\times 10^{-4}$	90%
$\Gamma_{11} \eta \pi^0$	< 1.0 $\times 10^{-3}$	90%
$\Gamma_{12} \eta \eta$	< 1.5 $\times 10^{-3}$	90%
$\Gamma_{13} \pi^+ K^-$	< 2.1 $\times 10^{-4}$	90%
$\Gamma_{14} K^+ K^-$	< 5.9 $\times 10^{-5}$	90%
$\Gamma_{15} p\bar{p}$	< 5.9 $\times 10^{-5}$	90%
$\Gamma_{16} \gamma\gamma$	< 1.48 $\times 10^{-4}$	90%
$\Gamma_{17} \phi\gamma$	< 7 $\times 10^{-4}$	90%

**Lepton Family number (*LF*) violating modes or
 $\Delta B = 1$ weak neutral current (*B1*) modes**

Γ_{18}	$\mu^+ \mu^-$	<i>B1</i>	< 2.0	$\times 10^{-6}$	90%
Γ_{19}	$e^+ e^-$	<i>B1</i>	< 5.4	$\times 10^{-5}$	90%
Γ_{20}	$e^\pm \mu^\mp$	<i>LF</i>	[<i>b</i>] < 4.1	$\times 10^{-5}$	90%
Γ_{21}	$\phi \nu \bar{\nu}$	<i>B1</i>	< 5.4	$\times 10^{-3}$	90%

[a] Not a pure measurement. See note at head of B_s^0 Decay Modes.

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

B_s^0 BRANCHING RATIOS

$\Gamma(D_s^- \text{ anything})/\Gamma_{\text{total}}$

Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.92 ± 0.33 OUR AVERAGE				
0.81 $\pm 0.24 \pm 0.24$	90	15 BUSKULIC	96E ALEP	$e^+ e^- \rightarrow Z$
1.56 $\pm 0.58 \pm 0.47$	147	16 ACTON	92N OPAL	$e^+ e^- \rightarrow Z$

¹⁵ BUSKULIC 96E separate $c\bar{c}$ and $b\bar{b}$ sources of D_s^+ mesons using a lifetime tag, subtract generic $\bar{b} \rightarrow W^+ \rightarrow D_s^+$ events, and obtain $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \text{ anything}) = 0.088 \pm 0.020 \pm 0.020$ assuming $B(D_s \rightarrow \phi\pi) = (3.5 \pm 0.4) \times 10^{-2}$ and PDG 1994 values for the relative partial widths to other D_s channels. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.105^{+0.018}_{-0.017}$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.

¹⁶ ACTON 92N assume that excess of 147 ± 48 D_s^0 events over that expected from B^0 , B^+ , and $c\bar{c}$ is all from B_s^0 decay. The product branching fraction is measured to be $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \text{ anything}) \times B(D_s^- \rightarrow \phi\pi^-) = (5.9 \pm 1.9 \pm 1.1) \times 10^{-3}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.105^{+0.018}_{-0.017}$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.

$\Gamma(D_s^- \ell^+ \nu \ell \text{ anything})/\Gamma_{\text{total}}$

Γ_2/Γ

The values and averages in this section serve only to show what values result if one assumes our $B(\bar{b} \rightarrow B_s^0)$. They cannot be thought of as measurements since the underlying product branching fractions were also used to determine $B(\bar{b} \rightarrow B_s^0)$ as described in the note on "Production and Decay of *b*-Flavored Hadrons."

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.081 ± 0.025 OUR AVERAGE				
0.076 $\pm 0.012 \pm 0.022$	134	17 BUSKULIC	95O ALEP	$e^+ e^- \rightarrow Z$
0.107 $\pm 0.043 \pm 0.032$		18 ABREU	92M DLPH	$e^+ e^- \rightarrow Z$
0.103 $\pm 0.036 \pm 0.031$	18	19 ACTON	92N OPAL	$e^+ e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.13 $\pm 0.04 \pm 0.04$	27	20 BUSKULIC	92E ALEP	$e^+ e^- \rightarrow Z$

- 17 BUSKULIC 950 use $D_s \ell$ correlations. The measured product branching ratio is $B(\bar{b} \rightarrow B_s) \times B(B_s \rightarrow D_s^- \ell^+ \nu_\ell \text{anything}) = (0.82 \pm 0.09^{+0.13}_{-0.14})\%$ assuming $B(D_s \rightarrow \phi\pi) = (3.5 \pm 0.4) \times 10^{-2}$ and PDG 1994 values for the relative partial widths to the six other D_s channels used in this analysis. Combined with results from $\Upsilon(4S)$ experiments this can be used to extract $B(\bar{b} \rightarrow B_s) = (11.0 \pm 1.2^{+2.5}_{-2.6})\%$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.105^{+0.018}_{-0.017}$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.
- 18 ABREU 92M measured muons only and obtained product branching ratio $B(Z \rightarrow b\bar{b}) \times B(\bar{b} \rightarrow B_s) \times B(B_s \rightarrow D_s \mu^+ \nu_\mu \text{anything}) \times B(D_s \rightarrow \phi\pi) = (18 \pm 8) \times 10^{-5}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.105^{+0.018}_{-0.017}$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$. We use $B(Z \rightarrow b\bar{b}) = 2B(Z \rightarrow b\bar{b}) = 2 \times (0.2212 \pm 0.0019)$.
- 19 ACTON 92N is measured using $D_s \rightarrow \phi\pi^+$ and $K^*(892)^0 K^+$ events. The product branching fraction measured is measured to be $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{anything}) \times B(D_s^- \rightarrow \phi\pi^-) = (3.9 \pm 1.1 \pm 0.8) \times 10^{-4}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.105^{+0.018}_{-0.017}$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.
- 20 BUSKULIC 92E is measured using $D_s \rightarrow \phi\pi^+$ and $K^*(892)^0 K^+$ events. They use $2.7 \pm 0.7\%$ for the $\phi\pi^+$ branching fraction. The average product branching fraction is measured to be $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{anything}) = 0.020 \pm 0.0055^{+0.005}_{-0.006}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.105^{+0.018}_{-0.017}$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$. Superseded by BUSKULIC 950.

$$\Gamma(D_s^- \pi^+)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<0.13	6	21 AKERS	94J OPAL	$e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 1 BUSKULIC 93G ALEP $e^+ e^- \rightarrow Z$

21 AKERS 94J sees ≤ 6 events and measures the limit on the product branching fraction $f(\bar{b} \rightarrow B_s^0) \cdot B(B_s^0 \rightarrow D_s^- \pi^+) < 1.3\%$ at CL = 90%. We divide by our current value $B(\bar{b} \rightarrow B_s^0) = 0.105$.

$$\Gamma(D_s^+ (*) + D_s^0 (-))/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.218	90	BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

$$\Gamma(J/\psi(1S)\phi)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.93 ± 0.28 ± 0.17	22 ABE	96Q CDF	$p\bar{p}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6	1	²³ AKERS	94J OPAL	$e^+ e^- \rightarrow Z$
seen	14	²⁴ ABE	93F CDF	$p\bar{p}$ at 1.8 TeV
seen	1	²⁵ ACTON	92N OPAL	Sup. by AKERS 94J

²² ABE 96Q assumes $f_u = f_d$ and $f_s/f_u = 0.40 \pm 0.06$. Uses $B \rightarrow J/\psi(1S)K$ and $B \rightarrow J/\psi(1S)K^*$ branching fractions from PDG 94. They quote two systematic errors, ± 0.10 and ± 0.14 where the latter is the uncertainty in f_s . We combine in quadrature.

²³ AKERS 94J sees one event and measures the limit on the product branching fraction $f(\bar{b} \rightarrow B_s^0) \cdot B(B_s^0 \rightarrow J/\psi(1S)\phi) < 7 \times 10^{-4}$ at CL = 90%. We divide by our current value $B(\bar{b} \rightarrow B_s^0) = 0.112$.

²⁴ ABE 93F measured using $J/\psi(1S) \rightarrow \mu^+ \mu^-$ and $\phi \rightarrow K^+ K^-$.

²⁵ In ACTON 92N a limit on the product branching fraction is measured to be

$$f(\bar{b} \rightarrow B_s^0) \cdot B(B_s^0 \rightarrow J/\psi(1S)\phi) \leq 0.22 \times 10^{-2}.$$

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$

Γ_6/Γ

VALUE	CL%	DOCUMENT ID	TECN
$<1.2 \times 10^{-3}$	90	²⁶ ACCIARRI	97C L3

²⁶ ACCIARRI 97C assumes B^0 production fraction ($39.5 \pm 4.0\%$) and B_s ($12.0 \pm 3.0\%$).

$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$

Γ_7/Γ

VALUE	CL%	DOCUMENT ID	TECN
$<3.8 \times 10^{-3}$	90	²⁷ ACCIARRI	97C L3

²⁷ ACCIARRI 97C assumes B^0 production fraction ($39.5 \pm 4.0\%$) and B_s ($12.0 \pm 3.0\%$).

$\Gamma(\psi(2S)\phi)/\Gamma_{\text{total}}$

Γ_8/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	1	BUSKULIC	93G ALEP	$e^+ e^- \rightarrow Z$

$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_9/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	²⁸ BUSKULIC	96V ALEP	$e^+ e^- \rightarrow Z$

²⁸ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

$\Gamma(\pi^0 \pi^0)/\Gamma_{\text{total}}$

Γ_{10}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-4}$	90	²⁹ ACCIARRI	95H L3	$e^+ e^- \rightarrow Z$

²⁹ ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

$\Gamma(\eta \pi^0)/\Gamma_{\text{total}}$

Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-3}$	90	³⁰ ACCIARRI	95H L3	$e^+ e^- \rightarrow Z$

³⁰ ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

$\Gamma(\eta \eta)/\Gamma_{\text{total}}$

Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-3}$	90	³¹ ACCIARRI	95H L3	$e^+ e^- \rightarrow Z$

³¹ ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

$\Gamma(\pi^+ K^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.1 × 10⁻⁴	90	32 BUSKULIC	96V ALEP	$e^+ e^- \rightarrow Z$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<2.6 \times 10^{-4}$	90	33 AKERS	94L OPAL	$e^+ e^- \rightarrow Z$
32 BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.				
33 Assumes $B(Z \rightarrow b\bar{b}) = 0.217$ and B_d^0 (B_s^0) fraction 39.5% (12%).				

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.9 × 10⁻⁵	90	34 BUSKULIC	96V ALEP	$e^+ e^- \rightarrow Z$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<1.4 \times 10^{-4}$	90	35 AKERS	94L OPAL	$e^+ e^- \rightarrow Z$
34 BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.				
35 Assumes $B(Z \rightarrow b\bar{b}) = 0.217$ and B_d^0 (B_s^0) fraction 39.5% (12%).				

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.9 × 10⁻⁵	90	36 BUSKULIC	96V ALEP	$e^+ e^- \rightarrow Z$
36 BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.				

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<14.8 × 10⁻⁵	90	37 ACCIARRI	95I L3	$e^+ e^- \rightarrow Z$
37 ACCIARRI 95I assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.				

$\Gamma(\phi\gamma)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7 × 10⁻⁴	90	38 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
38 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.				

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

Test for $\Delta B = 1$ weak neutral current.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.0 × 10⁻⁶	90	39 ABE	98 CDF	$p\bar{p}$ at 1.8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<3.8 \times 10^{-5}$	90	40 ACCIARRI	97B L3	$e^+ e^- \rightarrow Z$
$<8.4 \times 10^{-6}$	90	41 ABE	96L CDF	Repl. by ABE 98
39 ABE 98 assumes production of $\sigma(B^0) = \sigma(B^+)$ and $\sigma(B_s)/\sigma(B^0) = 1/3$. They normalize to their measured $\sigma(B^0, p_T(B) > 6, y < 1.0) = 2.39 \pm 0.32 \pm 0.44 \mu\text{b}$.				
40 ACCIARRI 97B assume PDG 96 production fractions for B^+ , B^0 , B_s , and Λ_b .				
41 ABE 96L assumes B^+/B_s production ratio 3/1. They normalize to their measured $\sigma(B^+, p_T(B) > 6 \text{ GeV}/c, y < 1) = 2.39 \pm 0.54 \mu\text{b}$.				

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

Test for $\Delta B = 1$ weak neutral current.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.4 \times 10^{-5}$	90	42 ACCIARRI	97B L3	$e^+ e^- \rightarrow Z$

42 ACCIARRI 97B assume PDG 96 production fractions for B^+ , B^0 , B_s , and Λ_b .

Γ_{19}/Γ

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$

test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.1 \times 10^{-5}$	90	43 ACCIARRI	97B L3	$e^+ e^- \rightarrow Z$

43 ACCIARRI 97B assume PDG 96 production fractions for B^+ , B^0 , B_s , and Λ_b .

Γ_{20}/Γ

$\Gamma(\phi \nu \bar{\nu})/\Gamma_{\text{total}}$

Test for $\Delta B = 1$ weak neutral current.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.4 \times 10^{-3}$	90	44 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$

44 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

Γ_{21}/Γ

POLARIZATION IN B_s^0 DECAY

Γ_L/Γ in $B_s^0 \rightarrow J/\psi(1S)\phi$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.56 \pm 0.21^{+0.02}_{-0.04}$	19	ABE	95Z CDF	$p\bar{p}$ at 1.8 TeV

B_s^0 - \bar{B}_s^0 MIXING

For a discussion of B_s^0 - \bar{B}_s^0 mixing see the note on “ B^0 - \bar{B}^0 Mixing” in the B^0 Particle Listings above.

χ_s is a measure of the time-integrated B_s^0 - \bar{B}_s^0 mixing probability that produced B_s^0 (\bar{B}_s^0) decays as a \bar{B}_s^0 (B_s^0). Mixing violates $\Delta B \neq 2$ rule.

$$\chi_s = \frac{x_s^2}{2(1+x_s^2)}$$

$$x_s = \frac{\Delta m_{B_s^0}}{\Gamma_{B_s^0}} = (m_{B_{sH}^0} - m_{B_{sL}^0}) \tau_{B_s^0},$$

where H, L stand for heavy and light states of two B_s^0 CP eigenstates and

$$\tau_{B_s^0} = \frac{1}{0.5(\Gamma_{B_{sH}^0} + \Gamma_{B_{sL}^0})}.$$

χ_B at high energy

This is a B - \bar{B} mixing measurement for an admixture of B^0 and B_s^0 at high energy.

$$\chi_B = f'_d \chi_d + f'_s \chi_s$$

where f'_d and f'_s are the branching ratio times production fractions of B_d^0 and B_s^0 mesons relative to all b -flavored hadrons which decay weakly. Mixing violates $\Delta B \neq 2$ rule.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.118 ± 0.006 OUR AVERAGE					
0.131 ± 0.020	± 0.016		45 ABE	97I CDF	$p\bar{p}$ 1.8 TeV
0.1107 ± 0.0062	± 0.0055		46 ALEXANDER	96 OPAL	$e^+e^- \rightarrow Z$
0.121 ± 0.016	± 0.006		47 ABREU	94J DLPH	$e^+e^- \rightarrow Z$
0.123 ± 0.012	± 0.008		ACCIARRI	94D L3	$e^+e^- \rightarrow Z$
0.114 ± 0.014	± 0.008		48 BUSKULIC	94G ALEP	$e^+e^- \rightarrow Z$
0.129 ± 0.022			49 BUSKULIC	92B ALEP	$e^+e^- \rightarrow Z$
0.176 ± 0.031	± 0.032	1112	50 ABE	91G CDF	$p\bar{p}$ 1.8 TeV
0.148 ± 0.029	± 0.017		51 ALBAJAR	91D UA1	$p\bar{p}$ 630 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.136 ± 0.037	± 0.040		52 UENO	96 AMY	e^+e^- at 57.9 GeV
0.144 ± 0.014	+ 0.017 - 0.011		53 ABREU	94F DLPH	Sup. by ABREU 94J
0.131 ± 0.014			54 ABREU	94J DLPH	$e^+e^- \rightarrow Z$
0.157 ± 0.020	± 0.032		55 ALBAJAR	94 UA1	$\sqrt{s} = 630$ GeV
0.121 + 0.044 - 0.040	± 0.017	1665	56 ABREU	93C DLPH	Sup. by ABREU 94J
0.143 + 0.022 - 0.021	± 0.007		57 AKERS	93B OPAL	Sup. by ALEXANDER 96
0.145 + 0.041 - 0.035	± 0.018		58 ACTON	92C OPAL	$e^+e^- \rightarrow Z$
0.121 ± 0.017	± 0.006		59 ADEVA	92C L3	Sup. by ACCIARRI 94D
0.132 ± 0.22	+ 0.015 - 0.012	823	60 DECAMP	91 ALEP	$e^+e^- \rightarrow Z$
0.178 + 0.049 - 0.040	± 0.020		61 ADEVA	90P L3	$e^+e^- \rightarrow Z$
0.17 + 0.15 - 0.08			62,63 WEIR	90 MRK2	e^+e^- 29 GeV
0.21 + 0.29 - 0.15			62 BAND	88 MAC	$E_{cm}^{ee} = 29$ GeV
>0.02		90	62 BAND	88 MAC	$E_{cm}^{ee} = 29$ GeV
0.121 ± 0.047			62,64 ALBAJAR	87C UA1	Repl. by ALBAJAR 91D
<0.12		90	62,65 SCHAAD	85 MRK2	$E_{cm}^{ee} = 29$ GeV

⁴⁵ Uses di-muon events.

⁴⁶ ALEXANDER 96 uses a maximum likelihood fit to simultaneously extract χ as well as the forward-backward asymmetries in $e^+e^- \rightarrow Z \rightarrow b\bar{b}$ and $c\bar{c}$.

⁴⁷ This ABREU 94J result is from 5182 $\ell\ell$ and 279 $A\ell$ events. The systematic error includes 0.004 for model dependence.

⁴⁸ BUSKULIC 94G data analyzed using ee , $e\mu$, and $\mu\mu$ events.

⁴⁹ BUSKULIC 92B uses a jet charge technique combined with electrons and muons.

⁵⁰ ABE 91G measurement of χ is done with $e\mu$ and ee events.

⁵¹ ALBAJAR 91D measurement of χ is done with dimuons.

- 52 UENO 96 extracted χ from the energy dependence of the forward-backward asymmetry.
 53 ABREU 94F uses the average electric charge sum of the jets recoiling against a b -quark jet tagged by a high p_T muon. The result is for $\bar{\chi} = f_d \chi_d + 0.9 f_s \chi_s$.
 54 This ABREU 94J result combines $\ell\ell$, $\Lambda\ell$, and jet-charge ℓ (ABREU 94F) analyses. It is for $\bar{\chi} = f_d \chi_d + 0.96 f_s \chi_s$.
 55 ALBAJAR 94 uses dimuon events. Not independent of ALBAJAR 91D.
 56 ABREU 93C data analyzed using $e\mu$, $\mu\mu$ events.
 57 AKERS 93B analysis performed using dilepton events.
 58 ACTON 92C uses electrons and muons. Superseded by AKERS 93B.
 59 ADEVA 92C uses electrons and muons.
 60 DECAMP 91 done with opposite and like-sign dileptons. Superseded by BUSKULIC 92B.
 61 ADEVA 90P measurement uses $e\mu$, $\mu\mu$, and $e\mu$ events from 118k events at the Z . Superseded by ADEVA 92C.
 62 These experiments are not in the average because the combination of B_s and B_d mesons which they see could differ from those at higher energy.
 63 The WEIR 90 measurement supersedes the limit obtained in SCHAAD 85. The 90% CL are 0.06 and 0.38.
 64 ALBAJAR 87C measured $\chi = (\overline{B}^0 \rightarrow B^0 \rightarrow \mu^+ X)$ divided by the average production weighted semileptonic branching fraction for B hadrons at 546 and 630 GeV.
 65 Limit is average probability for hadron containing B quark to produce a positive lepton.

$$\Delta m_{B_s^0} = m_{B_{sH}^0} - m_{B_{sL}^0}$$

$\Delta m_{B_s^0}$ is a measure of 2π times the B_s^0 - \overline{B}_s^0 oscillation frequency in time-dependent mixing experiments.

"OUR EVALUATION" is an average of the data listed below performed by the LEP B Oscillation Working Group as described in our review "Production and Decays of B -flavored Hadrons" in the B^\pm Section of these Listings. The averaging procedure takes into account correlations between the measurements.

VALUE (10^{12} s^{-1})	CL%	DOCUMENT ID	TECN	COMMENT
>9.1 (CL = 95%) OUR EVALUATION				
>7.9	95	66 BARATE 98C	ALEP	$e^+ e^- \rightarrow Z$
>3.1	95	67 ACKERSTAFF 97U	OPAL	$e^+ e^- \rightarrow Z$
>6.5	95	68 ADAM 97	DLPH	$e^+ e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>2.2	95	69 ACKERSTAFF 97V	OPAL	$e^+ e^- \rightarrow Z$
>6.6	95	70 BUSKULIC 96M	ALEP	Repl. by BARATE 98C
>2.2	95	69 AKERS 95J	OPAL	Sup. by ACKER-STAFF 97V
>5.7	95	71 BUSKULIC 95J	ALEP	$e^+ e^- \rightarrow Z$
>1.8	95	69 BUSKULIC 94B	ALEP	$e^+ e^- \rightarrow Z$

66 BARATE 98C combines results from $D_s h\ell/Q_{\text{hem}}$, $D_s hK$ in the same side, $D_s \ell\ell/Q_{\text{hem}}$ and $D_s \ell K$ in the same side.

67 Uses ℓQ_{hem} .

68 ADAM 97 combines results from $D_s \ell Q_{\text{hem}}$, ℓQ_{hem} , and $\ell\ell$.

69 Uses $\ell\ell$.

70 BUSKULIC 96M uses D_s lepton correlations and lepton, kaon, and jet charge tags.

71 BUSKULIC 95J uses ℓQ_{hem} . They find $\Delta m_s > 5.6$ [> 6.1] for $f_s = 10\%$ [12%]. We interpolate to our central value $f_s = 10.5\%$.

$$x_s = \Delta m_{B_s^0} / \Gamma_{B_s^0}$$

This is derived from "OUR EVALUATION" of $\Delta m_{B_s^0}$ measurements and $\tau_{B_s^0} = 1.54$ ps, our central value.

VALUE	CL%	DOCUMENT ID
>14.0 (CL = 95%) OUR EVALUATION		

χ_s

This B_s^0 - \bar{B}_s^0 integrated mixing parameter is derived from x_s above.

VALUE	CL%	DOCUMENT ID
>0.4975 (CL = 95%) OUR EVALUATION		

B_s^0 REFERENCES

ABE	98	PR D57 R3811	F. Abe+	(CDF Collab.)
ABE	98B	PR D57 5382	F. Abe+	(CDF Collab.)
ACCIARRI	98S	PL B438 417	M. Acciari+	(L3 Collab.)
ACKERSTAFF	98F	EPJ C2 407	K. Ackerstaff+	(OPAL Collab.)
ACKERSTAFF	98G	PL B426 161	K. Ackerstaff+	(OPAL Collab.)
BARATE	98C	EPJ C4 367	R. Barate+	(ALEPH Collab.)
BARATE	98Q	EPJ C4 387	R. Barate+	(ALEPH Collab.)
PDG	98	EPJ C3 1	C. Caso+	
ABE	97I	PR D55 2546	F. Abe+	(CDF Collab.)
ACCIARRI	97B	PL B391 474	M. Acciari+	(L3 Collab.)
ACCIARRI	97C	PL B391 481	M. Acciari+	(L3 Collab.)
ACKERSTAFF	97U	ZPHY C76 401	K. Ackerstaff+	(OPAL Collab.)
ACKERSTAFF	97V	ZPHY C76 417	K. Ackerstaff+	(OPAL Collab.)
ADAM	97	PL B414 382	W. Adam+	(DELPHI Collab.)
ABE	96B	PR D53 3496	+Albrow, Amendolia, Amidei+	(CDF Collab.)
ABE	96L	PRL 76 4675	+Akimoto, Akopian, Albrow+	(CDF Collab.)
ABE	96N	PRL 77 1945	+Akimoto, Akopian, Albrow+	(CDF Collab.)
ABE	96Q	PR D54 6596	+Akimoto, Akopian, Albrow+	(CDF Collab.)
ABREU	96F	ZPHY C71 11	+Adam, Adye, Agasi+	(DELPHI Collab.)
ADAM	96D	ZPHY C72 207	W. Adam+	(DELPHI Collab.)
ALEXANDER	96	ZPHY C70 357	+Allison, Altekamp+	(OPAL Collab.)
BUSKULIC	96E	ZPHY C69 585	+Casper, De Bonis, Decamp+	(ALEPH Collab.)
BUSKULIC	96M	PL B377 205	+De Bonis, Decamp, Ghez+	(ALEPH Collab.)
BUSKULIC	96V	PL B384 471	+De Bonis, Decamp, Ghez+	(ALEPH Collab.)
PDG	96	PR D54 1		
UENO	96	PL B381 365	+Kanda, Olsen, Kirk+	(AMY Collab.)
ABE	95R	PRL 74 4988	+Albrow, Amendolia, Amidei+	(CDF Collab.)
ABE	95Z	PRL 75 3068	+Albrow, Amendolia, Amidei+	(CDF Collab.)
ACCIARRI	95H	PL B363 127	+Adam, Adriani, Aguilar-Benitez+	(L3 Collab.)
ACCIARRI	95I	PL B363 137	+Adam, Adriani, Aguilar-Benitez+	(L3 Collab.)
AKERS	95G	PL B350 273	+Alexander, Allison, Ametewee+	(OPAL Collab.)
AKERS	95J	ZPHY C66 555	+Alexander, Allison, Ametewee+	(OPAL Collab.)
BUSKULIC	95J	PL B356 409	+Casper, De Bonis, Decamp+	(ALEPH Collab.)
BUSKULIC	95O	PL B361 221	+Casper, De Bonis, Decamp+	(ALEPH Collab.)
ABREU	94D	PL B324 500	+Adam, Adye, Agasi, Aleksan+	(DELPHI Collab.)
ABREU	94E	ZPHY C61 407	+Adam, Adye, Agasi, Aleksan+	(DELPHI Collab.)
Also	92M	PL B289 199	Abreu, Adam, Adye, Agasi, Alekseev+	(DELPHI Collab.)
ABREU	94F	PL B322 459	+Adam, Adye, Agasi, Ajinenko+	(DELPHI Collab.)
ABREU	94J	PL B332 488	+Adam, Adye, Agasi, Ajinenko+	(DELPHI Collab.)
ACCIARRI	94D	PL B335 542	+Adam, Adriani, Aguilar-Benitez, Ahlen+	(L3 Collab.)
AKERS	94J	PL B337 196	+Alexander, Allison, Anderson, Arcelli+	(OPAL Collab.)
AKERS	94L	PL B337 393	+Alexander, Allison, Anderson, Arcelli+	(OPAL Collab.)
ALBAJAR	94	ZPHY C61 41	+Ankoviak, Bartha, Bezaguet, Boehrer+	(UA1 Collab.)

BUSKULIC	94B	PL B322 441	+De Bonis, Decamp, Ghez, Goy, Lees+	(ALEPH Collab.)
BUSKULIC	94C	PL B322 275	+De Bonis, Decamp, Ghez, Goy, Lees+	(ALEPH Collab.)
BUSKULIC	94G	ZPHY C62 179	+Casper, De Bonis, Decamp, Ghez+	(ALEPH Collab.)
PDG	94	PR D50 1173	Montanet+	(CERN, LBL, BOST, IFIC+)
ABE	93F	PRL 71 1685	+Albrow, Amidei, Anway-Wiese+	(CDF Collab.)
ABREU	93C	PL B301 145	+Adam, Adye, Agasi, Aleksan+	(DELPHI Collab.)
ACTON	93H	PL B312 501	+Akers, Alexander, Allison, Anderson+	(OPAL Collab.)
AKERS	93B	ZPHY C60 199	+Alexander, Allison, Anderson, Arcelli+	(OPAL Collab.)
BUSKULIC	93G	PL B311 425	+De Bonis, Decamp, Ghez, Goy, Lees+	(ALEPH Collab.)
ABREU	92M	PL B289 199	+Adam, Adye, Agasi, Alekseev+	(DELPHI Collab.)
ACTON	92C	PL B276 379	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ACTON	92N	PL B295 357	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ADEVA	92C	PL B288 395	+Adriani, Aguilar-Benitez, Ahlen+	(L3 Collab.)
BUSKULIC	92B	PL B284 177	+Decamp, Goy, Lees, Minard+	(ALEPH Collab.)
BUSKULIC	92E	PL B294 145	+Decamp, Goy, Lees, Minard+	(ALEPH Collab.)
ABE	91G	PRL 67 3351	+Amidei, Apolinari, Atac, Auchincloss+	(CDF Collab.)
ALBAJAR	91D	PL B262 171	+Albrow, Allkofer, Ankoviak, Apsimon+	(UA1 Collab.)
DECAMP	91	PL B258 236	+Deschizeaux, Goy, Lees, Minard+	(ALEPH Collab.)
ADEVA	90P	PL B252 703	+Adriani, Aguilar-Benitez, Akbari, Alcaraz+	(L3 Collab.)
LEE-FRANZINI	90	PRL 65 2947	+Heintz, Lovelock, Narain, Schamberger+	(CUSB II Collab.)
WEIR	90	PL B240 289	+Abrams, Adolphsen, Alexander, Alvarez+	(Mark II Collab.)
BAND	88	PL B200 221	+Camporesi, Chadwick+	(MAC Collab.)
ALBAJAR	87C	PL B186 247	+Albrow, Allkofer, Arnison+	(UA1 Collab.)
SCHAAD	85	PL 160B 188	+Nelson, Abrams, Amidei+	(Mark II Collab.)