



$I(J^P) = 0(0^-)$

OMITTED FROM SUMMARY TABLE

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

B_c^\pm MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
6.4 $\pm 0.39 \pm 0.13$	¹ ABE	98M CDF	$p\bar{p}$ 1.8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
6.32 ± 0.06	² ACKERSTAFF	980 OPAL	$e^+ e^- \rightarrow Z$
¹ ABE 98M observed $20.4^{+6.2}_{-5.5}$ events in the $B_c^+ \rightarrow J/\psi(1S)\ell\nu_\ell$ with a significance of > 4.8 standard deviations. The mass value is estimated from $m(J/\psi(1S)\ell)$.			
² ACKERSTAFF 980 observed 2 candidate events in the $B_c \rightarrow J/\psi(1S)\pi^+$ channel with an estimated background of 0.63 ± 0.20 events.			

B_c^\pm MEAN LIFE

VALUE (10^{-12} s)	DOCUMENT ID	TECN	COMMENT
0.46 $\pm 0.18 \pm 0.03$	³ ABE	98M CDF	$p\bar{p}$ 1.8 TeV

³ The lifetime is measured from the $J/\psi(1S)\ell$ decay vertices.

B_c^+ DECAY MODES

B_c^- modes are charge conjugates of the modes below.

Mode

Γ_1	$J/\psi(1S)\ell^+\nu_\ell$ anything
Γ_2	$J/\psi(1S)\pi^+$
Γ_3	$J/\psi(1S)\pi^+\pi^-\pi^-$
Γ_4	$J/\psi(1S)a_1(1260)$
Γ_5	$D^*(2010)^+\overline{D}^0$

B_c^+ BRANCHING RATIOS

$\Gamma(J/\psi(1S)\ell^+\nu_\ell \text{anything}) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$	$\Gamma_1 / \Gamma \times B$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$(5.2^{+2.4}_{-2.1}) \times 10^{-5}$		⁴ ABE	98M CDF	$p\bar{p}$ 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.6	$\times 10^{-4}$	90	5 ACKERSTAFF 980 OPAL	$e^+ e^- \rightarrow Z$
< 1.9	$\times 10^{-4}$	90	6 ABREU 97E DLPH	$e^+ e^- \rightarrow Z$
< 1.2	$\times 10^{-4}$	90	7 BARATE 97H ALEP	$e^+ e^- \rightarrow Z$

⁴ ABE 98M result is derived from the measurement of $[\sigma(B_c) \times B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell)] / [\sigma(B^+) \times B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.132^{+0.041}_{-0.037}(\text{stat}) \pm 0.031(\text{sys})^{+0.032}_{-0.020}(\text{lifetime})$

by using PDG 98 values of $B(b \rightarrow B^+)$ and $B(B^+ \rightarrow J/\psi(1S)K^+)$.

⁵ ACKERSTAFF 980 reports $B(Z \rightarrow B_c X) / B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell) < 6.95 \times 10^{-5}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.

⁶ ABREU 97E value listed is for an assumed $\tau_{B_c} = 0.4$ ps and improves to 1.6×10^{-4} for $\tau_{B_c} = 1.4$ ps.

⁷ BARATE 97H reports $B(Z \rightarrow B_c X) / B(Z \rightarrow qq) \cdot B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell) < 5.2 \times 10^{-5}$ at 90%CL. We rescale to our PDG 96 values of $B(Z \rightarrow b\bar{b})$. A $B_c^+ \rightarrow J/\psi(1S)\mu^+\nu_\mu$ candidate event is found, compared to all the known background sources 2×10^{-3} , which gives $m_{B_c} = 5.96^{+0.25}_{-0.19}$ GeV and $\tau_{B_c} = 1.77 \pm 0.17$ ps.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$

$\Gamma_2/\Gamma \times B$

VALUE	CL %	DOCUMENT ID	TECN	COMMENT
$< 8.2 \times 10^{-5}$	90	8 BARATE 97H ALEP	$e^+ e^- \rightarrow Z$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.4 \times 10^{-4}$	90	9 ACKERSTAFF 980 OPAL	$e^+ e^- \rightarrow Z$
$< 3.4 \times 10^{-4}$	90	10 ABREU 97E DLPH	$e^+ e^- \rightarrow Z$
$< 2.0 \times 10^{-5}$	95	11 ABE 96R CDF	$p\bar{p}$ 1.8 TeV

⁸ BARATE 97H reports $B(Z \rightarrow B_c X) / B(Z \rightarrow qq) \cdot B(B_c \rightarrow J/\psi(1S)\pi) < 3.6 \times 10^{-5}$ at 90%CL. We rescale to our PDG 96 values of $B(Z \rightarrow b\bar{b})$.

⁹ ACKERSTAFF 980 reports $B(Z \rightarrow B_c X) / B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)\pi^+) < 1.06 \times 10^{-4}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.

¹⁰ ABREU 97E value listed is for an assumed $\tau_{B_c} = 0.4$ ps and improves to 2.7×10^{-4} for $\tau_{B_c} = 1.4$ ps.

¹¹ ABE 96R reports $B(b \rightarrow B_c X) / B(b \rightarrow B^+ X) \cdot B(B_c^+ \rightarrow J/\psi(1S)\pi^+) / B(B^+ \rightarrow J/\psi(1S)K^+) < 0.053$ at 95%CL for $\tau_{B_c} = 0.8$ ps. It changes from 0.15 to 0.04 for $0.17 \text{ ps} < \tau_{B_c} < 1.6 \text{ ps}$. We rescale to our PDG 96 values of $B(b \rightarrow B^+) = 0.378 \pm 0.022$ and $B(B^+ \rightarrow J/\psi(1S)K^+) = 0.00101 \pm 0.00014$.

$\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$

$\Gamma_3/\Gamma \times B$

VALUE	CL %	DOCUMENT ID	TECN	COMMENT
$< 5.7 \times 10^{-4}$	90	12 ABREU 97E DLPH	$e^+ e^- \rightarrow Z$	

¹² ABREU 97E value listed is independent of $0.4 \text{ ps} < \tau_{B_c} < 1.4 \text{ ps}$.

$\Gamma(J/\psi(1S)a_1(1260))/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$

$\Gamma_4/\Gamma \times B$

VALUE	CL %	DOCUMENT ID	TECN	COMMENT
$< 1.2 \times 10^{-3}$	90	13 ACKERSTAFF 980 OPAL	$e^+ e^- \rightarrow Z$	

¹³ ACKERSTAFF 980 reports $B(Z \rightarrow B_c X) / B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)a_1(1260)) < 5.29 \times 10^{-4}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.

$\Gamma(D^*(2010)^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_5/\Gamma \times B$

VALUE	CL %	DOCUMENT ID	TECN	COMMENT
$< 6.2 \times 10^{-3}$	90	¹⁴ BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$
¹⁴ BARATE 98Q reports $B(Z \rightarrow B_c X) \times B(B_c \rightarrow D^*(2010)^+ \bar{D}^0) < 1.9 \times 10^{-3}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.				

B_c^\pm REFERENCES

ABE	98M	PRL 81 2432	F. Abe+	(CDF Collab.)
Also	98R	PR D58 112004	F. Abe+	(CDF Collab.)
ACKERSTAFF	98O	PL B420 157	K. Ackerstaff+	(OPAL Collab.)
BARATE	98Q	EPJ C4 387	R. Barate+	(ALEPH Collab.)
PDG	98	EPJ C3 1	C. Caso+	
ABREU	97E	PL B398 207	P. Abreu+	(DELPHI Collab.)
BARATE	97H	PL B402 213	R. Barate+	(ALEPH Collab.)
ABE	96R	PRL 77 5176	+Akimoto, Akopian, Albrow+	(CDF Collab.)
PDG	96	PR D54 1		