$\Lambda(2000)$

$I(J^P) = 0(?^?)$ Status: *

OMITTED FROM SUMMARY TABLE

We list here all the ambiguous resonance possibilities with a mass around 2 GeV. The proposed quantum numbers are D_3 (BARBARO-GALTIERI 70 in $\Sigma\pi$), D_3+F_5, P_3+D_5, or P_1+D_3 (BRANDSTETTER 72 in $\Lambda\omega$), and S_1 (CAMERON 78B in $N\overline{K}^*$). The first two of the above analyses should now be considered obsolete. See also NAKKASYAN 75.

$\Lambda(2000)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2000 OUR ESTIMATE			
2030 ± 30	CAMERON 78B	DPWA	$K^- p \to N\overline{K}^*$
1935 to 1971	[1] BRANDSTET...72	DPWA	$K^- p \to \Lambda\omega$
1951 to 2034	[1] BRANDSTET...72	DPWA	$K^- p \to \Lambda\omega$
2010 ± 30	BARBARO-... 70	DPWA	$K^- p \to \Sigma\pi$

$\Lambda(2000)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
125 ± 25	CAMERON 78B	DPWA	$K^- p \to N\overline{K}^*$
180 to 240	[1] BRANDSTET...72	DPWA	(lower mass)
73 to 154	[1] BRANDSTET...72	DPWA	(higher mass)
130 ± 50	BARBARO-... 70	DPWA	$K^- p \to \Sigma\pi$

$\Lambda(2000)$ DECAY MODES

	Mode
Γ_1	$N\overline{K}$
Γ_2	$\Sigma\pi$
Γ_3	$\Lambda\omega$
Γ_4	$N\overline{K}^*(892)$, S=1/2, S-wave
Γ_5	$N\overline{K}^*(892)$, S=3/2, D-wave

$\Lambda(2000)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\overline{K} \to \Lambda(2000) \to \Sigma\pi$ $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
−0.20 ± 0.04	BARBARO-... 70	DPWA	$K^- p \to \Sigma\pi$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\overline{K} \rightarrow \Lambda(2000) \rightarrow \Lambda\omega$ $(\Gamma_1\Gamma_3)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
0.17 to 0.25	[1] BRANDSTET...72	DPWA	(lower mass)
0.04 to 0.15	[1] BRANDSTET...72	DPWA	(higher mass)

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\overline{K} \rightarrow \Lambda(2000) \rightarrow N\overline{K}^*(892)$, S=1/2, S-wave $(\Gamma_1\Gamma_4)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.12 ± 0.03	[2] CAMERON 78B	DPWA	$K^- p \rightarrow N\overline{K}^*$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\overline{K} \rightarrow \Lambda(2000) \rightarrow N\overline{K}^*(892)$, S=3/2, D-wave $(\Gamma_1\Gamma_5)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
$+0.09\pm0.03$	CAMERON 78B	DPWA	$K^- p \rightarrow N\overline{K}^*$

$\Lambda(2000)$ FOOTNOTES

[1] The parameters quoted here are ranges from the three best fits; the lower state probably has $J \leq 3/2$, and the higher one probably has $J \leq 5/2$.

[2] The published sign has been changed to be in accord with the baryon-first convention.

$\Lambda(2000)$ REFERENCES

CAMERON	78B	NP B146 327	W. Cameron *et al.*	(RHEL, LOIC) IJP
NAKKASYAN	75	NP B93 85	A. Nakkasyan	(CERN) IJP
BRANDSTET...	72	NP B39 13	A.A. Brandstetter *et al.*	(RHEL, CDEF+)
BARBARO-...	70	Duke Conf. 173	A. Barbaro-Galtieri	(LRL) IJP