

$f_2(1270)$ $I^G(J^{PC}) = 0^+(2^{++})$ **$f_2(1270)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1275.4 ± 1.2 OUR AVERAGE				
1283 \pm 5		ALDE 98	GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$
1278 \pm 5		1 BERTIN 97C	OBLX	$0.0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
1272 \pm 8	200k	PROKOSHKIN 94	GAM2	$38 \pi^- p \rightarrow \pi^0 \pi^0 n$
1269.7 \pm 5.2	5730	AUGUSTIN 89	DM2	$e^+ e^- \rightarrow 5\pi$
1283 \pm 8	400	2 ALDE 87	GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$
1274 \pm 5		2 AUGUSTIN 87	DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1283 \pm 6		3 LONGACRE 86	MPS	$22 \pi^- p \rightarrow n2K_S^0$
1276 \pm 7		COURAU 84	DLCO	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
1273.3 \pm 2.3		4 CHABAUD 83	ASPK	$17 \pi^- p$ polarized
1280 \pm 4		5 CASON 82	STRC	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
1281 \pm 7	11600	GIDAL 81	MRK2	J/ψ decay
1282 \pm 5		6 CORDEN 79	OMEG	$12-15 \pi^- p \rightarrow n2\pi$
1269 \pm 4	10k	APEL 75	NICE	$40 \pi^- p \rightarrow n2\pi^0$
1272 \pm 4	4600	ENGLER 74	DBC	$6 \pi^+ n \rightarrow \pi^+ \pi^- p$
1277 \pm 4	5300	FLATTE 71	HBC	$7.0 \pi^+ p$
1273 \pm 8		2 STUNTEBECK 70	HBC	$8 \pi^- p, 5.4 \pi^+ d$
1265 \pm 8		BOESEBECK 68	HBC	$8 \pi^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1260 \pm 10		7 ALDE 97	GAM2	$450 pp \rightarrow pp\pi^0\pi^0$
1278 \pm 6		7 GRYGOREV 96	SPEC	$40 \pi^- N \rightarrow K_S^0 K_S^0 X$
1262 \pm 11		AGUILAR-...	EHS	$400 pp$
1275 \pm 10		AKER 91	CBAR	$0.0 \bar{p} p \rightarrow 3\pi^0$
1220 \pm 10		BREAKSTONE 90	SFM	$pp \rightarrow pp\pi^+\pi^-$
1288 \pm 12		ABACHI 86B	HRS	$e^+ e^- \rightarrow \pi^+ \pi^- X$
1284 \pm 30	3k	BINON 83	GAM2	$38 \pi^- p \rightarrow n2\eta$
1280 \pm 20	3k	APEL 82	CNTR	$25 \pi^- p \rightarrow n2\pi^0$
1284 \pm 10	16000	DEUTSCH...	HBC	$16 \pi^+ p$
1258 \pm 10	600	TAKAHASHI 72	HBC	$8 \pi^- p \rightarrow n2\pi$
1275 \pm 13		ARMENISE 70	HBC	$9 \pi^+ n \rightarrow p\pi^+\pi^-$
1261 \pm 5	1960	2 ARMENISE 68	DBC	$5.1 \pi^+ n \rightarrow p\pi^+ MM^-$
1270 \pm 10	360	2 ARMENISE 68	DBC	$5.1 \pi^+ n \rightarrow p\pi^0 MM$
1268 \pm 6		8 JOHNSON 68	HBC	$3.7-4.2 \pi^- p$

¹ T-matrix pole.² Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.³ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.⁴ From an energy-independent partial-wave analysis.⁵ From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$.⁶ From an amplitude analysis of $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ scattering data.⁷ Systematic uncertainties not estimated.⁸ JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

$f_2(1270)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

185.1 $^{+3.4}_{-2.6}$ OUR FIT Error includes scale factor of 1.5.

184.3 $^{+4.0}_{-2.6}$ OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

171	± 10		ALDE	98	GAM4	100	$\pi^- p \rightarrow \pi^0 \pi^0 n$	
204	± 20		⁹ BERTIN	97C	OBLX	0.0	$\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$	
192	± 5	200k	PROKOSHKIN	94	GAM2	38	$\pi^- p \rightarrow \pi^0 \pi^0 n$	
180	± 24		AGUILAR-...	91	EHS	400	$p p$	
169	± 9	5730	AUGUSTIN	89	DM2	$e^+ e^- \rightarrow 5\pi$		
150	± 30	400	ALDE	87	GAM4	100	$\pi^- p \rightarrow 4\pi^0 n$	
186	$+9_{-2}$		LONGACRE	86	MPS	22	$\pi^- p \rightarrow n2K_S^0$	
179.2 $^{+6.9}_{-6.6}$			CHABAUD	83	ASPK	17	$\pi^- p$ polarized	
160	± 11		DENNEY	83	LASS	10	$\pi^+ N$	
196	± 10	3k	APEL	82	CNTR	25	$\pi^- p \rightarrow n2\pi^0$	
152	± 9		¹³ CASON	82	STRC	8	$\pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$	
186	± 27	11600	GIDAL	81	MRK2	J/ψ decay		
216	± 13		CORDEN	79	OMEG	12–15	$\pi^- p \rightarrow n2\pi$	
190	± 10	10k	APEL	75	NICE	40	$\pi^- p \rightarrow n2\pi^0$	
192	± 16	4600	ENGLER	74	DBC	6	$\pi^+ n \rightarrow \pi^+ \pi^- p$	
183	± 15	5300	FLATTE	71	HBC	7	$\pi^+ p \rightarrow \Delta^{++} f_2$	
196	± 30		STUNTEBECK	70	HBC	8	$\pi^- p$, 5.4 $\pi^+ d$	
216	± 20	1960	ARMENISE	68	DBC	5.1	$\pi^+ n \rightarrow p\pi^+ MM^-$	
128	± 27		BOESEBECK	68	HBC	8	$\pi^+ p$	
176	± 21		^{10,15} JOHNSON	68	HBC	3.7–4.2	$\pi^- p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

187	± 20		¹⁶ ALDE	97	GAM2	450	$p p \rightarrow p p \pi^0 \pi^0$	
184	± 10		¹⁶ GRYGOREV	96	SPEC	40	$\pi^- N \rightarrow K_S^0 K_S^0 X$	
200	± 10		AKER	91	CBAR	0.0	$\bar{p}p \rightarrow 3\pi^0$	
240	± 40	3k	BINON	83	GAM2	38	$\pi^- p \rightarrow n2\eta$	
187	± 30	650	¹⁰ ANTIPOV	77	CIBS	25	$\pi^- p \rightarrow p3\pi$	
225	± 38	16000	DEUTSCH...	76	HBC	16	$\pi^+ p$	
166	± 28	600	¹⁰ TAKAHASHI	72	HBC	8	$\pi^- p \rightarrow n2\pi$	
173	± 53		¹⁰ ARMENISE	70	HBC	9	$\pi^+ n \rightarrow p\pi^+ \pi^-$	

⁹T-matrix pole.

¹⁰Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

¹¹From a partial-wave analysis of data using a K-matrix formalism with 5 poles.


¹²From an energy-independent partial-wave analysis.

¹³From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$.

¹⁴From an amplitude analysis of $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ scattering data.

¹⁵JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

¹⁶Systematic uncertainties not estimated.



$f_2(1270)$ width (MeV)

$f_2(1270)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \pi\pi$	(84.7 $^{+2.4}_{-1.3}$) %	S=1.3
$\Gamma_2 \pi^+\pi^-2\pi^0$	(7.1 $^{+1.5}_{-2.6}$) %	S=1.3
$\Gamma_3 K\bar{K}$	(4.6 ± 0.5) %	S=2.8
$\Gamma_4 2\pi^+2\pi^-$	(2.8 ± 0.4) %	S=1.2
$\Gamma_5 \eta\eta$	(4.5 ± 1.0) $\times 10^{-3}$	S=2.4

Γ_6	$4\pi^0$	$(3.0 \pm 1.0) \times 10^{-3}$	
Γ_7	$\gamma\gamma$	$(1.41 \pm 0.13) \times 10^{-5}$	
Γ_8	$\eta\pi\pi$	$< 8 \times 10^{-3}$	CL=95%
Γ_9	$K^0 K^- \pi^+ + \text{c.c.}$	$< 3.4 \times 10^{-3}$	CL=95%
Γ_{10}	$e^+ e^-$	$< 9 \times 10^{-9}$	CL=90%

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 4 partial widths, a combination of partial widths obtained from integrated cross sections, and 6 branching ratios uses 41 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 73.5$ for 34 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-91						
x_3	11	-39					
x_4	11	-36	1				
x_5	2	-9	0	0			
x_6	0	-7	0	0	0		
x_7	11	-7	-9	1	0	0	
Γ	-79	73	-11	-8	-3	0	-15
	x_1	x_2	x_3	x_4	x_5	x_6	x_7

Mode	Rate (MeV)		Scale factor
Γ_1 $\pi\pi$	156.9	± 3.8 -1.3	
Γ_2 $\pi^+ \pi^- 2\pi^0$	13.1	± 3.0 -4.9	1.3
Γ_3 $K\bar{K}$	8.6	± 0.8	2.9
Γ_4 $2\pi^+ 2\pi^-$	5.2	± 0.7	1.2
Γ_5 $\eta\eta$	0.83	± 0.18	2.4
Γ_6 $4\pi^0$	0.55	± 0.19	
Γ_7 $\gamma\gamma$	0.00260	± 0.00024	

$f_2(1270)$ PARTIAL WIDTHS

$\Gamma(\pi\pi)$	Γ_1
<u>VALUE (MeV)</u>	
$156.9^{+3.8}_{-1.3}$ OUR FIT	
$157.0^{+6.0}_{-1.0}$	$^{18} \text{LONGACRE}$ 86 MPS $22 \pi^- p \rightarrow n 2K_S^0$

$\Gamma(K\bar{K})$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3
8.6 ±0.8 OUR FIT Error includes scale factor of 2.9.				
9.0 ±0.7 -0.3	18 LONGACRE	86 MPS	$22 \pi^- p \rightarrow n2K_S^0$	

 $\Gamma(\eta\eta)$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_5
0.83±0.18 OUR FIT Error includes scale factor of 2.4.				
1.0 ±0.1	18 LONGACRE	86 MPS	$22 \pi^- p \rightarrow n2K_S^0$	

 $\Gamma(\gamma\gamma)$

The value of this width depends on the theoretical model used. Unitarised models with scalars give values clustering around $\simeq 2.6$ keV; without an S -wave contribution, values are systematically higher (typically around 3 keV).

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_7
2.60±0.24 OUR FIT					

2.71 ±0.26 OUR AVERAGE

2.84±0.35		BOGLIONE	99 RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$	
2.58±0.13 ^{+0.36} _{-0.27}	19 BEHREND	92 CELL	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.93±0.23±0.32	17 YABUKI	95 VNS			
3.10±0.35±0.35	20 BLINOV	92 MD1	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^-$		
2.27±0.47±0.11	ADACHI	90D TOPZ	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^-$		
3.15±0.04±0.39	BOYER	90 MRK2	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^-$		
3.19±0.16 ^{+0.29} _{-0.28}	MARSISKE	90 CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$		
2.35±0.65	21 MORGAN	90 RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$		
3.19±0.09 ^{+0.22} _{-0.38}	2177 OEST	90 JADE	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$		
3.2 ±0.1 ±0.4	22 AIHARA	86B TPC	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^-$		
2.5 ±0.1 ±0.5	BEHREND	84B CELL	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^-$		
2.85±0.25±0.5	23 BERGER	84 PLUT	$e^+ e^- \rightarrow e^+ e^- 2\pi$		
2.70±0.05±0.20	COURAU	84 DLCO	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^-$		
2.52±0.13±0.38	24 SMITH	84C MRK2	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^-$		
2.7 ±0.2 ±0.6	EDWARDS	82F CBAL	$e^+ e^- \rightarrow e^+ e^- 2\pi^0$		
2.9 ^{+0.6} _{-0.4} ±0.6	25 EDWARDS	82F CBAL	$e^+ e^- \rightarrow e^+ e^- 2\pi^0$		
3.2 ±0.2 ±0.6	BRANDELIK	81B TASS	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^-$		
3.6 ±0.3 ±0.5	ROUSSARIE	81 MRK2	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^-$		
2.3 ±0.8	26 BERGER	80B PLUT	$e^+ e^- \rightarrow$ $e^+ e^- \pi^+ \pi^-$		

¹⁷ With a narrow scalar state around 1220 MeV.

$\Gamma(e^+ e^-)$	Γ_{10}
<u>VALUE (eV)</u>	<u>CL%</u>
<1.7	90
VOROBYEV	88 ND
	$e^+ e^- \rightarrow \pi^0 \pi^0$
18	From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
19	Using a unitarized model with a 300 - 500 keV wide scalar at 1100 MeV.
20	Using the unitarized model of LYTH 85.
21	Error includes spread of different solutions. Data of MARK2 and CRYSTAL BALL used in the analysis. Authors report strong correlations with $\gamma\gamma$ width of $f_0(1370)$: $\Gamma(f_2) + 1/4 \Gamma(f^0) = 3.6 \pm 0.3$ KeV.
22	Radiative corrections modify the partial widths; for instance the COURAU 84 value becomes 2.66 ± 0.21 in the calculation of LANDRO 86.
23	Using the MENNESSIER 83 model.
24	Superseded by BOYER 90.
25	If helicity = 2 assumption is not made.
26	Using mass, width and $B(f_2(1270) \rightarrow 2\pi)$ from PDG 78.

 $f_2(1270) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_7/\Gamma$
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
0.121 ± 0.015 OUR FIT	Error includes scale factor of 1.3.
0.091 ± 0.007 ± 0.027	27 ALBRECHT 90G ARG $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.104 ± 0.007 ± 0.072	28 ALBRECHT 90G ARG $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
27	Using an incoherent background.
28	Using a coherent background.

 $f_2(1270) \text{ BRANCHING RATIOS}$

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>EVTS</u>
0.847 $^{+0.024}_{-0.013}$ OUR FIT	Error includes scale factor of 1.3.
0.837 ± 0.020 OUR AVERAGE	
0.849 ± 0.025	CHABAUD 83 ASPK 17 $\pi^- p$ polarized
0.85 ± 0.05	BEAUPRE 71 HBC 8 $\pi^+ p \rightarrow \Delta^{++} f_2$
0.8 ± 0.04	OH 70 HBC 1.26 $\pi^- p \rightarrow \pi^+ \pi^- n$

$\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma(\pi\pi)$	Γ_2/Γ_1
Should be twice $\Gamma(2\pi^+ 2\pi^-)/\Gamma(\pi\pi)$ if decay is $\rho\rho$. (See ASCOLI 68D.)	
<u>VALUE</u>	<u>EVTS</u>
0.083 $^{+0.019}_{-0.033}$ OUR FIT	Error includes scale factor of 1.3.
0.15 ± 0.06	600 EISENBERG 74 HBC 4.9 $\pi^+ p \rightarrow \Delta^{++} f_2$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.07	EMMS 75D DBC 4 $\pi^+ n \rightarrow p f_2$

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$ Γ_3/Γ_1

We average only experiments which either take into account $f_2(1270)$ - $a_2(1320)$ interference explicitly or demonstrate that $a_2(1320)$ production is negligible.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.055^{+0.005}_{-0.006} OUR FIT				Error includes scale factor of 2.8.

0.040^{+0.005}_{-0.006} OUR AVERAGE

0.037 ^{+0.008} _{-0.021}	ETKIN	82B MPS	23 $\pi^- p \rightarrow n 2 K_S^0$	
0.045 ± 0.009	CHABAUD	81 ASPK	17 $\pi^- p$ polarized	
0.039 ± 0.008	LOVERRE	80 HBC	4 $\pi^- p \rightarrow K\bar{K}N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.036 ± 0.005	29 COSTA...	80 OMEG	1-2.2 $\pi^- p \rightarrow K^+ K^- n$	
0.030 ± 0.005	30 MARTIN	79 RVUE		
0.027 ± 0.009	31 POLYCHRO...	79 STRC	7 $\pi^- p \rightarrow n 2 K_S^0$	
0.025 ± 0.015	EMMS	75D DBC	4 $\pi^+ n \rightarrow p f_2$	
0.031 ± 0.012	20 ADERHOLZ	69 HBC	8 $\pi^+ p \rightarrow K^+ K^- \pi^+ p$	

29 Re-evaluated by CHABAUD 83.

30 Includes PAWLICKI 77 data.

31 Takes into account the $f_2(1270)$ - $f'_2(1525)$ interference.

 $\Gamma(2\pi^+ 2\pi^-)/\Gamma(\pi\pi)$ Γ_4/Γ_1

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.033± 0.005 OUR FIT				Error includes scale factor of 1.2.
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.033± 0.004 OUR AVERAGE				Error includes scale factor of 1.1.

0.024 ± 0.006	160	EMMS	75D DBC	4 $\pi^+ n \rightarrow p f_2$
0.051 ± 0.025	70	EISENBERG	74 HBC	4.9 $\pi^+ p \rightarrow \Delta^{++} f_2$
0.043 ^{+0.007} _{-0.011}	285	LOUIE	74 HBC	3.9 $\pi^- p \rightarrow n f_2$
0.037 ± 0.007	154	ANDERSON	73 DBC	6 $\pi^+ n \rightarrow p f_2$
0.047 ± 0.013	OH		70 HBC	1.26 $\pi^- p \rightarrow \pi^+ \pi^- n$

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_5/Γ_1

<u>VALUE (units 10^{-3})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.5± 1.0 OUR FIT				Error includes scale factor of 2.4.
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.1± 0.8 OUR AVERAGE				Error includes scale factor of 1.3.

2.8 ± 0.7	ALDE	86D GAM4	100 $\pi^- p \rightarrow 2\eta n$
5.2 ± 1.7	BINON	83 GAM2	38 $\pi^- p \rightarrow 2\eta n$

 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$ Γ_5/Γ_1

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.05	95	EDWARDS	82F CBAL	$e^+ e^- \rightarrow e^+ e^- 2\eta$
<0.016	95	EMMS	75D DBC	4 $\pi^+ n \rightarrow p f_2$
<0.09	95	EISENBERG	74 HBC	4.9 $\pi^+ p \rightarrow \Delta^{++} f_2$

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>EVTS</u>
0.0030 ± 0.0010 OUR FIT	
0.003 ± 0.001	400 \pm 50

 Γ_6/Γ $\Gamma(\eta\pi\pi)/\Gamma(\pi\pi)$

<u>VALUE</u>	<u>CL%</u>
<0.010	95

 Γ_8/Γ_1

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
EMMS	75D DBC	$4\pi^+ n \rightarrow p f_2$

 $\Gamma(K^0 K^- \pi^+ + \text{c.c.})/\Gamma(\pi\pi)$

<u>VALUE</u>	<u>CL%</u>
<0.004	95

 Γ_9/Γ_1

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
EMMS	75D DBC	$4\pi^+ n \rightarrow p f_2$

 $f_2(1270)$ REFERENCES

BOGLIONE 99	EPJ C9 11	M. Boglione, M.R. Pennington
ALDE 98	EPJ A3 361	D. Alde <i>et al.</i> (GAM4 Collab.)
Also 99	PAN 62 405	D. Alde <i>et al.</i> (GAMS Collab.)
ALDE 97	PL B397 350	D.M. Alde <i>et al.</i> (GAMS Collab.)
BERTIN 97C	PL B408 476	A. Bertin <i>et al.</i> (OBELIX Collab.)
GRYGOREV 96	PAN 59 2105	V.K. Grigoriev, O.N. Baloshin, B.P. Barkov (ITEP)
	Translated from YAF 59 2187.	
YABUKI 95	JPSJ 54 435	F. Yabuki <i>et al.</i> (VENUS Collab.)
PROKOSHKIN 94	SPD 39 420	Y.D. Prokoshkin, A.A. Kondashov (SERP)
	Translated from DANS 336 613.	
BEHREND 92	ZPHY C56 381	H.J. Behrend (CELLO Collab.)
BLINOV 92	ZPHY C53 33	A.E. Blinov <i>et al.</i> (NOVO)
AGUILAR-...	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i> (LEBC-EHS Collab.)
AKER 91	PL B260 249	E. Aker <i>et al.</i> (Crystal Barrel Collab.)
ADACHI 90D	PL B234 185	I. Adachi <i>et al.</i> (TOPAZ Collab.)
ALBRECHT 90G	ZPHY C48 183	H. Albrecht <i>et al.</i> (ARGUS Collab.)
BOYER 90	PR D42 1350	J. Boyer <i>et al.</i> (Mark II Collab.)
BREAKSTONE 90	ZPHY C48 569	A.M. Breakstone <i>et al.</i> (ISU, BGNA, CERN+)
MARSISKE 90	PR D41 3324	H. Marsiske <i>et al.</i> (Crystal Ball Collab.)
MORGAN 90	ZPHY C48 623	D. Morgan, M.R. Pennington (RAL, DURH)
OEST 90	ZPHY C47 343	T. Oest <i>et al.</i> (JADE Collab.)
AUGUSTIN 89	NP B320 1	J.E. Augustin, G. Cosme (DM2 Collab.)
VOROBIEV 88	SJNP 48 273	P.V. Vorobiev <i>et al.</i> (NOVO)
	Translated from YAF 48 436.	
ALDE 87	PL B198 286	D.M. Alde <i>et al.</i> (LANL, BRUX, SERP, LAPP)
AUGUSTIN 87	ZPHY C36 369	J.E. Augustin <i>et al.</i> (LALO, CLER, FRAS+)
ABACHI 86B	PRL 57 1990	S. Abachi <i>et al.</i> (PURD, ANL, IND, MICH+)
AIHARA 86B	PRL 57 404	H. Aihara <i>et al.</i> (TPC-2 γ Collab.)
ALDE 86D	NP B269 485	D.M. Alde <i>et al.</i> (BELG, LAPP, SERP, CERN+)
LANDRO 86	PL B172 445	M. Landro, K.J. Mork, H.A. Olsen (UTRO)
LONGACRE 86	PL B177 223	R.S. Longacre <i>et al.</i> (BNL, BRAN, CUNY+)
LYTH 85	JPG 11 459	D.H. Lyth
BEHREND 84B	ZPHY C23 223	H.J. Behrend <i>et al.</i> (CELLO Collab.)
BERGER 84	ZPHY C26 199	C. Berger <i>et al.</i> (PLUTO Collab.)
COURAU 84	PL 147B 227	A. Courau <i>et al.</i> (CIT, SLAC)
SMITH 84C	PR D30 851	J.R. Smith <i>et al.</i> (SLAC, LBL, HARV)
BINON 83	NC 78A 313	F.G. Binon <i>et al.</i> (BELG, LAPP, SERP+)
Also 83B	SJNP 38 561	F.G. Binon <i>et al.</i> (BELG, LAPP, SERP+)
	Translated from YAF 38 934.	

CHABAUD	83	NP B223 1	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)
MENNESSIER	83	ZPHY C16 241	G. Mennessier	(MONP)
APEL	82	NP B201 197	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)
EDWARDS	82F	PL 110B 82	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
BRANDELIK	81B	ZPHY C10 117	R. Brandelik <i>et al.</i>	(TASSO Collab.)
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)
ROUSSARIE	81	PL 105B 304	A. Roussarie <i>et al.</i>	(SLAC, LBL)
BERGER	80B	PL 94B 254	C. Berger <i>et al.</i>	(PLUTO Collab.)
COSTA...	80	NP B175 402	G. Costa de Beauregard <i>et al.</i>	(BARI, BONN+)
LOVERRE	80	ZPHY C6 187	P.F. Loverre <i>et al.</i>	(CERN, CDEF, MADR+)
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)
PDG	78	PL 75B	C. Bricman <i>et al.</i>	
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL)
DEUTSCH...	76	NP B103 426	M. Deutschmann <i>et al.</i>	(AACH3, BERL, BONN+)
APEL	75	PL 57B 398	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)
EMMS	75D	NP B96 155	M.J. Emms <i>et al.</i>	(BIRM, DURH, RHEL)
EISENBERG	74	PL 52B 239	Y. Eisenberg <i>et al.</i>	(REHO)
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)
LOUIE	74	PL 48B 385	J. Louie <i>et al.</i>	(SACL, CERN)
ANDERSON	73	PRL 31 562	J.C. Anderson <i>et al.</i>	(CMU, CASE)
TAKAHASHI	72	PR D6 1266	K. Takahashi <i>et al.</i>	(TOHOK, PENN, NDAM+)
BEAUPRE	71	NP B28 77	J.V. Beaupre <i>et al.</i>	(AACH, BERL, CERN)
FLATTE	71	PL 34B 551	S.M. Flatte <i>et al.</i>	(LBL)
ARMENISE	70	LNC 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)
OH	70	PR D1 2494	B.Y. Oh <i>et al.</i>	(WISC, TNTO) JP
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)
ASCOLI	68D	PRL 21 1712	G. Ascoli <i>et al.</i>	(ILL)
BOESEBECK	68	NP B4 501	K. Boesebeck <i>et al.</i>	(AACH, BERL, CERN)
JOHNSON	68	PR 176 1651	P.B. Johnson <i>et al.</i>	(NDAM, PURD, SLAC)
EISNER	67	PR 164 1699	R.L. Eisner <i>et al.</i>	(PURD)
DERADO	65	PRL 14 872	I. Derado <i>et al.</i>	(NDAM)
LEE	64	PRL 12 342	Y.Y. Lee <i>et al.</i>	(MICH)
BONDAR	63	PL 5 153	L. Bondar <i>et al.</i>	(AACH, BIRM, BONN, DESY+)