

$f_0(1370)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

SCALAR MESONS

Written April 2000 by S. Spanier (SLAC) and N.A. Törnqvist (Helsinki).

Introduction: In contrast to the vector and tensor mesons, the identification of the scalar mesons is a long-standing puzzle. The number of publications since our last issue indicates great activity in that field. Scalar resonances are difficult to resolve because of their large decay widths causing a strong overlap between resonances and background, and at the same time, several decay channels open up within a short mass interval. In addition, especially the $\bar{K}K$ and $\eta\eta$ thresholds produce important sharp cusps in the energy dependence of the resonant amplitude. Furthermore, one expects non- $\bar{q}q$ scalar objects, like glueballs and multiquark states, in the mass range below 1800 MeV.

Scalars are produced, for example, in $\bar{p}p$ annihilation (high statistics), πN scattering on polarized/unpolarized targets, central production, J/ψ decays, D - and K -meson decays, $\gamma\gamma$ formation, and ϕ radiative decays. Experiments are accompanied by the development of theoretical models for the reaction amplitudes, which are based on common fundamental principles of two-body unitarity, analyticity, Lorentz invariance, and chiral- and flavor-symmetry using different techniques (K -matrix formalism, N/D -method, Dalitz Tuan ansatz, unitarized quark models with coupled channels, effective chiral field theories like the linear sigma model, etc.).

The mass and width of a resonance are found from the position of the nearest pole in the T matrix (or equivalently, in the S matrix) at an unphysical sheet of the complex energy

plane: $(E - i\frac{\Gamma}{2})$. It is important to realize that only in the case of well-separated resonances, far away from the opening of decay channels, does the naive Breit-Wigner parameterization (or K -matrix pole parameterization) agree with the T -matrix pole position in the amplitude.

In this note, we discuss all light scalars organized in the listings under the entries $(I = 1/2) K^*(1430)$, $(I = 1) a_0(980)$, $a_0(1450)$, and $(I = 0) \sigma$ or $f_0(400-1200)$, $f_0(980)$, $f_0(1370)$, and $f_0(1500)$. This list is minimal and does not necessarily exhaust the list of actual resonances. The $(I = 2) \pi\pi$ and $(I = 3/2) K\pi$ phase shifts do not exhibit any resonant behavior.

See also our notes in previous issues for further comments on *e.g.*, scattering lengths and older papers.

The $I = 1/2$ States: The $K^*(1430)$ (ASTON 88) is perhaps the least controversial of the light scalar mesons. The $K\pi$ phase shift rises smoothly from the threshold, passes 90° at 1350 MeV, and continues to rise to about 170° at 1600 MeV, the first important inelastic threshold $K\eta'(958)$. Thus, it behaves like a single broad, nearly elastic resonance. ABELE 98, analyzing the $\overline{K}K\pi$ channel of $\overline{p}p$ annihilation at rest, finds the T -matrix pole parameters, $m \approx 1430$ MeV and $\Gamma \approx 290$ MeV, while the K -matrix pole of the same data is about 1340 MeV. This agrees with the LASS (ASTON 88) determination.

It should, however, be noted that several authors (BLACK 98, 99, DELBOURGO 98, ISHIDA 99, OLLER 99,99C, BEVEREN 99) have introduced a light “ $\kappa(900)$ ” meson, which in the model interferes destructively with a large background. This makes the existence of a such light state very model dependent.

The $I = 1$ States: Two isovector states are known, the established $a_0(980)$ and the $a_0(1450)$ found by the Crystal

Barrel experiment (AMSLER 94D). Independently of any model about the nature of the $a_0(980)$, the $\overline{K}K$ component in the wave function of this state must be large: the $a_0(980)$ lies close to the opening of the $\overline{K}K$ channel to which it couples strongly. This gives an important cusp-like behavior in the resonant amplitude. Hence, its mass and width parameters are strongly distorted. To reveal its true coupling constants, a coupled channel model with energy-dependent widths and mass shift contributions must be applied.

In our previous editions, the relative coupling $\overline{K}K/\pi\eta$ was only determined indirectly from $f_1(1285)$ (CORDEN 78, DEFOIX 72) or $\eta(1410)$ decays (BAI 90C, BOLTON 92B, AMSLER 95C), or from the line shape observed in the $\pi\eta$ decay mode (FLATTE 76, AMSLER 94D, BUGG 94, JANSSEN 95). From the analysis of $\pi\pi\eta$ and $\overline{K}K\pi$ final states of $\overline{p}p$ annihilation at rest, a relative production ratio $B(\overline{p}p \rightarrow \pi a_0; a_0 \rightarrow \overline{K}K)/B(\overline{p}p \rightarrow \pi a_0; a_0 \rightarrow \pi\eta) = 0.23 \pm 0.05$ is obtained by (ABELE 98). Tuning of the couplings in a coupled channel formula to reproduce the production ratio for the integrated mass distributions gives a relative branching ratio $\Gamma(\overline{K}K)/\Gamma(\pi\eta) = 1.03 \pm 0.14$. The analysis of the $\overline{p}p$ annihilation data also found that the width determined from the T -matrix pole is 92 ± 8 MeV, while the observed width of the peak in the $\pi\eta$ mass spectrum is about 45 MeV. In all measurements listed in our table, the mass position agrees on a value near 980 MeV, but the width takes values between 50 and 300 MeV due to the different applied models.

The $a_0(1450)$ is seen by the Crystal Barrel experiment in its $\pi\eta$, $\overline{K}K$, and $\pi\eta'(958)$ decay modes. The relative couplings to the different final states are found to be close to SU(3)-flavor predictions for an ordinary $\overline{q}q$ meson. The OBELIX experiment (BERTIN 98B) finds two solutions in the $K_S K^\pm \pi^\mp$ final state

of the $\bar{p}p$ annihilation, one at 1480 MeV and one with a mass value close to that of $a_2(1320)$, which is preferred by their fit, and by the low angular momentum in the production. The broad structure at about 1300 MeV observed in $\pi N \rightarrow \bar{K}KN$ reactions needs further confirmation in existence and isospin assignment.

The $I = 0$ States: The $I = 0 J^{PC} = 0^{++}$ sector is the most complex one, both experimentally and theoretically. The data have been obtained from $\pi\pi$, $\bar{K}K$, $\eta\eta$, 4π , and $\eta\eta'(958)$ systems produced in S wave. From the high-statistics data sets collected from $\bar{p}p$ annihilation at rest into $\pi^0 f_0$, where the f_0 decays into the channels mentioned above, one concludes that at least four poles are needed in the mass range from the $\pi\pi$ threshold to about 1600 MeV. The claimed isoscalar resonances are found under separate entries σ or $f_0(400-1200)$, $f_0(980)$, $f_0(1370)$, and $f_0(1500)$.

Below 1100 MeV, the important data come from the $\pi\pi$ and $\bar{K}K$ final states. Information on the $\pi\pi$ S -wave phase shift $\delta_J^I = \delta_0^0$ was already extracted 20 years ago from the πN scattering with unpolarized (GRAYER 74) and polarized targets (BECKER 79), and near threshold from the K_{e4} -decay (ROSSELET 77). The $\pi\pi$ S -wave inelasticity is not accurately known, and the reported $\pi\pi \rightarrow \bar{K}K$ cross sections (WETZEL 76, POLYCHRONAKOS 79, COHEN 80, and ETKIN 82B) may have large uncertainties. The πN data (GRAYER 74, BECKER 79) have been reanalyzed in combination with the $\bar{p}p$ annihilation data (KAMINSKI 97). Two out of four relevant solutions are found, with the S -wave phase shift rising slower than the P wave [$\rho(770)$], which is used as a reference. One of these corresponds to the well-known “down” solution of GRAYER 74. The other “up” solution shows a decrease of the modulus in the mass interval between 800–980 MeV. Both

solutions exhibit a sudden drop in the modulus and inelasticity at 1 GeV, due to the appearance of $f_0(980)$, which is very close to the opening of the $\bar{K}K$ threshold. The phase shift δ_0^0 rises smoothly up to this point, where it jumps by 120° (in the “up”) or 140° (in the “down”) solution to reach 230° , and then both continue to rise slowly.

SVEC 97 suggests the existence of a narrow state at 750 MeV, with a small width of 100 to 200 MeV in his analysis of the πN (polarized) data, from 600 to 900 MeV. Such a solution is also found by (KAMINSKI 97) using the CERN-Munich (-Cracow) data considering both the π - and $a_1(1260)$ -exchange in the reaction amplitudes. However, they show that unitarity is violated for this solution. Therefore, a narrow, light f_0 state below 900 MeV is excluded (KAMINSKI 97, 00). Also, the $2\pi^0$ invariant mass spectra of the $\bar{p}p$ annihilation at rest (AMSLER 95, ABELE 96), and the central collision (ALDE 97), do not show a narrow resonance below 900 MeV, and these data are consistently described with the standard “down” solution (GRAYNER 74, KAMINSKI 97), which allows for the existence of the broad ($\Gamma \approx 500$ MeV) σ listed under $f_0(400-1200)$. The σ is difficult to establish experimentally without models. It is expected to be very broad, and so can be easily distorted by large background from contact terms, crossed channel exchanges, the $f_0((1370))$, and other dynamical features. Further information on this object is expected from the analysis of three body decays of the D meson, *e.g.*, $D \rightarrow \sigma\pi \rightarrow 3\pi$ (E791 experiment).

The $f_0(980)$ interferes destructively with the background leading to a dip in the $\pi\pi$ spectrum at the $\bar{K}K$ threshold. It changes from a dip into a peak structure in the $\pi^0\pi^0$ invariant mass spectrum of the reaction $\pi^-p \rightarrow \pi^0\pi^0n$ (ACHASOV 98E), with increasing four-momentum transfer to the $\pi^0\pi^0$ system,

which means increasing the a_1 -exchange contribution in the amplitude, while the π -exchange decreases.

A meson resonance very well studied experimentally, is the $f_0(1500)$, seen by the Crystal Barrel experiment in five decay modes: $\pi\pi$, $\overline{K}K$, $\eta\eta$, $\eta\eta'(958)$, and 4π (AMSLER 95D, ABELE 96, and ABELE 98). Due to its interference with the $f_0(1370)$, the peak attributed to $f_0(1500)$ can appear shifted in mass to 1590 MeV, where it was observed by the GAMS Collaboration (BINON 83) in the $\eta\eta$ mass spectrum. For the dynamics in the resonant amplitude, they applied a sum of Breit-Wigner functions. In the central production (ANTINORI 95), a peak at a mass of 1450 MeV, having a width of 60 MeV, can be interpreted as the coherent sum of $f_0(1370)$ and $f_0(1500)$. The $\overline{p}p$ and $\overline{n}p/\overline{p}n$ reactions show a single enhancement at 1400 MeV in the invariant 4π mass (GASPERO 93, ADAMO 93, AMSLER 94, and ABELE 96). In the $5\pi^0$ channel (ABELE 96), this structure was resolved into $f_0(1500)$ and $f_0(1370)$, where the latter was found at somewhat lower mass at around 1300 MeV. An additional scalar had to be introduced in the reanalysis of the reaction $J/\psi(1S) \rightarrow \gamma 4\pi$ with a mass above 1700 MeV (BUGG 95). According to these investigations, the $f_0(1500)$ decay proceeds dominantly via $\sigma\sigma \rightarrow 4\pi$, where σ denotes the $\pi\pi$ S wave below the $\overline{K}K$ threshold. The $\overline{K}K$ decay of $f_0(1500)$ is suppressed (ABELE 98).

The determination of the $\pi\pi$ coupling of $f_0(1370)$ is aggravated by the strong overlap with the broad background from the $f_0(400-1200)$. Since it does not show up prominently in the 2π spectra, its mass and width are difficult to determine. As mentioned under the $I = 1$ states section, data on $\pi\pi \rightarrow \overline{K}K$ show an enhancement in the scalar partial wave at around 1300 MeV (WETZEL 76, COHEN 80, POLYCHRONAKOS 79,

COSTA 80, and LONGACRE 86). According to the phase shift, the resonance is found at about 1400 MeV (COHEN 80), while a reanalysis (BUGG 96) claims a trend towards lower mass. The recent three-channel approach (KAMINSKI 99) supports the Crystal Barrel findings, and yields a broad $f_0(1370)$ with a mass above 1400 MeV and a narrow $f_0(1500)$. Here, the $f_0(1370)$ couples more strongly to $\pi\pi$ than to $\bar{K}K$. The $f_0(1370)$ appears explicitly as $\eta\eta$ resonance in the $\pi^0\eta\eta$ final state of the $\bar{p}p$ annihilation at rest (AMSLER 95D). Further information about the $\bar{K}K$ decay of scalars are most welcome, in particular those that can clearly distinguish the $I = 0$ from the $I = 1$ system.

For numerical estimates of coupling constants of the lightest scalars to two pseudoscalars, see ACHASOV 89E,G,I, KAMINSKI 99, AKHMETSHIN 99C. For example, from these estimates, the $f_0(980)$ coupling to $K\bar{K}$ is much larger than its coupling to $\pi\pi$, which is an important constraint to model builders.

Interpretation: Almost every model on scalar states agrees that the $K^*(1430)$ is the quark model $s\bar{u}$ or $s\bar{d}$ state.

If one uses the naive quark model (which may be too naive because of lack of chiral symmetry constraints), it is natural to assume the $f_0(1370)$, $a_0(1450)$, and the $K^*(1430)$ are in the same SU(3) flavor nonet being the $(\bar{u}u + \bar{d}d)$, $u\bar{d}$ and $u\bar{s}$ state, respectively. In this picture, the choice of the ninth member of the nonet is ambiguous. The controversially discussed candidates are $f_0(1500)$ and $f_0(1710)$ (assuming $J = 0$). Compared to the above states, the $f_0(1500)$ is very narrow. Thus, it is unlikely to be their isoscalar partner. It is also too light to be the first radial excitation. Allowing for a gluonic admixture, one can come to an arrangement among these states. See our note on “Non- $\bar{q}q$ states.”

The $f_0(980)$ and $a_0(980)$ are often interpreted as being multiquark states (JAFFE 77), $\bar{K}K$ bound states (WEINSTEIN 90), or vacuum scalars (CLOSE 93A). These pictures are supported by the two-photon widths of these states, which are smaller than expected for naive $\bar{q}q$ mesons neglecting the large $\bar{K}K$ components in the wave function (BARNES 85, LI 91). The results from SND (ACHASOV 98I) reveal a much higher branching ratio for radiative $\phi \rightarrow \gamma f_0$ decays than expected for naive $\bar{q}q$ mesons, but also for $\bar{K}K$ molecules (CLOSE 93B).

On the other hand, the states $f_0(980)$ and $a_0(980)$ may form a low-mass state nonet with the σ as a central ingredient, and the $K^*(1430)$ (or “ $\kappa(900)$ ”). Attempts have been made to start directly from chiral symmetry or chiral Lagrangians (SCADRON 99, OLLER 98, 99, HANNAH 99, IGI 99, ISHIDA 99, and TORNQVIST 99), which all predict the existence of the σ meson near 500 MeV. Hence, *e.g.*, in the chiral linear sigma model, the σ is the $(\bar{u}u + \bar{d}d)$ state, and at the same time, also the chiral partner of the π . Hence, an experimental proof of its existence has become very important.

In the unitarized quark model with coupled channels, six of the light scalars are understood as different manifestations of bare quark model $\bar{q}q$ states (TORNQVIST 82,95,96, BEVEREN 86). The σ , $f_0(980)$, $f_0(1370)$, $a_0(980)$, $a_0(1450)$, and $K^*(1430)$ are described as unitarized remnants of strongly shifted and mixed $\bar{q}q$ 1^3P_0 states using six parameters. The $f_0(980)$ and $f_0(1370)$, as well as $a_0(980)$ and $a_0(1450)$, are two manifestations of the same $\bar{q}q$ state.

QCD sum rule techniques (ELIAS 99) generally find that the lightest scalars are nearly decoupled from $q\bar{q}$, which would suggest a non- $q\bar{q}$ structure. But this is also consistent with them being unitarized remnants of $q\bar{q}$ surrounded by large “clouds” of light mesons (forming part of the $q\bar{q}$ sea).

Other detailed models exist, which arrive at different groupings of the observed resonances. Further publications discussing the light scalar resonances are (see also our previous issues): AU 87, MORGAN 93, ZOU 94B, JANSSEN 95, KLEMPT 95, ANISOVICH 98, LOCHER 98, ACHASOV 98D, NARISON 98, and MINKOWSKI 99.

$f_0(1370)$ T-MATRIX POLE POSITION

Note that $\Gamma \approx 2 \text{Im}(\sqrt{s_{\text{pole}}})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1200–1500)–i(150–250) OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$(1312 \pm 25 \pm 10) - i(109 \pm 22 \pm 15)$	BARBERIS	99D OMEG	450 $p\bar{p} \rightarrow \pi^+ \pi^-$, $K^+ K^-$,
$(1406 \pm 27) - i(80 \pm 6)$	¹ KAMINSKI	99 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
$(1300 \pm 20) - i(120 \pm 20)$	ANISOVICH	98B RVUE	Compilation
$(1290 \pm 15) - i(145 \pm 15)$	BARBERIS	97B OMEG	450 $p\bar{p} \rightarrow p\bar{p} 2(\pi^+ \pi^-)$
$(1548 \pm 40) - i(560 \pm 40)$	BERTIN	97C OBLX	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
$(1380 \pm 40) - i(180 \pm 25)$	ABELE	96B CBAR	0.0 $\bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
$(1300 \pm 15) - i(115 \pm 8)$	BUGG	96 RVUE	
$(1330 \pm 50) - i(150 \pm 40)$	² AMSLER	95B CBAR	$\bar{p}p \rightarrow 3\pi^0$
$(1360 \pm 35) - i(150-300)$	² AMSLER	95C CBAR	$\bar{p}p \rightarrow \pi^0 \eta \eta$
$(1390 \pm 30) - i(190 \pm 40)$	³ AMSLER	95D CBAR	$\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta,$ $\pi^0 \pi^0 \eta$
1346 – i 249	^{4,5} JANSSEN	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
1214 – i 168	^{5,6} TORNQVIST	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi,$ $\eta\pi$
1364 – i 139	AMSLER	94D CBAR	$\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
$(1365^{+20}_{-55}) - i(134 \pm 35)$	ANISOVICH	94 CBAR	$\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta$
$(1340 \pm 40) - i(127^{+30}_{-20})$	⁷ BUGG	94 RVUE	$\bar{p}p \rightarrow 3\pi^0, \eta \eta \pi^0,$ $\eta \pi^0 \pi^0$
$(1430 \pm 5) - i(73 \pm 13)$	⁸ KAMINSKI	94 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
1515 – i 214	^{5,9} ZOU	93 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
1420 – i 220	¹⁰ AU	87 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$

¹ T-matrix pole on sheet –+.

² Supersedes ANISOVICH 94.

³ Coupled-channel analysis of $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta,$ and $\pi^0 \pi^0 \eta$ on sheet IV. Demonstrates explicitly that $f_0(400-1200)$ and $f_0(1370)$ are two different poles.

⁴ Analysis of data from FALVARD 88.

⁵ The pole is on Sheet III. Demonstrates explicitly that $f_0(400-1200)$ and $f_0(1370)$ are two different poles.

⁶ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

⁷ Reanalysis of ANISOVICH 94 data.

⁸T-matrix pole on sheet III.⁹Analysis of data from OCHS 73, GRAYER 74, and ROSSELET 77.¹⁰Analysis of data from OCHS 73, GRAYER 74, BECKER 79, and CASON 83. **$f_0(1370)$ BREIT-WIGNER MASS OR K-MATRIX POLE PARAMETER**VALUE (MeV)
1200 to 1500 OUR ESTIMATEDOCUMENT ID **$\pi\pi$ MODE**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1308±10	BARBERIS	99B OMEG	450 $p p \rightarrow p_S p_f \pi^+ \pi^-$
1315±50	BELLAZZINI	99 GAM4	450 $p p \rightarrow p p \pi^0 \pi^0$
1315±30	ALDE	98 GAM4	100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
1280±55	BERTIN	98 OBLX	0.05–0.405 $\bar{n} p \rightarrow$ $\pi^+ \pi^+ \pi^-$
1186	¹¹ TORNQVIST	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi,$ $\eta\pi$
1472±12	ARMSTRONG	91 OMEG	300 $p p \rightarrow p p \pi\pi,$ $p p K\bar{K}$
1275±20	BREAKSTONE	90 SFM	62 $p p \rightarrow p p \pi^+ \pi^-$
1420±20	AKESSON	86 SPEC	63 $p p \rightarrow p p \pi^+ \pi^-$
1256	FROGGATT	77 RVUE	$\pi^+ \pi^-$ channel

¹¹ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems. **$K\bar{K}$ MODE**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1440±50	BOLONKIN	88 SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1463±9	ETKIN	82B MPS	23 $\pi^- p \rightarrow n 2K_S^0$
1425±15	WICKLUND	80 SPEC	6 $\pi N \rightarrow K^+ K^- N$
~ 1300	POLYCHRO...	79 STRC	7 $\pi^- p \rightarrow n 2K_S^0$

 4π MODE $2(\pi\pi)_S + \rho\rho$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1374±38	AMSLER	94 CBAR	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- 3\pi^0$
1345±12	ADAMO	93 OBLX	$\bar{n} p \rightarrow 3\pi^+ 2\pi^-$
1386±30	GASPERO	93 DBC	0.0 $\bar{p} n \rightarrow 2\pi^+ 3\pi^-$

 $\eta\eta$ MODE

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1430	AMSLER	92 CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \eta\eta$
1220±40	ALDE	86D GAM4	100 $\pi^- p \rightarrow n 2\eta$

$f_0(1370)$ BREIT-WIGNER WIDTH

VALUE (MeV)
200 to 500 OUR ESTIMATE

DOCUMENT ID

$\pi\pi$ MODE

VALUE (MeV) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

222 ± 20	BARBERIS	99B	OMEG	450	$p p \rightarrow p_S p_f \pi^+ \pi^-$
255 ± 60	BELLAZZINI	99	GAM4	450	$p p \rightarrow p p \pi^0 \pi^0$
190 ± 50	ALDE	98	GAM4	100	$\pi^- p \rightarrow \pi^0 \pi^0 n$
323 ± 13	BERTIN	98	OBLX	0.05–0.405	$\bar{n} p \rightarrow$ $\pi^+ \pi^+ \pi^-$
350	¹² TORNQVIST	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi,$ $\eta\pi$	
195 ± 33	ARMSTRONG	91	OMEG	300	$p p \rightarrow p p \pi\pi,$ $p p K\bar{K}$
285 ± 60	BREAKSTONE	90	SFM	62	$p p \rightarrow p p \pi^+ \pi^-$
460 ± 50	AKESSON	86	SPEC	63	$p p \rightarrow p p \pi^+ \pi^-$
~ 400	¹³ FROGGATT	77	RVUE		$\pi^+ \pi^-$ channel

¹² Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CA-SON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

¹³ Width defined as distance between 45 and 135° phase shift.

$K\bar{K}$ MODE

VALUE (MeV) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

250 ± 80	BOLONKIN	88	SPEC	40	$\pi^- p \rightarrow K_S^0 K_S^0 n$
118 ⁺¹³⁸ - 16	ETKIN	82B	MPS	23	$\pi^- p \rightarrow n 2 K_S^0$
160 ± 30	WICKLUND	80	SPEC	6	$\pi N \rightarrow K^+ K^- N$
~ 150	POLYCHRO...	79	STRC	7	$\pi^- p \rightarrow n 2 K_S^0$

4π MODE $2(\pi\pi)_S + \rho\rho$

VALUE (MeV) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

375 ± 61	AMSLER	94	CBAR	0.0	$\bar{p} p \rightarrow \pi^+ \pi^- 3\pi^0$
398 ± 26	ADAMO	93	OBLX		$\bar{n} p \rightarrow 3\pi^+ 2\pi^-$
310 ± 50	GASPERO	93	DBC	0.0	$\bar{p} n \rightarrow 2\pi^+ 3\pi^-$

$\eta\eta$ MODE

VALUE (MeV) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

250	AMSLER	92	CBAR	0.0	$\bar{p} p \rightarrow \pi^0 \eta\eta$
320 ± 40	ALDE	86D	GAM4	100	$\pi^- p \rightarrow n 2\eta$

$f_0(1370)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen
Γ_2 4π	seen
Γ_3 $4\pi^0$	seen
Γ_4 $2\pi^+2\pi^-$	seen
Γ_5 $\pi^+\pi^-2\pi^0$	seen
Γ_6 $\rho\rho$	
Γ_7 $2(\pi\pi)_{S\text{-wave}}$	seen
Γ_8 $\eta\eta$	seen
Γ_9 $K\bar{K}$	seen
Γ_{10} $\gamma\gamma$	seen
Γ_{11} e^+e^-	not seen

$f_0(1370)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$				Γ_{10}
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	

••• We do not use the following data for averages, fits, limits, etc. •••

3.8 ± 1.5	¹⁴ BOGLIONE	99	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$
5.4 ± 2.3	MORGAN	90	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$

¹⁴ Supersedes MORGAN 90.

$\Gamma(e^+e^-)$				Γ_{11}
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT

<20	90	VOROBYEV	88	ND	$e^+e^- \rightarrow \pi^0\pi^0$
-----	----	----------	----	----	---------------------------------

$f_0(1370)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	

••• We do not use the following data for averages, fits, limits, etc. •••

0.26 ± 0.09	BUGG	96	RVUE	
<0.15	¹⁵ AMSLER	94	CBAR	$\bar{p}p \rightarrow \pi^+\pi^-3\pi^0$
<0.20	GASPERO	93	DBC	$0.0 \bar{p}n \rightarrow \text{hadrons}$

¹⁵ Using AMSLER 95B ($3\pi^0$).

$\Gamma(4\pi)/\Gamma_{\text{total}}$				$\Gamma_2/\Gamma = (\Gamma_3+\Gamma_4+\Gamma_5)/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	

••• We do not use the following data for averages, fits, limits, etc. •••

0.80 ± 0.04	GASPERO	93	DBC	$0.0 \bar{p}n \rightarrow \text{hadrons}$
-----------------	---------	----	-----	-------------------------------------------

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$				Γ_3/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	

••• We do not use the following data for averages, fits, limits, etc. •••

seen	ABELE	96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
------	-------	----	------	-----------------------------------

$\Gamma(2\pi^+2\pi^-)/\Gamma(4\pi)$ $\Gamma_4/\Gamma_2 = \Gamma_4/(\Gamma_3+\Gamma_4+\Gamma_5)$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.420±0.014 ¹⁶ GASPERO 93 DBC 0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$

¹⁶ Model-dependent evaluation.

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(4\pi)$ $\Gamma_5/\Gamma_2 = \Gamma_5/(\Gamma_3+\Gamma_4+\Gamma_5)$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.512±0.019 ¹⁷ GASPERO 93 DBC 0.0 $\bar{p}n \rightarrow$ hadrons

¹⁷ Model-dependent evaluation.

$\Gamma(\rho\rho)/\Gamma(2(\pi\pi)_{S\text{-wave}})$ Γ_6/Γ_7

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.6 ±0.2 AMSLER 94 CBAR $\bar{p}p \rightarrow \pi^+\pi^-3\pi^0$

0.58±0.16 GASPERO 93 DBC 0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE DOCUMENT ID TECN

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.35±0.13 BUGG 96 RVUE

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$ Γ_9/Γ_1

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.46±0.15±0.11 BARBERIS 99D OMEG 450 $pp \rightarrow K^+K^-, \pi^+\pi^-$

$f_0(1370)$ REFERENCES

BARBERIS 99B PL B453 316	D. Barberis <i>et al.</i>	(Omega expt.)
BARBERIS 99D PL B462 462	D. Barberis <i>et al.</i>	(Omega expt.)
BELLAZZINI 99 PL B467 296	R. Bellazzini <i>et al.</i>	
BOGLIONE 99 EPJ C9 11	M. Boggione, M.R. Pennington	
KAMINSKI 99 EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	
ALDE 98 EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also 99 PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
ANISOVICH 98B UFN 41 419	V.V. Anisovich <i>et al.</i>	
BERTIN 98 PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BARBERIS 97B PL B413 217	D. Barberis <i>et al.</i>	(WA102 Collab.)
BERTIN 97C PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE 96 PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE 96B PL B385 425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BUGG 96 NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)
AMSLER 95B PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 95C PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 95D PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
JANSEN 95 PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)
TORNQVIST 95 ZPHY C68 647	N.A. Tornqvist	(HELS)
AMSLER 94 PL B322 431	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) JPC
AMSLER 94D PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH 94 PL B323 233	V.V. Anisovich <i>et al.</i>	JPC
BUGG 94 PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
KAMINSKI 94 PR D50 3145	R. Kaminski <i>et al.</i>	(CRAC, IPN)
ADAMO 93 NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.) JPC

GASPERO	93	NP A562 407	M. Gaspero	(ROMAI) JPC
ZOU	93	PR D48 R3948	B.S. Zou, D.V. Bugg	(LOQM)
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)
AU	87	Translated from YAF 48 436.	K.L. Au, D. Morgan, M.R. Pennington	(DURH, RAL)
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
WICKLUND	80	PRL 45 1469	A.B. Wicklund <i>et al.</i>	(ANL)
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)
GRAYR	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)

OTHER RELATED PAPERS

AKHMETSHIN	00C	PL B476 33	R.R. Akhmetshin <i>et al.</i>	(CMD2 Collab.)
KAMINSKI	00	APP B31 895		
SADOVSKY	00	NP A655 131c	S.A. Sadovsky	
BEVEREN	99	EPJ C10 469	E. Van Beveren, G. Rupp	
GODFREY	99	RMP 71 1411	S. Godfrey, J. Napolitano	
ISHIDA	99	PTP 101 661		
MINKOWSKI	99	EPJ C9 283	P. Minkowski, W. Ochs	
TORNQVIST	99	EPJ C11 359	N. Tornqvist	
ACHASOV	98D	PAN 61 224		
ACHASOV	98E	PR D58 054011		
AMSLER	98	RMP 70 1293	C. Amsler	
ANISOVICH	98	PL B437 209	V.V. Anisovich <i>et al.</i>	
BLACK	98	PR D58 054012		
LOCHER	98	EPJ C4 317	M.P. Locher <i>et al.</i>	(PSI)
NARISON	98	NP B509 312		
ANISOVICH	97	PL B395 123	A.V. Anisovich, A.V. Sarantsev	(PNPI)
ANISOVICH	97B	ZPHY A357 123	A.V. Anisovich <i>et al.</i>	(PNPI)
ANISOVICH	97C	PL B413 137		
ANISOVICH	97E	PAN 60 1892	A.V. Anisovich <i>et al.</i>	(PNPI)
KAMINSKI	97	Translated from YAF 60 2065.	R. Kaminski <i>et al.</i>	(CRAC)
PROKOSHKIN	97	ZPHY C74 79	Y.D. Prokoshkin <i>et al.</i>	(SERP)
TORNQVIST	96	Translated from DANS 353 323.	N.A. Tornqvist, M. Roos	(HELS)
GASPERO	95	PRL 76 1575	M. Gaspero	(ROMA)
KLEMPPT	95	NP A588 861		
ZOU	94B	PL B361 160	B.S. Zou, D.V. Bugg	(LOQM)
CLOSE	93A	PR D50 591		
CLOSE	93B	PL B319 291		
MORGAN	93	NP B389 513		
LI	91	PR D48 1185	D. Morgan, M.R. Pennington	(RAL, DURH)
BARNES	85	PR D43 2161	Z.P. Li <i>et al.</i>	(TENN)
BIZZARRI	69	PL B165 434		
BETTINI	66	NP B14 169	R. Bizzarri <i>et al.</i>	(CERN, CDEF)
		NC 42A 695	A. Bettini <i>et al.</i>	(PADO, PISA)