

Index for Searches for Monopoles, Supersymmetry Technicolor, Compositeness, etc. Summary Table

- Page 2 Magnetic Monopole Searches
- Page 2 Supersymmetric Particle Searches
- Page 4 Technicolor
- Page 5 Quark and Lepton Compositeness, Searches for

Navigating PDF files: use CTRL-N to go to the desired page.

SEARCHES FOR MONOPOLES, SUPERSYMMETRY, TECHNICOLOR, COMPOSITENESS, etc.

Magnetic Monopole Searches

Isolated supermassive monopole candidate events have not been confirmed. The most sensitive experiments obtain negative results.

Best cosmic-ray supermassive monopole flux limit:

$$< 1.0 \times 10^{-15} \text{ cm}^{-2}\text{sr}^{-1}\text{s}^{-1} \quad \text{for } 1.1 \times 10^{-4} < \beta < 0.1$$

Supersymmetric Particle Searches

Limits are based on the Minimal Supersymmetric Standard Model.

Assumptions include: 1) $\tilde{\chi}_1^0$ (or $\tilde{\gamma}$) is lightest supersymmetric particle; 2) R -parity is conserved; 3) With the exception of \tilde{t} and \tilde{b} , all scalar quarks are assumed to be degenerate in mass and $m_{\tilde{q}_R} = m_{\tilde{q}_L}$. 4) Limits for sleptons refer to the $\tilde{\ell}_R$ states.

See the Particle Listings for a Note giving details of supersymmetry.

$\tilde{\chi}_i^0$ — neutralinos (mixtures of $\tilde{\gamma}$, \tilde{Z}^0 , and \tilde{H}_i^0)

Mass $m_{\tilde{\chi}_1^0} > 32.5$ GeV, CL = 95%

$$[\tan\beta > 0.7, m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} > 5 \text{ GeV}]$$

Mass $m_{\tilde{\chi}_2^0} > 55.9$ GeV, CL = 95%

$$[\tan\beta > 1.5, m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} > 10 \text{ GeV}]$$

Mass $m_{\tilde{\chi}_3^0} > 106.6$ GeV, CL = 95%

$$[\tan\beta > 1.5, m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} > 10 \text{ GeV}]$$

$\tilde{\chi}_i^\pm$ — charginos (mixtures of \tilde{W}^\pm and \tilde{H}_i^\pm)

Mass $m_{\tilde{\chi}_1^\pm} > 67.7$ GeV, CL = 95%

$$[\tan\beta > 0.7, m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} > 3 \text{ GeV}]$$

\tilde{e} — scalar electron (selectron)

Mass $m > 87.1$ GeV, CL = 95% $[m_{\tilde{e}_R} - m_{\tilde{\chi}_1^0} > 5 \text{ GeV}]$

$\tilde{\mu}$ — scalar muon (smuon)

Mass $m > 82.3$ GeV, CL = 95% $[m_{\tilde{\mu}_R} - m_{\tilde{\chi}_1^0} > 3 \text{ GeV}]$

$\tilde{\tau}$ — scalar tau (stau)

Mass $m > 81.0$ GeV, CL = 95% $[m_{\tilde{\tau}_R} - m_{\tilde{\chi}_1^0} > 8 \text{ GeV}]$

\tilde{q} — scalar quark (squark)

These limits include the effects of cascade decays, evaluated assuming a fixed value of the parameters μ and $\tan\beta$. The limits are weakly sensitive to these parameters over much of parameter space. Limits assume GUT relations between gaugino masses and the gauge coupling.

Mass $m > 250$ GeV, CL = 95% $[\tan\beta = 2, \mu < 0, A = 0]$

\tilde{b} — scalar bottom (sbottom)

Mass m none 40–75 GeV, CL = 95%

$[\tilde{b} \rightarrow b \tilde{\chi}_1^0, \text{ all } \theta_b, m_{\tilde{b}} - m_{\tilde{\chi}_1^0} > 10 \text{ GeV}]$

\tilde{t} — scalar top (stop)

Mass $m > 86.4$ GeV, CL = 95%

$[\tilde{t} \rightarrow t \tilde{\chi}_1^0, \text{ all } \theta_t, m_{\tilde{t}} - m_{\tilde{\chi}_1^0} > 5 \text{ GeV}]$

\tilde{g} — gluino

There is some controversy on whether gluinos in a low-mass window ($1 \lesssim m_{\tilde{g}} \lesssim 5$ GeV) are excluded or not. See the Supersymmetry Listings for details.

The limits summarised here refer to the high-mass region ($m_{\tilde{g}} \gtrsim 5$ GeV), and include the effects of cascade decays, evaluated assuming a fixed value of the parameters μ and $\tan\beta$. The limits are weakly sensitive to these parameters over much of parameter space. Limits assume GUT relations between gaugino masses and the gauge coupling,

Mass $m > 190$ GeV, CL = 95% $[\tan\beta = 2, \mu < 0, A = 0]$

Mass $m > 260$ GeV, CL = 95% $[m_{\tilde{q}} = m_{\tilde{g}}, \tan\beta = 2, \mu < 0, A = 0]$

Technicolor

Searches for a color-octet techni- ρ constrain its mass to be greater than 260 to 480 GeV, depending on allowed decay channels. Similar bounds exist on the color-octet techni- ω .

Quark and Lepton Compositeness, Searches for

Scale Limits Λ for Contact Interactions (the lowest dimensional interactions with four fermions)

If the Lagrangian has the form

$$\pm \frac{g^2}{2\Lambda^2} \bar{\psi}_L \gamma_\mu \psi_L \bar{\psi}_L \gamma^\mu \psi_L$$

(with $g^2/4\pi$ set equal to 1), then we define $\Lambda \equiv \Lambda_{LL}^\pm$. For the full definitions and for other forms, see the Note in the Listings on Searches for Quark and Lepton Compositeness in the full Review and the original literature.

$\Lambda_{LL}^+(eeee)$	> 3.5 TeV, CL = 95%
$\Lambda_{LL}^-(eeee)$	> 3.8 TeV, CL = 95%
$\Lambda_{LL}^+(ee\mu\mu)$	> 4.5 TeV, CL = 95%
$\Lambda_{LL}^-(ee\mu\mu)$	> 4.7 TeV, CL = 95%
$\Lambda_{LL}^+(ee\tau\tau)$	> 3.9 TeV, CL = 95%
$\Lambda_{LL}^-(ee\tau\tau)$	> 4.0 TeV, CL = 95%
$\Lambda_{LL}^+(\ell\ell\ell\ell)$	> 5.3 TeV, CL = 95%
$\Lambda_{LL}^-(\ell\ell\ell\ell)$	> 5.5 TeV, CL = 95%
$\Lambda_{LL}^+(eeqq)$	> 5.4 TeV, CL = 95%
$\Lambda_{LL}^-(eeqq)$	> 6.2 TeV, CL = 95%
$\Lambda_{LL}^+(eebb)$	> 5.6 TeV, CL = 95%
$\Lambda_{LL}^-(eebb)$	> 4.9 TeV, CL = 95%
$\Lambda_{LL}^+(\mu\mu qq)$	> 2.9 TeV, CL = 95%
$\Lambda_{LL}^-(\mu\mu qq)$	> 4.2 TeV, CL = 95%
$\Lambda_{LR}^\pm(\nu_\mu\nu_e\mu e)$	> 3.1 TeV, CL = 90%
$\Lambda_{LL}^+(qqqq)$	> 2.7 TeV, CL = 95%
$\Lambda_{LL}^-(qqqq)$	> 2.4 TeV, CL = 95%
$\Lambda_{LL}^+(\nu\nu qq)$	> 5.0 TeV, CL = 95%
$\Lambda_{LL}^-(\nu\nu qq)$	> 5.4 TeV, CL = 95%

Excited Leptons

The limits from $\ell^{*+} \ell^{*-}$ do not depend on λ (where λ is the $\ell\ell^*$ transition coupling). The λ -dependent limits assume chiral coupling, except for the third limit for e^* which is for nonchiral coupling. For chiral coupling, this limit corresponds to $\lambda_\gamma = \sqrt{2}$.

$e^{*\pm}$ — excited electron

Mass $m > 90.7$ GeV, CL = 95% (from $e^{*+} e^{*-}$)

Mass $m =$ none 30–200 GeV, CL = 95% (from $ep \rightarrow e^* X$)

Mass $m > 91$ GeV, CL = 95% (if $\lambda_Z > 1$)

Mass $m > 306$ GeV, CL = 95% (if $\lambda_\gamma = 1$)

$\mu^{*\pm}$ — excited muon

Mass $m > 90.7$ GeV, CL = 95% (from $\mu^{*+} \mu^{*-}$)

Mass $m > 91$ GeV, CL = 95% (if $\lambda_Z > 1$)

$\tau^{*\pm}$ — excited tau

Mass $m > 89.7$ GeV, CL = 95% (from $\tau^{*+} \tau^{*-}$)

Mass $m > 90$ GeV, CL = 95% (if $\lambda_Z > 0.18$)

ν^* — excited neutrino

Mass $m > 90.0$ GeV, CL = 95% (from $\nu^* \bar{\nu}^*$)

Mass $m > 91$ GeV, CL = 95% (if $\lambda_Z > 1$)

Mass $m =$ none 40–96 GeV, CL = 95% (from $ep \rightarrow \nu^* X$)

q^* — excited quark

Mass $m > 45.6$ GeV, CL = 95% (from $q^* \bar{q}^*$)

Mass $m > 88$ GeV, CL = 95% (if $\lambda_Z > 1$)

Mass $m > 570$ GeV, CL = 95% ($p\bar{p} \rightarrow q^* X$)

Color Sextet and Octet Particles

Color Sextet Quarks (q_6)

Mass $m > 84$ GeV, CL = 95% (Stable q_6)

Color Octet Charged Leptons (ℓ_8)

Mass $m > 86$ GeV, CL = 95% (Stable ℓ_8)

Color Octet Neutrinos (ν_8)

Mass $m > 110$ GeV, CL = 90% ($\nu_8 \rightarrow \nu g$)