



$$I(J^P) = \frac{1}{2}(0^-)$$

## $K_S^0$ MEAN LIFE

For earlier measurements, beginning with BOLDT 58B, see our our 1986 edition, Physics Letters **170B** 130 (1986).

OUR FIT is described in the note on "Fits for  $K_L^0$  CP-Violation Parameters" in the  $K_L^0$  Particle Listings.

VALUE ( $10^{-10}$ s)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.8935±0.0008 OUR FIT</b>				
<b>0.8940±0.0009 OUR AVERAGE</b>				
0.8971±0.0021		BERTANZA 97	NA31	
0.8941±0.0014±0.0009		SCHWINGEN... 95	E773	$\Delta m$ free, $\phi_{+-} = \phi_{SW}$
0.8929±0.0016		GIBBONS 93	E731	
0.8920±0.0044	214k	GROSSMAN 87	SPEC	
0.881 ± 0.009	26k	ARONSON 76	SPEC	
0.8924±0.0032		<sup>1</sup> CARITHERS 75	SPEC	
0.8937±0.0048	6M	GEWENIGER 74B	ASPK	
0.8958±0.0045	50k	<sup>2</sup> SKJEGGESTAD 72	HBC	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.905 ± 0.007		<sup>3</sup> ARONSON 82B	SPEC	
0.856 ± 0.008	19994	<sup>4</sup> DONALD 68B	HBC	
0.872 ± 0.009	20000	<sup>4,5</sup> HILL 68	DBC	

<sup>1</sup>CARITHERS 75 value is for  $m_{K_L^0} - m_{K_S^0}$ .  $\Delta m = 0.5301 \pm 0.0013$ . The  $\Delta m$  dependence of the total decay rate (inverse mean life) is  $\Gamma(K_S^0) = [(1.122 \pm 0.004) + 0.16(\Delta m - 0.5348)/\Delta m] 10^{10}/s$ , or, in terms of meanlife  $\tau_s = 0.8913 \pm 0.0032 - 0.238(\Delta m - 0.5348)$  where  $\Delta m$  and  $\tau_s$  are in units of  $10^{10}\text{fs}^{-1}$  and  $10^{-10}\text{s}$  respectively.

<sup>2</sup>HILL 68 has been changed by the authors from the published value ( $0.865 \pm 0.009$ ) because of a correction in the shift due to  $\eta_{+-}$ . SKJEGGESTAD 72 and HILL 68 give detailed discussions of systematics encountered in this type of experiment.

<sup>3</sup>ARONSON 82 find that  $K_S^0$  mean life may depend on the kaon energy.

<sup>4</sup>Pre-1971 experiments are excluded from the average because of disagreement with later more precise experiments.

<sup>5</sup>HILL 68 has been changed by the authors from the published value ( $0.865 \pm 0.009$ ) because of a correction in the shift due to  $\eta_{+-}$ . SKJEGGESTAD 72 and HILL 68 give detailed discussions of systematics encountered in this type of experiment.

## $K_S^0$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
------	--------------------------------	-----------------------------------

### Hadronic modes

$\Gamma_1$	$\pi^0 \pi^0$	$(31.40 \pm 0.27) \%$	S=1.2
$\Gamma_2$	$\pi^+ \pi^-$	$(68.60 \pm 0.27) \%$	S=1.2
$\Gamma_3$	$\pi^+ \pi^- \pi^0$	$(3.2 \pm 1.2) \times 10^{-7}$	

### Modes with photons or $\ell\bar{\ell}$ pairs

$\Gamma_4$	$\pi^+ \pi^- \gamma$	$[a,b] \quad (1.78 \pm 0.05) \times 10^{-3}$
$\Gamma_5$	$\pi^+ \pi^- e^+ e^-$	$(4.5 \pm 0.8) \times 10^{-5}$
$\Gamma_6$	$\gamma\gamma$	$(2.5 \pm 0.4) \times 10^{-6}$

### Semileptonic modes

$\Gamma_7$	$\pi^\pm e^\mp \nu_e$	$[c] \quad (7.2 \pm 1.4) \times 10^{-4}$
$\Gamma_8$	$\pi^\pm \mu^\mp \nu_\mu$	$[c]$

### CP violating (CP) and $\Delta S = 1$ weak neutral current (S1) modes

$\Gamma_9$	$3\pi^0$	$CP$	$< 1.4 \times 10^{-5}$	CL=90%
$\Gamma_{10}$	$\mu^+ \mu^-$	$S1$	$< 3.2 \times 10^{-7}$	CL=90%
$\Gamma_{11}$	$e^+ e^-$	$S1$	$< 1.4 \times 10^{-7}$	CL=90%
$\Gamma_{12}$	$\pi^0 e^+ e^-$	$S1$	$< 1.1 \times 10^{-6}$	CL=90%

[a] Most of this radiative mode, the low-momentum  $\gamma$  part, is also included in the parent mode listed without  $\gamma$ 's.

[b] See the Particle Listings below for the energy limits used in this measurement.

[c] The value is for the sum of the charge states or particle/antiparticle states indicated.

## CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 13 measurements and one constraint to determine 2 parameters. The overall fit has a  $\chi^2 = 15.5$  for 12 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$$\begin{array}{cc} x_2 & \boxed{-100} \\ & x_1 \end{array}$$


---

## $K_S^0$ DECAY RATES

### $\Gamma(\pi^\pm e^\mp \nu_e)$

<u>VALUE</u> ( $10^6$ s $^{-1}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.1 ±1.6</b>	75	6 AKHMETSHIN 99	CMD2	Tagged $K_S^0$ using $\phi \rightarrow K_L^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.50 ± 0.08	<sup>7</sup> PDG	98		
seen	BURGUN	72	HBC	$K^+ p \rightarrow K^0 p \pi^+$
9.3 ± 2.5	AUBERT	65	HLBC	$\Delta S = \Delta Q$ , $CP$ cons. not assumed

<sup>6</sup> AKHMETSHIN 99 is from a measured branching ratio  $B(K_S^0 \rightarrow \pi e \nu_e) = (7.2 \pm 1.4) \times 10^{-4}$  and  $\tau_{K_S^0} = (0.8934 \pm 0.0008) \times 10^{-10}$  s.

<sup>7</sup> PDG 98 from  $K_L^0$  measurements, assuming that  $\Delta S = \Delta Q$  in  $K^0$  decay so that  $\Gamma(K_S^0 \rightarrow \pi^\pm e^\mp \nu_e) = \Gamma(K_L^0 \rightarrow \pi^\pm e^\mp \nu_e)$ .

### $\Gamma(\pi^\pm \mu^\mp \nu_\mu)$

### $\Gamma_8$

<u>VALUE</u> ( $10^6$ s $^{-1}$ )	<u>DOCUMENT ID</u>
-----------------------------------	--------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.25 ± 0.07	<sup>8</sup> PDG	98
-------------	------------------	----

<sup>8</sup> PDG 98 from  $K_L^0$  measurements, assuming that  $\Delta S = \Delta Q$  in  $K^0$  decay so that  $\Gamma(K_S^0 \rightarrow \pi^\pm \mu^\mp \nu_\mu) = \Gamma(K_L^0 \rightarrow \pi^\pm \mu^\mp \nu_\mu)$ .

## $K_S^0$ BRANCHING RATIOS

### Hadronic modes

#### $\Gamma(\pi^0 \pi^0)/\Gamma_{\text{total}}$


#### $\Gamma_1/\Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
--------------	-------------	--------------------	-------------

**0.3140 ± 0.0027 OUR FIT** Error includes scale factor of 1.2.

**0.318 ± 0.015 OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.

0.335 ± 0.014	1066	BROWN	63	HLBC
0.288 ± 0.021	198	CHRETIEN	63	HLBC
0.30 ± 0.035		BROWN	61	HLBC



$$\Gamma(\pi^0 \pi^0)/\Gamma_{\text{total}}$$

$$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma$
<b>0.6860±0.0027 OUR FIT</b>				Error includes scale factor of 1.2.	
<b>0.670 ±0.010</b>	3447	DOYLE	69	HBC $\pi^- p \rightarrow \Lambda K^0$	

$$\Gamma(\pi^+ \pi^-)/\Gamma(\pi^0 \pi^0)$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma_1$
<b>2.185±0.027 OUR FIT</b>				Error includes scale factor of 1.2.	
<b>2.197±0.026 OUR AVERAGE</b>					

2.11 ±0.09	1315	EVERHART	76	WIRE $\pi^- p \rightarrow \Lambda K^0$	
2.169±0.094	16k	COWELL	74	OSPK $\pi^- p \rightarrow \Lambda K^0$	
2.16 ±0.08	4799	HILL	73	DBC $K^+ d \rightarrow K^0 pp$	
2.22 ±0.10	3068	<sup>9</sup> ALITTI	72	HBC $K^+ p \rightarrow \pi^+ p K^0$	
2.22 ±0.08	6380	MORSE	72B	DBC $K^+ n \rightarrow K^0 p$	
2.10 ±0.11	701	<sup>10</sup> NAGY	72	HLBC $K^+ n \rightarrow K^0 p$	
2.22 ±0.095	6150	<sup>11</sup> BALTAY	71	HBC $K p \rightarrow K^0$ neutrals	
2.282±0.043	7944	<sup>12</sup> MOFFETT	70	OSPK $K^+ n \rightarrow K^0 p$	
2.10 ±0.06	3700	MORFIN	69	HLBC $K^+ n \rightarrow K^0 p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.12 ±0.17	267	<sup>10</sup> BOZOKI	69	HLBC	
2.285±0.055	3016	<sup>12</sup> GOBBI	69	OSPK $K^+ n \rightarrow K^0 p$	

<sup>9</sup> The directly measured quantity is  $K_S^0 \rightarrow \pi^+ \pi^-$ /all  $K^0 = 0.345 \pm 0.005$ .

<sup>10</sup> NAGY 72 is a final result which includes BOZOKI 69.

<sup>11</sup> The directly measured quantity is  $K_S^0 \rightarrow \pi^+ \pi^-$ /all  $\bar{K}^0 = 0.345 \pm 0.005$ .

<sup>12</sup> MOFFETT 70 is a final result which includes GOBBI 69.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$				$\Gamma_3/\Gamma$
<u>VALUE (units <math>10^{-7}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3.2^{+1.2}_{-1.0}</math> OUR AVERAGE</b>				
$2.5^{+1.3+0.5}_{-1.0-0.6}$	500k	<sup>13</sup> ADLER	97B CPLR	
$4.8^{+2.2}_{-1.6}\pm 1.1$		<sup>14</sup> ZOU	96 E621	
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$4.1^{+2.5+0.5}_{-1.9-0.6}$		<sup>15</sup> ADLER	96E CPLR	Sup. by ADLER 97B
$3.9^{+5.4+0.9}_{-1.8-0.7}$		<sup>16</sup> THOMSON	94 E621	Sup. by ZOU 96
<sup>13</sup> ADLER 97B find the $CP$ -conserving parameters $\text{Re}(\lambda) = (28 \pm 7 \pm 3) \times 10^{-3}$ , $\text{Im}(\lambda) = (-10 \pm 8 \pm 2) \times 10^{-3}$ . They estimate $B(K_S^0 \rightarrow \pi^+\pi^-\pi^0)$ from $\text{Re}(\lambda)$ and the $K_L^0$ decay parameters. See also ANGELOPOULOS 98C.				
<sup>14</sup> ZOU 96 is from the measured quantities $ \rho_{+-0}  = 0.039^{+0.009}_{-0.006} \pm 0.005$ and $\phi_\rho = (-9 \pm 18)^\circ$ .				
<sup>15</sup> ADLER 96E is from the measured quantities $\text{Re}(\lambda) = 0.036 \pm 0.010^{+0.002}_{-0.003}$ and $\text{Im}(\lambda)$ consistent with zero. Note that the quantity $\lambda$ is the same as $\rho_{+-0}$ used in other footnotes.				
<sup>16</sup> THOMSON 94 calculates this branching ratio from their measurements $ \rho_{+-0}  = 0.035^{+0.019}_{-0.011} \pm 0.004$ and $\phi_\rho = (-59 \pm 48)^\circ$ where $ \rho_{+-0}  e^{i\phi_\rho} = A(K_S^0 \rightarrow \pi^+\pi^-\pi^0, I=2)/A(K_L^0 \rightarrow \pi^+\pi^-\pi^0)$ .				

**Modes with photons or  $\ell\bar{\ell}$  pairs**

$\Gamma(\pi^+\pi^-\gamma)/\Gamma(\pi^+\pi^-)$				$\Gamma_4/\Gamma_2$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.60 \pm 0.08</math> OUR AVERAGE</b>				
$2.56 \pm 0.09$	1286	RAMBERG	93 E731	$p_\gamma > 50 \text{ MeV}/c$
$2.68 \pm 0.15$		<sup>17</sup> TAUREG	76 SPEC	$p_\gamma > 50 \text{ MeV}/c$
$2.8 \pm 0.6$		<sup>18</sup> BURGUN	73 HBC	$p_\gamma > 50 \text{ MeV}/c$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$7.10 \pm 0.22$	3723	RAMBERG	93 E731	$p_\gamma > 20 \text{ MeV}/c$
$3.0 \pm 0.6$	29	<sup>19</sup> BOBISUT	74 HLBC	$p_\gamma > 40 \text{ MeV}/c$
<sup>17</sup> TAUREG 76 find direct emission contribution $< 0.06$ , CL = 90%.				
<sup>18</sup> BURGUN 73 estimates that direct emission contribution is $0.3 \pm 0.6$ .				
<sup>19</sup> BOBISUT 74 not included in average because $p_\gamma$ cut differs. Estimates direct emission contribution to be 0.5 or less, CL = 95%.				

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_5/\Gamma$
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	
<b><math>4.5 \pm 0.7 \pm 0.4</math></b>	56	LAI	00B NA48	

### $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

$\Gamma_6/\Gamma$

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN
<b>2.5 ±0.4 OUR AVERAGE</b>				
2.58±0.36±0.22	149	LAI	00	NA48
2.4 ±0.9	35	20 BARR	95B	NA31

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.2 ±1.1	16	21 BARR	95B	NA31
< 13	90	BALATS	89	SPEC
2.4 ±1.2	19	BURKHARDT	87	NA31
<133	90	BARMIN	86B	XEBC

20 BARR 95B quotes this as the combined BARR 95B + BURKHARDT 87 result after rescaling BURKHARDT 87 to use same branching ratios and lifetimes as BARR 95B.

21 BARR 95B result is calculated using  $B(K_L \rightarrow \gamma\gamma) = (5.86 \pm 0.17) \times 10^{-4}$ .

### Semileptonic modes

### $\Gamma(\pi^\pm e^\mp \nu_e)/\Gamma_{\text{total}}$

$\Gamma_7/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.2±1.4</b>	75	AKHMETSHIN 99	CMD2	Tagged $K_S^0$ using $\phi \rightarrow K_L^0 K_S^0$

### CP violating (CP) and $\Delta S = 1$ weak neutral current (S1) modes

### $\Gamma(3\pi^0)/\Gamma_{\text{total}}$

$\Gamma_9/\Gamma$

Violates CP conservation.

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN
<b>&lt;1.4</b>	90	7M	ACHASOV	99D SND

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.9	90	17300	22 ANGELOPO...	98B CPLR
<3.7	90		BARMIN	83 HLBC

22 ANGELOPOULOS 98B is from  $\text{Im}(\eta_{000}) = -0.05 \pm 0.12 \pm 0.05$ , assuming  $\text{Re}(\eta_{000}) = \text{Re}(\epsilon) = 1.635 \times 10^{-3}$  and using the value  $B(K_L^0 \rightarrow \pi^0 \pi^0 \pi^0) = 0.2112 \pm 0.0027$ .

### $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

$\Gamma_{10}/\Gamma$

Test for  $\Delta S = 1$  weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN
<b>&lt;0.032</b>	90		GJESDAL	73 ASPK

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.7	90		HYAMS	69B OSPK
------	----	--	-------	----------

### $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

$\Gamma_{11}/\Gamma$

Test for  $\Delta S = 1$  weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units $10^{-7}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt; 1.4</b>	90		ANGELOPO...	97 CPLR	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 28	90	0	BLICK	94 CNTR	Hyperon facility
<100	90		BARMIN	86 XEBC	

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

Test for  $\Delta S = 1$  weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
< 1.1	90	0	BARR	93B NA31
<45	90		GIBBONS	88 E731

• • • We do not use the following data for averages, fits, limits, etc. • • •

**CP-VIOLATION PARAMETERS IN  $K_S^0$  DECAY**

$$\text{Im}(\eta_{+-0})^2 = \Gamma(K_S^0 \rightarrow \pi^+ \pi^- \pi^0, \text{CP-violating}) / \Gamma(K_L^0 \rightarrow \pi^+ \pi^- \pi^0)$$

*CPT assumed valid (i.e.  $\text{Re}(\eta_{+-0}) \simeq 0$ ).*

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
--------------	------------	-------------	--------------------	-------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.23	90	601	23 BARMIN	85 HLBC
<0.12	90	384	METCALF	72 ASPK

<sup>23</sup> BARMIN 85 find  $\text{Re}(\eta_{+-0}) = (0.05 \pm 0.17)$  and  $\text{Im}(\eta_{+-0}) = (0.15 \pm 0.33)$ . Includes events of BALDO-CEOLIN 75.

$$\text{Im}(\eta_{+-0}) = \text{Im}(A(K_S^0 \rightarrow \pi^+ \pi^- \pi^0, \text{CP-violating}) / A(K_L^0 \rightarrow \pi^+ \pi^- \pi^0))$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.002 \pm 0.009^{+0.002}_{-0.001}$	500k	24 ADLER	97B CPLR	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.002 \pm 0.018 \pm 0.003$	137k	25 ADLER	96D CPLR	Sup. by ADLER 97B
$-0.015 \pm 0.017 \pm 0.025$	272k	26 ZOU	94 SPEC	

<sup>24</sup> ADLER 97B also find  $\text{Re}(\eta_{+-0}) = -0.002 \pm 0.007^{+0.004}_{-0.001}$ . See also ANGELOPOU-LOS 98C.

<sup>25</sup> The ADLER 96D fit also yields  $\text{Re}(\eta_{+-0}) = 0.006 \pm 0.013 \pm 0.001$  with a correlation  $+0.66$  between real and imaginary parts. Their results correspond to  $|\eta_{+-0}| < 0.037$  with 90% CL.

<sup>26</sup> ZOU 94 use theoretical constraint  $\text{Re}(\eta_{+-0}) = \text{Re}(\epsilon) = 0.0016$ . Without this constraint they find  $\text{Im}(\eta_{+-0}) = 0.019 \pm 0.061$  and  $\text{Re}(\eta_{+-0}) = 0.019 \pm 0.027$ .

$$\text{Im}(\eta_{000})^2 = \Gamma(K_S^0 \rightarrow 3\pi^0) / \Gamma(K_L^0 \rightarrow 3\pi^0)$$

*CPT assumed valid (i.e.  $\text{Re}(\eta_{000}) \simeq 0$ ). This limit determines branching ratio  $\Gamma(3\pi^0)/\Gamma_{\text{total}}$  above.*

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.1	90	632	27 BARMIN	83 HLBC
<0.28	90		28 GJESDAL	74B SPEC Indirect meas.

<sup>27</sup> BARMIN 83 find  $\text{Re}(\eta_{000}) = (-0.08 \pm 0.18)$  and  $\text{Im}(\eta_{000}) = (-0.05 \pm 0.27)$ . Assuming CPT invariance they obtain the limit quoted above.

<sup>28</sup> GJESDAL 74B uses  $K_2\pi$ ,  $K_{\mu 3}$ , and  $K_{e3}$  decay results, unitarity, and CPT. Calculates  $|\eta_{000}| = 0.26 \pm 0.20$ . We convert to upper limit.

$$\text{Im}(\eta_{000}) = \text{Im}(A(K_S^0 \rightarrow \pi^0 \pi^0 \pi^0)/A(K_L^0 \rightarrow \pi^0 \pi^0 \pi^0))$$

$K_S^0 \rightarrow \pi^0 \pi^0 \pi^0$  violates  $CP$  conservation, in contrast to  $K_S^0 \rightarrow \pi^+ \pi^- \pi^0$  which has a  $CP$ -conserving part.

VALUE	EVTS	DOCUMENT ID	TECN
<b><math>-0.05 \pm 0.12 \pm 0.05</math></b>	17300	29	ANGELOPO... 98B CPLR

29 ANGELOPOULOS 98B assumes  $\text{Re}(\eta_{000}) = \text{Re}(\epsilon) = 1.635 \times 10^{-3}$ . Without assuming  $CPT$  invariance, they obtain  $\text{Re}(\eta_{000}) = 0.18 \pm 0.14 \pm 0.06$  and  $\text{Im}(\eta_{000}) = 0.15 \pm 0.20 \pm 0.03$ .

## $K_S^0$ REFERENCES

LAI	00	PL B493 29	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
LAI	00B	PL B496 137	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
ACHASOV	99D	PL B459 674	M.N. Achasov <i>et al.</i>	
AKHMETSHIN	99	PL B456 90	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
ANGELOPO...	98B	PL B425 391	A. Angelopoulos <i>et al.</i>	(CLEAR Collab.)
ANGELOPO...	98C	EPJ C5 389	A. Angelopoulos <i>et al.</i>	(CLEAR Collab.)
PDG	98	EPJ C3 1	C. Caso <i>et al.</i>	
ADLER	97B	PL B407 193	R. Adler <i>et al.</i>	(CLEAR Collab.)
ANGELOPO...	97	PL B413 232	A. Angelopoulos <i>et al.</i>	(CLEAR Collab.)
BERTANZA	97	ZPHY C73 629	L. Bertanza (PISA, CERN, EDIN, MANZ, ORSAY+)	
ADLER	96D	PL B370 167	R. Adler <i>et al.</i>	(CLEAR Collab.)
ADLER	96E	PL B374 313	R. Adler <i>et al.</i>	(CLEAR Collab.)
ZOU	96	PL B369 362	Y. Zou <i>et al.</i>	(RUTG, MINN, MICH)
BARR	95B	PL B351 579	G.D. Barr <i>et al.</i>	(CERN, EDIN, MANZ, LALO+)
SCHWINGEN...	95	PRL 74 4376	B. Schwingerheuer <i>et al.</i>	(IFI, CHIC+)
BLICK	94	PL B334 234	A.M. Blick <i>et al.</i>	(SERP, JINR)
THOMSON	94	PL B337 411	G.B. Thomson <i>et al.</i>	(RUTG, MINN, MICH)
ZOU	94	PL B329 519	Y. Zou <i>et al.</i>	(RUTG, MINN, MICH)
BARR	93B	PL B304 381	G.D. Barr <i>et al.</i>	(CERN, EDIN, MANZ, LALO+)
GIBBONS	93	PRL 70 1199	L.K. Gibbons <i>et al.</i>	(FNAL E731 Collab.)
Also	97	PR D55 6625	L.K. Gibbons <i>et al.</i>	(FNAL E731 Collab.)
RAMBERG	93	PRL 70 2525	E. Ramberg <i>et al.</i>	(FNAL E731 Collab.)
BALATS	89	SJNP 49 828	M.Y. Balats <i>et al.</i>	(ITEP)
		Translated from YAF 49	1332.	
GIBBONS	88	PRL 61 2661	L.K. Gibbons <i>et al.</i>	(FNAL E731 Collab.)
BURKHARDT	87	PL B199 139	H. Burkhardt <i>et al.</i>	(CERN, EDIN, MANZ+)
GROSSMAN	87	PRL 59 18	N. Grossman <i>et al.</i>	(MINN, MICH, RUTG)
BARMIN	86	SJNP 44 622	V.V. Barmin <i>et al.</i>	(ITEP)
		Translated from YAF 44	965.	
BARMIN	86B	NC 96A 159	V.V. Barmin <i>et al.</i>	(ITEP, PADO)
PDG	86B	PL 170B 130	M. Aguilar-Benitez <i>et al.</i>	(CERN, CIT+)
BARMIN	85	NC 85A 67	V.V. Barmin <i>et al.</i>	(ITEP, PADO)
Also	85B	SJNP 41 759	V.V. Barmin <i>et al.</i>	(ITEP)
		Translated from YAF 41	1187.	
BARMIN	83	PL 128B 129	V.V. Barmin <i>et al.</i>	(ITEP, PADO)
Also	84	SJNP 39 269	V.V. Barmin <i>et al.</i>	(ITEP, PADO)
		Translated from YAF 39	428.	
ARONSON	82	PRL 48 1078	S.H. Aronson <i>et al.</i>	(BNL, CHIC, STAN+)
ARONSON	82B	PRL 48 1306	S.H. Aronson <i>et al.</i>	(BNL, CHIC, PURD)
Also	82B	PL 116B 73	E. Fischbach <i>et al.</i>	(PURD, BNL, CHIC)
Also	83	PR D28 476	S.H. Aronson <i>et al.</i>	(BNL, CHIC, PURD)
Also	83B	PR D28 495	S.H. Aronson <i>et al.</i>	(BNL, CHIC, PURD)
ARONSON	76	NC 32A 236	S.H. Aronson <i>et al.</i>	(WISC, EFI, UCSD+)
EVERHART	76	PR D14 661	G.C. Everhart <i>et al.</i>	(PENN)
TAUREG	76	PL 65B 92	H. Taureg <i>et al.</i>	(HEIDH, CERN, DORT)
BALDO-...	75	NC 25A 688	M. Baldo-Ceolin <i>et al.</i>	(PADO, WISC)
CARITHERS	75	PRL 34 1244	W.C.J. Carithers <i>et al.</i>	(COLU, NYU)
BOBISUT	74	LNC 11 646	F. Bobisut <i>et al.</i>	(PADO)
COWELL	74	PR D10 2083	P.L. Cowell <i>et al.</i>	(STON, COLU)
GEWENIGER	74B	PL 48B 487	C. Geweniger <i>et al.</i>	(CERN, HEIDH)

GJESDAL	74B	PL 52B 119	S. Gjesdal <i>et al.</i>	(CERN, HEIDH)
BURGUN	73	PL 46B 481	G. Burgun <i>et al.</i>	(SACL, CERN)
GJESDAL	73	PL 44B 217	S. Gjesdal <i>et al.</i>	(CERN, HEIDH)
HILL	73	PR D8 1290	D.G. Hill <i>et al.</i>	(BNL, CMU)
ALITTI	72	PL 39B 568	J. Alitti, E. Lesquoy, A. Muller	(SACL)
BURGUN	72	NP B50 194	G. Burgun <i>et al.</i>	(SACL, CERN, OSLO)
METCALF	72	PL 40B 703	M. Metcalf <i>et al.</i>	(CERN, IPN, WIEN)
MORSE	72B	PRL 28 388	R. Morse <i>et al.</i>	(COLO, PRIN, UMD)
NAGY	72	NP B47 94	E. Nagy, F. Telbisz, G. Vesztregombi	(BUDA)
Also	69	PL 30B 498	G. Bozoki <i>et al.</i>	(BUDA)
SKJEGGESTAD	72	NP B48 343	O. Skjeggestad <i>et al.</i>	(OSLO, CERN, SACL)
BALTAY	71	PRL 27 1678	C. Baltay <i>et al.</i>	(COLU)
Also	71	Thesis Nevis 187	W.A. Cooper	(COLU)
MOFFETT	70	BAPS 15 512	R. Moffett <i>et al.</i>	(ROCH)
BOZOKI	69	PL 30B 498	G. Bozoki <i>et al.</i>	(BUDA)
DOYLE	69	Thesis UCRL 18139	J.C. Doyle	(LRL)
GOBBI	69	PRL 22 682	B. Gobbi <i>et al.</i>	(ROCH)
HYAMS	69B	PL 29B 521	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
MORFIN	69	PRL 23 660	J.G. Morfin, D. Sinclair	(MICH)
DONALD	68B	PL 27B 58	R.A. Donald <i>et al.</i>	(LIVP, CERN, IPNP+)
HILL	68	PR 171 1418	D.G. Hill <i>et al.</i>	(BNL, CMU)
AUBERT	65	PL 17 59	B. Aubert <i>et al.</i>	(EPOL, ORSAY)
BROWN	63	PR 130 769	J.L. Brown <i>et al.</i>	(LRL, MICH)
CHRETIEN	63	PR 131 2208	M. Chretien <i>et al.</i>	(BRAN, BROW, HARV+)
BROWN	61	NC 19 1155	J.L. Brown <i>et al.</i>	(MICH)
BOLDT	58B	PRL 1 150	E. Boldt, D.O. Caldwell, Y. Pal	(MIT)

## OTHER RELATED PAPERS

LITTENBERG	93	ARNPS 43 729	L.S. Littenberg, G. Valencia	(BNL, FNAL)
Rare and Radiative Kaon Decays				
BATTISTON	92	PRPL 214 293	R. Battiston <i>et al.</i>	(PGIA, CERN, TRSTT)
Status and Perspectives of <i>K</i> Decay Physics				
TRILLING	65B	UCRL 16473	G.N. Trilling	(LRL)
Updated from 1965 Argonne Conference, page 115.				
CRAWFORD	62	CERN Conf. 827	F.S. Crawford	(LRL)
FITCH	61	NC 22 1160	V.L. Fitch, P.A. Piroue, R.B. Perkins	(PRIN+)
GOOD	61	PR 124 1223	R.H. Good <i>et al.</i>	(LRL)
BIRGE	60	Rochester Conf. 601	R.W. Birge <i>et al.</i>	(LRL, WISC)
MULLER	60	PRL 4 418	F. Muller <i>et al.</i>	(LRL, BNL)