



$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$ Status: ****

The parity has not actually been measured, but + is of course expected.

We have omitted some results that have been superseded by later experiments. See our earlier editions.

Ξ^- MASS

The fit uses the Ξ^- , Ξ^+ , and Ξ^0 mass and mass difference measurements. It assumes the Ξ^- and Ξ^+ masses are the same.

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1321.31±0.13 OUR FIT				
1321.34±0.14 OUR AVERAGE				
1321.46±0.34	632	DIBIANCA	75 DBC	4.9 GeV/c $K^- d$
1321.12±0.41	268	WILQUET	72 HLBC	
1321.87±0.51	195	¹ GOLDWASSER	70 HBC	5.5 GeV/c $K^- p$
1321.67±0.52	6	CHIEN	66 HBC	6.9 GeV/c $\bar{p} p$
1321.4 ±1.1	299	LONDON	66 HBC	
1321.3 ±0.4	149	PJERROU	65B HBC	
1321.1 ±0.3	241	² BADIER	64 HBC	
1321.4 ±0.4	517	² JAUNEAU	63D FBC	
1321.1 ±0.65	62	² SCHNEIDER	63 HBC	

¹ GOLDWASSER 70 uses $m_\Lambda = 1115.58$ MeV.
² These masses have been increased 0.09 MeV because the Λ mass increased.

Ξ^+ MASS

The fit uses the Ξ^- , Ξ^+ , and Ξ^0 mass and mass difference measurements. It assumes the Ξ^- and Ξ^+ masses are the same.

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1321.31±0.13 OUR FIT				
1321.20±0.33 OUR AVERAGE				
1321.6 ±0.8	35	VOTRUBA	72 HBC	10 GeV/c $K^+ p$
1321.2 ±0.4	34	STONE	70 HBC	
1320.69±0.93	5	CHIEN	66 HBC	6.9 GeV/c $\bar{p} p$

$(m_{\Xi^-} - m_{\Xi^+}) / m_{\Xi^-}$

A test of *CPT* invariance. We calculate this from the *average* Ξ^- and Ξ^+ masses above.

<u>VALUE</u>	<u>DOCUMENT ID</u>
$(1.1\pm2.7) \times 10^{-4}$ OUR EVALUATION	

Ξ^- MEAN LIFE

Measurements with an error $> 0.2 \times 10^{-10}$ s or with systematic errors not included have been omitted.

<u>VALUE (10^{-10} s)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.639 ± 0.015 OUR AVERAGE				
1.652 ± 0.051	32k	BOURQUIN	84 SPEC	Hyperon beam
1.665 ± 0.065	41k	BOURQUIN	79 SPEC	Hyperon beam
1.609 ± 0.028	4286	HEMINGWAY	78 HBC	4.2 GeV/c $K^- p$
1.67 ± 0.08		DIBIANCA	75 DBC	4.9 GeV/c $K^- d$
1.63 ± 0.03	4303	BALTAY	74 HBC	1.75 GeV/c $K^- p$
$1.73^{+0.08}_{-0.07}$	680	MAYEUR	72 HLBC	2.1 GeV/c K^-
1.61 ± 0.04	2610	DAUBER	69 HBC	
1.80 ± 0.16	299	LONDON	66 HBC	
1.70 ± 0.12	246	PJERROU	65B HBC	
1.69 ± 0.07	794	HUBBARD	64 HBC	
$1.86^{+0.15}_{-0.14}$	517	JAUNEAU	63D FBC	

Ξ^+ MEAN LIFE

<u>VALUE (10^{-10} s)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.6 ± 0.3	34	STONE	70 HBC	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.55^{+0.35}_{-0.20}$	35	³ VOTRUBA	72 HBC	10 GeV/c $K^+ p$
$1.9^{+0.7}_{-0.5}$	12	³ SHEN	67 HBC	
1.51 ± 0.55	5	³ CHIEN	66 HBC	6.9 GeV/c $\bar{p} p$
³ The error is statistical only.				

$$(\tau_{\Xi^-} - \tau_{\Xi^+}) / \tau_{\Xi^-}$$

A test of *CPT* invariance. Calculated from the Ξ^- and Ξ^+ mean lives, above.

<u>VALUE</u>	<u>DOCUMENT ID</u>
0.02 ± 0.18 OUR EVALUATION	

Ξ^- MAGNETIC MOMENT

See the “Note on Baryon Magnetic Moments” in the Λ Listings.

<u>VALUE (μ_N)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.6507 ± 0.0025 OUR AVERAGE				
-0.6505 ± 0.0025	4.36M	DURYEA	92 SPEC	800 GeV p Be
$-0.661 \pm 0.036 \pm 0.036$	44k	TROST	89 SPEC	$\Xi^- \sim 250$ GeV
-0.69 ± 0.04	218k	RAMEIKA	84 SPEC	400 GeV p Be

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

−0.674 ±0.021 ±0.020	122k	HO	90	SPEC	See DURYEA 92
−2.1 ±0.8	2436	COOL	74	OSPK	1.8 GeV/c $K^- p$
−0.1 ±2.1	2724	BINGHAM	70B	OSPK	1.8 GeV/c $K^- p$

Ξ^+ MAGNETIC MOMENT

See the “Note on Baryon Magnetic Moments” in the Λ Listings.

VALUE (μ_N)	EVTs	DOCUMENT ID	TECN	COMMENT
+0.657±0.028±0.020	70k	HO	90 SPEC	800 GeV p Be

$(\mu_{\Xi^-} + \mu_{\Xi^+}) / |\mu_{\Xi^-}|$

A test of CPT invariance. We calculate this from the Ξ^- and Ξ^+ magnetic moments above.

VALUE	DOCUMENT ID
+0.01±0.05 OUR EVALUATION	

Ξ^- DECAY MODES

Mode		Fraction (Γ_i/Γ)	Confidence level
Γ_1	$\Lambda\pi^-$	$(99.887\pm0.035)\%$	
Γ_2	$\Sigma^-\gamma$	$(1.27\pm0.23)\times10^{-4}$	
Γ_3	$\Lambda e^-\bar{\nu}_e$	$(5.63\pm0.31)\times10^{-4}$	
Γ_4	$\Lambda\mu^-\bar{\nu}_\mu$	$(3.5^{+3.5}_{-2.2})\times10^{-4}$	
Γ_5	$\Sigma^0 e^-\bar{\nu}_e$	$(8.7\pm1.7)\times10^{-5}$	
Γ_6	$\Sigma^0\mu^-\bar{\nu}_\mu$	$<8\times10^{-4}$	90%
Γ_7	$\Xi^0 e^-\bar{\nu}_e$	$<2.3\times10^{-3}$	90%

$\Delta S = 2$ forbidden ($S2$) modes

Γ_8	$n\pi^-$	$S2$	$<1.9\times10^{-5}$	90%
Γ_9	$ne^-\bar{\nu}_e$	$S2$	$<3.2\times10^{-3}$	90%
Γ_{10}	$n\mu^-\bar{\nu}_\mu$	$S2$	$<1.5\%$	90%
Γ_{11}	$p\pi^-\pi^-$	$S2$	$<4\times10^{-4}$	90%
Γ_{12}	$p\pi^-e^-\bar{\nu}_e$	$S2$	$<4\times10^{-4}$	90%
Γ_{13}	$p\pi^-\mu^-\bar{\nu}_\mu$	$S2$	$<4\times10^{-4}$	90%
Γ_{14}	$p\mu^-\mu^-$	L	$<4\times10^{-4}$	90%

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 5 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 1.0$ for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	−6			
x_3	−8	0		
x_4	−99	0	−1	
x_5	−5	0	0	0
	x_1	x_2	x_3	x_4

Ξ^- BRANCHING RATIOS

A number of early results have been omitted.

$\Gamma(\Sigma^- \gamma) / \Gamma(\Lambda \pi^-)$					Γ_2 / Γ_1
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.27 ± 0.24 OUR FIT					
1.27 ± 0.23 OUR AVERAGE					
1.22 ± 0.23 ± 0.06	211	⁴ DUBBS	94 E761	Ξ^- 375 GeV	
2.27 ± 1.02	9	BIAGI	87B SPEC	SPS hyperon beam	
⁴ DUBBS 94 also finds weak evidence that the asymmetry parameter α_γ is positive ($\alpha_\gamma = 1.0 \pm 1.3$).					

$\Gamma(\Lambda e^- \bar{\nu}_e) / \Gamma(\Lambda \pi^-)$					Γ_3 / Γ_1
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
0.564 ± 0.031 OUR FIT					
0.564 ± 0.031					
	2857	BOURQUIN	83 SPEC	SPS hyperon beam	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.30 ± 0.13	11	THOMPSON	80 ASPK	Hyperon beam	

$\Gamma(\Lambda \mu^- \bar{\nu}_\mu) / \Gamma(\Lambda \pi^-)$					Γ_4 / Γ_1
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.35^{+0.35}_{−0.22} OUR FIT					
0.35 ± 0.35					
	1	YEH	74 HBC	Effective denom.=2859	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 2.3	90	0	THOMPSON	80 ASPK	Effective denom.=1017
< 1.3			DAUBER	69 HBC	
< 12			BERGE	66 HBC	

$$\Gamma(\Sigma^0 e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-) \quad \Gamma_5/\Gamma_1$$

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.087 ± 0.017 OUR FIT					
0.087 ± 0.017		154	BOURQUIN	83	SPEC SPS hyperon beam

$$\Gamma(\Sigma^0 \mu^- \bar{\nu}_\mu)/\Gamma(\Lambda\pi^-) \quad \Gamma_6/\Gamma_1$$

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.76	90	0	YEH	74	HBC Effective denom.=3026
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<5			BERGE	66	HBC

$$[\Gamma(\Lambda e^- \bar{\nu}_e) + \Gamma(\Sigma^0 e^- \bar{\nu}_e)]/\Gamma(\Lambda\pi^-) \quad (\Gamma_3 + \Gamma_5)/\Gamma_1$$

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.651 ± 0.031		3011	⁵ BOURQUIN	83	SPEC SPS hyperon beam
0.68 ± 0.22		17	⁶ DUCLOS	71	OSPK

⁵ See the separate BOURQUIN 83 values for $\Gamma(\Lambda e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$ and $\Gamma(\Sigma^0 e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$ above.

⁶ DUCLOS 71 cannot distinguish Σ^0 's from Λ 's. The Cabibbo theory predicts the Σ^0 rate is about a factor 6 smaller than the Λ rate.

$$\Gamma(\Xi^0 e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-) \quad \Gamma_7/\Gamma_1$$

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<2.3	90	0	YEH	74	HBC Effective denom.=1000

$$\Gamma(n\pi^-)/\Gamma(\Lambda\pi^-) \quad \Gamma_8/\Gamma_1$$

$\Delta S=2$. Forbidden in first-order weak interaction.

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.019	90		BIAGI	82B	SPEC SPS hyperon beam
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.0	90	0	YEH	74	HBC Effective denom.=760
<1.1			DAUBER	69	HBC
<5.0			FERRO-LUZZI	63	HBC

$$\Gamma(ne^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-) \quad \Gamma_9/\Gamma_1$$

$\Delta S=2$. Forbidden in first-order weak interaction.

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.2	90	0	YEH	74	HBC Effective denom.=715
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<10	90		BINGHAM	65	RVUE

$$\Gamma(n\mu^- \bar{\nu}_\mu)/\Gamma(\Lambda\pi^-) \quad \Gamma_{10}/\Gamma_1$$

$\Delta S=2$. Forbidden in first-order weak interaction.

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<15.3	90	0	YEH	74	HBC Effective denom.=150

$\Gamma(p\pi^-\pi^-)/\Gamma(\Lambda\pi^-)$
 $\Delta S=2$. Forbidden in first-order weak interaction.

Γ_{11}/Γ_1

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.7	90	0	YEH	74	HBC Effective denom.=6200

$\Gamma(p\pi^-e^-\bar{\nu}_e)/\Gamma(\Lambda\pi^-)$
 $\Delta S=2$. Forbidden in first-order weak interaction.

Γ_{12}/Γ_1

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.7	90	0	YEH	74	HBC Effective denom.=6200

$\Gamma(p\pi^-\mu^-\bar{\nu}_\mu)/\Gamma(\Lambda\pi^-)$
 $\Delta S=2$. Forbidden in first-order weak interaction.

Γ_{13}/Γ_1

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.7	90	0	YEH	74	HBC Effective denom.=6200

$\Gamma(p\mu^-\mu^-)/\Gamma(\Lambda\pi^-)$
A $\Delta L=2$ decay, forbidden by total lepton number conservation.

Γ_{14}/Γ_1

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<3.7	90	⁷ LITTENBERG	92B	HBC Uses YEH 74 data

⁷ This LITTENBERG 92B limit and the identical YEH 74 limits for the preceding three modes all result from nonobservance of any 3-prong decays of the Ξ^- . One could as well apply the limit to the *sum* of the four modes.


Ξ^- DECAY PARAMETERS

See the "Note on Baryon Decay Parameters" in the neutron Listings.

$\alpha(\Xi^-)\alpha_-(\Lambda)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.294 ±0.005 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.		
-0.2963±0.0042	189k	LUK	00	E756 p Be, 800 GeV
-0.2894±0.0073	63k	⁸ LUK	00	E756 p Be, 800 GeV
-0.303 ±0.004 ±0.004	192k	RAMEIKA	86	SPEC 400 GeV p Be
-0.257 ±0.020	11k	ASTON	85B	LASS 11 GeV/ c $K^- p$
-0.260 ±0.017	21k	BENSINGER	85	MPS 5 GeV/ c $K^- p$
-0.299 ±0.007	150k	BIAGI	82	SPEC SPS hyperon beam
-0.315 ±0.026	9046	CLELAND	80C	ASPK BNL hyperon beam
-0.239 ±0.021	6599	HEMINGWAY	78	HBC 4.2 GeV/ c $K^- p$
-0.243 ±0.025	4303	BALTAY	74	HBC 1.75 GeV/ c $K^- p$
-0.252 ±0.032	2436	COOL	74	OSPK 1.8 GeV/ c $K^- p$
-0.253 ±0.028	2781	DAUBER	69	HBC

⁸ This LUK 00 value is for $\alpha(\Xi^+)\alpha_+(\bar{\Lambda})$. We assume CP conservation here by including it in the average for $\alpha(\Xi^-)\alpha_-(\Lambda)$. But see the second data block below for the CP test.



$\alpha(\Xi^-)\alpha_-(\Lambda)$

α FOR $\Xi^- \rightarrow \Lambda\pi^-$

The above average, $\alpha(\Xi^-)\alpha_-(\Lambda) = -0.293 \pm 0.007$, where the error includes a scale factor of 1.8, divided by our current average $\alpha_-(\Lambda) = 0.642 \pm 0.013$, gives the following value for $\alpha(\Xi^-)$.

VALUE	DOCUMENT ID
-0.458±0.012 OUR EVALUATION	Error includes scale factor of 1.8.

$[\alpha(\Xi^-)\alpha_-(\Lambda) - \alpha(\Xi^+)\alpha_+(\bar{\Lambda})]/[\alpha(\Xi^-)\alpha_-(\Lambda) + \alpha(\Xi^+)\alpha_+(\bar{\Lambda})]$

This is zero if CP is conserved. The α 's are the decay-asymmetry parameters for $\Xi^- \rightarrow \Lambda\pi^-$ and $\Lambda \rightarrow p\pi^-$ and for $\Xi^+ \rightarrow \bar{\Lambda}\pi^+$ and $\bar{\Lambda} \rightarrow \bar{p}\pi^-$.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
+0.012±0.014	252k	LUK	00 E756	p Be, 800 GeV

ϕ ANGLE FOR $\Xi^- \rightarrow \Lambda\pi^-$ **($\tan\phi = \beta/\gamma$)**

VALUE (°)	EVTs	DOCUMENT ID	TECN	COMMENT
4 ± 4 OUR AVERAGE				
5 ± 10	11k	ASTON	85B LASS	$K^- p$
14.7±16.0	21k	⁹ BENSINGER	85 MPS	5 GeV/c $K^- p$
11 ± 9	4303	BALTAY	74 HBC	1.75 GeV/c $K^- p$
5 ± 16	2436	COOL	74 OSPK	1.8 GeV/c $K^- p$
-26 ± 30	2724	BINGHAM	70B OSPK	
-14 ± 11	2781	DAUBER	69 HBC	Uses $\alpha_\Lambda = 0.647 \pm 0.020$
0 ± 12	1004	¹⁰ BERGE	66 HBC	
0 ± 20.4	364	¹⁰ LONDON	66 HBC	Using $\alpha_\Lambda = 0.62$
54 ± 30	356	¹⁰ CARMONY	64B HBC	

⁹ BENSINGER 85 used $\alpha_\Lambda = 0.642 \pm 0.013$.

¹⁰ The errors have been multiplied by 1.2 due to approximations used for the Ξ polarization; see DAUBER 69 for a discussion.

g_A / g_V FOR $\Xi^- \rightarrow \Lambda e^- \bar{\nu}_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.25 ± 0.05	1992	¹¹ BOURQUIN	83 SPEC	SPS hyperon beam

¹¹BOURQUIN 83 assumes that $g_2 = 0$. Also, the sign has been changed to agree with our conventions, given in the "Note on Baryon Decay Parameters" in the neutron Listings.

Ξ^- REFERENCES

We have omitted some papers that have been superseded by later experiments. See our earlier editions.

LUK	00	PRL 85 4860	K.B. Luk <i>et al.</i>	(FNAL E756 Collab.)
DUBBS	94	PRL 72 808	T. Dubbs <i>et al.</i>	(FNAL E761 Collab.)
DURYEA	92	PRL 68 768	J. Duryea <i>et al.</i>	(MINN, FNAL, MICH, RUTG)
LITTENBERG	92B	PR D46 R892	L.S. Littenberg, R.E. Shrock	(BNL, STON)
HO	90	PRL 65 1713	P.M. Ho <i>et al.</i>	(MICH, FNAL, MINN, RUTG)
Also	91	PR D44 3402	P.M. Ho <i>et al.</i>	(MICH, FNAL, MINN, RUTG)
TROST	89	PR D40 1703	L.H. Trost <i>et al.</i>	(FNAL-715 Collab.)
BIAGI	87B	ZPHY C35 143	S.F. Biagi <i>et al.</i>	(BRIS, CERN, GEVA+)
RAMEIKA	86	PR D33 3172	R. Rameika <i>et al.</i>	(RUTG, MICH, WISC+)
ASTON	85B	PR D32 2270	D. Aston <i>et al.</i>	(SLAC, CARL, CNRC, CINC)
BENSINGER	85	NP B252 561	J.R. Bensinger <i>et al.</i>	(CHIC, ELMT, FNAL+)
BOURQUIN	84	NP B241 1	M.H. Bourquin <i>et al.</i>	(BRIS, GEVA, HEIDP+)
RAMEIKA	84	PRL 52 581	R. Rameika <i>et al.</i>	(RUTG, MICH, WISC+)
BOURQUIN	83	ZPHY C21 1	M.H. Bourquin <i>et al.</i>	(BRIS, GEVA, HEIDP+)
BIAGI	82	PL 112B 265	S.F. Biagi <i>et al.</i>	(BRIS, CAVE, GEVA+)
BIAGI	82B	PL 112B 277	S.F. Biagi <i>et al.</i>	(LOQM, GEVA, RL+)
CLELAND	80C	PR D21 12	W.E. Cleland <i>et al.</i>	(PITT, BNL)
THOMPSON	80	PR D21 25	J.A. Thompson <i>et al.</i>	(PITT, BNL)
BOURQUIN	79	PL 87B 297	M.H. Bourquin <i>et al.</i>	(BRIS, GEVA, HEIDP+)
HEMINGWAY	78	NP B142 205	R.J. Hemingway <i>et al.</i>	(CERN, ZEEM, NIJM+)
DIBIANCA	75	NP B98 137	F.A. Dibianca, R.J. Endorf	(CMU)
BALTAY	74	PR D9 49	C. Baltay <i>et al.</i>	(COLU, BING) J
COOL	74	PR D10 792	R.L. Cool <i>et al.</i>	(BNL)
Also	72	PRL 29 1630	R.L. Cool <i>et al.</i>	(BNL)
YEH	74	PR D10 3545	N. Yeh <i>et al.</i>	(BING, COLU)
MAYEUR	72	NP B47 333	C. Mayeur <i>et al.</i>	(BRUX, CERN, TUFTS, LOUC)
VOTRUBA	72	NP B45 77	M.F. Votruba, A. Safder, T.M. Ratcliffe	(BIRM+)
WILQUET	72	PL 42B 372	G. Wilquet <i>et al.</i>	(BRUX, CERN, TUFTS+)
DUCLOS	71	NP B32 493	J. Duclos <i>et al.</i>	(CERN)
BINGHAM	70B	PR D1 3010	G.M. Bingham <i>et al.</i>	(UCSD, WASH)
GOLDWASSER	70	PR D1 1960	E.L. Goldwasser, P.F. Schultz	(ILL)
STONE	70	PL 32B 515	S.L. Stone <i>et al.</i>	(ROCH)
DAUBER	69	PR 179 1262	P.M. Dauber <i>et al.</i>	(LRL) J
SHEN	67	PL 25B 443	B.C. Shen, A. Firestone, G. Goldhaber	(UCB+)
BERGE	66	PR 147 945	J.P. Berge <i>et al.</i>	(LRL)
CHIEN	66	PR 152 1171	C.Y. Chien <i>et al.</i>	(YALE, BNL)
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA)
BINGHAM	65	PRSL 285 202	H.H. Bingham	(CERN)
PJERROU	65B	PRL 14 275	G.M. Pjerrou <i>et al.</i>	(UCLA)
Also	65	Thesis	G.M. Pjerrou	(UCLA)
BADIER	64	Dubna Conf. 1 593	J. Badier <i>et al.</i>	(EPOL, SACL, ZEEM)
CARMONY	64B	PRL 12 482	D.D. Carmony <i>et al.</i>	(UCLA) J
HUBBARD	64	PR 135B 183	J.R. Hubbard <i>et al.</i>	(LRL)
FERRO-LUZZI	63	PR 130 1568	M. Ferro-Luzzi <i>et al.</i>	(LRL)
JAUNEAU	63D	Siena Conf. 4	L. Jauneau <i>et al.</i>	(EPOL, CERN, LOUC+)
Also	63B	PL 5 261	L. Jauneau <i>et al.</i>	(EPOL, CERN, LOUC+)
SCHNEIDER	63	PL 4 360	J. Schneider	(CERN)