

τ

$$J = \frac{1}{2}$$

τ discovery paper was PERL 75. $e^+ e^- \rightarrow \tau^+ \tau^-$ cross-section threshold behavior and magnitude are consistent with pointlike spin-1/2 Dirac particle. BRANDELIK 78 ruled out pointlike spin-0 or spin-1 particle. FELDMAN 78 ruled out $J = 3/2$. KIRKBY 79 also ruled out $J=\text{integer}$, $J = 3/2$.

τ MASS

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----------|-------------|---------------------|---|
| 1776.99^{+0.29}_{-0.26} OUR AVERAGE | | | | |
| 1775.1 ± 1.6 | ± 1.0 | 13.3k | 1 ABBIENDI 00A OPAL | 1990–1995 LEP runs |
| 1778.2 ± 0.8 | ± 1.2 | | ANASTASSOV 97 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6$ GeV |
| 1776.96 ^{+0.18} _{-0.21} ^{+0.25} _{-0.17} | 65 | 2 BAI | 96 BES | $E_{\text{cm}}^{\text{ee}} = 3.54\text{--}3.57$ GeV |
| 1776.3 ± 2.4 | ± 1.4 | 11k | 3 ALBRECHT 92M ARG | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6$ GeV |
| 1783 $\begin{array}{l} +3 \\ -4 \end{array}$ | 692 | 4 BACINO | 78B DLCO | $E_{\text{cm}}^{\text{ee}} = 3.1\text{--}7.4$ GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1777.8 ± 0.7 | ± 1.7 | 35k | 5 BAlest | 93 CLEO Repl. by ANAS-TASSOV 97 |
| 1776.9 $\begin{array}{l} +0.4 \\ -0.5 \end{array}$ | ± 0.2 | 14 | 6 BAI | 92 BES Repl. by BAI 96 |

¹ ABBIENDI 00A fit τ pseudomass spectrum in $\tau \rightarrow \pi^\pm \leq 2\pi^0 \nu_\tau$ and $\tau \rightarrow \pi^\pm \pi^+ \pi^- \leq 1\pi^0 \nu_\tau$ decays. Result assumes $m_{\nu_\tau} = 0$.

² BAI 96 fit $\sigma(e^+ e^- \rightarrow \tau^+ \tau^-)$ at different energies near threshold.

³ ALBRECHT 92M fit τ pseudomass spectrum in $\tau^- \rightarrow 2\pi^- \pi^+ \nu_\tau$ decays. Result assumes $m_{\nu_\tau} = 0$.

⁴ BACINO 78B value comes from $e^\pm X^\mp$ threshold. Published mass 1782 MeV increased by 1 MeV using the high precision $\psi(2S)$ mass measurement of ZHOLENTZ 80 to eliminate the absolute SPEAR energy calibration uncertainty.

⁵ BAlest 93 fit spectra of minimum kinematically allowed τ mass in events of the type $e^+ e^- \rightarrow \tau^+ \tau^- \rightarrow (\pi^+ n\pi^0 \nu_\tau)(\pi^- m\pi^0 \nu_\tau)$ $n \leq 2$, $m \leq 2$, $1 \leq n+m \leq 3$. If $m_{\nu_\tau} \neq 0$, result increases by $(m_{\nu_\tau}^2 / 1100$ MeV).

⁶ BAI 92 fit $\sigma(e^+ e^- \rightarrow \tau^+ \tau^-)$ near threshold using $e\mu$ events.

$$(m_{\tau^+} - m_{\tau^-})/m_{\text{average}}$$

A test of *CPT* invariance.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-------------------|------|--------------------|
| <3.0 × 10⁻³ | 90 | ABBIENDI 00A OPAL | | 1990–1995 LEP runs |

τ MEAN LIFE

| <u>VALUE</u> (10^{-15} s) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--------------------------------------|
| 290.6 ± 1.1 OUR AVERAGE | | | | |
| 293.2 ± 2.0 ± 1.5 | | ACCIARRI | 00B L3 | 1991–1995 LEP runs |
| 290.1 ± 1.5 ± 1.1 | | BARATE | 97R ALEP | 1989–1994 LEP runs |
| 291.4 ± 3.0 | | ABREU | 96B DLPH | 1991–1993 LEP runs |
| 289.2 ± 1.7 ± 1.2 | | ALEXANDER | 96E OPAL | 1990–1994 LEP runs |
| 289.0 ± 2.8 ± 4.0 | 57.4k | BALEST | 96 CLEO | $E_{cm}^{ee} = 10.6$ GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 291.2 ± 2.0 ± 1.2 | | BARATE | 97I ALEP | Repl. by BARATE 97R |
| 290.1 ± 4.0 | 34k | ACCIARRI | 96K L3 | Repl. by ACCIARRI 00B |
| 297 ± 9 ± 5 | 1671 | ABE | 95Y SLD | 1992–1993 SLC runs |
| 304 ± 14 ± 7 | 4100 | BATTLE | 92 CLEO | $E_{cm}^{ee} = 10.6$ GeV |
| 301 ± 29 | 3780 | KLEINWORT | 89 JADE | $E_{cm}^{ee} = 35\text{--}46$ GeV |
| 288 ± 16 ± 17 | 807 | AMIDEI | 88 MRK2 | $E_{cm}^{ee} = 29$ GeV |
| 306 ± 20 ± 14 | 695 | BRAUNSCH... | 88C TASS | $E_{cm}^{ee} = 36$ GeV |
| 299 ± 15 ± 10 | 1311 | ABACHI | 87C HRS | $E_{cm}^{ee} = 29$ GeV |
| 295 ± 14 ± 11 | 5696 | ALBRECHT | 87P ARG | $E_{cm}^{ee} = 9.3\text{--}10.6$ GeV |
| 309 ± 17 ± 7 | 3788 | BAND | 87B MAC | $E_{cm}^{ee} = 29$ GeV |
| 325 ± 14 ± 18 | 8470 | BEBEK | 87C CLEO | $E_{cm}^{ee} = 10.5$ GeV |
| 460 ± 190 | 102 | FELDMAN | 82 MRK2 | $E_{cm}^{ee} = 29$ GeV |

τ MAGNETIC MOMENT ANOMALY

The q^2 dependence is expected to be small providing no thresholds are nearby.

$$\mu_\tau/(e\hbar/2m_\tau)-1 = (g_\tau-2)/2$$

For a theoretical calculation [$(g_\tau-2)/2 = 11773(3) \times 10^{-7}$], see SAMUEL 91B.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|------------------------|--|--------------------|
| > -0.052 and < 0.058 (CL = 95%) OUR LIMIT | | | | |
| > -0.052 and < 0.058 | 95 | ACCIARRI | 98E L3 | 1991–1995 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| > -0.007 and < 0.005 | 95 | 7 GONZALEZ-S...00 RVUE | $e^+ e^- \rightarrow \tau^+ \tau^-$ and $W \rightarrow \tau \nu_\tau$ | I |
| > -0.068 and < 0.065 | 95 | 8 ACKERSTAFF 98N OPAL | 1990–1995 LEP runs | |
| > -0.004 and < 0.006 | 95 | 9 ESCRIBANO 97 RVUE | $Z \rightarrow \tau^+ \tau^-$ at LEP | |
| < 0.01 | 95 | 10 ESCRIBANO 93 RVUE | $Z \rightarrow \tau^+ \tau^-$ at LEP | |
| < 0.12 | 90 | GRIFOLS 91 RVUE | $Z \rightarrow \tau \tau \gamma$ at LEP | |
| < 0.023 | 95 | 11 SILVERMAN 83 RVUE | $e^+ e^- \rightarrow \tau^+ \tau^-$ at PETRA | |

⁷ GONZALEZ-SPRINBERG 00 use data on tau lepton production at LEP1, SLC, and LEP2, and data from colliders and LEP2 to determine limits. Assume imaginary component is zero.

⁸ ACKERSTAFF 98N use $Z \rightarrow \tau^+ \tau^- \gamma$ events. The limit applies to an average of the form factor for off-shell τ 's having p^2 ranging from m_τ^2 to $(M_Z - m_\tau)^2$.

⁹ ESCRIBANO 97 use preliminary experimental results.

¹⁰ ESCRIBANO 93 limit derived from $\Gamma(Z \rightarrow \tau^+ \tau^-)$, and is on the absolute value of the magnetic moment anomaly.

¹¹ SILVERMAN 83 limit is derived from $e^+ e^- \rightarrow \tau^+ \tau^-$ total cross-section measurements for q^2 up to $(37 \text{ GeV})^2$.

τ ELECTRIC DIPOLE MOMENT (d_τ)

A nonzero value is forbidden by both T invariance and P invariance.

The q^2 dependence is expected to be small providing no thresholds are nearby.

$\text{Re}(d_\tau)$

| VALUE (10^{-16} e cm) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|----------------------------|----------|---|
| > -3.1 and < 3.1 (CL = 95%) OUR LIMIT | | | | |
| > -3.1 and < 3.1 | 95 | ACCIARRI | 98E L3 | 1991–1995 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <4.6 | 95 | ¹² ALBRECHT | 00 ARG | $E_{\text{cm}}^{\text{ee}} = 10.4 \text{ GeV}$ |
| > -3.8 and < 3.6 | 95 | ¹³ ACKERSTAFF | 98N OPAL | 1990–1995 LEP |
| <0.11 | 95 | ^{14,15} ESCRIBANO | 97 RVUE | $Z \rightarrow \tau^+ \tau^-$ at LEP runs |
| <0.5 | 95 | ¹⁶ ESCRIBANO | 93 RVUE | $Z \rightarrow \tau^+ \tau^-$ at LEP |
| <7 | 90 | GRIFOLS | 91 RVUE | $Z \rightarrow \tau \tau \gamma$ at LEP |
| <1.6 | 90 | DELAGUILA | 90 RVUE | $e^+ e^- \rightarrow \tau^+ \tau^-$ $E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$ |

¹² ALBRECHT 00 use $e^+ e^- \rightarrow \tau^+ \tau^-$ events. Limit is on the absolute value of $\text{Re}(d_\tau)$.

¹³ ACKERSTAFF 98N use $Z \rightarrow \tau^+ \tau^- \gamma$ events. The limit applies to an average of the form factor for off-shell τ 's having p^2 ranging from m_τ^2 to $(M_Z - m_\tau)^2$.

¹⁴ ESCRIBANO 97 derive the relationship $|d_\tau| = \cot \theta_W |d_\tau^W|$ using effective Lagrangian methods, and use a conference result $|d_\tau^W| < 5.8 \times 10^{-18} \text{ e cm}$ at 95% CL (L. Silvestris, ICHEP96) to obtain this result.

¹⁵ ESCRIBANO 97 use preliminary experimental results.

¹⁶ ESCRIBANO 93 limit derived from $\Gamma(Z \rightarrow \tau^+ \tau^-)$, and is on the absolute value of the electric dipole moment.

$\text{Im}(d_\tau)$

| VALUE (10^{-16} e cm) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------------|--------|--|
| <1.8 | 95 | ¹⁷ ALBRECHT | 00 ARG | $E_{\text{cm}}^{\text{ee}} = 10.4 \text{ GeV}$ |
| ¹⁷ ALBRECHT 00 use $e^+ e^- \rightarrow \tau^+ \tau^-$ events. Limit is on the absolute value of $\text{Im}(d_\tau)$. | | | | |

τ WEAK DIPOLE MOMENT (d_τ^w)

A nonzero value is forbidden by CP invariance.

The q^2 dependence is expected to be small providing no thresholds are nearby.

$\text{Re}(d_\tau^w)$

| VALUE (10^{-17} ecm) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------------|----------|--------------------------------------|
| <0.56 | 95 | ACKERSTAFF 97L | OPAL | 1991–1995 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <3.0 | 90 | ¹⁸ ACCIARRI | 98C L3 | 1991–1995 LEP runs |
| <0.78 | 95 | ¹⁹ AKERS | 95F OPAL | Repl. by ACKER-STAFF 97L |
| <1.5 | 95 | ¹⁹ BUSKULIC | 95C ALEP | 1990–1992 LEP runs |
| <7.0 | 95 | ¹⁹ ACTON | 92F OPAL | $Z \rightarrow \tau^+ \tau^-$ at LEP |
| <3.7 | 95 | ¹⁹ BUSKULIC | 92J ALEP | Repl. by BUSKULIC 95C |

¹⁸ ACCIARRI 98C limit is on the absolute value of the real part of the weak dipole moment.

¹⁹ Limit is on the absolute value of the real part of the weak dipole moment, and applies for $q^2 = m_Z^2$.

$\text{Im}(d_\tau^w)$

| VALUE (10^{-17} ecm) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|---------------------|----------|--------------------------|
| <1.5 | 95 | ACKERSTAFF 97L | OPAL | 1991–1995 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <4.5 | 95 | ²⁰ AKERS | 95F OPAL | Repl. by ACKER-STAFF 97L |
| ²⁰ Limit is on the absolute value of the imaginary part of the weak dipole moment, and applies for $q^2 = m_Z^2$. | | | | |

τ WEAK ANOMALOUS MAGNETIC DIPOLE MOMENT (α_τ^w)

Electroweak radiative corrections are expected to contribute at the 10^{-6} level. See BERNABEU 95.

The q^2 dependence is expected to be small providing no thresholds are nearby.

$\text{Re}(\alpha_\tau^w)$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------------------|--------|---|
| <4.5 × 10⁻³ | 90 | ²¹ ACCIARRI | 98C L3 | 1991–1995 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| > -0.0024 and < 0.0025 | 95 | ²² GONZALEZ-S..00 | RVUE | $e^+ e^- \rightarrow \tau^+ \tau^-$ and $W \rightarrow \tau \nu_\tau$ |

²¹ ACCIARRI 98C limit is on the absolute value of the real part of the weak anomalous magnetic dipole moment.

²² GONZALEZ-SPRINBERG 00 use data on tau lepton production at LEP1, SLC, and LEP2, and data from colliders and LEP2 to determine limits. Assume imaginary component is zero.

| Im(α_τ^w) | VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|--------------|------------|--------------------|-------------|--------------------|
| <9.9 × 10⁻³ | 90 | 23 | ACCIARRI | 98C L3 | 1991–1995 LEP runs |
| ²³ ACCIARRI 98C limit is on the absolute value of the imaginary part of the weak anomalous magnetic dipole moment. | | | | | |

τ^- DECAY MODES

τ^+ modes are charge conjugates of the modes below. “ h^\pm ” stands for π^\pm or K^\pm . “ ℓ ” stands for e or μ . “Neutral” means neutral hadron whose decay products include γ 's and/or π^0 's.

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|---|--|-----------------------------------|
| Modes with one charged particle | | |
| Γ_1 particle $^- \geq 0$ neutrals $\geq 0 K_L^0 \nu_\tau$ ("1-prong") | (84.71 \pm 0.13) % | S=1.2 |
| Γ_2 particle $^- \geq 0$ neutrals $\geq 0 K^0 \nu_\tau$ | (85.32 \pm 0.13) % | S=1.2 |
| Γ_3 $\mu^- \bar{\nu}_\mu \nu_\tau$ | [a] (17.37 \pm 0.07) % | |
| Γ_4 $\mu^- \bar{\nu}_\mu \nu_\tau \gamma$ | [b] (3.6 \pm 0.4) $\times 10^{-3}$ | |
| Γ_5 $e^- \bar{\nu}_e \nu_\tau$ | [a] (17.83 \pm 0.06) % | |
| Γ_6 $e^- \bar{\nu}_e \nu_\tau \gamma$ | [b] (1.75 \pm 0.18) % | |
| Γ_7 $h^- \geq 0$ neutrals $\geq 0 K_L^0 \nu_\tau$ | (49.51 \pm 0.15) % | S=1.2 |
| Γ_8 $h^- \geq 0 K_L^0 \nu_\tau$ | (12.35 \pm 0.12) % | S=1.4 |
| Γ_9 $h^- \nu_\tau$ | (11.79 \pm 0.12) % | S=1.4 |
| Γ_{10} $\pi^- \nu_\tau$ | [a] (11.09 \pm 0.12) % | S=1.4 |
| Γ_{11} $K^- \nu_\tau$ | [a] (6.99 \pm 0.27) $\times 10^{-3}$ | |
| Γ_{12} $h^- \geq 1$ neutrals ν_τ | (36.88 \pm 0.17) % | S=1.2 |
| Γ_{13} $h^- \pi^0 \nu_\tau$ | (25.86 \pm 0.14) % | S=1.1 |
| Γ_{14} $\pi^- \pi^0 \nu_\tau$ | [a] (25.40 \pm 0.14) % | S=1.1 |
| Γ_{15} $\pi^- \pi^0$ non- $\rho(770)$ ν_τ | (3.0 \pm 3.2) $\times 10^{-3}$ | |
| Γ_{16} $K^- \pi^0 \nu_\tau$ | [a] (4.54 \pm 0.33) $\times 10^{-3}$ | |
| Γ_{17} $h^- \geq 2 \pi^0 \nu_\tau$ | (10.73 \pm 0.16) % | S=1.2 |
| Γ_{18} $h^- 2 \pi^0 \nu_\tau$ | (9.36 \pm 0.14) % | S=1.2 |
| Γ_{19} $h^- 2 \pi^0 \nu_\tau$ (ex. K^0) | (9.19 \pm 0.14) % | S=1.2 |
| Γ_{20} $\pi^- 2 \pi^0 \nu_\tau$ (ex. K^0) | [a] (9.13 \pm 0.14) % | S=1.2 |
| Γ_{21} $\pi^- 2 \pi^0 \nu_\tau$ (ex. K^0), scalar | < 9 $\times 10^{-3}$ | CL=95% |
| Γ_{22} $\pi^- 2 \pi^0 \nu_\tau$ (ex. K^0), vector | < 7 $\times 10^{-3}$ | CL=95% |
| Γ_{23} $K^- 2 \pi^0 \nu_\tau$ (ex. K^0) | [a] (6.0 \pm 2.4) $\times 10^{-4}$ | |

| | | | |
|---------------|---|--|-------|
| Γ_{24} | $h^- \geq 3\pi^0 \nu_\tau$ | (1.37 ± 0.11) % | S=1.1 |
| Γ_{25} | $h^- 3\pi^0 \nu_\tau$ | (1.21 ± 0.10) % | S=1.1 |
| Γ_{26} | $\pi^- 3\pi^0 \nu_\tau$ (ex. K^0) | [a] (1.08 ± 0.10) % | S=1.1 |
| Γ_{27} | $K^- 3\pi^0 \nu_\tau$ (ex. K^0 , η) | [a] ($3.9 \begin{array}{l} +2.3 \\ -2.1 \end{array} \times 10^{-4}$) | |
| Γ_{28} | $h^- 4\pi^0 \nu_\tau$ (ex. K^0) | (1.6 ± 0.6) $\times 10^{-3}$ | |
| Γ_{29} | $h^- 4\pi^0 \nu_\tau$ (ex. K^0, η) | [a] ($1.0 \begin{array}{l} +0.6 \\ -0.5 \end{array} \times 10^{-3}$) | |
| Γ_{30} | $K^- \geq 0\pi^0 \geq 0K^0 \nu_\tau$ | (1.58 ± 0.06) % | |
| Γ_{31} | $K^- \geq 1 (\pi^0 \text{ or } K^0) \nu_\tau$ | (8.8 ± 0.5) $\times 10^{-3}$ | |

Modes with K^0 's

| | | | |
|---------------|---|--|--------|
| Γ_{32} | K^0 (particles) $-\nu_\tau$ | (1.71 ± 0.06) % | S=1.1 |
| Γ_{33} | $h^- \bar{K}^0 \geq 0$ neutrals $\geq 0 K_L^0 \nu_\tau$ | (1.67 ± 0.06) % | S=1.1 |
| Γ_{34} | $h^- \bar{K}^0 \nu_\tau$ | (1.06 ± 0.05) % | S=1.2 |
| Γ_{35} | $\pi^- \bar{K}^0 \nu_\tau$ | [a] (9.0 ± 0.4) $\times 10^{-3}$ | S=1.1 |
| Γ_{36} | $\pi^- \bar{K}^0$ (non- $K^*(892)^-$) ν_τ | < 1.7×10^{-3} | CL=95% |
| Γ_{37} | $K^- K^0 \nu_\tau$ | [a] (1.55 ± 0.17) $\times 10^{-3}$ | |
| Γ_{38} | $K^- \bar{K}^0 \geq 0 \pi^0 \nu_\tau$ | (3.12 ± 0.25) $\times 10^{-3}$ | |
| Γ_{39} | $h^- \bar{K}^0 \pi^0 \nu_\tau$ | (5.3 ± 0.4) $\times 10^{-3}$ | |
| Γ_{40} | $\pi^- \bar{K}^0 \pi^0 \nu_\tau$ | [a] (3.8 ± 0.4) $\times 10^{-3}$ | |
| Γ_{41} | $\bar{K}^0 \rho^- \nu_\tau$ | (2.2 ± 0.5) $\times 10^{-3}$ | |
| Γ_{42} | $K^- K^0 \pi^0 \nu_\tau$ | [a] (1.57 ± 0.21) $\times 10^{-3}$ | |
| Γ_{43} | $\pi^- \bar{K}^0 \geq 1 \pi^0 \nu_\tau$ | (3.2 ± 1.0) $\times 10^{-3}$ | |
| Γ_{44} | $\pi^- \bar{K}^0 \pi^0 \pi^0 \nu_\tau$ | (2.6 ± 2.4) $\times 10^{-4}$ | |
| Γ_{45} | $K^- K^0 \pi^0 \pi^0 \nu_\tau$ | < 1.6×10^{-4} | CL=95% |
| Γ_{46} | $\pi^- K^0 \bar{K}^0 \nu_\tau$ | [a] (1.19 ± 0.20) $\times 10^{-3}$ | S=1.2 |
| Γ_{47} | $\pi^- K_S^0 K_S^0 \nu_\tau$ | (3.0 ± 0.5) $\times 10^{-4}$ | S=1.2 |
| Γ_{48} | $\pi^- K_S^0 K_L^0 \nu_\tau$ | (6.0 ± 1.0) $\times 10^{-4}$ | S=1.2 |
| Γ_{49} | $\pi^- K^0 \bar{K}^0 \pi^0 \nu_\tau$ | (3.1 ± 2.3) $\times 10^{-4}$ | |
| Γ_{50} | $\pi^- K_S^0 K_S^0 \pi^0 \nu_\tau$ | < 2.0×10^{-4} | CL=95% |
| Γ_{51} | $\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau$ | (3.1 ± 1.2) $\times 10^{-4}$ | |
| Γ_{52} | $K^0 h^+ h^- h^- \geq 0$ neutrals ν_τ | < 1.7×10^{-3} | CL=95% |
| Γ_{53} | $K^0 h^+ h^- h^- \nu_\tau$ | (2.3 ± 2.0) $\times 10^{-4}$ | |

Modes with three charged particles

| | | | |
|---------------|--|------------------------|-------|
| Γ_{54} | $h^- h^- h^+ \geq 0$ neut. ν_τ ("3-prong") | (15.18 ± 0.13) % | S=1.2 |
| Γ_{55} | $h^- h^- h^+ \geq 0$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^+ \pi^-$) | (14.58 ± 0.13) % | S=1.2 |
| Γ_{56} | $\pi^- \pi^+ \pi^- \geq 0$ neutrals ν_τ | (14.49 ± 0.14) % | |
| Γ_{57} | $h^- h^- h^+ \nu_\tau$ | (9.97 ± 0.10) % | S=1.1 |
| Γ_{58} | $h^- h^- h^+ \nu_\tau$ (ex. K^0) | (9.61 ± 0.10) % | S=1.1 |
| Γ_{59} | $h^- h^- h^+ \nu_\tau$ (ex. K^0, ω) | (9.56 ± 0.10) % | S=1.1 |
| Γ_{60} | $\pi^- \pi^+ \pi^- \nu_\tau$ | (9.49 ± 0.11) % | S=1.1 |

| | | | |
|----------------|--|--|--------|
| Γ_{61} | $\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0) | (9.18 ± 0.11) % | S=1.1 |
| Γ_{62} | $\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0), non-axial vector | < 2.4 % | CL=95% |
| Γ_{63} | $\pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0, ω) | [a] (9.13 ± 0.11) % | S=1.1 |
| Γ_{64} | $h^- h^- h^+ \geq 1$ neutrals ν_τ | (5.17 ± 0.11) % | S=1.2 |
| Γ_{65} | $h^- h^- h^+ \geq 1$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^+ \pi^-$) | (4.97 ± 0.11) % | S=1.2 |
| Γ_{66} | $h^- h^- h^+ \pi^0 \nu_\tau$ | (4.49 ± 0.08) % | |
| Γ_{67} | $h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0) | (4.30 ± 0.08) % | |
| Γ_{68} | $h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0, ω) | (2.58 ± 0.08) % | |
| Γ_{69} | $\pi^- \pi^+ \pi^- \pi^0 \nu_\tau$ | (4.32 ± 0.08) % | |
| Γ_{70} | $\pi^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0) | (4.20 ± 0.08) % | |
| Γ_{71} | $\pi^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0, ω) | [a] (2.47 ± 0.08) % | |
| Γ_{72} | $h^- (\rho \pi)^0 \nu_\tau$ | | |
| Γ_{73} | $(a_1(1260) h)^- \nu_\tau$ | | |
| Γ_{74} | $h^- \rho \pi^0 \nu_\tau$ | | |
| Γ_{75} | $h^- \rho^+ h^- \nu_\tau$ | | |
| Γ_{76} | $h^- \rho^- h^+ \nu_\tau$ | | |
| Γ_{77} | $h^- h^- h^+ 2\pi^0 \nu_\tau$ | (5.4 ± 0.4) $\times 10^{-3}$ | |
| Γ_{78} | $h^- h^- h^+ 2\pi^0 \nu_\tau$ (ex. K^0) | (5.3 ± 0.4) $\times 10^{-3}$ | |
| Γ_{79} | $h^- h^- h^+ 2\pi^0 \nu_\tau$ (ex. K^0, ω, η) | [a] (1.1 ± 0.4) $\times 10^{-3}$ | |
| Γ_{80} | $h^- h^- h^+ \geq 3\pi^0 \nu_\tau$ | [a] ($1.3^{+0.8}_{-0.7}$) $\times 10^{-3}$ | S=1.3 |
| Γ_{81} | $h^- h^- h^+ 3\pi^0 \nu_\tau$ | (2.9 ± 0.8) $\times 10^{-4}$ | |
| Γ_{82} | $K^- h^+ h^- \geq 0$ neutrals ν_τ | (6.5 ± 0.5) $\times 10^{-3}$ | S=1.4 |
| Γ_{83} | $K^- h^+ \pi^- \nu_\tau$ (ex. K^0) | (4.3 ± 0.5) $\times 10^{-3}$ | S=1.5 |
| Γ_{84} | $K^- h^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0) | (1.07 ± 0.22) $\times 10^{-3}$ | |
| Γ_{85} | $K^- \pi^+ \pi^- \geq 0$ neutrals ν_τ | (4.4 ± 0.5) $\times 10^{-3}$ | S=1.4 |
| Γ_{86} | $K^- \pi^+ \pi^- \geq 0 \pi^0 \nu_\tau$ (ex. K^0) | (3.4 ± 0.5) $\times 10^{-3}$ | S=1.4 |
| Γ_{87} | $K^- \pi^+ \pi^- \nu_\tau$ | (3.2 ± 0.5) $\times 10^{-3}$ | S=1.5 |
| Γ_{88} | $K^- \pi^+ \pi^- \nu_\tau$ (ex. K^0) | [a] (2.7 ± 0.5) $\times 10^{-3}$ | S=1.5 |
| Γ_{89} | $K^- \rho^0 \nu_\tau \rightarrow$ $K^- \pi^+ \pi^- \nu_\tau$ | (1.3 ± 0.5) $\times 10^{-3}$ | |
| Γ_{90} | $K^- \pi^+ \pi^- \pi^0 \nu_\tau$ | (1.20 ± 0.25) $\times 10^{-3}$ | |
| Γ_{91} | $K^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0) | (6.7 ± 2.4) $\times 10^{-4}$ | |
| Γ_{92} | $K^- \pi^+ \pi^- \pi^0 \nu_\tau$ (ex. K^0, η) | [a] (6.0 ± 2.4) $\times 10^{-4}$ | |
| Γ_{93} | $K^- \pi^+ K^- \geq 0$ neut. ν_τ | < 9 $\times 10^{-4}$ | CL=95% |
| Γ_{94} | $K^- K^+ \pi^- \geq 0$ neut. ν_τ | (2.01 ± 0.23) $\times 10^{-3}$ | |
| Γ_{95} | $K^- K^+ \pi^- \nu_\tau$ | [a] (1.61 ± 0.18) $\times 10^{-3}$ | |
| Γ_{96} | $K^- K^+ \pi^- \pi^0 \nu_\tau$ | [a] (4.0 ± 1.6) $\times 10^{-4}$ | |
| Γ_{97} | $K^- K^+ K^- \geq 0$ neut. ν_τ | < 2.1 $\times 10^{-3}$ | CL=95% |
| Γ_{98} | $K^- K^+ K^- \nu_\tau$ | < 1.9 $\times 10^{-4}$ | CL=90% |
| Γ_{99} | $\pi^- K^+ \pi^- \geq 0$ neut. ν_τ | < 2.5 $\times 10^{-3}$ | CL=95% |
| Γ_{100} | $e^- e^- e^+ \bar{\nu}_e \nu_\tau$ | (2.8 ± 1.5) $\times 10^{-5}$ | |
| Γ_{101} | $\mu^- e^- e^+ \bar{\nu}_\mu \nu_\tau$ | < 3.6 $\times 10^{-5}$ | CL=90% |

Modes with five charged particles

| | | | |
|----------------|---|------------------------------------|--------|
| Γ_{102} | $3h^- 2h^+ \geq 0$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^- \pi^+$) ("5-prong") | $(9.9 \pm 0.7) \times 10^{-4}$ | |
| Γ_{103} | $3h^- 2h^+ \nu_\tau$ (ex. K^0) | [a] $(7.8 \pm 0.6) \times 10^{-4}$ | |
| Γ_{104} | $3h^- 2h^+ \pi^0 \nu_\tau$ (ex. K^0) | [a] $(2.2 \pm 0.5) \times 10^{-4}$ | |
| Γ_{105} | $3h^- 2h^+ 2\pi^0 \nu_\tau$ | $< 1.1 \times 10^{-4}$ | CL=90% |

Miscellaneous other allowed modes

| | | | |
|----------------|---|--------------------------------------|--------|
| Γ_{106} | $(5\pi)^- \nu_\tau$ | $(7.9 \pm 0.7) \times 10^{-3}$ | |
| Γ_{107} | $4h^- 3h^+ \geq 0$ neutrals ν_τ ("7-prong") | $< 2.4 \times 10^{-6}$ | CL=90% |
| Γ_{108} | $X^- (S=-1) \nu_\tau$ | $(2.89 \pm 0.09) \%$ | S=1.1 |
| Γ_{109} | $K^*(892)^- \geq 0 (h^0 \neq K_S^0) \nu_\tau$ | $(1.94 \pm 0.31) \%$ | |
| Γ_{110} | $K^*(892)^- \geq 0$ neutrals ν_τ | $(1.33 \pm 0.13) \%$ | |
| Γ_{111} | $K^*(892)^- \nu_\tau$ | $(1.29 \pm 0.05) \%$ | |
| Γ_{112} | $K^*(892)^0 K^- \geq 0$ neutrals ν_τ | $(3.2 \pm 1.4) \times 10^{-3}$ | |
| Γ_{113} | $K^*(892)^0 K^- \nu_\tau$ | $(2.1 \pm 0.4) \times 10^{-3}$ | |
| Γ_{114} | $\bar{K}^*(892)^0 \pi^- \geq 0$ neutrals ν_τ | $(3.8 \pm 1.7) \times 10^{-3}$ | |
| Γ_{115} | $\bar{K}^*(892)^0 \pi^- \nu_\tau$ | $(2.2 \pm 0.5) \times 10^{-3}$ | |
| Γ_{116} | $(\bar{K}^*(892)\pi)^- \nu_\tau \rightarrow \pi^- \bar{K}^0 \pi^0 \nu_\tau$ | $(1.0 \pm 0.4) \times 10^{-3}$ | |
| Γ_{117} | $K_1(1270)^- \nu_\tau$ | $(4.7 \pm 1.1) \times 10^{-3}$ | |
| Γ_{118} | $K_1(1400)^- \nu_\tau$ | $(1.7 \pm 2.6) \times 10^{-3}$ | S=1.7 |
| Γ_{119} | $K^*(1410)^- \nu_\tau$ | $(1.5 \pm 1.4) \times 10^{-3}$ | |
| Γ_{120} | $K_0^*(1430)^- \nu_\tau$ | $< 5 \times 10^{-4}$ | CL=95% |
| Γ_{121} | $K_2^*(1430)^- \nu_\tau$ | $< 3 \times 10^{-3}$ | CL=95% |
| Γ_{122} | $a_0(980)^- \geq 0$ neutrals ν_τ | | |
| Γ_{123} | $\eta \pi^- \nu_\tau$ | $< 1.4 \times 10^{-4}$ | CL=95% |
| Γ_{124} | $\eta \pi^- \pi^0 \nu_\tau$ | [a] $(1.74 \pm 0.24) \times 10^{-3}$ | |
| Γ_{125} | $\eta \pi^- \pi^0 \pi^0 \nu_\tau$ | $(1.4 \pm 0.7) \times 10^{-4}$ | |
| Γ_{126} | $\eta K^- \nu_\tau$ | [a] $(2.7 \pm 0.6) \times 10^{-4}$ | |
| Γ_{127} | $\eta K^*(892)^- \nu_\tau$ | $(2.9 \pm 0.9) \times 10^{-4}$ | |
| Γ_{128} | $\eta K^- \pi^0 \nu_\tau$ | $(1.8 \pm 0.9) \times 10^{-4}$ | |
| Γ_{129} | $\eta \bar{K}^0 \pi^- \nu_\tau$ | $(2.2 \pm 0.7) \times 10^{-4}$ | |
| Γ_{130} | $\eta \pi^+ \pi^- \pi^- \geq 0$ neutrals ν_τ | $< 3 \times 10^{-3}$ | CL=90% |
| Γ_{131} | $\eta \pi^- \pi^+ \pi^- \nu_\tau$ | $(3.4 \pm 0.8) \times 10^{-4}$ | |
| Γ_{132} | $\eta a_1(1260)^- \nu_\tau \rightarrow \eta \pi^- \rho^0 \nu_\tau$ | $< 3.9 \times 10^{-4}$ | CL=90% |
| Γ_{133} | $\eta \eta \pi^- \nu_\tau$ | $< 1.1 \times 10^{-4}$ | CL=95% |
| Γ_{134} | $\eta \eta \pi^- \pi^0 \nu_\tau$ | $< 2.0 \times 10^{-4}$ | CL=95% |
| Γ_{135} | $\eta'(958) \pi^- \nu_\tau$ | $< 7.4 \times 10^{-5}$ | CL=90% |

| | | | | |
|----------------|--|-------------------------|------------------|--------|
| Γ_{136} | $\eta'(958)\pi^-\pi^0\nu_\tau$ | < 8.0 | $\times 10^{-5}$ | CL=90% |
| Γ_{137} | $\phi\pi^-\nu_\tau$ | < 2.0 | $\times 10^{-4}$ | CL=90% |
| Γ_{138} | $\phi K^-\nu_\tau$ | < 6.7 | $\times 10^{-5}$ | CL=90% |
| Γ_{139} | $f_1(1285)\pi^-\nu_\tau$ | (5.8 \pm 2.3) | $\times 10^{-4}$ | |
| Γ_{140} | $f_1(1285)\pi^-\nu_\tau \rightarrow \eta\pi^-\pi^+\pi^-\nu_\tau$ | (1.9 \pm 0.7) | $\times 10^{-4}$ | |
| Γ_{141} | $\pi(1300)^-\nu_\tau \rightarrow (\rho\pi)^-\nu_\tau \rightarrow (3\pi)^-\nu_\tau$ | < 1.0 | $\times 10^{-4}$ | CL=90% |
| Γ_{142} | $\pi(1300)^-\nu_\tau \rightarrow ((\pi\pi)_{S-\text{wave}}\pi)^-\nu_\tau \rightarrow (3\pi)^-\nu_\tau$ | < 1.9 | $\times 10^{-4}$ | CL=90% |
| Γ_{143} | $h^-\omega \geq 0 \text{ neutrals } \nu_\tau$ | (2.36 \pm 0.08) | % | |
| Γ_{144} | $h^-\omega\nu_\tau$ | [a] (1.93 \pm 0.06) | % | |
| Γ_{145} | $h^-\omega\pi^0\nu_\tau$ | [a] (4.3 \pm 0.5) | $\times 10^{-3}$ | |
| Γ_{146} | $h^-\omega 2\pi^0\nu_\tau$ | (1.9 \pm 0.8) | $\times 10^{-4}$ | |

**Lepton Family number (*LF*), Lepton number (*L*),
or Baryon number (*B*) violating modes**

(In the modes below, ℓ means a sum over e and μ modes)

L means lepton number violation (e.g. $\tau^- \rightarrow e^+\pi^-\pi^-$). Following common usage, *LF* means lepton family violation *and not* lepton number violation (e.g. $\tau^- \rightarrow e^-\pi^+\pi^-$). *B* means baryon number violation.

| | | | | | |
|----------------|-------------------------|-----------|-------|------------------|--------|
| Γ_{147} | $e^-\gamma$ | <i>LF</i> | < 2.7 | $\times 10^{-6}$ | CL=90% |
| Γ_{148} | $\mu^-\gamma$ | <i>LF</i> | < 1.1 | $\times 10^{-6}$ | CL=90% |
| Γ_{149} | $e^-\pi^0$ | <i>LF</i> | < 3.7 | $\times 10^{-6}$ | CL=90% |
| Γ_{150} | $\mu^-\pi^0$ | <i>LF</i> | < 4.0 | $\times 10^{-6}$ | CL=90% |
| Γ_{151} | e^-K^0 | <i>LF</i> | < 1.3 | $\times 10^{-3}$ | CL=90% |
| Γ_{152} | μ^-K^0 | <i>LF</i> | < 1.0 | $\times 10^{-3}$ | CL=90% |
| Γ_{153} | $e^-\eta$ | <i>LF</i> | < 8.2 | $\times 10^{-6}$ | CL=90% |
| Γ_{154} | $\mu^-\eta$ | <i>LF</i> | < 9.6 | $\times 10^{-6}$ | CL=90% |
| Γ_{155} | $e^-\rho^0$ | <i>LF</i> | < 2.0 | $\times 10^{-6}$ | CL=90% |
| Γ_{156} | $\mu^-\rho^0$ | <i>LF</i> | < 6.3 | $\times 10^{-6}$ | CL=90% |
| Γ_{157} | $e^-K^*(892)^0$ | <i>LF</i> | < 5.1 | $\times 10^{-6}$ | CL=90% |
| Γ_{158} | $\mu^-K^*(892)^0$ | <i>LF</i> | < 7.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{159} | $e^-\bar{K}^*(892)^0$ | <i>LF</i> | < 7.4 | $\times 10^{-6}$ | CL=90% |
| Γ_{160} | $\mu^-\bar{K}^*(892)^0$ | <i>LF</i> | < 7.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{161} | $e^-\phi$ | <i>LF</i> | < 6.9 | $\times 10^{-6}$ | CL=90% |
| Γ_{162} | $\mu^-\phi$ | <i>LF</i> | < 7.0 | $\times 10^{-6}$ | CL=90% |
| Γ_{163} | $\pi^-\gamma$ | <i>L</i> | < 2.8 | $\times 10^{-4}$ | CL=90% |
| Γ_{164} | $\pi^-\pi^0$ | <i>L</i> | < 3.7 | $\times 10^{-4}$ | CL=90% |
| Γ_{165} | $e^-e^+e^-$ | <i>LF</i> | < 2.9 | $\times 10^{-6}$ | CL=90% |
| Γ_{166} | $e^-\mu^+\mu^-$ | <i>LF</i> | < 1.8 | $\times 10^{-6}$ | CL=90% |
| Γ_{167} | $e^+\mu^-\mu^-$ | <i>LF</i> | < 1.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{168} | $\mu^+e^+e^-$ | <i>LF</i> | < 1.7 | $\times 10^{-6}$ | CL=90% |

| | | | | | |
|----------------|----------------------|------------|-------|------------------|--------|
| Γ_{169} | $\mu^+ e^- e^-$ | <i>LF</i> | < 1.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{170} | $\mu^- \mu^+ \mu^-$ | <i>LF</i> | < 1.9 | $\times 10^{-6}$ | CL=90% |
| Γ_{171} | $e^- \pi^+ \pi^-$ | <i>LF</i> | < 2.2 | $\times 10^{-6}$ | CL=90% |
| Γ_{172} | $e^+ \pi^- \pi^-$ | <i>L</i> | < 1.9 | $\times 10^{-6}$ | CL=90% |
| Γ_{173} | $\mu^- \pi^+ \pi^-$ | <i>LF</i> | < 8.2 | $\times 10^{-6}$ | CL=90% |
| Γ_{174} | $\mu^+ \pi^- \pi^-$ | <i>L</i> | < 3.4 | $\times 10^{-6}$ | CL=90% |
| Γ_{175} | $e^- \pi^+ K^-$ | <i>LF</i> | < 6.4 | $\times 10^{-6}$ | CL=90% |
| Γ_{176} | $e^- \pi^- K^+$ | <i>LF</i> | < 3.8 | $\times 10^{-6}$ | CL=90% |
| Γ_{177} | $e^+ \pi^- K^-$ | <i>L</i> | < 2.1 | $\times 10^{-6}$ | CL=90% |
| Γ_{178} | $e^- K^+ K^-$ | <i>LF</i> | < 6.0 | $\times 10^{-6}$ | CL=90% |
| Γ_{179} | $e^+ K^- K^-$ | <i>L</i> | < 3.8 | $\times 10^{-6}$ | CL=90% |
| Γ_{180} | $\mu^- \pi^+ K^-$ | <i>LF</i> | < 7.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{181} | $\mu^- \pi^- K^+$ | <i>LF</i> | < 7.4 | $\times 10^{-6}$ | CL=90% |
| Γ_{182} | $\mu^+ \pi^- K^-$ | <i>L</i> | < 7.0 | $\times 10^{-6}$ | CL=90% |
| Γ_{183} | $\mu^- K^+ K^-$ | <i>LF</i> | < 1.5 | $\times 10^{-5}$ | CL=90% |
| Γ_{184} | $\mu^+ K^- K^-$ | <i>L</i> | < 6.0 | $\times 10^{-6}$ | CL=90% |
| Γ_{185} | $e^- \pi^0 \pi^0$ | <i>LF</i> | < 6.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{186} | $\mu^- \pi^0 \pi^0$ | <i>LF</i> | < 1.4 | $\times 10^{-5}$ | CL=90% |
| Γ_{187} | $e^- \eta \eta$ | <i>LF</i> | < 3.5 | $\times 10^{-5}$ | CL=90% |
| Γ_{188} | $\mu^- \eta \eta$ | <i>LF</i> | < 6.0 | $\times 10^{-5}$ | CL=90% |
| Γ_{189} | $e^- \pi^0 \eta$ | <i>LF</i> | < 2.4 | $\times 10^{-5}$ | CL=90% |
| Γ_{190} | $\mu^- \pi^0 \eta$ | <i>LF</i> | < 2.2 | $\times 10^{-5}$ | CL=90% |
| Γ_{191} | $\bar{p} \gamma$ | <i>L,B</i> | < 3.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{192} | $\bar{p} \pi^0$ | <i>L,B</i> | < 1.5 | $\times 10^{-5}$ | CL=90% |
| Γ_{193} | $\bar{p} 2\pi^0$ | <i>L,B</i> | < 3.3 | $\times 10^{-5}$ | CL=90% |
| Γ_{194} | $\bar{p} \eta$ | <i>L,B</i> | < 8.9 | $\times 10^{-6}$ | CL=90% |
| Γ_{195} | $\bar{p} \pi^0 \eta$ | <i>L,B</i> | < 2.7 | $\times 10^{-5}$ | CL=90% |
| Γ_{196} | e^- light boson | <i>LF</i> | < 2.7 | $\times 10^{-3}$ | CL=95% |
| Γ_{197} | μ^- light boson | <i>LF</i> | < 5 | $\times 10^{-3}$ | CL=95% |

[a] Basis mode for the τ .

[b] See the Particle Listings below for the energy limits used in this measurement.

CONSTRAINED FIT INFORMATION

An overall fit to 70 branching ratios uses 139 measurements and one constraint to determine 30 parameters. The overall fit has a $\chi^2 = 74.9$ for 110 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| x_5 | 8 | | | | | | | | | |
|-----------|-------|-------|----------|----------|----------|----------|----------|----------|----------|----------|
| x_{10} | -10 | -9 | | | | | | | | |
| x_{11} | 0 | 0 | -22 | | | | | | | |
| x_{14} | -13 | -12 | -20 | 1 | | | | | | |
| x_{16} | 0 | 0 | 1 | -4 | -24 | | | | | |
| x_{20} | -13 | -12 | -20 | 1 | -27 | 1 | | | | |
| x_{23} | 0 | 0 | 1 | -3 | 1 | -4 | -17 | | | |
| x_{26} | -8 | -7 | -12 | 1 | -12 | 1 | -19 | 1 | | |
| x_{27} | 0 | 0 | 1 | -3 | 1 | -3 | 0 | -3 | -23 | |
| x_{29} | -4 | -4 | -6 | 0 | -8 | 0 | -10 | 0 | -6 | 0 |
| x_{35} | -1 | -1 | -13 | 0 | -2 | 0 | -6 | 0 | -1 | 0 |
| x_{37} | 0 | 0 | -4 | -2 | 0 | -2 | -2 | -2 | 0 | -1 |
| x_{40} | -2 | -2 | -2 | 0 | -4 | 1 | -3 | 0 | -8 | 0 |
| x_{42} | -1 | -1 | 0 | -2 | -1 | -3 | -1 | -2 | -2 | -2 |
| x_{46} | -1 | -1 | -4 | 0 | -2 | 0 | -2 | 0 | -1 | 0 |
| x_{63} | -6 | -6 | -9 | 0 | -13 | 0 | -12 | 0 | -8 | 0 |
| x_{71} | -3 | -3 | -5 | 0 | -6 | 0 | -6 | 0 | -3 | 0 |
| x_{79} | -1 | -1 | -2 | 0 | -2 | 0 | -3 | 0 | -2 | 0 |
| x_{80} | -5 | -4 | -7 | 0 | -9 | 0 | -10 | 0 | -6 | 0 |
| x_{88} | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| x_{92} | 0 | 0 | 0 | 0 | -1 | 0 | -1 | 0 | 0 | 0 |
| x_{95} | 0 | 0 | 0 | 0 | -1 | 0 | -1 | 0 | 0 | 0 |
| x_{96} | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| x_{103} | 0 | 0 | -1 | 0 | -1 | 0 | -1 | 0 | 0 | 0 |
| x_{104} | 0 | 0 | 0 | 0 | -1 | 0 | -1 | 0 | 0 | 0 |
| x_{124} | -1 | -1 | -1 | 0 | -2 | 0 | -2 | 0 | -1 | 0 |
| x_{126} | 0 | 0 | 0 | -1 | 0 | -1 | 0 | 0 | -2 | 0 |
| x_{144} | -2 | -2 | -3 | 0 | -4 | 0 | -4 | 0 | -2 | 0 |
| x_{145} | -2 | -2 | -3 | 0 | -4 | 0 | -4 | 0 | -2 | 0 |
| | x_3 | x_5 | x_{10} | x_{11} | x_{14} | x_{16} | x_{20} | x_{23} | x_{26} | x_{27} |

| | x_{35} | x_{37} | x_{40} | x_{42} | x_{46} | x_{63} | x_{71} | x_{79} | x_{80} | x_{88} | x_{92} | x_{95} | x_{96} | x_{103} | x_{104} | x_{124} | x_{126} | x_{144} | x_{145} |
|-----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|-----------|-----------|-----------|-----------|-----------|-----------|
| | x_{29} | x_{35} | x_{37} | x_{40} | x_{42} | x_{46} | x_{63} | x_{71} | x_{79} | x_{80} | | | | | | | | | |
| x_{92} | | | | | | | | | | | -19 | | | | | | | | |
| x_{95} | | | | | | | | | | | -14 | 8 | | | | | | | |
| x_{96} | | | | | | | | | | | 10 | -47 | -14 | | | | | | |
| x_{103} | | | | | | | | | | | 0 | 0 | 0 | 0 | | | | | |
| x_{104} | | | | | | | | | | | 0 | 0 | 0 | 0 | -19 | | | | |
| x_{124} | | | | | | | | | | | 0 | 0 | 0 | 0 | 0 | 0 | | | |
| x_{126} | | | | | | | | | | | 0 | -6 | 0 | 0 | 0 | 0 | 0 | | |
| x_{144} | | | | | | | | | | | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | |
| x_{145} | | | | | | | | | | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | -1 | |

| | x_{88} | x_{92} | x_{95} | x_{96} | x_{103} | x_{104} | x_{124} | x_{126} | x_{144} |
|----------|----------|----------|----------|----------|-----------|-----------|-----------|-----------|-----------|
| x_{92} | | | | | | | | | |

A REVIEW GOES HERE – Check our WWW List of Reviews

τ^- BRANCHING RATIOS

$$\Gamma(\text{particle}^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau (\text{"1-prong"})) / \Gamma_{\text{total}} \quad \Gamma_1 / \Gamma$$

$$\Gamma_1 / \Gamma = (\Gamma_3 + \Gamma_5 + \Gamma_{10} + \Gamma_{11} + \Gamma_{14} + \Gamma_{16} + \Gamma_{20} + \Gamma_{23} + \Gamma_{26} + \Gamma_{27} + \Gamma_{29} + 0.6569 \Gamma_{35} + 0.6569 \Gamma_{37} + 0.6569 \Gamma_{40} + 0.6569 \Gamma_{42} + 0.4316 \Gamma_{46} + 0.708 \Gamma_{124} + 0.715 \Gamma_{126} + 0.09 \Gamma_{144} + 0.09 \Gamma_{145}) / \Gamma$$

The charged particle here can be e , μ , or hadron. In many analyses, the sum of the topological branching fractions (1, 3, and 5 prongs) is constrained to be unity. Since the 5-prong fraction is very small, the measured 1-prong and 3-prong fractions are highly correlated and cannot be treated as independent quantities in our overall fit. We arbitrarily choose to use the 3-prong fraction in our fit, and leave the 1-prong fraction out. We do, however, use these 1-prong measurements in our average below. The measurements used only for the average are marked "avg," whereas "f&a" marks a result used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|------|-------------------------------------|------|---------|
| 84.71 ± 0.13 OUR FIT | | Error includes scale factor of 1.2. | | |

85.1 ± 0.4 OUR AVERAGE

| | | | | |
|---|------|-----------------------|----------|---|
| 85.6 ± 0.6 ± 0.3 avg | 3300 | ²⁴ ADEVA | 91F L3 | $E_{\text{cm}}^{\text{ee}} = 88.3\text{--}94.3 \text{ GeV}$ |
| 84.9 ± 0.4 ± 0.3 avg | | BEHREND | 89B CELL | $E_{\text{cm}}^{\text{ee}} = 14\text{--}47 \text{ GeV}$ |
| 84.7 ± 0.8 ± 0.6 avg | | ²⁵ AIHARA | 87B TPC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 86.4 ± 0.3 ± 0.3 | | ABACHI | 89B HRS | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 87.1 ± 1.0 ± 0.7 | | ²⁶ BURCHAT | 87 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 87.2 ± 0.5 ± 0.8 | | SCHMIDKE | 86 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 84.7 ± 1.1 $^{+1.6}_{-1.3}$ | 169 | ²⁷ ALTHOFF | 85 TASS | $E_{\text{cm}}^{\text{ee}} = 34.5 \text{ GeV}$ |
| 86.1 ± 0.5 ± 0.9 | | BARTEL | 85F JADE | $E_{\text{cm}}^{\text{ee}} = 34.6 \text{ GeV}$ |
| 87.8 ± 1.3 ± 3.9 | | ²⁸ BERGER | 85 PLUT | $E_{\text{cm}}^{\text{ee}} = 34.6 \text{ GeV}$ |
| 86.7 ± 0.3 ± 0.6 | | FERNANDEZ | 85 MAC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

²⁴ Not independent of ADEVA 91F $\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \nu_\tau (\text{"3-prong"}) / \Gamma_{\text{total}}$ value.

²⁵ Not independent of AIHARA 87B $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma_{\text{total}}$, $\Gamma(e^- \bar{\nu}_e \nu_\tau) / \Gamma_{\text{total}}$, and $\Gamma(h^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$ values.

²⁶ Not independent of SCHMIDKE 86 value (also not independent of BURCHAT 87 value for $\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \nu_\tau (\text{"3-prong"}) / \Gamma_{\text{total}}$).

²⁷ Not independent of ALTHOFF 85 $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma_{\text{total}}$, $\Gamma(e^- \bar{\nu}_e \nu_\tau) / \Gamma_{\text{total}}$, $\Gamma(h^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}$, and $\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals} \nu_\tau (\text{"3-prong"}) / \Gamma_{\text{total}}$ values.

²⁸ Not independent of (1-prong + 0 π^0) and (1-prong + $\geq 1\pi^0$) values.

$$\Gamma(\text{particle}^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_2 / \Gamma$$

$$\Gamma_2 / \Gamma = (\Gamma_3 + \Gamma_5 + \Gamma_{10} + \Gamma_{11} + \Gamma_{14} + \Gamma_{16} + \Gamma_{20} + \Gamma_{23} + \Gamma_{26} + \Gamma_{27} + \Gamma_{29} + \Gamma_{35} + \Gamma_{37} + \Gamma_{40} + \Gamma_{42} + \Gamma_{46} + 0.708 \Gamma_{124} + 0.715 \Gamma_{126} + 0.09 \Gamma_{144} + 0.09 \Gamma_{145}) / \Gamma$$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------------------------------|------|---------|
| 85.32 ± 0.13 OUR FIT | Error includes scale factor of 1.2. | | |

84.59 ± 0.33 OUR AVERAGE

| | | | |
|---|--------|----------|--------------------|
| 84.48 ± 0.27 ± 0.23 avg | ACTON | 92H OPAL | 1990–1991 LEP runs |
| 85.45 $^{+0.69}_{-0.73}$ ± 0.65 f&a | DECAMP | 92C ALEP | 1989–1990 LEP runs |

$\Gamma(\mu^-\bar{\nu}_\mu\nu_\tau)/\Gamma_{\text{total}}$ Γ_3/Γ

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

To minimize the effect of experiments with large systematic errors, we exclude experiments which together would contribute 5% of the weight in the average.

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| 17.37 ±0.07 OUR FIT | | | | |
| 17.33 ±0.07 OUR AVERAGE | | | | |
| 17.325 ±0.095 ±0.077 | f&a 27.7k | ABREU | 99X DLPH | 1991–1995 LEP runs |
| 17.37 ±0.08 ±0.18 | avg | 29 ANASTASSOV 97 | CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| 17.31 ±0.11 ±0.05 | f&a 20.7k | BUSKULIC | 96C ALEP | 1991–1993 LEP runs |
| 17.36 ±0.27 | f&a 7941 | AKERS | 95I OPAL | 1990–1992 LEP runs |
| 17.6 ±0.4 ±0.4 | f&a 2148 | ADRIANI | 93M L3 | $E_{\text{cm}}^{\text{ee}} = 88–94 \text{ GeV}$ |
| 17.4 ±0.3 ±0.5 | avg | 30 ALBRECHT | 93G ARG | $E_{\text{cm}}^{\text{ee}} = 9.4–10.6 \text{ GeV}$ |
| 17.35 ±0.41 ±0.37 | f&a | DECAMP | 92C ALEP | 1989–1990 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 17.02 ±0.19 ±0.24 | 6586 | ABREU | 95T DLPH | Repl.. by ABREU 99X |
| 17.7 ±0.8 ±0.4 | 568 | BEHREND | 90 CELL | $E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$ |
| 17.4 ±1.0 | 2197 | ADEVA | 88 MRKJ | $E_{\text{cm}}^{\text{ee}} = 14–16 \text{ GeV}$ |
| 17.7 ±1.2 ±0.7 | | AIHARA | 87B TPC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 18.3 ±0.9 ±0.8 | | BURCHAT | 87 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 18.6 ±0.8 ±0.7 | 558 | 31 BARTEL | 86D JADE | $E_{\text{cm}}^{\text{ee}} = 34.6 \text{ GeV}$ |
| 12.9 ±1.7 ±0.7 | | ALTHOFF | 85 TASS | $E_{\text{cm}}^{\text{ee}} = 34.5 \text{ GeV}$ |
| 18.0 ±0.9 ±0.5 | 473 | 31 ASH | 85B MAC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 18.0 ±1.0 ±0.6 | | 32 BALTRUSAIT..85 | MRK3 | $E_{\text{cm}}^{\text{ee}} = 3.77 \text{ GeV}$ |
| 19.4 ±1.6 ±1.7 | 153 | BERGER | 85 PLUT | $E_{\text{cm}}^{\text{ee}} = 34.6 \text{ GeV}$ |
| 17.6 ±2.6 ±2.1 | 47 | BEHREND | 83C CELL | $E_{\text{cm}}^{\text{ee}} = 34 \text{ GeV}$ |
| 17.8 ±2.0 ±1.8 | | BERGER | 81B PLUT | $E_{\text{cm}}^{\text{ee}} = 9–32 \text{ GeV}$ |

29 This ANASTASSOV 97 result is not independent of $\Gamma(\mu^-\bar{\nu}_\mu\nu_\tau)/\Gamma(e^-\bar{\nu}_e\nu_\tau)$ and $\Gamma(e^-\bar{\nu}_e\nu_\tau)/\Gamma_{\text{total}}$ values.

30 Not independent of ALBRECHT 92D $\Gamma(\mu^-\bar{\nu}_\mu\nu_\tau)/\Gamma(e^-\bar{\nu}_e\nu_\tau)$ and ALBRECHT 93G $\Gamma(\mu^-\bar{\nu}_\mu\nu_\tau) \times \Gamma(e^-\bar{\nu}_e\nu_\tau)/\Gamma_{\text{total}}^2$ values.

31 Modified using $B(e^-\bar{\nu}_e\nu_\tau)/B(\text{"1 prong"})$ and $B(\text{"1 prong"}) = 0.855$.

32 Error correlated with BALTRUSAITIS 85 $e\nu\bar{\nu}$ value.

 $\Gamma(\mu^-\bar{\nu}_\mu\nu_\tau\gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| 0.361±0.016±0.035 | | 33 BERGFELD 00 | CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.30 ±0.04 ±0.05 | 116 | 34 ALEXANDER 96S | OPAL | 1991–1994 LEP runs |
| 0.23 ±0.10 | 10 | 35 WU | 90 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

³³ BERGFELD 00 impose requirements on detected γ 's corresponding to a τ -rest-frame energy cutoff $E_{\gamma}^* > 10$ MeV. For $E_{\gamma}^* > 20$ MeV, they quote $(3.04 \pm 0.14 \pm 0.30) \times 10^{-3}$.

³⁴ ALEXANDER 96S impose requirements on detected γ 's corresponding to a τ -rest-frame energy cutoff $E_{\gamma} > 20$ MeV.

³⁵ WU 90 reports $\Gamma(\mu^- \bar{\nu}_{\mu} \nu_{\tau} \gamma) / \Gamma(\mu^- \bar{\nu}_{\mu} \nu_{\tau}) = 0.013 \pm 0.006$, which is converted to $\Gamma(\mu^- \bar{\nu}_{\mu} \nu_{\tau} \gamma) / \Gamma_{\text{total}}$ using $\Gamma(\mu^- \bar{\nu}_{\mu} \nu_{\tau} \gamma) / \Gamma_{\text{total}} = 17.35\%$. Requirements on detected γ 's correspond to a τ rest frame energy cutoff $E_{\gamma} > 37$ MeV.

$\Gamma(e^- \bar{\nu}_e \nu_{\tau}) / \Gamma_{\text{total}}$

Γ_5 / Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

To minimize the effect of experiments with large systematic errors, we exclude experiments which together would contribute 5% of the weight in the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----------|-------------|----------|--|
| 17.83 ± 0.06 OUR FIT | | | | |
| 17.81 ± 0.07 OUR AVERAGE | | | | |
| 17.81 ± 0.09 ± 0.06 | 33.1k | ABBIENDI | 99H OPAL | 1991–1995 LEP runs |
| 17.877 ± 0.109 ± 0.110 | f&a 23.3k | ABREU | 99X DLPH | 1991–1995 LEP runs |
| 17.76 ± 0.06 ± 0.17 | f&a | ANASTASSOV | 97 CLEO | $E_{\text{cm}}^{ee} = 10.6$ GeV |
| 17.79 ± 0.12 ± 0.06 | f&a 20.6k | BUSKULIC | 96C ALEP | 1991–1993 LEP runs |
| 17.9 ± 0.4 ± 0.4 | f&a 2892 | ADRIANI | 93M L3 | $E_{\text{cm}}^{ee} = 88\text{--}94$ GeV |
| 17.5 ± 0.3 ± 0.5 | avg | 36 ALBRECHT | 93G ARG | $E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV |
| 18.09 ± 0.45 ± 0.45 | f&a | | 92C ALEP | 1989–1990 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 17.78 ± 0.10 ± 0.09 | f&a 25.3k | ALEXANDER | 96D OPAL | Repl. by ABBIENDI 99H |
| 17.51 ± 0.23 ± 0.31 | 5059 | ABREU | 95T DLPH | Repl. by ABREU 99X |
| 17.97 ± 0.14 ± 0.23 | 3970 | AKERIB | 92 CLEO | Repl. by ANASTASSOV 97 |
| 19.1 ± 0.4 ± 0.6 | 2960 | 37 AMMAR | 92 CLEO | $E_{\text{cm}}^{ee} = 10.5\text{--}10.9$ GeV |
| 17.0 ± 0.5 ± 0.6 | 1.7k | | 90 HRS | $E_{\text{cm}}^{ee} = 29$ GeV |
| 18.4 ± 0.8 ± 0.4 | 644 | BEHREND | 90 CELL | $E_{\text{cm}}^{ee} = 35$ GeV |
| 16.3 ± 0.3 ± 3.2 | | JANSSEN | 89 CBAL | $E_{\text{cm}}^{ee} = 9.4\text{--}10.6$ GeV |
| 18.4 ± 1.2 ± 1.0 | | AIHARA | 87B TPC | $E_{\text{cm}}^{ee} = 29$ GeV |
| 19.1 ± 0.8 ± 1.1 | | BURCHAT | 87 MRK2 | $E_{\text{cm}}^{ee} = 29$ GeV |
| 16.8 ± 0.7 ± 0.9 | 515 | 37 BARTEL | 86D JADE | $E_{\text{cm}}^{ee} = 34.6$ GeV |
| 20.4 ± 3.0 ± 1.4 | –0.9 | | 85 TASS | $E_{\text{cm}}^{ee} = 34.5$ GeV |
| 17.8 ± 0.9 ± 0.6 | 390 | 37 ASH | 85B MAC | $E_{\text{cm}}^{ee} = 29$ GeV |
| 18.2 ± 0.7 ± 0.5 | | | 85 MRK3 | $E_{\text{cm}}^{ee} = 3.77$ GeV |
| 13.0 ± 1.9 ± 2.9 | | BERGER | 85 PLUT | $E_{\text{cm}}^{ee} = 34.6$ GeV |
| 18.3 ± 2.4 ± 1.9 | 60 | BEHREND | 83C CELL | $E_{\text{cm}}^{ee} = 34$ GeV |
| 16.0 ± 1.3 | 459 | 39 BACINO | 78B DLCO | $E_{\text{cm}}^{ee} = 3.1\text{--}7.4$ GeV |

- ³⁶ Not independent of ALBRECHT 92D $\Gamma(\mu^-\bar{\nu}_\mu\nu_\tau)/\Gamma(e^-\bar{\nu}_e\nu_\tau)$ and ALBRECHT 93G $\Gamma(\mu^-\bar{\nu}_\mu\nu_\tau) \times \Gamma(e^-\bar{\nu}_e\nu_\tau)/\Gamma_{\text{total}}^2$ values.
³⁷ Modified using $B(e^-\bar{\nu}_e\nu_\tau)/B(\text{"1 prong"})$ and $B(\text{"1 prong"}) = 0.855$.
³⁸ Error correlated with BALTRUSAITIS 85 $\Gamma(\mu^-\bar{\nu}_\mu\nu_\tau)/\Gamma_{\text{total}}$.
³⁹ BACINO 78B value comes from fit to events with e^\pm and one other nonelectron charged prong.

| $\Gamma(e^-\bar{\nu}_e\nu_\tau\gamma)/\Gamma_{\text{total}}$ | Γ_6/Γ | | |
|--|-------------------|------|--|
| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
| 1.75 ± 0.06 ± 0.17 | 40 BERGFELD 00 | CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

40 BERGFELD 00 impose requirements on detected γ 's corresponding to a τ -rest-frame energy cutoff $E_\gamma^* > 10 \text{ MeV}$.

| $\Gamma(\mu^-\bar{\nu}_\mu\nu_\tau) \times \Gamma(e^-\bar{\nu}_e\nu_\tau)/\Gamma_{\text{total}}^2$ | $\Gamma_3\Gamma_5/\Gamma^2$ | | | |
|--|-----------------------------|------------------|--|---------|
| VALUE | EVTs | DOCUMENT ID | TECN | COMMENT |
| 0.03097 ± 0.00018 OUR FIT | | | | |
| 0.0306 ± 0.0005 ± 0.0013 | 3230 | ALBRECHT 93G ARG | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.0288 ± 0.0017 ± 0.0019 | ASH | 85B MAC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ | |

| $\Gamma(\mu^-\bar{\nu}_\mu\nu_\tau)/\Gamma(e^-\bar{\nu}_e\nu_\tau)$ | Γ_3/Γ_5 |
|--|---------------------|
| PREDICTED | COMMENT |
| Predicted to be 1 for sequential lepton, 1/2 for para-electron, and 2 for para-muon. | |
| Para-electron also ruled out by HEILE 78. | |

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------------------|------|--|
| 0.974 ± 0.005 OUR FIT | | | |
| 0.978 ± 0.011 OUR AVERAGE | | | |
| 0.9777 ± 0.0063 ± 0.0087 f&a | ANASTASSOV 97 | CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| 0.997 ± 0.035 ± 0.040 f&a | ALBRECHT 92D ARG | | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ |

| $\Gamma(h^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$ | Γ_7/Γ |
|--|-------------------|
| $\Gamma_7/\Gamma = (\Gamma_{10} + \Gamma_{11} + \Gamma_{14} + \Gamma_{16} + \Gamma_{20} + \Gamma_{23} + \Gamma_{26} + \Gamma_{27} + \Gamma_{29} + 0.6569\Gamma_{35} + 0.6569\Gamma_{37} + 0.6569\Gamma_{40} + 0.6569\Gamma_{42} + 0.4316\Gamma_{46} + 0.708\Gamma_{124} + 0.715\Gamma_{126} + 0.09\Gamma_{144} + 0.09\Gamma_{145})/\Gamma$ | |

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------------------------------|------|--|
| 49.51 ± 0.15 OUR FIT | Error includes scale factor of 1.2. | | |
| 48.6 ± 1.2 ± 0.9 avg | 41 AIHARA 87B TPC | | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

41 Not independent of AIHARA 87B $e\nu\bar{\nu}$, $\mu\nu\bar{\nu}$, and $\pi^+\pi^-(\geq 0\pi^0)\nu$ values.

$\Gamma(h^- \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_8/Γ

$$\Gamma_8/\Gamma = (\Gamma_{10} + \Gamma_{11} + \frac{1}{2}\Gamma_{35} + \frac{1}{2}\Gamma_{37} + \frac{1}{4}\Gamma_{46})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|------|-------------|----------|--|
| 12.35±0.12 OUR FIT | Error includes scale factor of 1.4. | | | | |
| 12.43±0.14 OUR AVERAGE | | | | | |
| 12.44±0.11±0.11 | f&a | 15k | 42 BUSKULIC | 96 ALEP | 1991–1993 LEP run |
| 12.47±0.26±0.43 | f&a | 2967 | 43 ACCIARRI | 95 L3 | 1992 LEP run |
| 12.4 ± 0.7 ± 0.7 | f&a | 283 | 44 ABREU | 92N DLPH | 1990 LEP run |
| 11.7 ± 0.6 ± 0.8 | avg | | 45 ALBRECHT | 92D ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |
| 12.98±0.44±0.33 | f&a | | 46 DECOMP | 92C ALEP | 1989–1990 LEP runs |
| 12.1 ± 0.7 ± 0.5 | f&a | 309 | ALEXANDER | 91D OPAL | 1990 LEP run |
| 11.3 ± 0.5 ± 0.8 | avg | 798 | 47 FORD | 87 MAC | $E_{cm}^{ee} = 29 \text{ GeV}$ |
| 12.3 ± 0.6 ± 1.1 | avg | 328 | 48 BARTEL | 86D JADE | $E_{cm}^{ee} = 34.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 12.3 ± 0.9 ± 0.5 | | 1338 | BEHREND | 90 CELL | $E_{cm}^{ee} = 35 \text{ GeV}$ |
| 11.1 ± 1.1 ± 1.4 | | | 49 BURCHAT | 87 MRK2 | $E_{cm}^{ee} = 29 \text{ GeV}$ |
| 13.0 ± 2.0 ± 4.0 | | | BERGER | 85 PLUT | $E_{cm}^{ee} = 34.6 \text{ GeV}$ |
| 11.2 ± 1.7 ± 1.2 | | 34 | 50 BEHREND | 83C CELL | $E_{cm}^{ee} = 34 \text{ GeV}$ |

42 BUSKULIC 96 quote $11.78 \pm 0.11 \pm 0.13$ We add 0.66 to undo their correction for unseen K_L^0 and modify the systematic error accordingly.

43 ACCIARRI 95 with 0.65% added to remove their correction for $\pi^- K_L^0$ backgrounds.

44 ABREU 92N with 0.5% added to remove their correction for $K^*(892)^-$ backgrounds.

45 Not independent of ALBRECHT 92D $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$, $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \times \Gamma(e^- \bar{\nu}_e \nu_\tau)$, and $\Gamma(h^- \geq 0 K_L^0 \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ values.

46 DECOMP 92C quote $B(h^- \geq 0 K_S^0 \geq 0 (K_S^0 \rightarrow \pi^+ \pi^-) \nu_\tau) = 13.32 \pm 0.44 \pm 0.33$.

We subtract 0.35 to correct for their inclusion of the K_S^0 decays.

47 FORD 87 result for $B(\pi^- \nu_\tau)$ with 0.67% added to remove their K^- correction and adjusted for 1992 $B(\text{"1 prong"})$.

48 BARTEL 86D result for $B(\pi^- \nu_\tau)$ with 0.59% added to remove their K^- correction and adjusted for 1992 $B(\text{"1 prong"})$.

49 BURCHAT 87 with 1.1% added to remove their correction for K^- and $K^*(892)^-$ backgrounds.

50 BEHREND 83C quote $B(\pi^- \nu_\tau) = 9.9 \pm 1.7 \pm 1.3$ after subtracting 1.3 ± 0.5 to correct for $B(K^- \nu_\tau)$.

 $\Gamma(h^- \geq 0 K_L^0 \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ Γ_8/Γ_5

$$\Gamma_8/\Gamma_5 = (\Gamma_{10} + \Gamma_{11} + \frac{1}{2}\Gamma_{35} + \frac{1}{2}\Gamma_{37} + \frac{1}{4}\Gamma_{46})/\Gamma_5$$

| VALUE | | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|-------------|--|---------|
| 0.692±0.007 OUR FIT | Error includes scale factor of 1.3. | | | |
| 0.678±0.037±0.044 | ALBRECHT | 92D ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |

0.647±0.039±0.061 51 BARTEL 86D JADE $E_{cm}^{ee} = 34.6 \text{ GeV}$

51 Combined result of BARTEL 86D $e\nu\bar{\nu}$, $\mu\nu\bar{\nu}$, and $\pi^-\nu$ assuming $B(\mu\nu\bar{\nu})/B(e\nu\bar{\nu}) = 0.973$.

$\Gamma(h^-\nu_\tau)/\Gamma_{\text{total}}$

$\Gamma_9/\Gamma = (\Gamma_{10} + \Gamma_{11})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------------------------------|---------------------|--|---------|
| 11.79 ± 0.12 OUR FIT | Error includes scale factor of 1.4. | | | |
| 11.65 ± 0.21 OUR AVERAGE | Error includes scale factor of 1.9. | | | |
| 11.98 ± 0.13 ± 0.16 | f&a | ACKERSTAFF 98M OPAL | 1991–1995 LEP runs | |
| 11.52 ± 0.05 ± 0.12 | f&a | ANASTASSOV 97 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ | |

$\Gamma(h^-\nu_\tau)/\Gamma(e^-\bar{\nu}_e\nu_\tau)$

$\Gamma_9/\Gamma_5 = (\Gamma_{10} + \Gamma_{11})/\Gamma_5$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE | | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------|-------------------------------------|--|------|---------|
| 0.661 ± 0.007 OUR FIT | Error includes scale factor of 1.4. | | | |
| 0.6484 ± 0.0041 ± 0.0060 avg | 52 ANASTASSOV 97 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ | | |

52 Not independent of ANASTASSOV 97 $\Gamma(h^-\nu_\tau)/\Gamma_{\text{total}}$ value.

$\Gamma(\pi^-\nu_\tau)/\Gamma_{\text{total}}$

Γ_{10}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------------------------------|------|-------------|------|---------|
| 11.09 ± 0.12 OUR FIT | Error includes scale factor of 1.4. | | | | |
| 11.07 ± 0.18 OUR AVERAGE | | | | | |

| | | | | |
|---------------------|-----|--------------|----------|---|
| 11.06 ± 0.11 ± 0.14 | avg | 53 BUSKULIC | 96 ALEP | LEP 1991–1993 data |
| 11.7 ± 0.4 ± 1.8 | f&a | 1138 BLOCKER | 82D MRK2 | $E_{\text{cm}}^{\text{ee}} = 3.5\text{--}6.7 \text{ GeV}$ |

53 Not independent of BUSKULIC 96 $B(h^-\nu_\tau)$ and $B(K^-\nu_\tau)$ values.

$\Gamma(K^-\nu_\tau)/\Gamma_{\text{total}}$

Γ_{11}/Γ

| VALUE (%) | | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------|-------------|---|---------|
| 0.699 ± 0.027 OUR FIT | | | | | |
| 0.697 ± 0.027 OUR AVERAGE | | | | | |
| 0.696 ± 0.025 ± 0.014 | 2032 | BARATE | 99K ALEP | 1991–1995 LEP runs | |
| 0.85 ± 0.18 | 27 | ABREU | 94K DLPH | LEP 1992 Z data | |
| 0.66 ± 0.07 ± 0.09 | 99 | BATTLE | 94 CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ | |
| 0.59 ± 0.18 | 16 | MILLS | 84 DLCO | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ | |
| 1.3 ± 0.5 | 15 | BLOCKER | 82B MRK2 | $E_{\text{cm}}^{\text{ee}} = 3.9\text{--}6.7 \text{ GeV}$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 0.72 ± 0.04 ± 0.04 | 728 | BUSKULIC | 96 ALEP | Repl. by BARATE 99K | |

$\Gamma(h^- \geq 1 \text{ neutrals} \nu_\tau)/\Gamma_{\text{total}}$ Γ_{12}/Γ

$$\Gamma_{12}/\Gamma = (\Gamma_{14} + \Gamma_{16} + \Gamma_{20} + \Gamma_{23} + \Gamma_{26} + \Gamma_{27} + \Gamma_{29} + 0.157\Gamma_{35} + 0.157\Gamma_{37} + 0.157\Gamma_{40} + 0.157\Gamma_{42} + 0.0246\Gamma_{46} + 0.708\Gamma_{124} + 0.715\Gamma_{126} + 0.09\Gamma_{144} + 0.09\Gamma_{145})/\Gamma$$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|-----------|-------------|------|---------|
|-----------|-------------|------|---------|

36.88±0.17 OUR FIT Error includes scale factor of 1.2.

36.14±0.33±0.58 AKERS 94E OPAL 1991–1992 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

38.4 ± 1.2 ± 1.0 54 BURCHAT 87 MRK2 $E_{\text{cm}}^{\text{ee}} = 29$ GeV

42.7 ± 2.0 ± 2.9 BERGER 85 PLUT $E_{\text{cm}}^{\text{ee}} = 34.6$ GeV

54 BURCHAT 87 quote for $B(\pi^\pm \geq 1 \text{ neutral} \nu_\tau) = 0.378 \pm 0.012 \pm 0.010$. We add 0.006 to account for contribution from $(K^{*-} \nu_\tau)$ which they fixed at BR = 0.013.

 $\Gamma(h^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{13}/\Gamma = (\Gamma_{14} + \Gamma_{16})/\Gamma$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

25.86±0.14 OUR FIT Error includes scale factor of 1.1.

25.76±0.15 OUR AVERAGE

25.89 ± 0.17 ± 0.29 ACKERSTAFF 98M OPAL 1991–1995 LEP runs

25.76 ± 0.15 ± 0.13 31k BUSKULIC 96 ALEP LEP 1991–1993 data

25.05 ± 0.35 ± 0.50 6613 ACCIARRI 95 L3 1992 LEP run

25.87 ± 0.12 ± 0.42 51k ARTUSO 94 CLEO $E_{\text{cm}}^{\text{ee}} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

25.98 ± 0.36 ± 0.52 56 AKERS 94E OPAL Repl. by ACKER-STAFF 98M

22.9 ± 0.8 ± 1.3 283 57 ABREU 92N DLPH $E_{\text{cm}}^{\text{ee}} = 88.2\text{--}94.2$ GeV

23.1 ± 0.4 ± 0.9 1249 58 ALBRECHT 92Q ARG $E_{\text{cm}}^{\text{ee}} = 10$ GeV

25.02 ± 0.64 ± 0.88 1849 DECAMP 92C ALEP 1989–1990 LEP runs

22.0 ± 0.8 ± 1.9 779 ANTREASYAN 91 CBAL $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6$ GeV

22.6 ± 1.5 ± 0.7 1101 BEHREND 90 CELL $E_{\text{cm}}^{\text{ee}} = 35$ GeV

23.1 ± 1.9 ± 1.6 BEHREND 84 CELL $E_{\text{cm}}^{\text{ee}} = 14.22$ GeV

55 ARTUSO 94 reports the combined result from three independent methods, one of which (23% of the $\tau^- \rightarrow h^- \pi^0 \nu_\tau$) is normalized to the inclusive one-prong branching fraction, taken as 0.854 ± 0.004 . Renormalization to the present value causes negligible change.

56 AKERS 94E quote $(26.25 \pm 0.36 \pm 0.52) \times 10^{-2}$; we subtract 0.27% from their number to correct for $\tau^- \rightarrow h^- K_L^0 \nu_\tau$.

57 ABREU 92N with 0.5% added to remove their correction for $K^*(892)^-$ backgrounds.

58 ALBRECHT 92Q with 0.5% added to remove their correction for $\tau^- \rightarrow K^*(892)^- \nu_\tau$ background.

 $\Gamma(\pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{14}/Γ

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

25.40±0.14 OUR FIT Error includes scale factor of 1.1.

25.31±0.18 OUR AVERAGE

25.30 ± 0.15 ± 0.13 avg 59 BUSKULIC 96 ALEP LEP 1991–1993 data

25.36 ± 0.44 avg 60 ARTUSO 94 CLEO $E_{\text{cm}}^{\text{ee}} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------|------|---------------------------|----------|--|
| $21.5 \pm 0.4 \pm 1.9$ | 4400 | ^{61,62} ALBRECHT | 88L ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $23.0 \pm 1.3 \pm 1.7$ | 582 | ADLER | 87B MRK3 | $E_{\text{cm}}^{\text{ee}} = 3.77 \text{ GeV}$ |
| $25.8 \pm 1.7 \pm 2.5$ | | ⁶³ BURCHAT | 87 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| $22.3 \pm 0.6 \pm 1.4$ | 629 | ⁶² YELTON | 86 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

⁵⁹ Not independent of BUSKULIC 96 $B(h^- \pi^0 \nu_\tau)$ and $B(K^- \pi^0 \nu_\tau)$ values.

⁶⁰ Not independent of ARTUSO 94 $B(h^- \pi^0 \nu_\tau)$ and BATTLE 94 $B(K^- \pi^0 \nu_\tau)$ values.

⁶¹ The authors divide by $(\Gamma_3 + \Gamma_5 + \Gamma_{10} + \Gamma_{11})/\Gamma = 0.467$ to obtain this result.

⁶² Experiment had no hadron identification. Kaon corrections were made, but insufficient information is given to permit their removal.

⁶³ BURCHAT 87 value is not independent of YELTON 86 value. Nonresonant decays included.

$\Gamma(\pi^- \pi^0 \text{non-}\rho(770)\nu_\tau)/\Gamma_{\text{total}}$

Γ_{15}/Γ

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|---------|---|
| 0.3 ± 0.1 ± 0.3 | 64 BEHREND | 84 CELL | $E_{\text{cm}}^{\text{ee}} = 14,22 \text{ GeV}$ |

⁶⁴ BEHREND 84 assume a flat nonresonant mass distribution down to the $\rho(770)$ mass, using events with mass above 1300 to set the level.

$\Gamma(K^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$

Γ_{16}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

0.454±0.033 OUR FIT

0.449±0.034 OUR AVERAGE

$0.444 \pm 0.026 \pm 0.024$ 923 BARATE 99K ALEP 1991–1995 LEP runs

$0.51 \pm 0.10 \pm 0.07$ 37 BATTLE 94 CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.52 \pm 0.04 \pm 0.05$ 395 BUSKULIC 96 ALEP Repl. by BARATE 99K

$\Gamma(h^- \geq 2\pi^0 \nu_\tau)/\Gamma_{\text{total}}$

Γ_{17}/Γ

$$\Gamma_{17}/\Gamma = (\Gamma_{20} + \Gamma_{23} + \Gamma_{26} + \Gamma_{27} + \Gamma_{29} + 0.157\Gamma_{35} + 0.157\Gamma_{37} + 0.157\Gamma_{40} + 0.157\Gamma_{42} + 0.0246\Gamma_{46} + 0.319\Gamma_{124} + 0.322\Gamma_{126})/\Gamma$$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

10.73±0.16 OUR FIT Error includes scale factor of 1.2.

10.0 ± 0.4 OUR AVERAGE

$9.91 \pm 0.31 \pm 0.27$ f&a ACKERSTAFF 98M OPAL 1991–1995 LEP runs

$12.0 \pm 1.4 \pm 2.5$ f&a ⁶⁵ BURCHAT 87 MRK2 $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.89 \pm 0.34 \pm 0.55$ ⁶⁶ AKERS 94E OPAL Repl. by ACKER-STAFF 98M

$14.0 \pm 1.2 \pm 0.6$ 938 ⁶⁷ BEHREND 90 CELL $E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$

$13.9 \pm 2.0 \begin{matrix} +1.9 \\ -2.2 \end{matrix}$ ⁶⁸ AIHARA 86E TPC $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

⁶⁵ Error correlated with BURCHAT 87 $\Gamma(\rho^- \nu_e)/\Gamma(\text{total})$ value.

⁶⁶ AKERS 94E not independent of AKERS 94E $B(h^- \geq 1\pi^0 \nu_\tau)$ and $B(h^- \pi^0 \nu_\tau)$ measurements.

⁶⁷ No independent of BEHREND 90 $\Gamma(h^- 2\pi^0 \nu_\tau \text{ (exp. } K^0\text{)})$ and $\Gamma(h^- \geq 3\pi^0 \nu_\tau)$.

⁶⁸ AIHARA 86E (TPC) quote $B(2\pi^0 \pi^- \nu_\tau) + 1.6B(3\pi^0 \pi^- \nu_\tau) + 1.1B(\pi^0 \eta \pi^- \nu_\tau)$.

$$\Gamma(h^- 2\pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{18} / \Gamma = (\Gamma_{20} + \Gamma_{23} + 0.157\Gamma_{35} + 0.157\Gamma_{37}) / \Gamma$$

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|--|
| 9.36±0.14 OUR FIT | | | | Error includes scale factor of 1.2. |
| 9.48±0.13±0.10 | 12k | 69 BUSKULIC | 96 ALEP | LEP 1991–1993 data 69 BUSKULIC 96 quote $9.29 \pm 0.13 \pm 0.10$. We add 0.19 to undo their correction for $\tau^- \rightarrow h^- K^0 \nu_\tau$. |

$$\Gamma(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{19} / \Gamma = (\Gamma_{20} + \Gamma_{23}) / \Gamma$$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. f&a marks results used for the fit and the average.

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| 9.19±0.14 OUR FIT | | | | Error includes scale factor of 1.2. |
| 8.9 ±0.4 OUR AVERAGE | | | | Error includes scale factor of 1.1. |
| 8.88±0.37±0.42 f&a | 1060 | ACCIARRI | 95 L3 | 1992 LEP run |
| 8.96±0.16±0.44 avg | | 70 PROCARIO | 93 CLEO | $E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$ |
| 10.38±0.66±0.82 f&a | 809 | 71 DECOMP | 92C ALEP | 1989–1990 LEP runs |
| 5.7 ±0.5 ^{+1.7} _{-1.0} f&a | 133 | 72 ANTREASYAN | 91 CBAL | $E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |
| 8.7 ±0.4 ±1.1 f&a | 815 | 73 BAND | 87 MAC | $E_{\text{cm}}^{ee} = 29 \text{ GeV}$ |
| 6.0 ±3.0 ±1.8 f&a | | BEHREND | 84 CELL | $E_{\text{cm}}^{ee} = 14,22 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|----------------|-----|------------|---------|---------------------------------------|
| 10.0 ±1.5 ±1.1 | 333 | 74 BEHREND | 90 CELL | $E_{\text{cm}}^{ee} = 35 \text{ GeV}$ |
| 6.2 ±0.6 ±1.2 | | 75 GAN | 87 MRK2 | $E_{\text{cm}}^{ee} = 29 \text{ GeV}$ |

70 PROCARIO 93 entry is obtained from $B(h^- 2\pi^0 \nu_\tau) / B(h^- \pi^0 \nu_\tau)$ using ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$.

71 We subtract 0.0015 to account for $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.

72 ANTREASYAN 91 subtract 0.001 to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.

73 BAND 87 assume $B(\pi^- 3\pi^0 \nu_\tau) = 0.01$ and $B(\pi^- \pi^0 \eta \nu_\tau) = 0.005$.

74 BEHREND 90 subtract 0.002 to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.

75 GAN 87 analysis use photon multiplicity distribution.

$$\Gamma(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma(h^- \pi^0 \nu_\tau) \quad \Gamma_{19} / \Gamma_{13} = (\Gamma_{20} + \Gamma_{23}) / (\Gamma_{14} + \Gamma_{16})$$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|--------------------|-------------|---|
| 0.355±0.006 OUR FIT | | | Error includes scale factor of 1.2. |
| 0.342±0.006±0.016 | 76 PROCARIO | 93 CLEO | $E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$ |

76 PROCARIO 93 quote $0.345 \pm 0.006 \pm 0.016$ after correction for 2 kaon backgrounds assuming $B(K^* \nu_\tau) = 1.42 \pm 0.18\%$ and $B(h^- K^0 \pi^0 \nu_\tau) = 0.48 \pm 0.48\%$. We multiply by 0.990 ± 0.010 to remove these corrections to $B(h^- \pi^0 \nu_\tau)$.

$\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$ Γ_{20} / Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------------------------|-------------|------|---------|
| 9.13±0.14 OUR FIT | Error includes scale factor of 1.2. | | | |

9.21±0.13±0.11 **avg** 77 BUSKULIC 96 ALEP LEP 1991–1993 data

77 Not independent of BUSKULIC 96 $B(h^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ and $B(K^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ values.

 $\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0), \text{scalar}) / \Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ $\Gamma_{21} / \Gamma_{20}$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|-------------|---------|--|
| <0.094 | 95 | 78 BROWDER | 00 CLEO | $4.7 \text{ fb}^{-1} E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

78 Model-independent limit from structure function analysis on contribution to $B(\tau^- \rightarrow \pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ from scalars.

 $\Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0), \text{vector}) / \Gamma(\pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ $\Gamma_{22} / \Gamma_{20}$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|-------------|---------|--|
| <0.073 | 95 | 79 BROWDER | 00 CLEO | $4.7 \text{ fb}^{-1} E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

79 Model-independent limit from structure function analysis on contribution to $B(\tau^- \rightarrow \pi^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$ from vectors.

 $\Gamma(K^- 2\pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$ Γ_{23} / Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|---------|
| 0.060±0.024 OUR FIT | | | | |

0.058±0.024 OUR AVERAGE

0.056±0.020±0.015 131 BARATE 99K ALEP 1991–1995 LEP runs
 0.09 ± 0.10 ± 0.03 3 80 BATTLE 94 CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.08 ± 0.02 ± 0.02 59 BUSKULIC 96 ALEP Repl. by BARATE 99K

80 BATTLE 94 quote $0.14 \pm 0.10 \pm 0.03$ or $< 0.3\%$ at 90% CL. We subtract $(0.05 \pm 0.02)\%$ to account for $\tau^- \rightarrow K^- (K^0 \rightarrow \pi^0 \pi^0) \nu_\tau$ background.

 $\Gamma(h^- \geq 3\pi^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_{24} / Γ

$$\Gamma_{24} / \Gamma = (\Gamma_{26} + \Gamma_{27} + \Gamma_{29} + 0.157\Gamma_{40} + 0.157\Gamma_{42} + 0.0246\Gamma_{46} + 0.319\Gamma_{124} + 0.322\Gamma_{126}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------------------------|-------------|------|---------|
| 1.37±0.11 OUR FIT | Error includes scale factor of 1.1. | | | |

1.53±0.40±0.46 **f&a** 186 DECAMP 92C ALEP 1989–1990 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.2 ± 1.0 ± 1.0 BEHREND 90 CELL $E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$

$$\Gamma(h^- 3\pi^0 \nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{25}/\Gamma$$

$$\Gamma_{25}/\Gamma = (\Gamma_{26} + \Gamma_{27} + 0.157\Gamma_{40} + 0.157\Gamma_{42} + 0.322\Gamma_{126})/\Gamma$$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

| VALUE (%) | | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|--|------|-------------|------|---------|
|-----------|--|------|-------------|------|---------|

1.21±0.10 OUR FIT Error includes scale factor of 1.1.

1.22±0.10 OUR AVERAGE

| | | | | | |
|----------------|-----|------|-------------|---------|--|
| 1.24±0.09±0.11 | f&a | 2.3k | 81 BUSKULIC | 96 ALEP | LEP 1991–1993 data |
| 1.70±0.24±0.38 | f&a | 293 | ACCIARRI | 95 L3 | 1992 LEP run |
| 1.15±0.08±0.13 | avg | | 82 PROCARIO | 93 CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----|------|------|--------|---------|--|
| 0.0 | +1.4 | +1.1 | 83 GAN | 87 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
|-----|------|------|--------|---------|--|

81 BUSKULIC 96 quote $B(h^- 3\pi^0 \nu_\tau \text{ (ex. } K^0)) = 1.17 \pm 0.09 \pm 0.11$. We add 0.07 to remove their correction for K^0 backgrounds.

82 PROCARIO 93 entry is obtained from $B(h^- 3\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau)$ using ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$.

83 Highly correlated with GAN 87 $\Gamma(\eta \pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ value. Authors quote $B(\pi^\pm 3\pi^0 \nu_\tau) + 0.67B(\pi^\pm \eta \pi^0 \nu_\tau) = 0.047 \pm 0.010 \pm 0.011$.

$$\Gamma(h^- 3\pi^0 \nu_\tau)/\Gamma(h^- \pi^0 \nu_\tau) \quad \Gamma_{25}/\Gamma_{13}$$

$$\Gamma_{25}/\Gamma_{13} = (\Gamma_{26} + \Gamma_{27} + 0.157\Gamma_{40} + 0.157\Gamma_{42} + 0.322\Gamma_{126})/(\Gamma_{14} + \Gamma_{16})$$

| VALUE | | DOCUMENT ID | TECN | COMMENT |
|-------|--|-------------|------|---------|
|-------|--|-------------|------|---------|

0.047±0.004 OUR FIT Error includes scale factor of 1.1.

| | | | |
|--------------------------|-------------|---------|--|
| 0.044±0.003±0.005 | 84 PROCARIO | 93 CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |
|--------------------------|-------------|---------|--|

84 PROCARIO 93 quote $0.041 \pm 0.003 \pm 0.005$ after correction for 2 kaon backgrounds assuming $B(K^{*-} \nu_\tau) = 1.42 \pm 0.18\%$ and $B(h^- K^0 \pi^0 \nu_\tau) = 0.48 \pm 0.48\%$. We add 0.003 ± 0.003 and multiply the sum by 0.990 ± 0.010 to remove these corrections.

$$\Gamma(\pi^- 3\pi^0 \nu_\tau \text{ (ex. } K^0))/\Gamma_{\text{total}} \quad \Gamma_{26}/\Gamma$$

| VALUE (%) | | DOCUMENT ID |
|-----------|--|-------------|
|-----------|--|-------------|

1.08±0.10 OUR FIT Error includes scale factor of 1.1.

$$\Gamma(K^- 3\pi^0 \nu_\tau \text{ (ex. } K^0, \eta))/\Gamma_{\text{total}} \quad \Gamma_{27}/\Gamma$$

| VALUE (%) | | DOCUMENT ID | TECN | COMMENT |
|-----------|--|-------------|------|---------|
|-----------|--|-------------|------|---------|

**0.039+0.023
-0.021 OUR FIT**

0.037±0.021±0.011 22 BARATE 99K ALEP 1991–1995 LEP runs

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------------|-------------|----------|---------------------|
| 0.05 ± 0.13 | 85 BUSKULIC | 94E ALEP | Repl. by BARATE 99K |
|-------------|-------------|----------|---------------------|

85 BUSKULIC 94E quote $B(K^- \geq 0 \pi^0 \geq 0 K^0 \nu_\tau) = [B(K^- \nu_\tau) + B(K^- \pi^0 \nu_\tau) + B(K^- K^0 \nu_\tau) + B(K^- \pi^0 \pi^0 \nu_\tau) + B(K^- \pi^0 K^0 \nu_\tau)] = 0.05 \pm 0.13\%$ accounting for common systematic errors in BUSKULIC 94E and BUSKULIC 94F measurements of these modes. We assume $B(K^- \geq 2K^0 \nu_\tau)$ and $B(K^- \geq 4\pi^0 \nu_\tau)$ are negligible.

$$\Gamma(h^- 4\pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{28}/\Gamma = (\Gamma_{29} + 0.319\Gamma_{124})/\Gamma$$

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|-------------|--------------------|-------------|----------------|
|------------------|-------------|--------------------|-------------|----------------|

0.16±0.06 OUR FIT**0.16±0.06 OUR AVERAGE**

0.16±0.04±0.09 232 86 BUSKULIC 96 ALEP LEP 1991–1993 data

0.16±0.05±0.05 87 PROCARIO 93 CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$

86 BUSKULIC 96 quote result for $\tau^- \rightarrow h^- \geq 4\pi^0 \nu_\tau$. We assume $B(h^- \geq 5\pi^0 \nu_\tau)$ is negligible.

87 PROCARIO 93 quotes $B(h^- 4\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau) = 0.006 \pm 0.002 \pm 0.002$. We multiply by the ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$ to obtain $B(h^- 4\pi^0 \nu_\tau)$. PROCARIO 93 assume $B(h^- \geq 5 \pi^0 \nu_\tau)$ is small and do not correct for it.

$$\Gamma(h^- 4\pi^0 \nu_\tau (\text{ex. } K^0, \eta)) / \Gamma_{\text{total}} \quad \Gamma_{29}/\Gamma$$

| <u>VALUE (%)</u> | <u>DOCUMENT ID</u> |
|------------------|--------------------|
|------------------|--------------------|

0.10^{+0.06}_{-0.05} OUR FIT

$$\Gamma(K^- \geq 0\pi^0 \geq 0 K^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{30}/\Gamma$$

$\Gamma_{30}/\Gamma = (\Gamma_{11} + \Gamma_{16} + \Gamma_{23} + \Gamma_{27} + \Gamma_{37} + \Gamma_{42} + 0.715\Gamma_{126})/\Gamma$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|-------------|--------------------|-------------|----------------|
|------------------|-------------|--------------------|-------------|----------------|

1.58 ±0.06 OUR FIT**1.54 ±0.05 OUR AVERAGE**

1.520±0.040±0.041 avg 4006 88 BARATE 99K ALEP 1991–1995 LEP runs

1.54 ±0.24 f&a ABREU 94K DLPH LEP 1992 Z data

1.70 ±0.12 ±0.19 f&a 202 89 BATTLE 94 CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ 1.6 ±0.4 ±0.2 f&a 35 AIHARA 87B TPC $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ 1.71 ±0.29 f&a 53 MILLS 84 DLCO $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.70 ±0.05 ±0.06 1610 90 BUSKULIC 96 ALEP Repl. by
BARATE 99K

88 Not independent of BARATE 99K $B(K^- \nu_\tau)$, $B(K^- \pi^0 \nu_\tau)$, $B(K^- 2\pi^0 \nu_\tau (\text{ex. } K^0))$, $B(K^- 3\pi^0 \nu_\tau (\text{ex. } K^0))$, $B(K^- K^0 \nu_\tau)$, and $B(K^- K^0 \pi^0 \nu_\tau)$ values.

89 BATTLE 94 quote $1.60 \pm 0.12 \pm 0.19$. We add 0.10 ± 0.02 to correct for their rejection of $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

90 Not independent of BUSKULIC 96 $B(K^- \nu_\tau)$, $B(K^- \pi^0 \nu_\tau)$, $B(K^- 2\pi^0 \nu_\tau)$, $B(K^- K^0 \nu_\tau)$, and $B(K^- K^0 \pi^0 \nu_\tau)$ values.

$$\Gamma(K^- \geq 1 (\pi^0 \text{ or } K^0) \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{31}/\Gamma$$

$$\Gamma_{31}/\Gamma = (\Gamma_{16} + \Gamma_{23} + \Gamma_{27} + \Gamma_{37} + \Gamma_{42} + 0.715\Gamma_{126})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------------|----------------------|---------------------|---|
| 0.88 ± 0.05 OUR FIT | | | | |
| 0.76 ± 0.23 OUR AVERAGE | | | | |
| 0.69 ± 0.25 1.2 ± 0.5 ^{+0.2} _{-0.4} | avg f&a | 91 ABREU 9 AIHARA | 94K DLPH 87B TPC | LEP 1992 Z data $E_{\text{cm}}^{\text{ee}} = 29$ GeV |

⁹¹ Not independent of ABREU 94K $B(K^- \nu_\tau)$ and $B(K^- \geq 0 \text{ neutrals} \nu_\tau)$ measurements.

$$\Gamma(K^0(\text{particles})^- \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{32}/\Gamma$$

$$\Gamma_{32}/\Gamma = (\Gamma_{35} + \Gamma_{37} + \Gamma_{40} + \Gamma_{42} + \Gamma_{46})/\Gamma$$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------------------|-----------------------|----------------------|---|
| 1.71 ± 0.06 OUR FIT | Error includes scale factor of 1.1. | | | |
| 1.94 ± 0.13 OUR AVERAGE | | | | |
| 1.94 ± 0.12 ± 0.12 1.94 ± 0.18 ± 0.12 | 929 141 | 92 BARATE 93 AKERS | 98E ALEP 94G OPAL | 1991–1995 LEP runs $E_{\text{cm}}^{\text{ee}} = 88–94$ GeV |

⁹² BARATE 98E measure $\Gamma(K_S^0(\text{particles})^- \nu_\tau) / \Gamma_{\text{total}} = (0.970 \pm 0.058 \pm 0.062)\%$. We multiply this by 2 to obtain the listed value.

⁹³ AKERS 94G measure $\Gamma(K_S^0(\text{particles})^- \nu_\tau) / \Gamma_{\text{total}} = 0.97 \pm 0.09 \pm 0.06$.

$$\Gamma(h^- \bar{K}^0 \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{33}/\Gamma$$

$$\Gamma_{33}/\Gamma = (\Gamma_{35} + \Gamma_{37} + \Gamma_{40} + \Gamma_{42} + 0.657\Gamma_{46})/\Gamma$$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------------------------------|-------------|--------|--------------------------------------|
| 1.67 ± 0.06 OUR FIT | Error includes scale factor of 1.1. | | | |
| 1.3 ± 0.3 | 44 | TSCHIRHART | 88 HRS | $E_{\text{cm}}^{\text{ee}} = 29$ GeV |

$$\Gamma(h^- \bar{K}^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{34}/\Gamma = (\Gamma_{35} + \Gamma_{37})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|-------------|-------------------|--|
| 1.06 ± 0.05 OUR FIT | Error includes scale factor of 1.2. | | | |
| 0.90 ± 0.07 OUR AVERAGE | | | | |
| 1.01 ± 0.11 ± 0.07 0.855 ± 0.036 ± 0.073 | avg f&a | 555 1242 | 94 BARATE COAN | 98E ALEP 96 CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6$ GeV |

⁹⁴ Not independent of BARATE 98E $B(\tau^- \rightarrow \pi^- \bar{K}^0 \nu_\tau)$ and $B(\tau^- \rightarrow K^- K^0 \nu_\tau)$ values.

$\Gamma(\pi^- \bar{K}^0 \nu_\tau)/\Gamma_{\text{total}}$

Γ_{35}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------------------|-----------|-------------|--------------------------------|---------|
| 0.90 ± 0.04 OUR FIT | Error includes scale factor of 1.1. | | | | |
| 0.88 ± 0.05 OUR AVERAGE | Error includes scale factor of 1.2. | | | | |
| 0.933 ± 0.068 ± 0.049 | 377 | ABBIENDI | 00C OPAL | 1991–1995 LEP runs | |
| 0.928 ± 0.045 ± 0.034 | f&a 937 | 95 BARATE | 99K ALEP | 1991–1995 LEP runs | |
| 0.855 ± 0.117 ± 0.066 | avg 509 | 96 BARATE | 98E ALEP | 1991–1995 LEP runs | |
| 0.704 ± 0.041 ± 0.072 | avg 97 COAN | | 96 CLEO | $E_{cm}^{ee} \approx 10.6$ GeV | |
| 0.95 ± 0.15 ± 0.06 | f&a 98 ACCIARRI | | 95F L3 | 1991–1993 LEP runs | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 0.79 ± 0.10 ± 0.09 | 98 BUSKULIC | 99 | 96 ALEP | Repl. by BARATE 99K | |

95 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

96 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays. Not independent of BARATE 98E $B(K^0 \text{ particles} - \nu_\tau)$ value.

97 Not independent of COAN 96 $B(h^- K^0 \nu_\tau)$ and $B(K^- K^0 \nu_\tau)$ measurements.

98 ACCIARRI 95F do not identify π^- / K^- and assume $B(K^- K^0 \nu_\tau) = (0.29 \pm 0.12)\%$.

99 BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

$\Gamma(\pi^- \bar{K}^0 (\text{non-}K^*(892)^-) \nu_\tau)/\Gamma_{\text{total}}$

Γ_{36}/Γ

| VALUE (%) | CL% | | DOCUMENT ID | TECN | COMMENT |
|-----------|-----|--|-------------|--------|--------------------|
| <0.17 | 95 | | ACCIARRI | 95F L3 | 1991–1993 LEP runs |

$\Gamma(K^- K^0 \nu_\tau)/\Gamma_{\text{total}}$

Γ_{37}/Γ

| VALUE (%) | | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|--------------|-------------|--------------------------------|---------|
| 0.155 ± 0.017 OUR FIT | | | | | |
| 0.158 ± 0.017 OUR AVERAGE | | | | | |
| 0.162 ± 0.021 ± 0.011 | 150 | 100 BARATE | 99K ALEP | 1991–1995 LEP runs | |
| 0.158 ± 0.042 ± 0.017 | 46 | 101 BARATE | 98E ALEP | 1991–1995 LEP runs | |
| 0.151 ± 0.021 ± 0.022 | 111 | COAN | 96 CLEO | $E_{cm}^{ee} \approx 10.6$ GeV | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 0.26 ± 0.09 ± 0.02 | 13 | 102 BUSKULIC | 96 ALEP | Repl. by BARATE 99K | |
| 100 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter. | | | | | |
| 101 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays. | | | | | |
| 102 BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter. | | | | | |

$\Gamma(K^- \bar{K}^0 \geq 0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$

$\Gamma_{38}/\Gamma = (\Gamma_{37} + \Gamma_{42})/\Gamma$

| VALUE (%) | EVTS | | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|----------|-------------|--------------------|---------|
| 0.312 ± 0.025 OUR FIT | | | | | |
| 0.330 ± 0.055 ± 0.039 | 124 | ABBIENDI | 00C OPAL | 1991–1995 LEP runs | |

$\Gamma(h^-\bar{K}^0\pi^0\nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{39}/\Gamma = (\Gamma_{40} + \Gamma_{42})/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------------------|----------|--|
| 0.53 ±0.04 OUR FIT | | | | |
| 0.50 ±0.06 OUR AVERAGE | | Error includes scale factor of 1.2. | | |
| 0.446±0.052±0.046 avg | 157 | 103 BARATE | 98E ALEP | 1991–1995 LEP runs |
| 0.562±0.050±0.048 f&a | 264 | COAN | 96 CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |
| 103 Not independent of BARATE 98E $B(\tau^- \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau)$ and $B(\tau^- \rightarrow K^-K^0\pi^0\nu_\tau)$ values. | | | | |

 $\Gamma(\pi^-\bar{K}^0\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{40}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|--------------|----------|--|
| 0.38 ±0.04 OUR FIT | | | | |
| 0.36 ±0.04 OUR AVERAGE | | | | |
| 0.347±0.053±0.037 f&a | 299 | 104 BARATE | 99K ALEP | 1991–1995 LEP runs |
| 0.294±0.073±0.037 f&a | 142 | 105 BARATE | 98E ALEP | 1991–1995 LEP runs |
| 0.417±0.058±0.044 avg | | 106 COAN | 96 CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |
| 0.41 ±0.12 ±0.03 f&a | | 107 ACCIARRI | 95F L3 | 1991–1993 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.32 ±0.11 ±0.05 | 23 | 108 BUSKULIC | 96 ALEP | Repl. by BARATE 99K |

104 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

105 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+\pi^-$ decays.

106 Not independent of COAN 96 $B(h^-\bar{K}^0\pi^0\nu_\tau)$ and $B(K^-\bar{K}^0\pi^0\nu_\tau)$ measurements.

107 ACCIARRI 95F do not identify π^-/K^- and assume $B(K^-\bar{K}^0\pi^0\nu_\tau) = (0.05 \pm 0.05)\%$.

108 BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

 $\Gamma(\bar{K}^0\rho^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{41}/Γ

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|--|-------------|----------|--------------------|
| 0.22 ±0.05 OUR AVERAGE | | | |
| 0.250±0.057±0.044 | 109 BARATE | 99K ALEP | 1991–1995 LEP runs |
| 0.188±0.054±0.038 | 110 BARATE | 98E ALEP | 1991–1995 LEP runs |
| 109 BARATE 99K measure K^0 's by detecting K_L^0 's in hadron calorimeter. They determine the $\bar{K}^0\rho^-$ fraction in $\tau^- \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau$ decays to be $(0.72 \pm 0.12 \pm 0.10)$ and multiply their $B(\pi^-\bar{K}^0\pi^0\nu_\tau)$ measurement by this fraction to obtain the quoted result. | | | |
| 110 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+\pi^-$ decays. They determine the $\bar{K}^0\rho^-$ fraction in $\tau^- \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau$ decays to be $(0.64 \pm 0.09 \pm 0.10)$ and multiply their $B(\pi^-\bar{K}^0\pi^0\nu_\tau)$ measurement by this fraction to obtain the quoted result. | | | |

$\Gamma(K^- K^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{42}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|--------------|----------|--|
| 0.157 ± 0.021 OUR FIT | | | | |
| 0.144 ± 0.023 OUR AVERAGE | | | | |
| 0.143 ± 0.025 ± 0.015 | 78 | 111 BARATE | 99K ALEP | 1991–1995 LEP runs |
| 0.152 ± 0.076 ± 0.021 | 15 | 112 BARATE | 98E ALEP | 1991–1995 LEP runs |
| 0.145 ± 0.036 ± 0.020 | 32 | COAN | 96 CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.10 ± 0.05 ± 0.03 | 5 | 113 BUSKULIC | 96 ALEP | Repl. by BARATE 99K |
| 111 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter. | | | | |
| 112 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays. | | | | |
| 113 BUSKULIC 96 measure K^0 's by detecting K_L^0 's in their hadron calorimeter. | | | | |

$\Gamma(\pi^- \bar{K}^0 \geq 1 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{43}/\Gamma = (\Gamma_{40} + \Gamma_{44})/\Gamma$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|----------|--------------------|
| 0.324 ± 0.074 ± 0.066 | 148 | ABBIENDI | 00C OPAL | 1991–1995 LEP runs |

$\Gamma(\pi^- \bar{K}^0 \pi^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{44}/Γ

| VALUE (units 10^{-3}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|-------------|----------|--------------------|
| 0.26 ± 0.24 | | | 114 BARATE | 99R ALEP | 1991–1995 LEP runs |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.66 95 17 115 BARATE 99K ALEP 1991–1995 LEP runs

0.58 ± 0.33 ± 0.14 5 116 BARATE 98E ALEP 1991–1995 LEP runs

114 BARATE 99R combine the BARATE 98E and BARATE 99K measurements to obtain this value.

115 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter.

116 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

$\Gamma(K^- K^0 \pi^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{45}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|-----|-------------|----------|--------------------|
| <0.16 × 10⁻³ | 95 | 117 BARATE | 99R ALEP | 1991–1995 LEP runs |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.18 × 10⁻³ 95 118 BARATE 99K ALEP 1991–1995 LEP runs

<0.39 × 10⁻³ 95 119 BARATE 98E ALEP 1991–1995 LEP runs

117 BARATE 99R combine the BARATE 98E and BARATE 99K bounds to obtain this value.

118 BARATE 99K measure K^0 's by detecting K_L^0 's in hadron calorimeter.

119 BARATE 98E reconstruct K^0 's by using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

$\Gamma(\pi^- K^0 \bar{K}^0 \nu_\tau)/\Gamma_{\text{total}}$

Γ_{46}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

0.119±0.020 OUR FIT Error includes scale factor of 1.2.

0.116±0.028 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.


$0.153 \pm 0.030 \pm 0.016$ f&a 74 120 BARATE 98E ALEP 1991–1995 LEP

$0.092 \pm 0.020 \pm 0.012$ avg 42 121 COAN 96 CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$
runs

$0.31 \pm 0.12 \pm 0.04$ f&a ACCIARRI 95F L3 1991–1993 LEP
runs

120 BARATE 98E obtain this value by adding twice their $B(\pi^- K_S^0 \bar{K}_S^0 \nu_\tau)$ value to their $B(\pi^- K_L^0 \bar{K}_L^0 \nu_\tau)$ value.

121 We multiply the COAN 96 measurement $B(h^- K_S^0 \bar{K}_S^0 \nu_\tau) = (0.023 \pm 0.005 \pm 0.003)\%$ by 4 to obtain the listed value. This factor of 1/4 is uncertain, and might be as large as 1/2, due to Bose-Einstein correlations and the resonant parentage of this state.



$\Gamma(\pi^- K_S^0 \bar{K}_S^0 \nu_\tau)/\Gamma_{\text{total}} (\%)$

$\Gamma(\pi^- K_S^0 \bar{K}_S^0 \nu_\tau)/\Gamma_{\text{total}}$

$\Gamma_{47}/\Gamma = \frac{1}{4}\Gamma_{46}/\Gamma$

Bose-Einstein correlations might make the mixing fraction different than 1/4.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

0.030±0.005 OUR FIT Error includes scale factor of 1.2.

0.024±0.005 OUR AVERAGE

$0.026 \pm 0.010 \pm 0.005$ 6 BARATE 98E ALEP 1991–1995 LEP runs

$0.023 \pm 0.005 \pm 0.003$ 42 COAN 96 CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$

$\Gamma(\pi^- K_S^0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$ $\Gamma_{48}/\Gamma = \frac{1}{2}\Gamma_{46}/\Gamma$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------------|-------------|--------|-------------------------------------|
| 0.060 ± 0.010 OUR FIT | | | | Error includes scale factor of 1.2. |
| 0.101 ± 0.023 ± 0.013 | avg | 68 | BARATE | 98E ALEP 1991–1995 LEP runs |

 $\Gamma(\pi^- K^0 \bar{K}^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{49}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|----------|--------------------|
| (0.31 ± 0.23) × 10⁻³ | 122 BARATE | 99R ALEP | 1991–1995 LEP runs |
| 122 BARATE 99R combine BARATE 98E $\Gamma(\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ and $\Gamma(\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ measurements to obtain this value. | | | |

 $\Gamma(\pi^- K_S^0 K_S^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{50}/Γ

| VALUE (%) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|-------------|----------|--------------------|
| <0.020 | 95 | BARATE | 98E ALEP | 1991–1995 LEP runs |

 $\Gamma(\pi^- K_S^0 K_L^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{51}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|----------|--------------------|
| 0.031 ± 0.011 ± 0.005 | 11 | BARATE | 98E ALEP | 1991–1995 LEP runs |

 $\Gamma(K^0 h^+ h^- h^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{52}/Γ

| VALUE (%) | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|---------------|--------|--|
| <0.17 | 95 | TSCHIRHART 88 | HRS | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.27 | 90 | BELTRAMI | 85 HRS | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

 $\Gamma(K^0 h^+ h^- h^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{53}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|----------|--------------------|
| 0.023 ± 0.019 ± 0.007 | 6 | 123 BARATE | 98E ALEP | 1991–1995 LEP runs |

123 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+ \pi^-$ decays.

 $\Gamma(h^- h^- h^+ \geq 0 \text{ neut. } \nu_\tau \text{ ("3-prong")})/\Gamma_{\text{total}}$ Γ_{54}/Γ

$$\begin{aligned} \Gamma_{54}/\Gamma &= (0.3431\Gamma_{35} + 0.3431\Gamma_{37} + 0.3431\Gamma_{40} + 0.3431\Gamma_{42} + 0.4508\Gamma_{46} + \Gamma_{63} + \Gamma_{71} + \\ &\Gamma_{79} + \Gamma_{80} + \Gamma_{88} + \Gamma_{92} + \Gamma_{95} + \Gamma_{96} + 0.285\Gamma_{124} + 0.285\Gamma_{126} + 0.9101\Gamma_{144} + \\ &0.9101\Gamma_{145})/\Gamma \end{aligned}$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|------|-------------|----------|---|
| 15.18 ± 0.13 OUR FIT | | | | Error includes scale factor of 1.2. |
| 14.8 ± 0.4 OUR AVERAGE | | | | |
| 14.4 ± 0.6 ± 0.3 | f&a | ADEVA | 91F L3 | $E_{\text{cm}}^{\text{ee}} = 88.3\text{--}94.3 \text{ GeV}$ |
| 15.0 ± 0.4 ± 0.3 | f&a | BEHREND | 89B CELL | $E_{\text{cm}}^{\text{ee}} = 14\text{--}47 \text{ GeV}$ |
| 15.1 ± 0.8 ± 0.6 | f&a | AIHARA | 87B TPC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------------------------|------|---------------|----------|---|
| 13.5 ± 0.3 ± 0.3 | | ABACHI | 89B HRS | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 12.8 ± 1.0 ± 0.7 | 124 | BURCHAT | 87 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 12.1 ± 0.5 ± 1.2 | | RUCKSTUHL | 86 DLCO | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 12.8 ± 0.5 ± 0.8 | 1420 | SCHMIDKE | 86 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 15.3 ± 1.1 +1.3 -1.6 | 367 | ALTHOFF | 85 TASS | $E_{\text{cm}}^{\text{ee}} = 34.5 \text{ GeV}$ |
| 13.6 ± 0.5 ± 0.8 | | BARTEL | 85F JADE | $E_{\text{cm}}^{\text{ee}} = 34.6 \text{ GeV}$ |
| 12.2 ± 1.3 ± 3.9 | 125 | BERGER | 85 PLUT | $E_{\text{cm}}^{\text{ee}} = 34.6 \text{ GeV}$ |
| 13.3 ± 0.3 ± 0.6 | | FERNANDEZ | 85 MAC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 24 ± 6 | 35 | BRANDELIK | 80 TASS | $E_{\text{cm}}^{\text{ee}} = 30 \text{ GeV}$ |
| 32 ± 5 | 692 | 126 BACINO | 78B DLCO | $E_{\text{cm}}^{\text{ee}} = 3.1\text{--}7.4 \text{ GeV}$ |
| 35 ± 11 | | 126 BRANDELIK | 78 DASP | Assumes $V-A$ decay |
| 18 ± 6.5 | 33 | 126 JAROS | 78 MRK1 | $E_{\text{cm}}^{\text{ee}} > 6 \text{ GeV}$ |

124 BURCHAT 87 value is not independent of SCHMIDKE 86 value.

125 Not independent of BERGER 85 $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma_{\text{total}}$, $\Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}$, $\Gamma(h^- \geq 1 \text{ neutrals} \nu_\tau)/\Gamma_{\text{total}}$, and $\Gamma(h^- \geq 0 K_L^0 \nu_\tau)/\Gamma_{\text{total}}$, and therefore not used in the fit.

126 Low energy experiments are not in average or fit because the systematic errors in background subtraction are judged to be large.

$$\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{55}/\Gamma$$

$$\Gamma_{55}/\Gamma = (\Gamma_{63} + \Gamma_{71} + \Gamma_{79} + \Gamma_{80} + \Gamma_{88} + \Gamma_{92} + \Gamma_{95} + \Gamma_{96} + 0.285\Gamma_{124} + 0.285\Gamma_{126} + 0.9101\Gamma_{144} + 0.9101\Gamma_{145}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|---|-------|-------------|----------|--|
| 14.58 ± 0.13 OUR FIT | Error includes scale factor of 1.2. | | | | |
| 14.63 ± 0.25 OUR AVERAGE | Error includes scale factor of 1.4. See the ideogram below. | | | | |
| 14.96 ± 0.09 ± 0.22 | f&a | 10.4k | AKERS | 95Y OPAL | 1991–1994 LEP runs |
| 14.22 ± 0.10 ± 0.37 | avg | 127 | BALEST | 95C CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |
| 13.3 ± 0.3 ± 0.8 | f&a | 128 | ALBRECHT | 92D ARG | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ |
| 14.35 +0.40 -0.45 | ± 0.24 | f&a | DECAMP | 92C ALEP | 1989–1990 LEP runs |


• • • We do not use the following data for averages, fits, limits, etc. • • •

15.26 ± 0.26 ± 0.22

ACTON 92H OPAL Repl. by AKERS 95Y

127 Not independent of BALEST 95C $B(h^- h^- h^+ \nu_\tau)$ and $B(h^- h^- h^+ \pi^0 \nu_\tau)$ values, and BORTOLETTO 93 $B(h^- h^- h^+ 2\pi^0 \nu_\tau)/B(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau)$ value.

128 This ALBRECHT 92D value is not independent of their $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)\Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}^2$ value.



$$\Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}} (\%)$$

$$\Gamma(\pi^- \pi^+ \pi^- \geq 0 \text{ neutrals } \nu_\tau) / \Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{"3-prong"})) \quad \Gamma_{56}/\Gamma_{54}$$

$$\Gamma_{56}/\Gamma_{54} = (0.3431\Gamma_{35} + 0.3431\Gamma_{40} + 0.1078\Gamma_{46} + \Gamma_{63} + \Gamma_{71} + \Gamma_{79} + \Gamma_{80} + 0.285\Gamma_{124} + 0.9101\Gamma_{144} + 0.9101\Gamma_{145}) / (0.3431\Gamma_{35} + 0.3431\Gamma_{37} + 0.3431\Gamma_{40} + 0.3431\Gamma_{42} + 0.4508\Gamma_{46} + \Gamma_{63} + \Gamma_{71} + \Gamma_{79} + \Gamma_{80} + \Gamma_{88} + \Gamma_{92} + \Gamma_{95} + \Gamma_{96} + 0.285\Gamma_{124} + 0.285\Gamma_{126} + 0.9101\Gamma_{144} + 0.9101\Gamma_{145})$$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|--------|--|
| 0.9547±0.0035 OUR FIT | | | | Error includes scale factor of 1.4. |
| 0.945 ±0.019 | 490 | 129 BAUER | 94 TPC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

129 BAUER 94 quote $B(\pi^- \pi^+ \pi^- \geq 0 \text{ neutrals } \nu_\tau) = 0.1329 \pm 0.0027$. We divide by 0.1406, their assumed value for $B(\text{"3prong"})$.

$$\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{57}/\Gamma$$

$$\Gamma_{57}/\Gamma = (0.3431\Gamma_{35} + 0.3431\Gamma_{37} + \Gamma_{63} + \Gamma_{88} + \Gamma_{95} + 0.0221\Gamma_{144}) / \Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------|--------------|----------|---|
| 9.97±0.10 OUR FIT | | | | Error includes scale factor of 1.1. |
| 9.8 ±0.6 OUR AVERAGE | | | | Error includes scale factor of 4.4. See the ideogram below. |
| 7.6 ±0.1 ±0.5 avg | 7.5k | 130 ALBRECHT | 96E ARG | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ |
| 9.92±0.10±0.09 f&a | 11.2k | 131 BUSKULIC | 96 ALEP | LEP 1991–1993 data |
| 9.49±0.36±0.63 f&a | | DECAMP | 92C ALEP | 1989–1990 LEP runs |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|------|---------------|---------|---|
| $8.7 \pm 0.7 \pm 0.3$ | 694 | 132 BEHREND | 90 CELL | $E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$ |
| $7.0 \pm 0.3 \pm 0.7$ | 1566 | 133 BAND | 87 MAC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| $6.7 \pm 0.8 \pm 0.9$ | | 134 BURCHAT | 87 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| $6.4 \pm 0.4 \pm 0.9$ | | 135 RUCKSTUHL | 86 DLCO | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| $7.8 \pm 0.5 \pm 0.8$ | 890 | SCHMIDKE | 86 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| $8.4 \pm 0.4 \pm 0.7$ | 1255 | 135 FERNANDEZ | 85 MAC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| $9.7 \pm 2.0 \pm 1.3$ | | BEHREND | 84 CELL | $E_{\text{cm}}^{\text{ee}} = 14,22 \text{ GeV}$ |

130 ALBRECHT 96E not independent of ALBRECHT 93C $\Gamma(h^- h^- h^+ \nu_\tau)$ (ex. K^0) $\times \Gamma(\text{particle}^- \geq 0 \text{ neutrals} \geq 0 K_L^0 \nu_\tau) / \Gamma_{\text{total}}^2$ value.


131 BUSKULIC 96 quote $B(h^- h^- h^+ \nu_\tau)$ (ex. K^0) = $9.50 \pm 0.10 \pm 0.11$. We add 0.42 to remove their K^0 correction and reduce the systematic error accordingly.

132 BEHREND 90 subtract 0.3% to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution to measured events.

133 BAND 87 subtract for charged kaon modes; not independent of FERNANDEZ 85 value.

134 BURCHAT 87 value is not independent of SCHMIDKE 86 value.

135 Value obtained by multiplying paper's $R = B(h^- h^- h^+ \nu_\tau) / B(3\text{-prong})$ by $B(3\text{-prong}) = 0.143$ and subtracting 0.3% for $K^*(892)$ background.



$$\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma_{\text{total}} (\%)$$

$$\Gamma(h^- h^- h^+ \nu_\tau(\text{ex. } K^0)) / \Gamma_{\text{total}} = \Gamma_{58} / \Gamma = (\Gamma_{63} + \Gamma_{88} + \Gamma_{95} + 0.0221\Gamma_{144}) / \Gamma$$

Γ_{58}/Γ

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------------------------------|--------------|----------------------------|
| 9.61±0.10 OUR FIT | | Error includes scale factor of 1.1. | | |
| 9.57±0.11 OUR AVERAGE | | | | |
| 9.50±0.10±0.11 | avg | 11.2k | 136 BUSKULIC | 96 ALEP LEP 1991–1993 data |

9.87±0.10±0.24 avg 137 AKERS 95Y OPAL 1991–1994 LEP runs

9.51±0.07±0.20 f&a 37.7k BAlest 95C CLEO $E_{\text{cm}}^{ee} \approx 10.6$ GeV

136 Not independent of BUSKULIC 96 B($h^- h^- h^+ \nu_\tau$) value.

137 Not independent of AKERS 95Y B($h^- h^- h^+ \geq 0$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^+ \pi^-$)) and B($h^- h^- h^+ \nu_\tau$ (ex. K^0))/B($h^- h^- h^+ \geq 0$ neutrals ν_τ (ex. $K_S^0 \rightarrow \pi^+ \pi^-$)) values.

$$\Gamma(h^- h^- h^+ \nu_\tau(\text{ex. } K^0)) / \Gamma(h^- h^- h^+ \geq 0 \text{ neutrals } \nu_\tau(\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-)) = \Gamma_{58}/\Gamma_{55}$$

$$\Gamma_{58}/\Gamma_{55} = (\Gamma_{63} + \Gamma_{88} + \Gamma_{95} + 0.0221\Gamma_{144}) / (\Gamma_{63} + \Gamma_{71} + \Gamma_{79} + \Gamma_{80} + \Gamma_{88} + \Gamma_{92} + \Gamma_{95} + \Gamma_{96} + 0.285\Gamma_{124} + 0.285\Gamma_{126} + 0.9101\Gamma_{144} + 0.9101\Gamma_{145})$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------|-------------------------------------|--------------------|
| 0.659±0.006 OUR FIT | | Error includes scale factor of 1.1. | |
| 0.660±0.004±0.014 | AKERS | 95Y OPAL | 1991–1994 LEP runs |

$$\Gamma(h^- h^- h^+ \nu_\tau(\text{ex. } K^0, \omega)) / \Gamma_{\text{total}} = \Gamma_{59}/\Gamma = (\Gamma_{63} + \Gamma_{88} + \Gamma_{95}) / \Gamma$$

| VALUE (%) | DOCUMENT ID |
|--------------------------|-------------------------------------|
| 9.56±0.10 OUR FIT | Error includes scale factor of 1.1. |

$$\Gamma(\pi^- \pi^+ \pi^- \nu_\tau) / \Gamma_{\text{total}} = \Gamma_{60}/\Gamma = (0.3431\Gamma_{35} + \Gamma_{63} + 0.0221\Gamma_{144}) / \Gamma$$

| VALUE (%) | DOCUMENT ID |
|--------------------------|-------------------------------------|
| 9.49±0.11 OUR FIT | Error includes scale factor of 1.1. |

$$\Gamma(\pi^- \pi^+ \pi^- \nu_\tau(\text{ex. } K^0)) / \Gamma_{\text{total}} = \Gamma_{61}/\Gamma = (\Gamma_{63} + 0.0221\Gamma_{144}) / \Gamma$$

| VALUE (%) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|----------------|----------|--------------------|
| 9.18±0.11 OUR FIT | 95 | 138 ACKERSTAFF | 97R OPAL | 1992–1994 LEP runs |

138 Model-independent limit from structure function analysis on contribution to B($\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau$ (ex. K^0)) from non-axial vectors.

$$\Gamma(\pi^- \pi^+ \pi^- \nu_\tau(\text{ex. } K^0, \omega)) / \Gamma_{\text{total}} = \Gamma_{63}/\Gamma$$

| VALUE (%) | DOCUMENT ID |
|--------------------------|-------------------------------------|
| 9.13±0.11 OUR FIT | Error includes scale factor of 1.1. |

$\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals} \nu_\tau)/\Gamma_{\text{total}}$ Γ_{64}/Γ

$$\Gamma_{64}/\Gamma = (0.3431\Gamma_{40} + 0.3431\Gamma_{42} + 0.1077\Gamma_{46} + \Gamma_{71} + \Gamma_{79} + \Gamma_{80} + \Gamma_{92} + \Gamma_{96} + 0.285\Gamma_{124} + 0.285\Gamma_{126} + 0.888\Gamma_{144} + 0.9101\Gamma_{145})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

5.17±0.11 OUR FIT Error includes scale factor of 1.2.

4.4 ±1.0 OUR AVERAGE

| | | | | | |
|-----------------|-----|-----|--------------|---------|---|
| 4.2 ± 0.5 ± 0.9 | f&a | 203 | 139 ALBRECHT | 87L ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| 6.2 ± 2.3 ± 1.7 | f&a | | BEHREND | 84 CELL | $E_{\text{cm}}^{\text{ee}} = 14,22 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------|-----|-------------------|---------|--|
| 5.6 ± 0.7 ± 0.3 | 352 | 140 BEHREND | 90 CELL | $E_{\text{cm}}^{\text{ee}} = 35 \text{ GeV}$ |
| 6.1 ± 0.8 ± 0.9 | | 141 BURCHAT | 87 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 7.6 ± 0.4 ± 0.9 | | 142,143 RUCKSTUHL | 86 DLCO | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 4.7 ± 0.5 ± 0.8 | 530 | 144 SCHMIDKE | 86 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 5.6 ± 0.4 ± 0.7 | | 143 FERNANDEZ | 85 MAC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

139 ALBRECHT 87L measure the product of branching ratios $B(3\pi^\pm \pi^0 \nu_\tau)$ $B((e\bar{\nu} \text{ or } \mu\bar{\nu} \text{ or } \pi \text{ or } K \text{ or } \rho)\nu_\tau) = 0.029$ and use the PDG 86 values for the second branching ratio which sum to 0.69 ± 0.03 to get the quoted value.

140 BEHREND 90 value is not independent of BEHREND 90 $B(3h\nu_\tau \geq 1 \text{ neutrals}) + B(5\text{-prong})$.

141 BURCHAT 87 value is not independent of SCHMIDKE 86 value.

142 Contributions from kaons and from $>1\pi^0$ are subtracted. Not independent of (3-prong + $0\pi^0$) and (3-prong + $\geq 0\pi^0$) values.

143 Value obtained using paper's $R = B(h^- h^- h^+ \nu_\tau)/B(3\text{-prong})$ and current $B(3\text{-prong}) = 0.143$.

144 Not independent of SCHMIDKE 86 $h^- h^- h^+ \nu_\tau$ and $h^- h^- h^+(\geq 0\pi^0)\nu_\tau$ values.

 $\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals} \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{65}/Γ

$$\Gamma_{65}/\Gamma = (\Gamma_{71} + \Gamma_{79} + \Gamma_{80} + \Gamma_{92} + \Gamma_{96} + 0.285\Gamma_{124} + 0.285\Gamma_{126} + 0.888\Gamma_{144} + 0.9101\Gamma_{145})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

4.97±0.11 OUR FIT Error includes scale factor of 1.2.

5.07±0.24 OUR AVERAGE

| | | | | |
|--------------------|-----|------------|----------|--------------------|
| 5.09 ± 0.10 ± 0.23 | avg | 145 AKERS | 95Y OPAL | 1991–1994 LEP runs |
| 4.95 ± 0.29 ± 0.65 | f&a | 570 DECAMP | 92C ALEP | 1989–1990 LEP runs |

145 Not independent of AKERS 95Y $B(h^- h^- h^+ \geq 0 \text{ neutrals} \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))$ and $B(h^- h^- h^+ \geq 0 \text{ neutrals} \nu_\tau (\text{ex. } K^0))/B(h^- h^- h^+ \geq 0 \text{ neutrals} \nu_\tau (\text{ex. } K_S^0 \rightarrow \pi^+ \pi^-))$ values.

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{66}/\Gamma$$

$$\Gamma_{66}/\Gamma = (0.3431\Gamma_{40} + 0.3431\Gamma_{42} + \Gamma_{71} + \Gamma_{92} + \Gamma_{96} + 0.231\Gamma_{126} + 0.888\Gamma_{144} + 0.0221\Gamma_{145})/\Gamma$$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------|------|-------------------------|
| 4.49±0.08 OUR FIT | | | | |
| 4.45±0.09±0.07 | 6.1k | ¹⁴⁶ BUSKULIC | 96 | ALEP LEP 1991–1993 data |
| ¹⁴⁶ BUSKULIC 96 quote $B(h^- h^- h^+ \pi^0 \nu_\tau (\text{ex. } K^0)) = 4.30 \pm 0.09 \pm 0.09$. We add 0.15 to remove their K^0 correction and reduce the systematic error accordingly. | | | | |

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{67}/\Gamma$$

$$\Gamma_{67}/\Gamma = (\Gamma_{71} + \Gamma_{92} + \Gamma_{96} + 0.231\Gamma_{126} + 0.888\Gamma_{144} + 0.0221\Gamma_{145})/\Gamma$$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---|
| 4.30±0.08 OUR FIT | | | | |
| 4.23±0.06±0.22 | 7.2k | BALEST | 95C | CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |

$$\Gamma(h^- h^- h^+ \pi^0 \nu_\tau (\text{ex. } K^0, \omega)) / \Gamma_{\text{total}} \quad \Gamma_{68}/\Gamma = (\Gamma_{71} + \Gamma_{92} + \Gamma_{96} + 0.231\Gamma_{126})/\Gamma$$

$$\Gamma_{68}/\Gamma = (\Gamma_{71} + \Gamma_{92} + \Gamma_{96} + 0.231\Gamma_{126})/\Gamma$$

| VALUE (%) | DOCUMENT ID |
|--------------------------|-------------|
| 2.58±0.08 OUR FIT | |

$$\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{69}/\Gamma = (0.3431\Gamma_{40} + \Gamma_{71} + 0.888\Gamma_{144} + 0.0221\Gamma_{145})/\Gamma$$

$$\Gamma_{69}/\Gamma = (0.3431\Gamma_{40} + \Gamma_{71} + 0.888\Gamma_{144} + 0.0221\Gamma_{145})/\Gamma$$

| VALUE (%) | DOCUMENT ID |
|--------------------------|-------------|
| 4.32±0.08 OUR FIT | |

$$\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}} \quad \Gamma_{70}/\Gamma = (\Gamma_{71} + 0.888\Gamma_{144} + 0.0221\Gamma_{145})/\Gamma$$

$$\Gamma_{70}/\Gamma = (\Gamma_{71} + 0.888\Gamma_{144} + 0.0221\Gamma_{145})/\Gamma$$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------------------------|------|---|
| 4.20±0.08 OUR FIT | | | |
| 4.19±0.10±0.21 | ¹⁴⁷ EDWARDS | 00A | CLEO $4.7 \text{ fb}^{-1} E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

¹⁴⁷ EDWARDS 00A quote $(4.19 \pm 0.10) \times 10^{-2}$ with a 5% systematic error.

$$\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex. } K^0, \omega)) / \Gamma_{\text{total}} \quad \Gamma_{71}/\Gamma$$

$$\Gamma_{71}/\Gamma = (\Gamma_{71} + 0.888\Gamma_{144} + 0.0221\Gamma_{145})/\Gamma$$

| VALUE (%) | DOCUMENT ID |
|--------------------------|-------------|
| 2.47±0.08 OUR FIT | |

$$\Gamma(h^- (\rho\pi)^0 \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau) \quad \Gamma_{72}/\Gamma_{66}$$

$$\Gamma_{72}/\Gamma_{66} = (\Gamma_{74} + \Gamma_{75} + \Gamma_{76})/\Gamma_{66}$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |

0.64±0.07±0.03 ¹⁴⁸ ALBRECHT 91D ARG $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$

¹⁴⁸ ALBRECHT 91D not independent of their $\Gamma(h^- \rho^+ h^- \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$, $\Gamma(h^- \rho^- h^+ \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$, and $\Gamma(h^- \rho \pi^0 \nu_\tau) / \Gamma(h^- h^- h^+ \pi^0 \nu_\tau)$ values.

$\Gamma((a_1(1260)h)^-\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ Γ_{73}/Γ_{66}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|--------------|---------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.44 | 95 | 149 ALBRECHT | 91D ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |
| 149 ALBRECHT 91D not independent of their $\Gamma(h^-\omega\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ (ex. K^0), $\Gamma(h^-\rho^0\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$, $\Gamma(h^-\rho^+h^-\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$, and $\Gamma(h^-\rho^-h^+\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ values. | | | | |

$\Gamma(h^-\rho\pi^0\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ Γ_{74}/Γ_{66}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|---------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $0.30 \pm 0.04 \pm 0.02$ | 393 | ALBRECHT | 91D ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |

$\Gamma(h^-\rho^+h^-\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ Γ_{75}/Γ_{66}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|---------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $0.10 \pm 0.03 \pm 0.04$ | 142 | ALBRECHT | 91D ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |

$\Gamma(h^-\rho^-h^+\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ Γ_{76}/Γ_{66}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|---------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $0.26 \pm 0.05 \pm 0.01$ | 370 | ALBRECHT | 91D ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |

$[\Gamma(h^-\rho^+h^-\nu_\tau) + \Gamma(h^-\rho^-h^+\nu_\tau)]/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ $(\Gamma_{75} + \Gamma_{76})/\Gamma_{66}$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|--------------|---------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $0.33 \pm 0.06 \pm 0.01$ | 475 | 150 ALBRECHT | 91D ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |
| 150 ALBRECHT 91D not independent of their $\Gamma(h^-\rho^+h^-\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ and $\Gamma(h^-\rho^-h^+\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ values. | | | | |

$\Gamma(h^-h^-h^+2\pi^0\nu_\tau)/\Gamma_{total}$ Γ_{77}/Γ

| VALUE (%) | DOCUMENT ID |
|----------------------------|-------------|
| 0.54 ± 0.04 OUR FIT | |

$\Gamma(h^-h^-h^+2\pi^0\nu_\tau(\text{ex. } K^0))/\Gamma_{total}$ Γ_{78}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|---------|--------------------|
| 0.53 ± 0.04 OUR FIT | | | | |
| 0.50 ± 0.07 ± 0.07 | 1.8k | BUSKULIC | 96 ALEP | LEP 1991–1993 data |

$$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau (\text{ex.} K^0)) / \Gamma(h^- h^- h^+ \geq 1 \text{neut. } \nu_\tau (\text{"3-prong"}) \quad \Gamma_{78}/\Gamma_{54}$$

$$\Gamma_{78}/\Gamma_{54} = (\Gamma_{79} + 0.236\Gamma_{124} + 0.888\Gamma_{145}) / (0.3431\Gamma_{35} + 0.3431\Gamma_{37} + 0.3431\Gamma_{40} + 0.3431\Gamma_{42} + 0.4508\Gamma_{46} + \Gamma_{63} + \Gamma_{71} + \Gamma_{79} + \Gamma_{80} + \Gamma_{88} + \Gamma_{92} + \Gamma_{95} + \Gamma_{96} + 0.285\Gamma_{124} + 0.285\Gamma_{126} + 0.9101\Gamma_{144} + 0.9101\Gamma_{145})$$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------|--------------------|-------------|--|
| 0.0347 ± 0.0028 OUR FIT | | | | |
| 0.034 ± 0.002 ± 0.003 | 668 | BORTOLETTO93 | CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |

$$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau (\text{ex.} K^0, \omega, \eta)) / \Gamma_{\text{total}} \quad \Gamma_{79}/\Gamma$$

| <u>VALUE (%)</u> | <u>DOCUMENT ID</u> |
|----------------------------|--------------------|
| 0.11 ± 0.04 OUR FIT | |

$$\Gamma(h^- h^- h^+ \geq 3\pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{80}/\Gamma$$

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|-------------|--------------------|-------------|-------------------------------------|
| 0.13 ± 0.08 OUR FIT | | | | Error includes scale factor of 1.3. |
| 0.11 ± 0.04 ± 0.05 | 440 | BUSKULIC | 96 | ALEP LEP 1991–1993 data |

$$\Gamma(h^- h^- h^+ 3\pi^0 \nu_\tau) / \Gamma_{\text{total}} \quad \Gamma_{81}/\Gamma$$

| <u>VALUE (units 10⁻⁴)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------------|-------------|--------------------|-------------|---|
| 2.85 ± 0.56 ± 0.51 | 57 | ANDERSON | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

$$\Gamma(K^- h^+ h^- \geq 0 \text{ neutrals } \nu_\tau) / \Gamma_{\text{total}}$$

$$\Gamma_{82}/\Gamma = (0.3431\Gamma_{37} + 0.3431\Gamma_{42} + \Gamma_{88} + \Gamma_{92} + \Gamma_{95} + \Gamma_{96} + 0.285\Gamma_{126}) / \Gamma$$

| <u>VALUE (%)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|------------|--------------------|-------------|--|
| 0.65 ± 0.05 OUR FIT | | | | Error includes scale factor of 1.4. |
| <0.6 | 90 | AIHARA | 84C TPC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

$$\Gamma(K^- h^+ \pi^- \nu_\tau (\text{ex.} K^0)) / \Gamma_{\text{total}} \quad \Gamma_{83}/\Gamma = (\Gamma_{88} + \Gamma_{95}) / \Gamma$$

| <u>VALUE (%)</u> | <u>DOCUMENT ID</u> |
|----------------------------|--------------------|
| 0.43 ± 0.05 OUR FIT | |

Error includes scale factor of 1.5.

$$\Gamma(K^- h^+ \pi^- \nu_\tau (\text{ex.} K^0)) / \Gamma(\pi^- \pi^+ \pi^- \nu_\tau (\text{ex.} K^0))$$

$$\Gamma_{83}/\Gamma_{61} = (\Gamma_{88} + \Gamma_{95}) / (\Gamma_{63} + 0.0221\Gamma_{144})$$

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|-------------|--------------------|-------------|---|
| 4.7 ± 0.6 OUR FIT | | | | Error includes scale factor of 1.5. |
| 5.44 ± 0.21 ± 0.53 | 7.9k | RICHICHI | 99 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

$$\Gamma(K^- h^+ \pi^- \pi^0 \nu_\tau (\text{ex.} K^0)) / \Gamma_{\text{total}} \quad \Gamma_{84}/\Gamma = (\Gamma_{92} + \Gamma_{96} + 0.231\Gamma_{126}) / \Gamma$$

| <u>VALUE (%)</u> | <u>DOCUMENT ID</u> |
|------------------------------|--------------------|
| 0.107 ± 0.022 OUR FIT | |

$$\Gamma(K^- h^+ \pi^- \pi^0 \nu_\tau (\text{ex.} K^0)) / \Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau (\text{ex.} K^0))$$

$$\Gamma_{84}/\Gamma_{70} = (\Gamma_{92} + \Gamma_{96} + 0.231\Gamma_{126}) / (\Gamma_{71} + 0.888\Gamma_{144} + 0.0221\Gamma_{145})$$

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|-------------|--------------------|-------------|---|
| 2.5 ± 0.5 OUR FIT | | | | |
| 2.61 ± 0.45 ± 0.42 | 719 | RICHICHI | 99 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

$$\Gamma(K^-\pi^+\pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$$

$$\Gamma_{85}/\Gamma = (0.3431\Gamma_{37} + 0.3431\Gamma_{42} + \Gamma_{88} + \Gamma_{92} + 0.285\Gamma_{126})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------------------------------|------|-------------|------|---------|
| 0.44 ± 0.05 OUR FIT | Error includes scale factor of 1.4. | | | | |

0.39 ± 0.19 OUR AVERAGE Error includes scale factor of 1.5.

| | | | | | |
|--|--------|-----|----|-----------|--|
| 0.58 ^{+0.15} _{-0.13} | ± 0.12 | f&a | 20 | 151 BAUER | 94 TPC $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 0.22 ^{+0.16} _{-0.13} | ± 0.05 | f&a | 9 | 152 MILLS | 85 DLCO $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

151 We multiply 0.58% by 0.20, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

152 Error correlated with MILLS 85 ($K K \pi \nu$) value. We multiply 0.22% by 0.23, the relative systematic error quoted by MILLS 85, to obtain the systematic error.

$$\Gamma(K^-\pi^+\pi^- \geq 0 \pi^0 \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}} \quad \Gamma_{86}/\Gamma = (\Gamma_{88} + \Gamma_{92} + 0.231\Gamma_{126})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------------------------------|-------------|------|---------|
| 0.34 ± 0.05 OUR FIT | Error includes scale factor of 1.4. | | | |

0.30 ± 0.05 OUR AVERAGE

| | | | | |
|-----------------------|-----|------------|----------|--------------------|
| 0.343 ± 0.073 ± 0.031 | f&a | ABBIENDI | 00D OPAL | 1990–1995 LEP runs |
| 0.275 ± 0.064 | avg | 153 BARATE | 98 ALEP | 1991–1995 LEP runs |

153 Not independent of BARATE 98 $\Gamma(\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$ and $\Gamma(\tau^- \rightarrow K^-\pi^+\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ values.

$$\Gamma(K^-\pi^+\pi^-\nu_\tau)/\Gamma_{\text{total}} \quad \Gamma_{87}/\Gamma = (0.3431\Gamma_{37} + \Gamma_{88})/\Gamma$$

| VALUE (%) | | DOCUMENT ID |
|----------------------------|-------------------------------------|-------------|
| 0.32 ± 0.05 OUR FIT | Error includes scale factor of 1.5. | |

$$\Gamma(K^-\pi^+\pi^-\nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}$$

$$\Gamma_{88}/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.


| VALUE (%) | | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------------------------------|------|-------------|------|---------|
| 0.27 ± 0.05 OUR FIT | Error includes scale factor of 1.5. | | | | |

0.28 ± 0.05 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

| | | | |
|---------------------------|------------------|--|--------------------|
| 0.360 ± 0.082 ± 0.048 avg | ABBIENDI | 00D OPAL | 1990–1995 LEP runs |
| 0.346 ± 0.023 ± 0.056 avg | 158 154 RICHICHI | 99 CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ | |
| 0.214 ± 0.037 ± 0.029 f&a | BARATE | 98 ALEP | 1991–1995 LEP runs |

154 Not independent of RICHICHI 99

$\Gamma(\tau^- \rightarrow K^- h^+ \pi^- \nu_\tau (\text{ex. } K^0))/\Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$, $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau (\text{ex. } K^0))$ and BAEST 95C $\Gamma(\tau^- \rightarrow h^- h^- h^+ \nu_\tau (\text{ex. } K^0))/\Gamma_{\text{total}}$ values.



$$\Gamma(K^-\pi^+\pi^-\nu_\tau(\text{ex. } K^0))/\Gamma_{\text{total}} (\%)$$

| $\Gamma(K^-\rho^0\nu_\tau \rightarrow K^-\pi^+\pi^-\nu_\tau)/\Gamma(K^-\pi^+\pi^-\nu_\tau(\text{ex. } K^0))$ | Γ_{89}/Γ_{88} |
|--|---|
| <u>VALUE</u> | <u>DOCUMENT ID</u> |
| 0.48±0.14±0.10 | 155 ASNER |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | 00B CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| 0.39±0.14 | 156 BARATE 99R ALEP 1991–1995 LEP runs |
| 155 ASNER 00B assume $\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau$ (ex. K^0) decays proceed only through $K\rho$ and $K^*\pi$ intermediate states. They assume the resonance structure of $\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau$ (ex. K^0) decays is dominated by $K_1(1270)^-$ and $K_1(1400)^-$ resonances, and assume $B(K_1(1270) \rightarrow K^*(892)\pi) = (16 \pm 5)\%$, $B(K_1(1270) \rightarrow K\rho) = (42 \pm 6)\%$, and $B(K_1(1400) \rightarrow K\rho) = 0$. | |
| 156 BARATE 99R assume $\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau$ (ex. K^0) decays proceed only through $K\rho$ and $K^*\pi$ intermediate states. The quoted error is statistical only. | |

$$\Gamma(K^-\pi^+\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$$

VALUE (units 10^{-4})
 12.0 ± 2.5 OUR FIT

$$\Gamma_{90}/\Gamma = (0.3431\Gamma_{42} + \Gamma_{92} + 0.231\Gamma_{126})/\Gamma$$

DOCUMENT ID

$$\Gamma(K^-\pi^+\pi^-\pi^0\nu_\tau(\text{ex. } K^0))/\Gamma_{\text{total}}$$

$$\Gamma_{91}/\Gamma = (\Gamma_{92} + 0.231\Gamma_{126})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|----------------|
| 6.7 ± 2.4 OUR FIT | | | | |
| 7.0 ± 2.5 OUR AVERAGE | | | | |

| | | | | |
|-----------------------|-----|--------------|---------|---------------------------------|
| $7.5 \pm 2.6 \pm 1.8$ | avg | 157 RICHICHI | 99 CLEO | $E_{\text{cm}}^{ee} = 10.6$ GeV |
| $6.1 \pm 3.9 \pm 1.8$ | f&a | BARATE | 98 ALEP | 1991–1995 LEP runs |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----|----|----------|----------|--------------------|
| <17 | 95 | ABBIENDI | 00D OPAL | 1990–1995 LEP runs |
|-----|----|----------|----------|--------------------|

157 Not independent of RICHICHI 99

$\Gamma(\tau^- \rightarrow K^- h^+ \pi^- \nu_\tau(\text{ex. } K^0))/\Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau(\text{ex. } K^0))$, $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau(\text{ex. } K^0))$ and BALEST 95C $\Gamma(\tau^- \rightarrow h^- h^- h^+ \nu_\tau(\text{ex. } K^0))/\Gamma_{\text{total}}$ values.

$$\Gamma(K^-\pi^+K^-\geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$$

$$\Gamma_{93}/\Gamma$$

| <u>VALUE (%)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|------------|--------------------|-------------|-------------------------------|
| <0.09 | 95 | BAUER | 94 TPC | $E_{\text{cm}}^{ee} = 29$ GeV |

$$\Gamma(K^-\bar{K}^+\pi^-\geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$$

$$\Gamma_{94}/\Gamma = (\Gamma_{95} + \Gamma_{96})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|----------------|
| 0.201 ± 0.023 OUR FIT | | | | |
| 0.203 ± 0.031 OUR AVERAGE | | | | |

| | | | | |
|-----------------------------|-----|-------------|----------|-------------------------------|
| $0.159 \pm 0.053 \pm 0.020$ | f&a | ABBIENDI | 00D OPAL | 1990–1995 LEP runs |
| 0.238 ± 0.042 | avg | 158 BARATE | 98 ALEP | 1991–1995 LEP runs |
| $0.15 \pm 0.09 \pm 0.03$ | f&a | 4 159 BAUER | 94 TPC | $E_{\text{cm}}^{ee} = 29$ GeV |

158 Not independent of BARATE 98 $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$ and $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ values.

159 We multiply 0.15% by 0.20, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

$\Gamma(K^- K^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{95}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-----|------|-------------------------|----------|--|
| 0.161 ± 0.018 OUR FIT | | | | | |
| 0.151 ± 0.019 OUR AVERAGE | | | | | |
| 0.087 $\pm 0.056 \pm 0.040$ | avg | | ABBIENDI | 00D OPAL | 1990–1995 LEP runs |
| 0.145 $\pm 0.013 \pm 0.028$ | avg | 2.3k | ¹⁶⁰ RICHICHI | 99 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6$ GeV |
| 0.163 $\pm 0.021 \pm 0.017$ | f&a | | BARATE | 98 ALEP | 1991–1995 LEP runs |
| 0.22 $^{+0.17}_{-0.11} \pm 0.05$ | f&a | 9 | ¹⁶¹ MILLS | 85 DLCO | $E_{\text{cm}}^{\text{ee}} = 29$ GeV |
| 160 Not independent of RICHICHI 99 $\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau \text{ (ex. } K^0))$ and BAEST 95C $\Gamma(\tau^- \rightarrow h^- h^- h^+ \nu_\tau \text{ (ex. } K^0))/\Gamma_{\text{total}}$ values. | | | | | |
| 161 Error correlated with MILLS 85 ($K\pi\pi\pi^0\nu$) value. We multiply 0.22% by 0.23, the relative systematic error quoted by MILLS 85, to obtain obtain the systematic error. | | | | | |

 $\Gamma(K^- K^+ \pi^- \nu_\tau)/\Gamma(\pi^- \pi^+ \pi^- \nu_\tau \text{ (ex. } K^0))$ $\Gamma_{95}/\Gamma_{61} = \Gamma_{95}/(\Gamma_{63} + 0.0221\Gamma_{144})$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|---------|--|
| 1.75 ± 0.20 OUR FIT | | | | |
| $1.60 \pm 0.15 \pm 0.30$ | 2.3k | RICHICHI | 99 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6$ GeV |

 $\Gamma(K^- K^+ \pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{96}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------------------|---------|--|
| 4.0 ± 1.6 OUR FIT | | | | | |
| 4.4 ± 1.8 OUR AVERAGE Error includes scale factor of 1.1. | | | | | |
| 3.3 $\pm 1.8 \pm 0.7$ | avg | 158 | ¹⁶² RICHICHI | 99 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6$ GeV |
| 7.5 $\pm 2.9 \pm 1.5$ | f&a | | BARATE | 98 ALEP | 1991–1995 LEP runs |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----|----|----------|----------|--------------------|
| <27 | 95 | ABBIENDI | 00D OPAL | 1990–1995 LEP runs |
|-----|----|----------|----------|--------------------|

162 Not independent of RICHICHI 99

$\Gamma(\tau^- \rightarrow K^- K^+ \pi^- \nu_\tau)/\Gamma(\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau \text{ (ex. } K^0))$ and BAEST 95C $\Gamma(\tau^- \rightarrow h^- h^- h^+ \nu_\tau \text{ (ex. } K^0))/\Gamma_{\text{total}}$ values.

 $\Gamma(K^- K^+ \pi^- \pi^0 \nu_\tau)/\Gamma(\pi^- \pi^+ \pi^- \pi^0 \nu_\tau \text{ (ex. } K^0))$

$\Gamma_{96}/\Gamma_{70} = \Gamma_{96}/(\Gamma_{71} + 0.888\Gamma_{144} + 0.0221\Gamma_{145})$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------------------|---------|--|
| 1.0 ± 0.4 OUR FIT | | | | |
| $0.79 \pm 0.44 \pm 0.16$ | 158 | ¹⁶³ RICHICHI | 99 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6$ GeV |

163 RICHICHI 99 also quote a 95%CL upper limit of 0.0157 for this measurement.

$\Gamma(K^- K^+ K^- \geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{97}/Γ

| <u>VALUE (%)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|------------|--------------------|-------------|---|
| <0.21 | 95 | BAUER | 94 | TPC $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

 $\Gamma(K^- K^+ K^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{98}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|------------|--------------------|-------------|----------------------------|
| $<1.9 \times 10^{-4}$ | 90 | BARATE | 98 | ALEP 1991–1995 LEP runs |

 $\Gamma(\pi^- K^+ \pi^- \geq 0 \text{ neut. } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{99}/Γ

| <u>VALUE (%)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|------------|--------------------|-------------|---|
| <0.25 | 95 | BAUER | 94 | TPC $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

 $\Gamma(e^- e^- e^+ \bar{\nu}_e \nu_\tau)/\Gamma_{\text{total}}$ Γ_{100}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| $2.8 \pm 1.4 \pm 0.4$ | 5 | ALAM | 96 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 $\Gamma(\mu^- e^- e^+ \bar{\nu}_\mu \nu_\tau)/\Gamma_{\text{total}}$ Γ_{101}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| <3.6 | 90 | ALAM | 96 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 $\Gamma(3h^- 2h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ (ex. } K_S^0 \rightarrow \pi^- \pi^+ \text{ ("5-prong"))})/\Gamma_{\text{total}}$ Γ_{102}/Γ

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average. $\Gamma_{102}/\Gamma = (\Gamma_{103} + \Gamma_{104})/\Gamma$

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|---------------------------|---|
| 0.099 ± 0.007 OUR FIT | | | | |
| 0.107 ± 0.009 OUR AVERAGE | | | | |
| 0.119 $\pm 0.013 \pm 0.008$ | avg | 119 | ¹⁶⁴ ACKERSTAFF | 99E OPAL 1991–1995 LEP runs |
| 0.097 $\pm 0.005 \pm 0.011$ | f&a | 419 | GIBAUT | 94B CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| 0.26 $\pm 0.06 \pm 0.05$ | f&a | | ACTON | 92H OPAL $E_{\text{cm}}^{\text{ee}} = 88.2\text{--}94.2$ GeV |
| 0.10 $\begin{array}{l} +0.05 \\ -0.04 \end{array} \pm 0.03$ | f&a | | DECAMP | 92C ALEP 1989–1990 LEP runs |
| 0.102 ± 0.029 | f&a | 13 | BYLSMA | 87 HRS $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 0.16 $\pm 0.08 \pm 0.04$ | f&a | 4 | BURCHAT | 85 MRK2 $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.16 $\pm 0.13 \pm 0.04$ | | | BEHREND | 89B CELL $E_{\text{cm}}^{\text{ee}} = 14\text{--}47 \text{ GeV}$ |
| 0.3 $\pm 0.1 \pm 0.2$ | | | BARTEL | 85F JADE $E_{\text{cm}}^{\text{ee}} = 34.6 \text{ GeV}$ |
| 0.13 ± 0.04 | | 10 | BELTRAMI | 85 HRS Repl. by BYLSMA 87 |
| 1.0 ± 0.4 | | 10 | BEHREND | 82 CELL Repl. by BEHREND 89B |

¹⁶⁴ Not independent of ACKERSTAFF 99E $B(\tau^- \rightarrow 3h^- 2h^+ \nu_\tau \text{ (ex. } K^0))$ and $B(\tau^- \rightarrow 3h^- 2h^+ \pi^0 \nu_\tau \text{ (ex. } K^0))$ measurements.

$\Gamma(3h^- 2h^+ \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$ Γ_{103}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|----------------|--------|--|
| 0.078±0.006 OUR FIT | | | | |
| 0.076±0.007 OUR AVERAGE | | | | |
| 0.091±0.014±0.006 | 97 | ACKERSTAFF 99E | OPAL | 1991–1995 LEP runs |
| 0.080±0.011±0.013 | 58 | BUSKULIC 96 | ALEP | LEP 1991–1993 data |
| 0.077±0.005±0.009 | 295 | GIBAUT 94B | CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| 0.064±0.023±0.01 | 12 | ALBRECHT 88B | ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| 0.051±0.020 | 7 | BYLSMA 87 | HRS | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.067±0.030 | 5 | 165 BELTRAMI | 85 HRS | Repl. by BYLSMA 87 |

165 The error quoted is statistical only.

$\Gamma(3h^- 2h^+ \pi^0 \nu_\tau (\text{ex. } K^0)) / \Gamma_{\text{total}}$ Γ_{104}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|----------------|--------|--|
| 0.022±0.005 OUR FIT | | | | |
| 0.021±0.005 OUR AVERAGE | | | | |
| 0.027±0.018±0.009 | 23 | ACKERSTAFF 99E | OPAL | 1991–1995 LEP runs |
| 0.018±0.007±0.012 | 18 | BUSKULIC 96 | ALEP | LEP 1991–1993 data |
| 0.019±0.004±0.004 | 31 | GIBAUT 94B | CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| 0.051±0.022 | 6 | BYLSMA 87 | HRS | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.067±0.030 | 5 | 166 BELTRAMI | 85 HRS | Repl. by BYLSMA 87 |

166 The error quoted is statistical only.

$\Gamma(3h^- 2h^+ 2\pi^0 \nu_\tau) / \Gamma_{\text{total}}$ Γ_{105}/Γ

| VALUE (%) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|-------------|------|--|
| <0.011 | 90 | GIBAUT 94B | CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

$\Gamma((5\pi)^- \nu_\tau) / \Gamma_{\text{total}}$ Γ_{106}/Γ

$$\Gamma_{106}/\Gamma = (\Gamma_{29} + \frac{1}{4}\Gamma_{46} + \Gamma_{79} + \Gamma_{103} + 0.553\Gamma_{124} + 0.888\Gamma_{145})/\Gamma$$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

| VALUE (%) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-----|-------------|------|--|
| 0.79±0.07 OUR FIT | | | | |
| 0.61±0.06±0.08 avg | 167 | GIBAUT 94B | CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

167 Not independent of GIBAUT 94B $B(3h^- 2h^+ \nu_\tau)$, PROCARIO 93 $B(h^- 4\pi^0 \nu_\tau)$, and BORTOLETTO 93 $B(2h^- h^+ 2\pi^0 \nu_\tau)/B(\text{"3prong"})$ measurements. Result is corrected for η contributions.

$\Gamma(4h^- 3h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{"7-prong"}) / \Gamma_{\text{total}}$ Γ_{107}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|----------------|------|--|
| <2.4 × 10⁻⁶ | 90 | EDWARDS 97J | CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <1.8 × 10 ⁻⁵ | 95 | ACKERSTAFF 97J | OPAL | 1990–1995 LEP runs |
| <2.9 × 10 ⁻⁴ | 90 | BYLSMA 87 | HRS | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

$\Gamma(X^-(S=-1)\nu_\tau)/\Gamma_{\text{total}}$

$$\Gamma_{108}/\Gamma = (\Gamma_{11} + \Gamma_{16} + \Gamma_{23} + \Gamma_{27} + \Gamma_{35} + \Gamma_{40} + \Gamma_{88} + \Gamma_{92} + \Gamma_{126})/\Gamma$$

Data marked “avg” are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. “f&a” marks results used for the fit and the average.

| VALUE (%) | | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------------------------|-------------|------|---------|
| 2.89±0.09 OUR FIT | Error includes scale factor of 1.1. | | | |

2.87±0.12 avg 168 BARATE 99R ALEP 1991–1995 LEP runs

168 BARATE 99R perform a combined analysis of all ALEPH LEP 1 data on τ branching fraction measurements for decay modes having total strangeness equal to -1 .

$\Gamma(K^*(892)^- \geq 0 (h^0 \neq K_S^0)\nu_\tau)/\Gamma_{\text{total}}$

$$\Gamma_{109}/\Gamma$$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 1.94±0.27±0.15 | 74 | AKERS | 94G OPAL | $E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$ |

$\Gamma(K^*(892)^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$

$$\Gamma_{110}/\Gamma$$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|------|---------|
| 1.33±0.13 OUR AVERAGE | | | | |

$1.19 \pm 0.15^{+0.13}_{-0.18}$ 104 ALBRECHT 95H ARG $E_{\text{cm}}^{ee} = 9.4\text{--}10.6 \text{ GeV}$

$1.43 \pm 0.11 \pm 0.13$ 475 169 GOLDBERG 90 CLEO $E_{\text{cm}}^{ee} = 9.4\text{--}10.9 \text{ GeV}$

169 GOLDBERG 90 estimates that 10% of observed $K^*(892)$ are accompanied by a π^0 .

$\Gamma(K^*(892)^-\nu_\tau)/\Gamma_{\text{total}}$

$$\Gamma_{111}/\Gamma$$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-------------|------|---------|
| 1.29 ± 0.05 OUR AVERAGE | | | | |

1.326 ± 0.063 BARATE 99R ALEP 1991–1995 LEP runs

1.11 ± 0.12 170 COAN 96 CLEO $E_{\text{cm}}^{ee} \approx 10.6 \text{ GeV}$

$1.42 \pm 0.22 \pm 0.09$ 171 ACCIARRI 95F L3 1991–1993 LEP runs

$1.23 \pm 0.21^{+0.11}_{-0.21}$ 54 172 ALBRECHT 88L ARG $E_{\text{cm}}^{ee} = 10 \text{ GeV}$

$1.9 \pm 0.3 \pm 0.4$ 44 173 TSCHIRHART 88 HRS $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

$1.5 \pm 0.4 \pm 0.4$ 15 174 AIHARA 87C TPC $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

$1.3 \pm 0.3 \pm 0.3$ 31 YELTON 86 MRK2 $E_{\text{cm}}^{ee} = 29 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.39 \pm 0.09 \pm 0.10$ 175 BUSKULIC 96 ALEP Repl. by BARATE 99R

$1.45 \pm 0.13 \pm 0.11$ 273 176 BUSKULIC 94F ALEP Repl. by BUSKULIC 96

1.7 ± 0.7 11 DORFAN 81 MRK2 $E_{\text{cm}}^{ee} = 4.2\text{--}6.7 \text{ GeV}$

170 Not independent of COAN 96 $B(\pi^-\bar{K}^0\nu_\tau)$ and BATTLE 94 $B(K^-\pi^0\nu_\tau)$ measurements. $K\pi$ final states are consistent with and assumed to originate from $K^*(892)^-$ production.

171 This result is obtained from their $B(\pi^-\bar{K}^0\nu_\tau)$ assuming all those decays originate in $K^*(892)^-$ decays.

172 The authors divide by $\Gamma_1/\Gamma = 0.865$ to obtain this result.

173 Not independent of TSCHIRHART 88 $\Gamma(\tau^- \rightarrow h^-\bar{K}^0 \geq 0 \text{ neutrals} \geq 0 K_L^0\nu_\tau)/\Gamma(\text{total})$.

174 Decay π^- identified in this experiment, is assumed in the others.

175 Not independent of BUSKULIC 96 $B(\pi^-\bar{K}^0\nu_\tau)$ and $B(K^-\pi^0\nu_\tau)$ measurements.

176 BUSKULIC 94F obtain this result from BUSKULIC 94F $B(\bar{K}^0\pi^-\nu_\tau)$ and BUSKULIC 94E $B(K^-\pi^0\nu_\tau)$ assuming all of those decays originate in $K^*(892)^-$ decays.

$\Gamma(K^*(892)^-\nu_\tau)/\Gamma(\pi^-\pi^0\nu_\tau)$ Γ_{111}/Γ_{14}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|----------|-----------------|
| 0.075±0.027 | 177 ABREU | 94K DLPH | LEP 1992 Z data |
| 177 ABREU 94K quote $B(\tau^- \rightarrow K^*(892)^-\nu_\tau)B(K^*(892)^-\rightarrow K^-\pi^0)/B(\tau^- \rightarrow \rho^-\nu_\tau) = 0.025 \pm 0.009$. We divide by $B(K^*(892)^-\rightarrow K^-\pi^0) = 0.333$ to obtain this result. | | | |

 $\Gamma(K^*(892)^0 K^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{112}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|------|---|
| 0.32±0.08±0.12 | 119 | GOLDBERG | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.9 \text{ GeV}$ |

 $\Gamma(K^*(892)^0 K^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{113}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|---|
| 0.21 ± 0.04 OUR AVERAGE | | | | |
| 0.213±0.048 | | | | 178 BARATE 98 ALEP 1991–1995 LEP runs |
| 0.20 ± 0.05 ± 0.04 | | | | ALBRECHT 95H ARG $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ |
| 178 BARATE 98 measure the $K^- (\rho^0 \rightarrow \pi^+\pi^-)$ fraction in $\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau$ decays to be $(35 \pm 11)\%$ and derive this result from their measurement of $\Gamma(\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$ assuming the intermediate states are all $K^-\rho$ and $K^-K^*(892)^0$. | | | | |

 $\Gamma(\bar{K}^*(892)^0 \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{114}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|------|---|
| 0.38±0.11±0.13 | 105 | GOLDBERG | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.9 \text{ GeV}$ |

 $\Gamma(\bar{K}^*(892)^0 \pi^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{115}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| 0.22 ± 0.05 OUR AVERAGE | | | | |
| 0.209±0.058 | | | | 179 BARATE 98 ALEP 1991–1995 LEP runs |
| 0.25 ± 0.10 ± 0.05 | | | | ALBRECHT 95H ARG $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ |
| 179 BARATE 98 measure the $K^-K^*(892)^0$ fraction in $\tau^- \rightarrow K^-K^+\pi^-\nu_\tau$ decays to be $(87 \pm 13)\%$ and derive this result from their measurement of $\Gamma(\tau^- \rightarrow K^-K^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$. | | | | |

 $\Gamma((\bar{K}^*(892)\pi)^-\nu_\tau \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{116}/Γ

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------------|----------|--------------------|
| 0.10 ± 0.04 OUR AVERAGE | | | |
| 0.097±0.044±0.036 | 180 BARATE | 99K ALEP | 1991–1995 LEP runs |
| 0.106±0.037±0.032 | 181 BARATE | 98E ALEP | 1991–1995 LEP runs |

180 BARATE 99K measure K^0 's by detecting K_L^0 's in their hadron calorimeter. They determine the $\bar{K}^0\rho^-$ fraction in $\tau^- \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau$ decays to be $(0.72 \pm 0.12 \pm 0.10)$ and multiply their $B(\pi^-\bar{K}^0\pi^0\nu_\tau)$ measurement by one minus this fraction to obtain the quoted result.

181 BARATE 98E reconstruct K^0 's using $K_S^0 \rightarrow \pi^+\pi^-$ decays. They determine the $\bar{K}^0\rho^-$ fraction in $\tau^- \rightarrow \pi^-\bar{K}^0\pi^0\nu_\tau$ decays to be $(0.64 \pm 0.09 \pm 0.10)$ and multiply their $B(\pi^-\bar{K}^0\pi^0\nu_\tau)$ measurement by one minus this fraction to obtain the quoted result.

| $\Gamma(K_1(1270)^-\nu_\tau)/\Gamma_{\text{total}}$ | | Γ_{117}/Γ | | |
|---|------|-----------------------|----------|--------------------|
| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
| 0.47±0.11 OUR AVERAGE | | | | |
| 0.48±0.11 | | BARATE | 99R ALEP | 1991–1995 LEP runs |

182 We multiply 0.41% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

| $\Gamma(K_1(1400)^-\nu_\tau)/\Gamma_{\text{total}}$ | | Γ_{118}/Γ | | |
|---|------|-------------------------------------|----------|--|
| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
| 0.17±0.26 OUR AVERAGE | | Error includes scale factor of 1.7. | | |
| 0.05±0.17 | | BARATE | 99R ALEP | 1991–1995 LEP runs |
| 0.76 ^{+0.40} _{-0.33} ±0.20 | 11 | 183 BAUER | 94 TPC | $E_{\text{cm}}^{\text{ee}}=29 \text{ GeV}$ |

183 We multiply 0.76% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error.

| $[\Gamma(K_1(1270)^-\nu_\tau) + \Gamma(K_1(1400)^-\nu_\tau)]/\Gamma_{\text{total}}$ | | $(\Gamma_{117}+\Gamma_{118})/\Gamma$ | | |
|---|------|--------------------------------------|--------|--|
| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
| 1.17^{+0.41}_{-0.37}±0.29 | 16 | 184 BAUER | 94 TPC | $E_{\text{cm}}^{\text{ee}}=29 \text{ GeV}$ |

184 We multiply 1.17% by 0.25, the relative systematic error quoted by BAUER 94, to obtain the systematic error. Not independent of BAUER 94 $B(K_1(1270)^-\nu_\tau)$ and BAUER 94 $B(K_1(1400)^-\nu_\tau)$ measurements.

| $\Gamma(K_1(1270)^-\nu_\tau)/[\Gamma(K_1(1270)^-\nu_\tau) + \Gamma(K_1(1400)^-\nu_\tau)]$ | | $\Gamma_{117}/(\Gamma_{117}+\Gamma_{118})$ | | |
|---|-------------|--|---------|--|
| VALUE | DOCUMENT ID | TECN | COMMENT | |
| 0.69±0.15 OUR AVERAGE | | | | |

185 ABBIENDI 00D assume the resonance structure of $\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau$ decays is dominated by the $K_1(1270)^-$ and $K_1(1400)^-$ resonances.

186 ASNER 00B assume the resonance structure of $\tau^- \rightarrow K^-\pi^+\pi^-\nu_\tau$ (ex. K^0) decays is dominated by $K_1(1270)^-$ and $K_1(1400)^-$ resonances.

| $\Gamma(K^*(1410)^-\nu_\tau)/\Gamma_{\text{total}}$ | | Γ_{119}/Γ | | |
|---|-------------|-----------------------|--------------------|--|
| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT | |
| 1.5^{+1.4}_{-1.0} | BARATE | 99R ALEP | 1991–1995 LEP runs | |

| $\Gamma(K_0^*(1430)^-\nu_\tau)/\Gamma_{\text{total}}$ | | Γ_{120}/Γ | | |
|---|------|-----------------------|----------|--------------------|
| VALUE (units 10^{-3}) | CL % | DOCUMENT ID | TECN | COMMENT |
| <0.5 | 95 | BARATE | 99R ALEP | 1991–1995 LEP runs |

$\Gamma(K_2^*(1430)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{121}/Γ

| VALUE (%) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|-----|------|-------------|--------|--|
| <0.3 | 95 | | TSCHIRHART | 88 HRS | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------|----|-----|----------|---------|---|
| <0.33 | 95 | 187 | ACCIARRI | 95F L3 | 1991–1993 LEP runs |
| <0.9 | 95 | 0 | DORFAN | 81 MRK2 | $E_{\text{cm}}^{\text{ee}} = 4.2\text{--}6.7 \text{ GeV}$ |

187 ACCIARRI 95F quote $B(\tau^- \rightarrow K^*(1430)^- \rightarrow \pi^-\bar{K}^0\nu_\tau) < 0.11\%$. We divide by $B(K^*(1430)^- \rightarrow \pi^-\bar{K}^0) = 0.33$ to obtain the limit shown.

 $\Gamma(a_0(980)^-\geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}} \times B(a_0(980) \rightarrow K^0 K^-)$ $\Gamma_{122}/\Gamma \times B$

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|-------------|---------|--|
| <2.8 | 90 | | GOLDBERG | 90 CLEO | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.9 \text{ GeV}$ |

 $\Gamma(\eta\pi^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{123}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|-------------|---------|--|
| < 1.4 | 95 | 0 | BARTELT | 96 CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------------|----|---|----------|----------|---|
| < 6.2 | 95 | | BUSKULIC | 97C ALEP | 1991–1994 LEP runs |
| < 3.4 | 95 | | ARTUSO | 92 CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |
| < 90 | 95 | | ALBRECHT | 88M ARG | $E_{\text{cm}}^{\text{ee}} \approx 10 \text{ GeV}$ |
| <140 | 90 | | BEHREND | 88 CELL | $E_{\text{cm}}^{\text{ee}} = 14\text{--}46.8 \text{ GeV}$ |
| <180 | 95 | | BARINGER | 87 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.5 \text{ GeV}$ |
| <250 | 90 | 0 | COFFMAN | 87 MRK3 | $E_{\text{cm}}^{\text{ee}} = 3.77 \text{ GeV}$ |
| 510 $\pm 100 \pm 120$ | 65 | | DERRICK | 87 HRS | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| <100 | 95 | | GAN | 87B MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

 $\Gamma(\eta\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{124}/Γ

| VALUE (%) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------|------|---------|
| 0.174\pm0.024 OUR FIT | | | | | |
| 0.173\pm0.024 OUR AVERAGE | | | | | |

| | | | | | |
|--------------------------|-----|--|----------|----------|--|
| 0.18 $\pm 0.04 \pm 0.02$ | | | BUSKULIC | 97C ALEP | 1991–1994 LEP runs |
| 0.17 $\pm 0.02 \pm 0.02$ | 125 | | ARTUSO | 92 CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|----------------------------------|----|---------|----------|---------|--|
| <1.10 | 95 | | ALBRECHT | 88M ARG | $E_{\text{cm}}^{\text{ee}} \approx 10 \text{ GeV}$ |
| <2.10 | 95 | | BARINGER | 87 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.5 \text{ GeV}$ |
| 4.20 $^{+0.70}_{-1.20} \pm 1.60$ | | 188 GAN | | 87 MRK2 | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

188 Highly correlated with GAN 87 $\Gamma(\pi^- 3\pi^0\nu_\tau)/\Gamma(\text{total})$ value.

$\Gamma(\eta\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{125}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|-------------|--------------------|-------------|---|
| $1.4 \pm 0.6 \pm 0.3$ | | 15 | BERGFELD | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| < 4.3 | 95 | | ARTUSO | 92 | CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |
| <120 | 95 | | ALBRECHT | 88M ARG | $E_{\text{cm}}^{\text{ee}} \approx 10 \text{ GeV}$ |

$\Gamma(\eta K^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{126}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|-------------|--------------------|-------------|--|
| $2.7 \pm 0.6 \text{ OUR FIT}$ | | | | | |
| $2.7 \pm 0.6 \text{ OUR AVERAGE}$ | | | | | |
| $2.9^{+1.3}_{-1.2} \pm 0.7$ | | | BUSKULIC | 97C ALEP | 1991–1994 LEP runs |
| $2.6 \pm 0.5 \pm 0.5$ | 85 | | BARTEL | 96 CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <4.7 | 95 | | ARTUSO | 92 CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |

$\Gamma(\eta K^*(892)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{127}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| $2.90 \pm 0.80 \pm 0.42$ | 25 | BISHAI | 99 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

$\Gamma(\eta K^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{128}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| $1.77 \pm 0.56 \pm 0.71$ | 36 | BISHAI | 99 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

$\Gamma(\eta \bar{K}^0 \pi^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{129}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| $2.20 \pm 0.70 \pm 0.22$ | 15 | 189 BISHAI | 99 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

189 We multiply the BISHAI 99 measurement $B(\tau^- \rightarrow \eta K_S^0 \pi^- \nu_\tau) = (1.10 \pm 0.35 \pm 0.11) \times 10^{-4}$ by 2 to obtain the listed value.

$\Gamma(\eta\pi^+\pi^-\pi^-\geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{130}/Γ

| <u>VALUE (%)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|------------|--------------------|-------------|--|
| <0.3 | 90 | ABACHI | 87B HRS | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |

$\Gamma(\eta\pi^-\pi^+\pi^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{131}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| $3.4^{+0.6}_{-0.5} \pm 0.6$ | 89 | BERGFELD | 97 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

$\Gamma(\eta a_1(1260)^-\nu_\tau \rightarrow \eta\pi^-\rho^0\nu_\tau)/\Gamma_{\text{total}}$ Γ_{132}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|--------------------|-------------|--|
| $<3.9 \times 10^{-4}$ | 90 | BERGFELD | 97 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

$\Gamma(\eta\eta\pi^-\nu_\tau)/\Gamma_{\text{total}}$

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| < 1.1 | 95 | ARTUSO | 92 | CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <83 | 95 | ALBRECHT | 88M ARG | $E_{\text{cm}}^{\text{ee}} \approx 10 \text{ GeV}$ |

 Γ_{133}/Γ $\Gamma(\eta\eta\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| < 2.0 | 95 | ARTUSO | 92 | CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <90 | 95 | ALBRECHT | 88M ARG | $E_{\text{cm}}^{\text{ee}} \approx 10 \text{ GeV}$ |

 Γ_{134}/Γ $\Gamma(\eta'(958)\pi^-\nu_\tau)/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|------------|--------------------|-------------|---|
| < 7.4×10^{-5} | 90 | BERGFELD | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 Γ_{135}/Γ $\Gamma(\eta'(958)\pi^-\pi^0\nu_\tau)/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|------------|--------------------|-------------|---|
| < 8.0×10^{-5} | 90 | BERGFELD | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 Γ_{136}/Γ $\Gamma(\phi\pi^-\nu_\tau)/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| < 2.0×10^{-4} | 90 | 190 AVERY | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |

 Γ_{137}/Γ $\Gamma(\phi K^-\nu_\tau)/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| < 6.7×10^{-5} | 90 | 191 AVERY | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| 191 Avery 97 limit varies from $(5.4\text{--}6.7) \times 10^{-5}$ depending on decay model assumptions. | | | | |

 Γ_{138}/Γ $\Gamma(f_1(1285)\pi^-\nu_\tau)/\Gamma_{\text{total}}$

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|---|
| $5.8^{+1.4}_{-1.3} \pm 1.8$ | 54 | BERGFELD | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 Γ_{139}/Γ $\Gamma(f_1(1285)\pi^-\nu_\tau \rightarrow \eta\pi^-\pi^+\pi^-\nu_\tau)/\Gamma(\eta\pi^-\pi^+\pi^-\nu_\tau)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|--------------------|-------------|---|
| 0.55 ± 0.14 | BERGFELD | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 $\Gamma_{140}/\Gamma_{131}$ $\Gamma(\pi(1300)^-\nu_\tau \rightarrow (\rho\pi)^-\nu_\tau \rightarrow (3\pi)^-\nu_\tau)/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|------------|--------------------|-------------|---|
| < 1.0×10^{-4} | 90 | ASNER | 00 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 Γ_{141}/Γ

$\Gamma(\pi(1300)^-\nu_\tau \rightarrow ((\pi\pi)_S\text{-wave}\pi)^-\nu_\tau \rightarrow (3\pi)^-\nu_\tau)/\Gamma_{\text{total}}$ Γ_{142}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|------|---|
| $<1.9 \times 10^{-4}$ | 90 | ASNER | 00 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

$\Gamma(h^-\omega \geq 0 \text{ neutrals} \nu_\tau)/\Gamma_{\text{total}}$ Γ_{143}/Γ

$$\Gamma_{143}/\Gamma = (\Gamma_{144} + \Gamma_{145})/\Gamma$$

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

2.36±0.08 OUR FIT

| | | | | |
|-----------------|-------------|------------|------|---|
| 1.65±0.3 | ±0.2 | avg | 1513 | ALBRECHT 88M ARG $E_{\text{cm}}^{\text{ee}} \approx 10 \text{ GeV}$ |
|-----------------|-------------|------------|------|---|

$\Gamma(h^-\omega\nu_\tau)/\Gamma_{\text{total}}$ Γ_{144}/Γ

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

1.93±0.06 OUR FIT

1.92±0.07 OUR AVERAGE

| | | | | |
|----------------|-----|------|------------|---|
| 1.91±0.07±0.06 | f&a | 5803 | BUSKULIC | 97C ALEP LEP runs |
| 1.95±0.07±0.11 | avg | 2223 | 192 BALEST | 95C CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |
| 1.60±0.27±0.41 | f&a | 139 | BARINGER | 87 CLEO $E_{\text{cm}}^{\text{ee}} = 10.5 \text{ GeV}$ |

192 Not independent of BALEST 95C $B(\tau^- \rightarrow h^-\omega\nu_\tau)/B(\tau^- \rightarrow h^-h^-h^+\pi^0\nu_\tau)$ value.

$$\frac{[\Gamma(h^-\rho\pi^0\nu_\tau) + \Gamma(h^-\rho^+h^-\nu_\tau) + \Gamma(h^-\rho^-h^+\nu_\tau) + \Gamma(h^-\omega\nu_\tau)]}{\Gamma(h^-h^-h^+\pi^0\nu_\tau)} / (\Gamma_{74} + \Gamma_{75} + \Gamma_{76} + \Gamma_{144})/\Gamma_{66}$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------|-----|--------------|---------|--|
| >0.81 | 95 | 193 ALBRECHT | 91D ARG | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ |

193 ALBRECHT 91D not independent of their $\Gamma(h^-\omega\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ (ex. K^0), $\Gamma(h^-\rho\pi^0\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$, $\Gamma(h^-\rho^+h^-\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$, and $\Gamma(h^-\rho^-h^+\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau)$ values.

$\Gamma(h^-\omega\nu_\tau)/\Gamma(h^-h^-h^+\pi^0\nu_\tau \text{ (ex. } K^0\text{)})$ Γ_{144}/Γ_{67}

$$\Gamma_{144}/\Gamma_{67} = \Gamma_{144}/(\Gamma_{71} + \Gamma_{92} + \Gamma_{96} + 0.231\Gamma_{126} + 0.888\Gamma_{144} + 0.0221\Gamma_{145})$$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.448±0.015 OUR FIT

0.453±0.019 OUR AVERAGE

| | | | | |
|-------------------|------|--------------|----------|--|
| 0.431±0.033 | 2350 | 194 BUSKULIC | 96 ALEP | LEP 1991–1993 data |
| 0.464±0.016±0.017 | 2223 | 195 BALEST | 95C CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------|-----|--------------|---------|--|
| 0.37 ± 0.05 ± 0.02 | 458 | 196 ALBRECHT | 91D ARG | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ |
|--------------------|-----|--------------|---------|--|

194 BUSKULIC 96 quote the fraction of $\tau^- \rightarrow h^-h^-h^+\pi^0\nu_\tau$ (ex. K^0) decays which originate in a $h^-\omega$ final state = 0.383 ± 0.029 . We divide this by the $\omega(782) \rightarrow \pi^+\pi^-\pi^0$ branching fraction (0.888).

195 BAEST 95C quote the fraction of $\tau^- \rightarrow h^- h^- h^+ \pi^0 \nu_\tau$ (ex. K^0) decays which originate in a $h^- \omega$ final state equals $0.412 \pm 0.014 \pm 0.015$. We divide this by the $\omega(782) \rightarrow \pi^+ \pi^- \pi^0$ branching fraction (0.888).

196 ALBRECHT 91D quote the fraction of $\tau^- \rightarrow h^- h^- h^+ \pi^0 \nu_\tau$ decays which originate in a $\pi^- \omega$ final state equals $0.33 \pm 0.04 \pm 0.02$. We divide this by the $\omega(782) \rightarrow \pi^+ \pi^- \pi^0$ branching fraction (0.888).

| $\Gamma(h^- \omega \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ | Γ_{145}/Γ | | | |
|---|-----------------------|--------------------|-------------|--------------------|
| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.43 ± 0.05 OUR FIT | | | | |
| 0.43 ± 0.06 ± 0.05 | 7283 | BUSKULIC | 97C ALEP | 1991–1994 LEP runs |

| $\Gamma(h^- \omega 2\pi^0 \nu_\tau)/\Gamma_{\text{total}}$ | Γ_{146}/Γ | | | |
|--|-----------------------|--------------------|-------------|--|
| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| $1.89^{+0.74}_{-0.67} \pm 0.40$ | 19 | ANDERSON | 97 CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

| $\Gamma(h^- \omega \pi^0 \nu_\tau)/\Gamma(h^- h^- h^+ \geq 1 \text{ neut. } \nu_\tau \text{ ("3-prong")})$ | Γ_{145}/Γ_{54} | | | |
|--|----------------------------|--------------------|------------------|---|
| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.0282 ± 0.0031 OUR FIT | | | | |
| 0.028 ± 0.003 ± 0.003 | avg | 430 | 197 BORTOLETTO93 | CLEO $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |

197 Not independent of BORTOLETTO 93 $\Gamma(\tau^- \rightarrow h^- \omega \pi^0 \nu_\tau)/\Gamma(\tau^- \rightarrow h^- h^- h^+ 2\pi^0 \nu_\tau \text{ (ex. } K^0\text{)})$ value.

| $\Gamma(h^- \omega \pi^0 \nu_\tau)/\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau \text{ (ex. } K^0\text{)})$ | Γ_{145}/Γ_{78} | | | |
|---|----------------------------|--------------------|-------------|--|
| <u>VALUE</u> | <u>CL %</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.81 ± 0.08 OUR FIT | | | | |
| 0.81 ± 0.06 ± 0.06 | | BORTOLETTO93 | CLEO | $E_{\text{cm}}^{\text{ee}} \approx 10.6 \text{ GeV}$ |

| $\Gamma(e^- \gamma)/\Gamma_{\text{total}}$ | Γ_{147}/Γ | | | |
|---|-----------------------|--------------------|-------------|---|
| Test of lepton family number conservation. | | | | |
| <u>VALUE</u> | <u>CL %</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| $<2.7 \times 10^{-6}$ | 90 | EDWARDS | 97 CLEO | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $<1.1 \times 10^{-4}$ | 90 | ABREU | 95U DLPH | 1990–1993 LEP runs |
| $<1.2 \times 10^{-4}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $<2.0 \times 10^{-4}$ | 90 | KEH | 88 CBAL | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $<6.4 \times 10^{-4}$ | 90 | HAYES | 82 MRK2 | $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$ |

$\Gamma(\mu^- \gamma)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

 Γ_{148}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|--|
| $< 1.1 \times 10^{-6}$ | 90 | AHMED | 00 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 3.0 \times 10^{-6}$ | 90 | EDWARDS | 97 | CLEO |
| $< 6.2 \times 10^{-5}$ | 90 | ABREU | 95U | DLPH 1990–1993 LEP runs |
| $< 0.42 \times 10^{-5}$ | 90 | BEAN | 93 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $< 3.4 \times 10^{-5}$ | 90 | ALBRECHT | 92K | ARG $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 55 \times 10^{-5}$ | 90 | HAYES | 82 | MRK2 $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$ |

 $\Gamma(e^- \pi^0)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

 Γ_{149}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|--|
| $< 3.7 \times 10^{-6}$ | 90 | BONVICINI | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 17 \times 10^{-5}$ | 90 | ALBRECHT | 92K | ARG $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 14 \times 10^{-5}$ | 90 | KEH | 88 | CBAL $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 210 \times 10^{-5}$ | 90 | HAYES | 82 | MRK2 $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$ |

 $\Gamma(\mu^- \pi^0)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

 Γ_{150}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|--|
| $< 4.0 \times 10^{-6}$ | 90 | BONVICINI | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 4.4 \times 10^{-5}$ | 90 | ALBRECHT | 92K | ARG $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 82 \times 10^{-5}$ | 90 | HAYES | 82 | MRK2 $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$ |

 $\Gamma(e^- K^0)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

 Γ_{151}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|-------------|------|--|
| $< 1.3 \times 10^{-3}$ | 90 | HAYES | 82 | MRK2 $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$ |

 $\Gamma(\mu^- K^0)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

 Γ_{152}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|-------------|------|--|
| $< 1.0 \times 10^{-3}$ | 90 | HAYES | 82 | MRK2 $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$ |

 $\Gamma(e^- \eta)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

 Γ_{153}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|---|
| $< 8.2 \times 10^{-6}$ | 90 | BONVICINI | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 6.3 \times 10^{-5}$ | 90 | ALBRECHT | 92K | ARG $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 24 \times 10^{-5}$ | 90 | KEH | 88 | CBAL $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |

$\Gamma(\mu^- \eta)/\Gamma_{\text{total}}$ Γ_{154}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| <9.6 × 10⁻⁶ | 90 | BONVICINI | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $<7.3 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |

 $\Gamma(e^- \rho^0)/\Gamma_{\text{total}}$ Γ_{155}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| < 2.0 × 10⁻⁶ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $< 0.42 \times 10^{-5}$ | 90 | 198 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 1.9 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 37 \times 10^{-5}$ | 90 | HAYES | 82 | MRK2 $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$ |

198 BARTEL 94 assume phase space decays.

 $\Gamma(\mu^- \rho^0)/\Gamma_{\text{total}}$ Γ_{156}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| < 6.3 × 10⁻⁶ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $< 0.57 \times 10^{-5}$ | 90 | 199 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 2.9 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 44 \times 10^{-5}$ | 90 | HAYES | 82 | MRK2 $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$ |

199 BARTEL 94 assume phase space decays.

 $\Gamma(e^- K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{157}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| <5.1 × 10⁻⁶ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $<0.63 \times 10^{-5}$ | 90 | 200 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $<3.8 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |

200 BARTEL 94 assume phase space decays.

 $\Gamma(\mu^- K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{158}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| <7.5 × 10⁻⁶ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $<0.94 \times 10^{-5}$ | 90 | 201 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $<4.5 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |

201 BARTEL 94 assume phase space decays.

$\Gamma(e^- \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{159}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| $<7.4 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $<1.1 \times 10^{-5}$ | 90 | 202 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| 202 BARTEL 94 assume phase space decays. | | | | |

 $\Gamma(\mu^- \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{160}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| $<7.5 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $<0.87 \times 10^{-5}$ | 90 | 203 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| 203 BARTEL 94 assume phase space decays. | | | | |

 $\Gamma(e^- \phi)/\Gamma_{\text{total}}$ Γ_{161}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|------------|--------------------|-------------|---|
| $<6.9 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 $\Gamma(\mu^- \phi)/\Gamma_{\text{total}}$ Γ_{162}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|------------|--------------------|-------------|---|
| $<7.0 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 $\Gamma(\pi^- \gamma)/\Gamma_{\text{total}}$ Γ_{163}/Γ

Test of lepton number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|--------------------|-------------|--|
| $<28 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |

 $\Gamma(\pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{164}/Γ

Test of lepton number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|--------------------|-------------|--|
| $<37 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |

 $\Gamma(e^- e^+ e^-)/\Gamma_{\text{total}}$ Γ_{165}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| $< 2.9 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 0.33 \times 10^{-5}$ | 90 | 204 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 1.3 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |

 $< 2.7 \times 10^{-5}$ 90 BOWCOCK 90 CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ $< 40 \times 10^{-5}$ 90 HAYES 82 MRK2 $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$

204 BARTEL 94 assume phase space decays.

$\Gamma(e^- \mu^+ \mu^-)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

 Γ_{166}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| $< 1.8 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 0.36 \times 10^{-5}$ | 90 | 205 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 1.9 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 2.7 \times 10^{-5}$ | 90 | BOWCOCK | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ |
| $< 33 \times 10^{-5}$ | 90 | HAYES | 82 | MRK2 $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$ |

205 BARTEL 94 assume phase space decays.

 $\Gamma(e^+ \mu^- \mu^-)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

 Γ_{167}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| $< 1.5 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 0.35 \times 10^{-5}$ | 90 | 206 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 1.8 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 1.6 \times 10^{-5}$ | 90 | BOWCOCK | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ |

206 BARTEL 94 assume phase space decays.

 $\Gamma(\mu^- e^+ e^-)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

 Γ_{168}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| $< 1.7 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 0.34 \times 10^{-5}$ | 90 | 207 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 1.4 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 2.7 \times 10^{-5}$ | 90 | BOWCOCK | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ |
| $< 44 \times 10^{-5}$ | 90 | HAYES | 82 | MRK2 $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$ |

207 BARTEL 94 assume phase space decays.

 $\Gamma(\mu^+ e^- e^-)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

 Γ_{169}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| $< 1.5 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 0.34 \times 10^{-5}$ | 90 | 208 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 1.4 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 1.6 \times 10^{-5}$ | 90 | BOWCOCK | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ |

208 BARTEL 94 assume phase space decays.

$\Gamma(\mu^- \mu^+ \mu^-)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

Γ_{170}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|---------|--|
| $< 1.9 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 0.43 \times 10^{-5}$ | 90 | 209 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 1.9 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 1.7 \times 10^{-5}$ | 90 | BOWCOCK | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ |
| $< 49 \times 10^{-5}$ | 90 | HAYES | 82 | MRK2 $E_{\text{cm}}^{\text{ee}} = 3.8\text{--}6.8 \text{ GeV}$ |

209 BARTEL 94 assume phase space decays.

$\Gamma(e^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

Γ_{171}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|---------|--|
| $< 2.2 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 0.44 \times 10^{-5}$ | 90 | 210 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 2.7 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 6.0 \times 10^{-5}$ | 90 | BOWCOCK | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ |

210 BARTEL 94 assume phase space decays.

$\Gamma(e^+ \pi^- \pi^-)/\Gamma_{\text{total}}$

Test of lepton number conservation.

Γ_{172}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|---------|--|
| $< 1.9 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 0.44 \times 10^{-5}$ | 90 | 211 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 1.8 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 1.7 \times 10^{-5}$ | 90 | BOWCOCK | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ |

211 BARTEL 94 assume phase space decays.

$\Gamma(\mu^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

Γ_{173}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|---------|--|
| $< 8.2 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 0.74 \times 10^{-5}$ | 90 | 212 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 3.6 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 3.9 \times 10^{-5}$ | 90 | BOWCOCK | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ |

212 BARTEL 94 assume phase space decays.

$\Gamma(\mu^+ \pi^- \pi^-)/\Gamma_{\text{total}}$

Test of lepton number conservation.

Γ_{174}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|-------------|------|---|
| $< 3.4 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------|----|--------------|---------|-----------------------------------|
| $<0.69 \times 10^{-5}$ | 90 | 213 BARTELTT | 94 CLEO | Repl. by BLISS 98 |
| $<6.3 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{cm}^{ee} = 10$ GeV |
| $<3.9 \times 10^{-5}$ | 90 | BOWCOCK | 90 CLEO | $E_{cm}^{ee} = 10.4\text{--}10.9$ |

213 BARTELTT 94 assume phase space decays.

$\Gamma(e^- \pi^+ K^-)/\Gamma_{total}$

Γ_{175}/Γ

Test of lepton family number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|---------|--------------------------|
| $<6.4 \times 10^{-6}$ | 90 | BLISS | 98 CLEO | $E_{cm}^{ee} = 10.6$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------|----|--------------|---------|-----------------------------------|
| $<0.77 \times 10^{-5}$ | 90 | 214 BARTELTT | 94 CLEO | Repl. by BLISS 98 |
| $<2.9 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{cm}^{ee} = 10$ GeV |
| $<5.8 \times 10^{-5}$ | 90 | BOWCOCK | 90 CLEO | $E_{cm}^{ee} = 10.4\text{--}10.9$ |

214 BARTELTT 94 assume phase space decays.

$\Gamma(e^- \pi^- K^+)/\Gamma_{total}$

Γ_{176}/Γ

Test of lepton family number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|---------|--------------------------|
| $<3.8 \times 10^{-6}$ | 90 | BLISS | 98 CLEO | $E_{cm}^{ee} = 10.6$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------|----|--------------|---------|-----------------------------------|
| $<0.46 \times 10^{-5}$ | 90 | 215 BARTELTT | 94 CLEO | Repl. by BLISS 98 |
| $<5.8 \times 10^{-5}$ | 90 | BOWCOCK | 90 CLEO | $E_{cm}^{ee} = 10.4\text{--}10.9$ |

215 BARTELTT 94 assume phase space decays.

$\Gamma(e^+ \pi^- K^-)/\Gamma_{total}$

Γ_{177}/Γ

Test of lepton number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|---------|--------------------------|
| $<2.1 \times 10^{-6}$ | 90 | BLISS | 98 CLEO | $E_{cm}^{ee} = 10.6$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------|----|--------------|---------|-----------------------------------|
| $<0.45 \times 10^{-5}$ | 90 | 216 BARTELTT | 94 CLEO | Repl. by BLISS 98 |
| $<2.0 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{cm}^{ee} = 10$ GeV |
| $<4.9 \times 10^{-5}$ | 90 | BOWCOCK | 90 CLEO | $E_{cm}^{ee} = 10.4\text{--}10.9$ |

216 BARTELTT 94 assume phase space decays.

$\Gamma(e^- K^+ K^-)/\Gamma_{total}$

Γ_{178}/Γ

Test of lepton family number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|---------|--------------------------|
| $<6.0 \times 10^{-6}$ | 90 | BLISS | 98 CLEO | $E_{cm}^{ee} = 10.6$ GeV |

$\Gamma(e^+ K^- K^-)/\Gamma_{total}$

Γ_{179}/Γ

Test of lepton number conservation.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|---------|--------------------------|
| $<3.8 \times 10^{-6}$ | 90 | BLISS | 98 CLEO | $E_{cm}^{ee} = 10.6$ GeV |

$\Gamma(\mu^- \pi^+ K^-)/\Gamma_{\text{total}}$ Γ_{180}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| $< 7.5 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 0.87 \times 10^{-5}$ | 90 | 217 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 11 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 7.7 \times 10^{-5}$ | 90 | BOWCOCK | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ |

217 BARTEL 94 assume phase space decays.

 $\Gamma(\mu^- \pi^- K^+)/\Gamma_{\text{total}}$ Γ_{181}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| $< 7.4 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 1.5 \times 10^{-5}$ | 90 | 218 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 7.7 \times 10^{-5}$ | 90 | BOWCOCK | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ |

218 BARTEL 94 assume phase space decays.

 $\Gamma(\mu^+ \pi^- K^-)/\Gamma_{\text{total}}$ Γ_{182}/Γ

Test of lepton number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| $< 7.0 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $< 2.0 \times 10^{-5}$ | 90 | 219 BARTEL | 94 | CLEO Repl. by BLISS 98 |
| $< 5.8 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |
| $< 4.0 \times 10^{-5}$ | 90 | BOWCOCK | 90 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.4\text{--}10.9$ |

219 BARTEL 94 assume phase space decays.

 $\Gamma(\mu^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{183}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|------------|--------------------|-------------|---|
| $< 15 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 $\Gamma(\mu^+ K^- K^-)/\Gamma_{\text{total}}$ Γ_{184}/Γ

Test of lepton number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|------------|--------------------|-------------|---|
| $< 6.0 \times 10^{-6}$ | 90 | BLISS | 98 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 $\Gamma(e^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{185}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|------------|--------------------|-------------|---|
| $< 6.5 \times 10^{-6}$ | 90 | BONVICINI | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 $\Gamma(\mu^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{186}/Γ

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|------------|--------------------|-------------|---|
| $< 14 \times 10^{-6}$ | 90 | BONVICINI | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

$\Gamma(e^-\eta\eta)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|--------------------|-------------|---|
| $<35 \times 10^{-6}$ | 90 | BONVICINI | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 Γ_{187}/Γ $\Gamma(\mu^-\eta\eta)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|--------------------|-------------|---|
| $<60 \times 10^{-6}$ | 90 | BONVICINI | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 Γ_{188}/Γ $\Gamma(e^-\pi^0\eta)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|--------------------|-------------|---|
| $<24 \times 10^{-6}$ | 90 | BONVICINI | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 Γ_{189}/Γ $\Gamma(\mu^-\pi^0\eta)/\Gamma_{\text{total}}$

Test of lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|--------------------|-------------|---|
| $<22 \times 10^{-6}$ | 90 | BONVICINI | 97 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 Γ_{190}/Γ $\Gamma(\bar{p}\gamma)/\Gamma_{\text{total}}$

Test of lepton number and baryon number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| $< 3.5 \times 10^{-6}$ | 90 | GODANG | 99 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $<29 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |

 Γ_{191}/Γ $\Gamma(\bar{p}\pi^0)/\Gamma_{\text{total}}$

Test of lepton number and baryon number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| $<15 \times 10^{-6}$ | 90 | GODANG | 99 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $<66 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |

 Γ_{192}/Γ $\Gamma(\bar{p}2\pi^0)/\Gamma_{\text{total}}$

Test of lepton number and baryon number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|--------------------|-------------|---|
| $<33 \times 10^{-6}$ | 90 | GODANG | 99 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 Γ_{193}/Γ $\Gamma(\bar{p}\eta)/\Gamma_{\text{total}}$

Test of lepton number and baryon number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| $< 8.9 \times 10^{-6}$ | 90 | GODANG | 99 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $<130 \times 10^{-5}$ | 90 | ALBRECHT | 92K ARG | $E_{\text{cm}}^{\text{ee}} = 10 \text{ GeV}$ |

 Γ_{194}/Γ $\Gamma(\bar{p}\pi^0\eta)/\Gamma_{\text{total}}$

Test of lepton number and baryon number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|--------------------|-------------|---|
| $<27 \times 10^{-6}$ | 90 | GODANG | 99 | CLEO $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |

 Γ_{195}/Γ

$\Gamma(e^- \text{ light boson})/\Gamma(e^- \bar{\nu}_e \nu_\tau)$

Test of lepton family number conservation.

 Γ_{196}/Γ_5

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|--------------------|---------|--|
| <0.015 | 95 | 220 ALBRECHT | 95G ARG | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <0.018 | 95 | 221 ALBRECHT | 90E ARG | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ |
| <0.040 | 95 | 222 BALTRUSAIT..85 | MRK3 | $E_{\text{cm}}^{\text{ee}} = 3.77 \text{ GeV}$ |
| 220 ALBRECHT 95G limit holds for bosons with mass < 0.4 GeV. The limit rises to 0.036 for a mass of 1.0 GeV, then falls to 0.006 at the upper mass limit of 1.6 GeV. | | | | |
| 221 ALBRECHT 90E limit applies for spinless boson with mass < 100 MeV, and rises to 0.050 for mass = 500 MeV. | | | | |
| 222 BALTRUSAITIS 85 limit applies for spinless boson with mass < 100 MeV. | | | | |

 $\Gamma(\mu^- \text{ light boson})/\Gamma(e^- \bar{\nu}_e \nu_\tau)$

Test of lepton family number conservation.

 Γ_{197}/Γ_5

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|--------------------|---------|--|
| <0.026 | 95 | 223 ALBRECHT | 95G ARG | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| <0.033 | 95 | 224 ALBRECHT | 90E ARG | $E_{\text{cm}}^{\text{ee}} = 9.4\text{--}10.6 \text{ GeV}$ |
| <0.125 | 95 | 225 BALTRUSAIT..85 | MRK3 | $E_{\text{cm}}^{\text{ee}} = 3.77 \text{ GeV}$ |
| 223 ALBRECHT 95G limit holds for bosons with mass < 1.3 GeV. The limit rises to 0.034 for a mass of 1.4 GeV, then falls to 0.003 at the upper mass limit of 1.6 GeV. | | | | |
| 224 ALBRECHT 90E limit applies for spinless boson with mass < 100 MeV, and rises to 0.071 for mass = 500 MeV. | | | | |
| 225 BALTRUSAITIS 85 limit applies for spinless boson with mass < 100 MeV. | | | | |

 τ -DECAY PARAMETERS

A REVIEW GOES HERE – Check our WWW List of Reviews

 $\rho^\tau(e \text{ or } \mu)$ PARAMETER(V-A) theory predicts $\rho = 0.75$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|--------------|----------|--|
| 0.747±0.009 OUR FIT | | | | |
| 0.751±0.009 OUR AVERAGE | | | | |
| 0.775±0.023±0.020 | 36k | ABREU | 00L DLPH | 1992–1995 runs |
| 0.781±0.028±0.018 | 46k | ACKERSTAFF | 99D OPAL | 1990–1995 LEP runs |
| 0.762±0.035 | 54k | ACCIARRI | 98R L3 | 1991–1995 LEP runs |
| 0.731±0.031 | | 226 ALBRECHT | 98 ARG | $E_{\text{cm}}^{\text{ee}} = 9.5\text{--}10.6 \text{ GeV}$ |
| 0.72 ± 0.09 ± 0.03 | | 227 ABE | 970 SLD | 1993–1995 SLC runs |
| 0.747±0.010±0.006 | 55k | ALEXANDER | 97F CLEO | $E_{\text{cm}}^{\text{ee}} = 10.6 \text{ GeV}$ |
| 0.751±0.039±0.022 | | BUSKULIC | 95D ALEP | 1990–1992 LEP runs |
| 0.79 ± 0.10 ± 0.10 | 3732 | FORD | 87B MAC | $E_{\text{cm}}^{\text{ee}} = 29 \text{ GeV}$ |
| 0.71 ± 0.09 ± 0.03 | 1426 | BEHRENDS | 85 CLEO | $e^+ e^-$ near $\gamma(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------------|------|--------------|----------|--|
| $0.735 \pm 0.013 \pm 0.008$ | 31k | AMMAR | 97B CLEO | Repl. by ALEXANDER 97F |
| $0.794 \pm 0.039 \pm 0.031$ | 18k | ACCIARRI | 96H L3 | Repl. by ACCIARRI 98R |
| $0.732 \pm 0.034 \pm 0.020$ | 8.2k | 228 ALBRECHT | 95 ARG | $E_{cm}^{ee} = 9.5\text{--}10.6 \text{ GeV}$ |
| 0.738 ± 0.038 | | 229 ALBRECHT | 95C ARG | Repl. by ALBRECHT 98 |
| $0.742 \pm 0.035 \pm 0.020$ | 8000 | ALBRECHT | 90E ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |

226 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

227 ABE 970 assume $\eta^\tau = 0$ in their fit. Letting η^τ vary in the fit gives a ρ^τ value of $0.69 \pm 0.13 \pm 0.05$.

228 Value is from a simultaneous fit for the ρ^τ and η^τ decay parameters to the lepton energy spectrum. Not independent of ALBRECHT 90E $\rho^\tau(e \text{ or } \mu)$ value which assumes $\eta^\tau=0$. Result is strongly correlated with ALBRECHT 95C.

229 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E.

$\rho^\tau(e)$ PARAMETER

(V-A) theory predicts $\rho = 0.75$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|--------------|----------|--|
| 0.749±0.011 OUR FIT | | | | |
| 0.745±0.011 OUR AVERAGE | | | | |
| $0.744 \pm 0.036 \pm 0.037$ | 17k | ABREU | 00L DLPH | 1992–1995 runs |
| $0.779 \pm 0.047 \pm 0.029$ | 25k | ACKERSTAFF | 99D OPAL | 1990–1995 LEP runs |
| $0.68 \pm 0.04 \pm 0.07$ | | 230 ALBRECHT | 98 ARG | $E_{cm}^{ee} = 9.5\text{--}10.6 \text{ GeV}$ |
| $0.71 \pm 0.14 \pm 0.05$ | | ABE | 970 SLD | 1993–1995 SLC runs |
| $0.747 \pm 0.012 \pm 0.004$ | 34k | ALEXANDER | 97F CLEO | $E_{cm}^{ee} = 10.6 \text{ GeV}$ |
| $0.735 \pm 0.036 \pm 0.020$ | 4.7k | 231 ALBRECHT | 95 ARG | $E_{cm}^{ee} = 9.5\text{--}10.6 \text{ GeV}$ |
| $0.793 \pm 0.050 \pm 0.025$ | | BUSKULIC | 95D ALEP | 1990–1992 LEP runs |
| $0.79 \pm 0.08 \pm 0.06$ | 3230 | 232 ALBRECHT | 93G ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |
| $0.64 \pm 0.06 \pm 0.07$ | 2753 | JANSSEN | 89 CBAL | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |
| $0.62 \pm 0.17 \pm 0.14$ | 1823 | FORD | 87B MAC | $E_{cm}^{ee} = 29 \text{ GeV}$ |
| 0.60 ± 0.13 | 699 | BEHRENDS | 85 CLEO | $e^+ e^-$ near $\Upsilon(4S)$ |
| $0.72 \pm 0.10 \pm 0.11$ | 594 | BACINO | 79B DLCO | $E_{cm}^{ee} = 3.5\text{--}7.4 \text{ GeV}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------------|------|----------|----------|------------------------|
| $0.732 \pm 0.014 \pm 0.009$ | 19k | AMMAR | 97B CLEO | Repl. by ALEXANDER 97F |
| $0.747 \pm 0.045 \pm 0.028$ | 5106 | ALBRECHT | 90E ARG | Repl. by ALBRECHT 95 |

230 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

231 ALBRECHT 95 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)$ ($h^+ h^- h^+ (\pi^0) \bar{\nu}_\tau$) and their charged conjugates.

232 ALBRECHT 93G use tau pair events of the type $\tau^- \tau^+ \rightarrow (\mu^- \bar{\nu}_\mu \nu_\tau)(e^+ \nu_e \bar{\nu}_\tau)$ and their charged conjugates.

$\rho^\tau(\mu)$ PARAMETER(V-A) theory predicts $\rho = 0.75$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------|------|--|
| 0.752±0.021 OUR FIT | | | | |
| 0.758±0.023 OUR AVERAGE | | | | |
| 0.999±0.098±0.045 | 22k | ABREU 00L | DLPH | 1992–1995 runs |
| 0.777±0.044±0.016 | 27k | ACKERSTAFF 99D | OPAL | 1990–1995 LEP runs |
| 0.69 ± 0.06 ± 0.06 | 233 | ALBRECHT 98 | ARG | $E_{cm}^{ee} = 9.5\text{--}10.6 \text{ GeV}$ |
| 0.54 ± 0.28 ± 0.14 | | ABE 970 | SLD | 1993–1995 SLC runs |
| 0.750±0.017±0.045 | 22k | ALEXANDER 97F | CLEO | $E_{cm}^{ee} = 10.6 \text{ GeV}$ |
| 0.693±0.057±0.028 | | BUSKULIC 95D | ALEP | 1990–1992 LEP runs |
| 0.76 ± 0.07 ± 0.08 | 3230 | ALBRECHT 93G | ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |
| 0.734±0.055±0.027 | 3041 | ALBRECHT 90E | ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |
| 0.89 ± 0.14 ± 0.08 | 1909 | FORD 87B | MAC | $E_{cm}^{ee} = 29 \text{ GeV}$ |
| 0.81 ± 0.13 | 727 | BEHRENDS 85 | CLEO | $e^+ e^-$ near $\gamma(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.747±0.048±0.044 | 13k | AMMAR 97B | CLEO | Repl. by ALEXANDER 97F |

233 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

 $\xi^\tau(e \text{ or } \mu)$ PARAMETER(V-A) theory predicts $\xi = 1$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|------------------|------|--|
| 0.997±0.032 OUR FIT | | | | |
| 0.984±0.034 OUR AVERAGE | | | | |
| 0.929±0.070±0.030 | 36k | ABREU 00L | DLPH | 1992–1995 runs |
| 0.98 ± 0.22 ± 0.10 | 46k | ACKERSTAFF 99D | OPAL | 1990–1995 LEP runs |
| 0.70 ± 0.16 | 54k | ACCIARRI 98R | L3 | 1991–1995 LEP runs |
| 1.03 ± 0.11 | 234 | ALBRECHT 98 | ARG | $E_{cm}^{ee} = 9.5\text{--}10.6 \text{ GeV}$ |
| 1.05 ± 0.35 ± 0.04 | 235 | ABE 970 | SLD | 1993–1995 SLC runs |
| 1.007±0.040±0.015 | 55k | ALEXANDER 97F | CLEO | $E_{cm}^{ee} = 10.6 \text{ GeV}$ |
| 1.18 ± 0.15 ± 0.16 | | BUSKULIC 95D | ALEP | 1990–1992 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.94 ± 0.21 ± 0.07 | 18k | ACCIARRI 96H | L3 | Repl. by ACCIARRI 98R |
| 0.97 ± 0.14 | 236 | ALBRECHT 95C | ARG | Repl. by ALBRECHT 98 |
| 0.90 ± 0.15 ± 0.10 | 3230 | 237 ALBRECHT 93G | ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |

234 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

235 ABE 970 assume $\eta^\tau = 0$ in their fit. Letting η^τ vary in the fit gives a ξ^τ value of $1.02 \pm 0.36 \pm 0.05$.

236 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 95C uses events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(h^+ h^- h^+ \bar{\nu}_\tau)$ and their charged conjugates.

237 ALBRECHT 93G measurement determines $|\xi^\tau|$ for the case $\xi^\tau(e) = \xi^\tau(\mu)$, but the authors point out that other LEP experiments determine the sign to be positive.

$\xi^\tau(e)$ PARAMETER(V-A) theory predicts $\xi = 1$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|------|-----------------|------|--|
| 0.996±0.044 OUR FIT | | | | |
| 0.99 ±0.04 OUR AVERAGE | | | | |
| 1.01 ±0.12 ±0.05 | 17k | ABREU 00L | DLPH | 1992–1995 runs |
| 1.13 ±0.39 ±0.14 | 25k | ACKERSTAFF 99D | OPAL | 1990–1995 LEP runs |
| 1.11 ±0.20 ±0.08 | | 238 ALBRECHT 98 | ARG | $E_{cm}^{ee} = 9.5\text{--}10.6 \text{ GeV}$ |
| 1.16 ±0.52 ±0.06 | | ABE 970 | SLD | 1993–1995 SLC runs |
| 0.979±0.048±0.016 | 34k | ALEXANDER 97F | CLEO | $E_{cm}^{ee} = 10.6 \text{ GeV}$ |
| 1.03 ±0.23 ±0.09 | | BUSKULIC 95D | ALEP | 1990–1992 LEP runs |

238 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

 $\xi^\tau(\mu)$ PARAMETER(V-A) theory predicts $\xi = 1$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|------|-----------------|------|--|
| 1.046±0.065 OUR FIT | | | | |
| 1.08 ±0.07 OUR AVERAGE | | | | |
| 1.16 ±0.19 ±0.06 | 22k | ABREU 00L | DLPH | 1992–1995 runs |
| 0.79 ±0.41 ±0.09 | 27k | ACKERSTAFF 99D | OPAL | 1990–1995 LEP runs |
| 1.26 ±0.27 ±0.14 | | 239 ALBRECHT 98 | ARG | $E_{cm}^{ee} = 9.5\text{--}10.6 \text{ GeV}$ |
| 0.75 ±0.50 ±0.14 | | ABE 970 | SLD | 1993–1995 SLC runs |
| 1.054±0.069±0.047 | 22k | ALEXANDER 97F | CLEO | $E_{cm}^{ee} = 10.6 \text{ GeV}$ |
| 1.23 ±0.22 ±0.10 | | BUSKULIC 95D | ALEP | 1990–1992 LEP runs |

239 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

 $\eta^\tau(e \text{ or } \mu)$ PARAMETER(V-A) theory predicts $\eta = 0$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------|------|--|
| 0.011±0.031 OUR FIT | | | | |
| 0.019±0.033 OUR AVERAGE | | | | |
| -0.005±0.036±0.037 | | ABREU 00L | DLPH | 1992–1995 runs |
| 0.027±0.055±0.005 | 46k | ACKERSTAFF 99D | OPAL | 1990–1995 LEP runs |
| 0.27 ±0.14 | 54k | ACCIARRI 98R | L3 | 1991–1995 LEP runs |
| -0.13 ±0.47 ±0.15 | | ABE 970 | SLD | 1993–1995 SLC runs |
| -0.015±0.061±0.062 | 31k | AMMAR 97B | CLEO | $E_{cm}^{ee} = 10.6 \text{ GeV}$ |
| 0.03 ±0.18 ±0.12 | 8.2k | ALBRECHT 95 | ARG | $E_{cm}^{ee} = 9.5\text{--}10.6 \text{ GeV}$ |
| -0.04 ±0.15 ±0.11 | | BUSKULIC 95D | ALEP | 1990–1992 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.25 ±0.17 ±0.11 | 18k | ACCIARRI 96H | L3 | Repl. by ACCIARRI 98R |

$\eta^\tau(\mu)$ PARAMETER

($V-A$) theory predicts $\eta = 0$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|------|-------------|------|---------|
| -0.013±0.097 OUR FIT | | | | |

0.06 ± 0.20 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

| | | | |
|-----------------------|---------------|----------|--------------------------|
| 0.72 ± 0.32 ± 0.15 | ABREU | 00L DLPH | 1992–1995 runs |
| -0.59 ± 0.82 ± 0.45 | 240 ABE | 97O SLD | 1993–1995 SLC runs |
| 0.010 ± 0.149 ± 0.171 | 13k 241 AMMAR | 97B CLEO | $E_{cm}^{ee} = 10.6$ GeV |
| -0.24 ± 0.23 ± 0.18 | BUSKULIC | 95D ALEP | 1990–1992 LEP runs |


• • • We do not use the following data for averages, fits, limits, etc. • • •

0.010 ± 0.065 ± 0.001 27k 242 ACKERSTAFF 99D OPAL 1990–1995 LEP runs

240 Highly correlated (corr. = 0.92) with ABE 97O $\rho^\tau(\mu)$ measurement.

241 Highly correlated (corr. = 0.949) with AMMAR 97B $\rho^\tau(\mu)$ value.

242 ACKERSTAFF 99D result is dominated by a constraint on η^τ from the OPAL measurements of the τ lifetime and $B(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau)$ assuming lepton universality for the total coupling strength.



$\eta^\tau(\mu)$ parameter

$(\delta\xi)^T(e \text{ or } \mu)$ PARAMETER $(V-A)$ theory predicts $(\delta\xi) = 0.75$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------|------|--------------------------------------|
| 0.746±0.023 OUR FIT | | | | |
| 0.742±0.023 OUR AVERAGE | | | | |
| 0.779±0.070±0.028 | 36k | ABREU 00L | DLPH | 1992–1995 runs |
| 0.65 ± 0.14 ± 0.07 | 46k | ACKERSTAFF 99D | OPAL | 1990–1995 LEP runs |
| 0.70 ± 0.11 | 54k | ACCIARRI 98R | L3 | 1991–1995 LEP runs |
| 0.63 ± 0.09 | 243 | ALBRECHT 98 | ARG | $E_{cm}^{ee} = 9.5\text{--}10.6$ GeV |
| 0.88 ± 0.27 ± 0.04 | 244 | ABE 970 | SLD | 1993–1995 SLC runs |
| 0.745±0.026±0.009 | 55k | ALEXANDER 97F | CLEO | $E_{cm}^{ee} = 10.6$ GeV |
| 0.88 ± 0.11 ± 0.07 | | BUSKULIC 95D | ALEP | 1990–1992 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.81 ± 0.14 ± 0.06 | 18k | ACCIARRI 96H | L3 | Repl. by ACCIARRI 98R |
| 0.65 ± 0.12 | 245 | ALBRECHT 95C | ARG | Repl. by ALBRECHT 98 |

243 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 98, ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.


244 ABE 970 assume $\eta^T = 0$ in their fit. Letting η^T vary in the fit gives a $(\rho\xi)^T$ value of $0.87 \pm 0.27 \pm 0.04$.

245 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 93G, and ALBRECHT 94E. ALBRECHT 95C uses events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(h^+ h^- h^+ \bar{\nu}_\tau)$ and their charged conjugates.

 $(\delta\xi)^T(e)$ PARAMETER $(V-A)$ theory predicts $(\delta\xi) = 0.75$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|---|------|--------------------------------------|
| 0.735±0.030 OUR FIT | | | | |
| 0.73 ± 0.05 OUR AVERAGE | | | | |
| | | Error includes scale factor of 1.5. See the ideogram below. | | |
| 0.85 ± 0.12 ± 0.04 | 17k | ABREU 00L | DLPH | 1992–1995 runs |
| 0.72 ± 0.31 ± 0.14 | 25k | ACKERSTAFF 99D | OPAL | 1990–1995 LEP runs |
| 0.56 ± 0.14 ± 0.06 | 246 | ALBRECHT 98 | ARG | $E_{cm}^{ee} = 9.5\text{--}10.6$ GeV |
| 0.85 ± 0.43 ± 0.08 | | ABE 970 | SLD | 1993–1995 SLC runs |
| 0.720±0.032±0.010 | 34k | ALEXANDER 97F | CLEO | $E_{cm}^{ee} = 10.6$ GeV |
| 1.11 ± 0.17 ± 0.07 | | BUSKULIC 95D | ALEP | 1990–1992 LEP runs |

246 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.



$(\delta\xi)^\tau(e)$ PARAMETER

$(\delta\xi)^\tau(\mu)$ PARAMETER

(V-A) theory predicts $(\delta\xi) = 0.75$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------|-------------|----------|--|
| 0.774 +/- 0.043 OUR FIT | | | | |
| 0.78 +/- 0.04 OUR AVERAGE | | | | |
| 0.86 +/- 0.13 +/- 0.04 | 22k | ABREU | 00L DLPH | 1992–1995 runs |
| 0.63 +/- 0.23 +/- 0.05 | 27k | ACKERSTAFF | 99D OPAL | 1990–1995 LEP runs |
| 0.73 +/- 0.18 +/- 0.10 | 247 | ALBRECHT | 98 ARG | $E_{cm}^{ee} = 9.5\text{--}10.6 \text{ GeV}$ |
| 0.82 +/- 0.32 +/- 0.07 | | ABE | 97O SLD | 1993–1995 SLC runs |
| 0.786 +/- 0.041 +/- 0.032 | 22k | ALEXANDER | 97F CLEO | $E_{cm}^{ee} = 10.6 \text{ GeV}$ |
| 0.71 +/- 0.14 +/- 0.06 | | BUSKULIC | 95D ALEP | 1990–1992 LEP runs |

247 ALBRECHT 98 use tau pair events of the type $\tau^- \tau^+ \rightarrow (\ell^- \bar{\nu}_\ell \nu_\tau)(\pi^+ \pi^0 \bar{\nu}_\tau)$, and their charged conjugates.

$\xi^\tau(\pi)$ PARAMETER

(V-A) theory predicts $\xi^\tau(\pi) = 1$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|----------|----------------------------------|
| 0.992 +/- 0.046 OUR FIT | | | | |
| 0.99 +/- 0.05 OUR AVERAGE | | | | |
| 0.81 +/- 0.17 +/- 0.02 | | ABE | 97O SLD | 1993–1995 SLC runs |
| 1.03 +/- 0.06 +/- 0.04 | 2.0k | COAN | 97 CLEO | $E_{cm}^{ee} = 10.6 \text{ GeV}$ |
| 0.987 +/- 0.057 +/- 0.027 | | BUSKULIC | 95D ALEP | 1990–1992 LEP runs |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.95 +/- 0.11 +/- 0.05 | 248 | BUSKULIC | 94D ALEP | 1990+1991 LEP run |

248 Superseded by BUSKULIC 95D.

$\xi^T(\rho)$ PARAMETER

($V-A$) theory predicts $\xi^T(\rho) = 1$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|--------------|--------------------|--------------------------|------------------------------|
| 0.998±0.010 OUR FIT | | | | |
| 0.998±0.010 OUR AVERAGE | | | | |
| 0.99 ± 0.12 ± 0.04 | ABE | 970 SLD | 1993–1995 SLC runs | |
| 0.995 ± 0.010 ± 0.003 | 66k | ALEXANDER 97F CLEO | $E_{cm}^{ee} = 10.6$ GeV | |
| 1.045 ± 0.058 ± 0.032 | BUSKULIC | 95D ALEP | 1990–1992 LEP runs | |
| 1.022 ± 0.028 ± 0.030 | 1.7k | 249 ALBRECHT | 94E ARG | $E_{cm}^{ee} = 9.4–10.6$ GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.03 ± 0.11 ± 0.05 | 250 BUSKULIC | 94D ALEP | 1990+1991 LEP run | |
| 249 ALBRECHT 94E measure the square of this quantity and use the sign determined by ALBRECHT 90I to obtain the quoted result. | | | | |
| 250 Superseded by BUSKULIC 95D. | | | | |

$\xi^T(a_1)$ PARAMETER

($V-A$) theory predicts $\xi^T(a_1) = 1$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------|-------------------------|------------------------------|---------|
| 0.998±0.077 OUR FIT | | | | |
| 1.00 ± 0.08 OUR AVERAGE | | | | |
| 1.02 ± 0.13 ± 0.03 | 17.2k | ASNER 00 CLEO | $E_{cm}^{ee} = 10.6$ GeV | |
| 1.29 ± 0.26 ± 0.11 | 7.4k | 251 ACKERSTAFF 97R OPAL | 1992–1994 LEP runs | |
| 0.85 $^{+0.15}_{-0.17}$ ± 0.05 | | ALBRECHT 95C ARG | $E_{cm}^{ee} = 9.5–10.6$ GeV | |
| 0.937 ± 0.116 ± 0.064 | | BUSKULIC 95D ALEP | 1990–1992 LEP runs | |
| 1.25 ± 0.23 $^{+0.15}_{-0.08}$ | 7.5k | ALBRECHT 93C ARG | $E_{cm}^{ee} = 9.4–10.6$ GeV | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.08 $^{+0.46}_{-0.41}$ $^{+0.14}_{-0.25}$ | 2.6k | 252 AKERS 95P OPAL | Repl. by ACKER-STAFF 97R | |
| 251 ACKERSTAFF 97R obtain this result with a model independent fit to the hadronic structure functions. Fitting with the model of Kuhn and Santamaria (ZPHY C48 , 445 (1990)) gives $0.87 \pm 0.16 \pm 0.04$, and with the model of Isgur <i>et al.</i> (PR D39 , 1357 (1989)) they obtain $1.20 \pm 0.21 \pm 0.14$. | | | | |
| 252 AKERS 95P obtain this result with a model independent fit to the hadronic structure functions. Fitting with the model of Kuhn and Santamaria (ZPHY C48 , 445 (1990)) gives $0.87 \pm 0.27 ^{+0.05}_{-0.06}$, and with the model of Isgur <i>et al.</i> (PR D39 , 1357 (1989)) they obtain $1.10 \pm 0.31 ^{+0.13}_{-0.14}$. | | | | |

$\xi^T(\text{all hadronic modes})$ PARAMETER

($V-A$) theory predicts $\xi^T = 1$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------|-------------------------|--------------------------|---------|
| 1.000±0.008 OUR FIT | | | | |
| 1.001±0.009 OUR AVERAGE | | | | |
| 0.997 ± 0.027 ± 0.011 | 39k | 253 ABREU 00L DLPH | 1992–1995 runs | |
| 1.02 ± 0.13 ± 0.03 | 17.2k | 254 ASNER 00 CLEO | $E_{cm}^{ee} = 10.6$ GeV | |
| 1.032 ± 0.031 | 37k | 255 ACCIARRI 98R L3 | 1991–1995 LEP runs | |
| 0.93 ± 0.10 ± 0.04 | | ABE 970 SLD | 1993–1995 SLC runs | |
| 1.29 ± 0.26 ± 0.11 | 7.4k | 256 ACKERSTAFF 97R OPAL | 1992–1994 LEP runs | |

| | | | | |
|---|------|---------------|----------|--|
| 0.995±0.010±0.003 | 66k | 257 ALEXANDER | 97F CLEO | $E_{cm}^{ee} = 10.6 \text{ GeV}$ |
| 1.03 ± 0.06 ± 0.04 | 2.0k | 258 COAN | 97 CLEO | $E_{cm}^{ee} = 10.6 \text{ GeV}$ |
| 1.017±0.039 | | 259 ALBRECHT | 95C ARG | $E_{cm}^{ee} = 9.5\text{--}10.6 \text{ GeV}$ |
| 1.006±0.032±0.019 | | 260 BUSKULIC | 95D ALEP | 1990–1992 LEP runs |
| 1.25 ± 0.23 +0.15 -0.08 | 7.5k | 261 ALBRECHT | 93C ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.970±0.053±0.011 | 14k | 262 ACCIARRI | 96H L3 | Repl. by ACCIARRI 98R |
| 1.08 +0.46 +0.14 -0.41 -0.25 | 2.6k | 263 AKERS | 95P OPAL | Repl. by ACKER-STAFF 97R |
| 1.022±0.028±0.030 | 1.7k | 264 ALBRECHT | 94E ARG | $E_{cm}^{ee} = 9.4\text{--}10.6 \text{ GeV}$ |
| 0.99 ± 0.07 ± 0.04 | | 265 BUSKULIC | 94D ALEP | 1990+1991 LEP run |
| 253 ABREU 00L use $\tau^- \rightarrow h^- \geq 0\pi^0\nu_\tau$ decays. | | | | |
| 254 ASNER 00 use $\tau^- \rightarrow \pi^- 2\pi^0\nu_\tau$ decays. | | | | |
| 255 ACCIARRI 98R use $\tau \rightarrow \pi\nu_\tau$, $\tau \rightarrow K\nu_\tau$, and $\tau \rightarrow \rho\nu_\tau$ decays. | | | | |
| 256 ACKERSTAFF 97R use $\tau \rightarrow a_1\nu_\tau$ decays. | | | | |
| 257 ALEXANDER 97F use $\tau \rightarrow \rho\nu_\tau$ decays. | | | | |
| 258 COAN 97 use $h^+ h^-$ energy correlations. | | | | |
| 259 Combined fit to ARGUS tau decay parameter measurements in ALBRECHT 95C, ALBRECHT 95G, and ALBRECHT 94E. | | | | |
| 260 BUSKULIC 95D use $\tau \rightarrow \pi\nu_\tau$, $\tau \rightarrow \rho\nu_\tau$, and $\tau \rightarrow a_1\nu_\tau$ decays. | | | | |
| 261 Uses $\tau \rightarrow a_1\nu_\tau$ decays. Replaced by ALBRECHT 95C. | | | | |
| 262 ACCIARRI 96H use $\tau \rightarrow \pi\nu_\tau$, $\tau \rightarrow K\nu_\tau$, and $\tau \rightarrow \rho\nu_\tau$ decays. | | | | |
| 263 AKERS 95P use $\tau \rightarrow a_1\nu_\tau$ decays. | | | | |
| 264 ALBRECHT 94E measure the square of this quantity and use the sign determined by ALBRECHT 90I to obtain the quoted result. Uses $\tau \rightarrow a_1\nu_\tau$ decays. Replaced by ALBRECHT 95C. | | | | |
| 265 BUSKULIC 94D use $\tau \rightarrow \pi\nu_\tau$ and $\tau \rightarrow \rho\nu_\tau$ decays. Superseded by BUSKULIC 95D. | | | | |

τ REFERENCES

| | | | | |
|---------------|-----|----------------|---------------------------------------|------------------|
| ABBIENDI | 00A | PL B492 23 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI | 00C | EPJ C13 213 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABBIENDI | 00D | EPJ C13 197 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABREU | 00L | EPJ C16 229 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 00B | PL B479 67 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| AHMED | 00 | PR D61 071101R | S. Ahmed <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 00 | PL B485 37 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ASNER | 00 | PR D61 012002 | D.M. Asner <i>et al.</i> | (CLEO Collab.) |
| ASNER | 00B | PR D62 072006 | D.M. Asner <i>et al.</i> | (CLEO Collab.) |
| BERGFELD | 00 | PR L84 830 | T. Bergfeld <i>et al.</i> | (CLEO Collab.) |
| BROWDER | 00 | PR D61 052004 | T.E. Browder <i>et al.</i> | (CLEO Collab.) |
| EDWARDS | 00A | PR D61 072003 | K.W. Edwards <i>et al.</i> | (CLEO Collab.) |
| GONZALEZ-S... | 00 | NP B582 3 | G.A. Gonzalez-Sprinberg <i>et al.</i> | |
| ABBIENDI | 99H | PL B447 134 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ABREU | 99X | EPJ C10 201 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACKERSTAFF | 99D | EPJ C8 3 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKERSTAFF | 99E | EPJ C8 183 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| BARATE | 99K | EPJ C10 1 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BARATE | 99R | EPJ C11 599 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BISHAI | 99 | PRL 82 281 | M. Bishai <i>et al.</i> | (CLEO Collab.) |
| GODANG | 99 | PR D59 091303 | R. Godang <i>et al.</i> | (CLEO Collab.) |
| RICHICHI | 99 | PR D60 112002 | S.J. Richichi <i>et al.</i> | (CLEO Collab.) |
| ACCIARRI | 98C | PL B426 207 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 98E | PL B434 169 | M. Aciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 98R | PL B438 405 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACKERSTAFF | 98M | EPJ C4 193 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKERSTAFF | 98N | PL B431 188 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |

| | | | | |
|------------|-----|-------------------------|------------------------------|----------------------|
| ALBRECHT | 98 | PL B431 179 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BARATE | 98 | EPJ C1 65 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BARATE | 98E | EPJ C4 29 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BLISS | 98 | PR D57 5903 | D.W. Bliss <i>et al.</i> | (CLEO Collab.) |
| ABE | 97O | PRL 78 4691 | K. Abe <i>et al.</i> | (SLD Collab.) |
| ACKERSTAFF | 97J | PL B404 213 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKERSTAFF | 97L | ZPHY C74 403 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ACKERSTAFF | 97R | ZPHY C75 593 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| ALEXANDER | 97F | PR D56 5320 | J.P. Alexander <i>et al.</i> | (CLEO Collab.) |
| AMMAR | 97B | PRL 78 4686 | R. Ammar <i>et al.</i> | (CLEO Collab.) |
| ANASTASSOV | 97 | PR D55 2559 | A. Anastassov <i>et al.</i> | (CLEO Collab.) |
| Also | 98B | PR D58 119903 (erratum) | A. Anastassov <i>et al.</i> | (CLEO Collab.) |
| ANDERSON | 97 | PRL 79 3814 | S. Anderson <i>et al.</i> | (CLEO Collab.) |
| AVERY | 97 | PR D55 R1119 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| BARATE | 97I | ZPHY C74 387 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BARATE | 97R | PL B414 362 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| BERGFELD | 97 | PRL 79 2406 | T. Bergfeld <i>et al.</i> | (CLEO Collab.) |
| BONVICINI | 97 | PRL 79 1221 | G. Bonvicini <i>et al.</i> | (CLEO Collab.) |
| BUSKULIC | 97C | ZPHY C74 263 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| COAN | 97 | PR D55 7291 | T.E. Coan <i>et al.</i> | (CLEO Collab.) |
| EDWARDS | 97 | PR D55 R3919 | K.W. Edwards <i>et al.</i> | (CLEO Collab.) |
| EDWARDS | 97B | PR D56 R5297 | K.W. Edwards <i>et al.</i> | (CLEO Collab.) |
| ESCRIBANO | 97 | PL B395 369 | R. Escribano, E. Masso | (BARC, PARIT) |
| ABREU | 96B | PL B365 448 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 96H | PL B377 313 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 96K | PL B389 187 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ALAM | 96 | PRL 76 2637 | M.S. Alam <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 96E | PRPL 276 223 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALEXANDER | 96D | PL B369 163 | G. Alexander <i>et al.</i> | (OPAL Collab.) |
| ALEXANDER | 96E | PL B374 341 | G. Alexander <i>et al.</i> | (OPAL Collab.) |
| ALEXANDER | 96S | PL B388 437 | G. Alexander <i>et al.</i> | (OPAL Collab.) |
| BAI | 96 | PR D53 20 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BALEST | 96 | PL B388 402 | R. Balest <i>et al.</i> | (CLEO Collab.) |
| BARTELT | 96 | PRL 76 4119 | J.E. Bartelt <i>et al.</i> | (CLEO Collab.) |
| BUSKULIC | 96 | ZPHY C70 579 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 96C | ZPHY C70 561 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| COAN | 96 | PR D53 6037 | T.E. Coan <i>et al.</i> | (CLEO Collab.) |
| ABE | 95Y | PR D52 4828 | K. Abe <i>et al.</i> | (SLD Collab.) |
| ABREU | 95T | PL B357 715 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 95U | PL B359 411 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACCIARRI | 95 | PL B345 93 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 95F | PL B352 487 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| AKERS | 95F | ZPHY C66 31 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| AKERS | 95I | ZPHY C66 543 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| AKERS | 95P | ZPHY C67 45 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| AKERS | 95Y | ZPHY C68 555 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| ALBRECHT | 95 | PL B341 441 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 95C | PL B349 576 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 95G | ZPHY C68 25 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 95H | ZPHY C68 215 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BALEST | 95C | PRL 75 3809 | R. Balest <i>et al.</i> | (CLEO Collab.) |
| BERNABEU | 95 | NP B436 474 | J. Bernabeu <i>et al.</i> | |
| BUSKULIC | 95C | PL B346 371 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 95D | PL B346 379 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| Also | 95P | PL B363 265 erratum | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| ABREU | 94K | PL B334 435 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| AKERS | 94E | PL B328 207 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| AKERS | 94G | PL B339 278 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| ALBRECHT | 94E | PL B337 383 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ARTUSO | 94 | PRL 72 3762 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| BARTELT | 94 | PRL 73 1890 | J.E. Bartelt <i>et al.</i> | (CLEO Collab.) |
| BATTLE | 94 | PRL 73 1079 | M. Battle <i>et al.</i> | (CLEO Collab.) |
| BAUER | 94 | PR D50 R13 | D.A. Bauer <i>et al.</i> | (TPC/2gamma Collab.) |
| BUSKULIC | 94D | PL B321 168 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 94E | PL B332 209 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 94F | PL B332 219 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| GIBAUT | 94B | PRL 73 934 | D. Gibaut <i>et al.</i> | (CLEO Collab.) |
| ADRIANI | 93M | PRPL 236 1 | O. Adriani <i>et al.</i> | (L3 Collab.) |
| ALBRECHT | 93C | ZPHY C58 61 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 93G | PL B316 608 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BALEST | 93 | PR D47 R3671 | R. Balest <i>et al.</i> | (CLEO Collab.) |

| | | | | |
|-------------|-----|-----------------------|---------------------------------|------------------------|
| BEAN | 93 | PRL 70 138 | A. Bean <i>et al.</i> | (CLEO Collab.) |
| BORTOLETTO | 93 | PRL 71 1791 | D. Bortoletto <i>et al.</i> | (CLEO Collab.) |
| ESCRIBANO | 93 | PL B301 419 | R. Escribano, E. Masso | (BARC) |
| PROCARIO | 93 | PRL 70 1207 | M. Procario <i>et al.</i> | (CLEO Collab.) |
| ABREU | 92N | ZPHY C55 555 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACTON | 92F | PL B281 405 | D.P. Acton <i>et al.</i> | (OPAL Collab.) |
| ACTON | 92H | PL B288 373 | P.D. Acton <i>et al.</i> | (OPAL Collab.) |
| AKERIB | 92 | PRL 69 3610 | D.S. Akerib <i>et al.</i> | (CLEO Collab.) |
| Also | 93B | PRL 71 3395 (erratum) | D.S. Akerib <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 92D | ZPHY C53 367 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 92K | ZPHY C55 179 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 92M | PL B292 221 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 92Q | ZPHY C56 339 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| AMMAR | 92 | PR D45 3976 | R. Ammar <i>et al.</i> | (CLEO Collab.) |
| ARTUSO | 92 | PRL 69 3278 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| BAI | 92 | PRL 69 3021 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BATTLE | 92 | PL B291 488 | M. Battle <i>et al.</i> | (CLEO Collab.) |
| BUSKULIC | 92J | PL B297 459 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| DECAMP | 92C | ZPHY C54 211 | D. Decamp <i>et al.</i> | (ALEPH Collab.) |
| ADEVA | 91F | PL B265 451 | B. Adeva <i>et al.</i> | (L3 Collab.) |
| ALBRECHT | 91D | PL B260 259 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALEXANDER | 91D | PL B266 201 | G. Alexander <i>et al.</i> | (OPAL Collab.) |
| ANTREASYAN | 91 | PL B259 216 | D. Antreasyan <i>et al.</i> | (Crystal Ball Collab.) |
| GRIFOLS | 91 | PL B255 611 | J.A. Grifols, A. Mendez | (BARC) |
| SAMUEL | 91B | PRL 67 668 | M.A. Samuel, G.W. Li, R. Mendel | (OKSU, WONT) |
| Also | 92B | PRL 69 995 | M.A. Samuel, G.W. Li, R. Mendel | (OKSU, WONT) |
| Erratum. | | | | |
| ABACHI | 90 | PR D41 1414 | S. Abachi <i>et al.</i> | (HRS Collab.) |
| ALBRECHT | 90E | PL B246 278 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 90I | PL B250 164 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BEHREND | 90 | ZPHY C46 537 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| BOWCOCK | 90 | PR D41 805 | T.J.V. Bowcock <i>et al.</i> | (CLEO Collab.) |
| DELAGUILA | 90 | PL B252 116 | F. del Aguila, M. Sher | (BARC, WILL) |
| GOLDBERG | 90 | PL B251 223 | M. Goldberg <i>et al.</i> | (CLEO Collab.) |
| WU | 90 | PR D41 2339 | D.Y. Wu <i>et al.</i> | (Mark II Collab.) |
| ABACHI | 89B | PR D40 902 | S. Abachi <i>et al.</i> | (HRS Collab.) |
| BEHREND | 89B | PL B222 163 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| JANSSEN | 89 | PL B228 273 | H. Janssen <i>et al.</i> | (Crystal Ball Collab.) |
| KLEINWORT | 89 | ZPHY C42 7 | C. Kleinwort <i>et al.</i> | (JADE Collab.) |
| ADEVA | 88 | PR D38 2665 | B. Adeva <i>et al.</i> | (Mark-J Collab.) |
| ALBRECHT | 88B | PL B202 149 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 88L | ZPHY C41 1 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 88M | ZPHY C41 405 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| AMIDEI | 88 | PR D37 1750 | D. Amidei <i>et al.</i> | (Mark II Collab.) |
| BEHREND | 88 | PL B200 226 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| BRAUNSCH... | 88C | ZPHY C39 331 | W. Braunschweig <i>et al.</i> | (TASSO Collab.) |
| KEH | 88 | PL B212 123 | S. Keh <i>et al.</i> | (Crystal Ball Collab.) |
| TSCHIRHART | 88 | PL B205 407 | R. Tschirhart <i>et al.</i> | (HRS Collab.) |
| ABACHI | 87B | PL B197 291 | S. Abachi <i>et al.</i> | (HRS Collab.) |
| ABACHI | 87C | PRL 59 2519 | S. Abachi <i>et al.</i> | (HRS Collab.) |
| ADLER | 87B | PRL 59 1527 | J. Adler <i>et al.</i> | (Mark III Collab.) |
| AIHARA | 87B | PR D35 1553 | H. Aihara <i>et al.</i> | (TPC Collab.) |
| AIHARA | 87C | PRL 59 751 | H. Aihara <i>et al.</i> | (TPC Collab.) |
| ALBRECHT | 87L | PL B185 223 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALBRECHT | 87P | PL B199 580 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BAND | 87 | PL B198 297 | H.R. Band <i>et al.</i> | (MAC Collab.) |
| BAND | 87B | PRL 59 415 | H.R. Band <i>et al.</i> | (MAC Collab.) |
| BARINGER | 87 | PRL 59 1993 | P. Baringer <i>et al.</i> | (CLEO Collab.) |
| BEBEK | 87C | PR D36 690 | C. Bebek <i>et al.</i> | (CLEO Collab.) |
| BURCHAT | 87 | PR D35 27 | P.R. Burchat <i>et al.</i> | (Mark II Collab.) |
| BYLSMA | 87 | PR D35 2269 | B.G. Bylsma <i>et al.</i> | (HRS Collab.) |
| COFFMAN | 87 | PR D36 2185 | D.M. Coffman <i>et al.</i> | (Mark III Collab.) |
| DERRICK | 87 | PL B189 260 | M. Derrick <i>et al.</i> | (HRS Collab.) |
| FORD | 87 | PR D35 408 | W.T. Ford <i>et al.</i> | (MAC Collab.) |
| FORD | 87B | PR D36 1971 | W.T. Ford <i>et al.</i> | (MAC Collab.) |
| GAN | 87 | PRL 59 411 | K.K. Gan <i>et al.</i> | (Mark II Collab.) |
| GAN | 87B | PL B197 561 | K.K. Gan <i>et al.</i> | (Mark II Collab.) |
| AIHARA | 86E | PRL 57 1836 | H. Aihara <i>et al.</i> | (TPC Collab.) |
| BARTEL | 86D | PL B182 216 | W. Bartel <i>et al.</i> | (JADE Collab.) |

| | | | | |
|-----------------------------------|-----|------------------------|----------------------------------|-------------------------|
| PDG | 86 | PL 170B | M. Aguilar-Benitez <i>et al.</i> | (CERN, CIT+) |
| RUCKSTUHL | 86 | PRL 56 2132 | W. Ruckstuhl <i>et al.</i> | (DELCO Collab.) |
| SCHMIDKE | 86 | PRL 57 527 | W.B. Schmidke <i>et al.</i> | (Mark II Collab.) |
| YELTON | 86 | PRL 56 812 | J.M. Yelton <i>et al.</i> | (Mark II Collab.) |
| ALTHOFF | 85 | ZPHY C26 521 | M. Althoff <i>et al.</i> | (TASSO Collab.) |
| ASH | 85B | PRL 55 2118 | W.W. Ash <i>et al.</i> | (MAC Collab.) |
| BALTRUSAIT... | 85 | PRL 55 1842 | R.M. Baltrusaitis <i>et al.</i> | (Mark III Collab.) |
| BARTEL | 85F | PL 161B 188 | W. Bartel <i>et al.</i> | (JADE Collab.) |
| BEHRENDS | 85 | PR D32 2468 | S. Behrends <i>et al.</i> | (CLEO Collab.) |
| BELTRAMI | 85 | PRL 54 1775 | I. Beltrami <i>et al.</i> | (HRS Collab.) |
| BERGER | 85 | ZPHY C28 1 | C. Berger <i>et al.</i> | (PLUTO Collab.) |
| BURCHAT | 85 | PRL 54 2489 | P.R. Burchat <i>et al.</i> | (Mark II Collab.) |
| FERNANDEZ | 85 | PRL 54 1624 | E. Fernandez <i>et al.</i> | (MAC Collab.) |
| MILLS | 85 | PRL 54 624 | G.B. Mills <i>et al.</i> | (DELCO Collab.) |
| AIHARA | 84C | PR D30 2436 | H. Aihara <i>et al.</i> | (TPC Collab.) |
| BEHREND | 84 | ZPHY C23 103 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| MILLS | 84 | PRL 52 1944 | G.B. Mills <i>et al.</i> | (DELCO Collab.) |
| BEHREND | 83C | PL 127B 270 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| SILVERMAN | 83 | PR D27 1196 | D.J. Silverman, G.L. Shaw | (UCI) |
| BEHREND | 82 | PL 114B 282 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| BLOCKER | 82B | PRL 48 1586 | C.A. Blocker <i>et al.</i> | (Mark II Collab.) |
| BLOCKER | 82D | PL 109B 119 | C.A. Blocker <i>et al.</i> | (Mark II Collab.) J |
| FELDMAN | 82 | PRL 48 66 | G.J. Feldman <i>et al.</i> | (Mark II Collab.) |
| HAYES | 82 | PR D25 2869 | K.G. Hayes <i>et al.</i> | (Mark II Collab.) |
| BERGER | 81B | PL 99B 489 | C. Berger <i>et al.</i> | (PLUTO Collab.) |
| DORFAN | 81 | PRL 46 215 | J.M. Dorfan <i>et al.</i> | (Mark II Collab.) |
| BRANDELIK | 80 | PL 92B 199 | R. Brandelik <i>et al.</i> | (TASSO Collab.) |
| ZHOLENTZ | 80 | PL 96B 214 | A.A. Zholents <i>et al.</i> | (NOVO) |
| Also | 81 | SJNP 34 814 | A.A. Zholents <i>et al.</i> | (NOVO) |
| | | Translated from YAF 34 | 1471. | |
| BACINO | 79B | PRL 42 749 | W.J. Bacino <i>et al.</i> | (DELCO Collab.) |
| KIRKBY | 79 | SLAC-PUB-2419 | J. Kirkby | (SLAC) J |
| Batavia Lepton Photon Conference. | | | | |
| BACINO | 78B | PRL 41 13 | W.J. Bacino <i>et al.</i> | (DELCO Collab.) J |
| Also | 78 | Tokyo Conf. 249 | J. Kirz | (STON) |
| Also | 80 | PL 96B 214 | A.A. Zholents <i>et al.</i> | (NOVO) |
| BRANDELIK | 78 | PL 73B 109 | R. Brandelik <i>et al.</i> | (DASP Collab.) J |
| FELDMAN | 78 | Tokyo Conf. 777 | G.J. Feldman | (SLAC) J |
| HEILE | 78 | NP B138 189 | F.B. Heile <i>et al.</i> | (SLAC, LBL) |
| JAROS | 78 | PRL 40 1120 | J. Jaros <i>et al.</i> | (SLAC, LBL, NWES, HAWA) |
| PERL | 75 | PRL 35 1489 | M.L. Perl <i>et al.</i> | (LBL, SLAC) |

OTHER RELATED PAPERS

| | | | | |
|--------------|----|--------------|--------------------------------|--------------|
| RAHAL-CAL... | 98 | IJMP A13 695 | G. Rahal-Callot | (ETH) |
| GENTILE | 96 | PRPL 274 287 | S. Gentile, M. Pohl | (ROMAI, ETH) |
| WEINSTEIN | 93 | ARNPS 43 457 | A.J. Weinstein, R. Stroynowski | (CIT, SMU) |
| PERL | 92 | RPP 55 653 | M.L. Perl | (SLAC) |
| PICH | 90 | MPL A5 1995 | A. Pich | (VALE) |
| BARISH | 88 | PRPL 157 1 | B.C. Barish, R. Stroynowski | (CIT) |
| GAN | 88 | IJMP A3 531 | K.K. Gan, M.L. Perl | (SLAC) |
| HAYES | 88 | PR D38 3351 | K.G. Hayes, M.L. Perl | (SLAC) |
| PERL | 80 | ARNPS 30 299 | M.L. Perl | (SLAC) |