

# CHARMED BARYONS ( $C=+1$ )

$$\begin{aligned}\Lambda_c^+ &= u d c, \quad \Sigma_c^{++} = u u c, \quad \Sigma_c^+ = u d c, \quad \Sigma_c^0 = d d c, \\ \Xi_c^+ &= u s c, \quad \Xi_c^0 = d s c, \quad \Omega_c^0 = s s c\end{aligned}$$

$\Lambda_c^+$

$$I(J^P) = 0(\frac{1}{2}^+)$$

$J$  is not well measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2284.9 \pm 0.6$  MeV

Mean life  $\tau = (200 \pm 6) \times 10^{-15}$  s ( $S = 1.6$ )

$c\tau = 59.9 \mu\text{m}$

### Decay asymmetry parameters

$$\Lambda\pi^+ \quad \alpha = -0.98 \pm 0.19$$

$$\Sigma^+\pi^0 \quad \alpha = -0.45 \pm 0.32$$

$$\Lambda\ell^+\nu_\ell \quad \alpha = -0.82^{+0.11}_{-0.07}$$

Nearly all branching fractions of the  $\Lambda_c^+$  are measured relative to the  $pK^-\pi^+$  mode, but there are no model-independent measurements of this branching fraction. We explain how we arrive at our value of  $B(\Lambda_c^+ \rightarrow pK^-\pi^+)$  in a Note at the beginning of the branching-ratio measurements in the Listings. When this branching fraction is eventually well determined, all the other branching fractions will slide up or down proportionally as the true value differs from the value we use here.

$\Lambda_c^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level (MeV/c)
<b>Hadronic modes with a <math>p</math>: <math>S = -1</math> final states</b>		
$p\bar{K}^0$	( 2.3 $\pm$ 0.6 ) %	872
$pK^-\pi^+$	[ <i>I</i> ] ( 5.0 $\pm$ 1.3 ) %	822
$p\bar{K}^*(892)^0$	[ <i>m</i> ] ( 1.6 $\pm$ 0.5 ) %	681
$\Delta(1232)^{++}K^-$	( 8.6 $\pm$ 3.0 ) $\times 10^{-3}$	709
$\Lambda(1520)\pi^+$	[ <i>m</i> ] ( 5.9 $\pm$ 2.1 ) $\times 10^{-3}$	626
$pK^-\pi^+$ nonresonant	( 2.8 $\pm$ 0.8 ) %	822
$p\bar{K}^0\pi^0$	( 3.3 $\pm$ 1.0 ) %	822

$p\bar{K}^0\eta$	( 1.2 ± 0.4 ) %	567
$p\bar{K}^0\pi^+\pi^-$	( 2.6 ± 0.7 ) %	753
$pK^-\pi^+\pi^0$	( 3.4 ± 1.0 ) %	758
$pK^*(892)^-\pi^+$	[m] ( 1.1 ± 0.5 ) %	579
$p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	( 3.6 ± 1.2 ) %	758
$\Delta(1232)\bar{K}^*(892)$	seen	416
$pK^-\pi^+\pi^+\pi^-$	( 1.1 ± 0.8 ) × 10 <sup>-3</sup>	670
$pK^-\pi^+\pi^0\pi^0$	( 8 ± 4 ) × 10 <sup>-3</sup>	676

**Hadronic modes with a  $p$ :  $S = 0$  final states**

$p\pi^+\pi^-$	( 3.5 ± 2.0 ) × 10 <sup>-3</sup>	926
$p f_0(980)$	[m] ( 2.8 ± 1.9 ) × 10 <sup>-3</sup>	621
$p\pi^+\pi^+\pi^-\pi^-$	( 1.8 ± 1.2 ) × 10 <sup>-3</sup>	851
$pK^+K^-$	( 7.7 ± 3.5 ) × 10 <sup>-4</sup>	615
$p\phi$	[m] ( 8.2 ± 2.7 ) × 10 <sup>-4</sup>	589
$pK^+K^- \text{ non-}\phi$	( 3.5 ± 1.7 ) × 10 <sup>-4</sup>	615

**Hadronic modes with a hyperon:  $S = -1$  final states**

$\Lambda\pi^+$	( 9.0 ± 2.8 ) × 10 <sup>-3</sup>	863
$\Lambda\pi^+\pi^0$	( 3.6 ± 1.3 ) %	843
$\Lambda\rho^+$	< 5 %	CL=95% 638
$\Lambda\pi^+\pi^+\pi^-$	( 3.3 ± 1.0 ) %	806
$\Lambda\pi^+\eta$	( 1.8 ± 0.6 ) %	690
$\Sigma(1385)^+\eta$	[m] ( 8.5 ± 3.3 ) × 10 <sup>-3</sup>	569
$\Lambda K^+\bar{K}^0$	( 6.0 ± 2.1 ) × 10 <sup>-3</sup>	441
$\Xi(1690)^0 K^+, \Xi(1690)^0 \rightarrow \Lambda\bar{K}^0$	( 1.6 ± 0.8 ) × 10 <sup>-3</sup>	286
$\Sigma^0\pi^+$	( 9.9 ± 3.2 ) × 10 <sup>-3</sup>	824
$\Sigma^+\pi^0$	( 1.00 ± 0.34 ) %	826
$\Sigma^+\eta$	( 5.5 ± 2.3 ) × 10 <sup>-3</sup>	712
$\Sigma^+\pi^+\pi^-$	( 3.6 ± 1.0 ) %	803
$\Sigma^+\rho^0$	< 1.4 %	CL=95% 578
$\Sigma^-\pi^+\pi^+$	( 1.9 ± 0.8 ) %	798
$\Sigma^0\pi^+\pi^0$	( 1.8 ± 0.8 ) %	802
$\Sigma^0\pi^+\pi^+\pi^-$	( 1.1 ± 0.4 ) %	762
$\Sigma^+\pi^+\pi^-\pi^0$	—	766
$\Sigma^+\omega$	[m] ( 2.7 ± 1.0 ) %	568
$\Sigma^+K^+K^-$	( 2.9 ± 0.9 ) × 10 <sup>-3</sup>	346
$\Sigma^+\phi$	[m] ( 3.1 ± 1.0 ) × 10 <sup>-3</sup>	292
$\Xi(1690)^0 K^+, \Xi(1690)^0 \rightarrow \Sigma^+ K^-$	( 8.3 ± 3.5 ) × 10 <sup>-4</sup>	286
$\Sigma^+K^+K^- \text{ nonresonant}$	< 7 × 10 <sup>-4</sup>	CL=90% 346
$\Xi^0K^+$	( 3.9 ± 1.4 ) × 10 <sup>-3</sup>	652
$\Xi^-K^+\pi^+$	( 4.9 ± 1.7 ) × 10 <sup>-3</sup>	564
$\Xi(1530)^0 K^+$	[m] ( 2.6 ± 1.0 ) × 10 <sup>-3</sup>	471

### **Hadronic modes with a hyperon: $S = 0$ final states**

$\Lambda K^+$	$( 6.7 \pm 2.5 ) \times 10^{-4}$	780
$\Sigma^0 K^+$	$( 5.6 \pm 2.4 ) \times 10^{-4}$	734
$\Sigma^+ K^+ \pi^-$	$( 1.7 \pm 0.7 ) \times 10^{-3}$	668

### **Semileptonic modes**

$\Lambda \ell^+ \nu_\ell$	[n] $( 2.0 \pm 0.6 ) \%$	—
$\Lambda e^+ \nu_e$	$( 2.1 \pm 0.6 ) \%$	870
$\Lambda \mu^+ \nu_\mu$	$( 2.0 \pm 0.7 ) \%$	866

### **Inclusive modes**

$e^+$ anything	$( 4.5 \pm 1.7 ) \%$	—
$p e^+$ anything	$( 1.8 \pm 0.9 ) \%$	—
$p$ anything	$( 50 \pm 16 ) \%$	—
$p$ anything (no $\Lambda$ )	$( 12 \pm 19 ) \%$	—
$n$ anything	$( 50 \pm 16 ) \%$	—
$n$ anything (no $\Lambda$ )	$( 29 \pm 17 ) \%$	—
$\Lambda$ anything	$( 35 \pm 11 ) \%$	S=1.4
$\Sigma^\pm$ anything	[o] $( 10 \pm 5 ) \%$	—

### **$\Delta C = 1$ weak neutral current ( $C1$ ) modes, or Lepton number ( $L$ ) violating modes**

$p \mu^+ \mu^-$	$C1 < 3.4 \times 10^{-4}$	CL=90%	936
$\Sigma^- \mu^+ \mu^+$	$L < 7.0 \times 10^{-4}$	CL=90%	811

---

**$\Lambda_c(2593)^+$** 

$$I(J^P) = 0(\frac{1}{2}^-)$$

The spin-parity follows from the fact that  $\Sigma_c(2455)\pi$  decays, with little available phase space, are dominant.

Mass  $m = 2593.9 \pm 0.8$  MeV

$m - m_{\Lambda_c^+} = 308.9 \pm 0.6$  MeV ( $S = 1.1$ )

Full width  $\Gamma = 3.6^{+2.0}_{-1.3}$  MeV

$\Lambda_c^+\pi\pi$  and its submode  $\Sigma_c(2455)\pi$  — the latter just barely — are the only strong decays allowed to an excited  $\Lambda_c^+$  having this mass; and the submode seems to dominate.

 **$\Lambda_c(2593)^+$  DECAY MODES**

Fraction ( $\Gamma_i/\Gamma$ )

$p$  (MeV/c)

$\Lambda_c^+\pi^+\pi^-$	$[p] \approx 67\%$	124
$\Sigma_c(2455)^{++}\pi^-$	$24 \pm 7\%$	21
$\Sigma_c(2455)^0\pi^+$	$24 \pm 7\%$	24
$\Lambda_c^+\pi^+\pi^-$ 3-body	$18 \pm 10\%$	124
$\Lambda_c^+\pi^0$	$[q]$ not seen	261
$\Lambda_c^+\gamma$	not seen	291

 **$\Lambda_c(2625)^+$** 

$$I(J^P) = 0(\frac{3}{2}^-)$$

$J^P$  has not been measured;  $\frac{3}{2}^-$  is the quark-model prediction.

Mass  $m = 2626.6 \pm 0.8$  MeV ( $S = 1.2$ )

$m - m_{\Lambda_c^+} = 341.7 \pm 0.6$  MeV ( $S = 1.6$ )

Full width  $\Gamma < 1.9$  MeV, CL = 90%

$\Lambda_c^+\pi\pi$  and its submode  $\Sigma(2455)\pi$  are the only strong decays allowed to an excited  $\Lambda_c^+$  having this mass.

 **$\Lambda_c(2625)^+$  DECAY MODES**

Fraction ( $\Gamma_i/\Gamma$ )

$p$

		Confidence level	(MeV/c)
$\Lambda_c^+\pi^+\pi^-$	$[p] \approx 67\%$		184
$\Sigma_c(2455)^{++}\pi^-$	<5	90%	100
$\Sigma_c(2455)^0\pi^+$	<5	90%	101
$\Lambda_c^+\pi^+\pi^-$ 3-body	large		184
$\Lambda_c^+\pi^0$	$[q]$ not seen		293
$\Lambda_c^+\gamma$	not seen		319

## $\Sigma_c(2455)$

$I(J^P) = 1(\frac{1}{2}^+)$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

$\Sigma_c(2455)^{++}$  mass  $m = 2452.6 \pm 0.6$  MeV

$\Sigma_c(2455)^+$  mass  $m = 2451.3 \pm 0.7$  MeV

$\Sigma_c(2455)^0$  mass  $m = 2452.2 \pm 0.6$  MeV

$m_{\Sigma_c^{++}} - m_{\Lambda_c^+} = 167.67 \pm 0.15$  MeV

$m_{\Sigma_c^+} - m_{\Lambda_c^+} = 166.4 \pm 0.4$  MeV

$m_{\Sigma_c^0} - m_{\Lambda_c^+} = 167.32 \pm 0.15$  MeV

$m_{\Sigma_c^{++}} - m_{\Sigma_c^0} = 0.35 \pm 0.18$  MeV

$m_{\Sigma_c^+} - m_{\Sigma_c^0} = -0.9 \pm 0.4$  MeV

$\Sigma_c(2455)^{++}$  full width  $\Gamma = 2.0 \pm 0.5$  MeV

$\Sigma_c(2455)^+$  full width  $\Gamma < 4.6$  MeV, CL = 90%

$\Sigma_c(2455)^0$  full width  $\Gamma = 1.6 \pm 0.5$  MeV

$\Lambda_c^+ \pi$  is the only strong decay allowed to a  $\Sigma_c$  having this mass.

$\Sigma_c(2455)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100$ %	90

## **$\Sigma_c(2520)$**

$$I(J^P) = 1(\frac{3}{2}^+)$$

$J^P$  has not been measured;  $\frac{3}{2}^+$  is the quark-model prediction.

$\Sigma_c(2520)^{++}$  mass  $m = 2519.4 \pm 1.5$  MeV

$\Sigma_c(2520)^+$  mass  $m = 2515.9 \pm 2.4$  MeV

$\Sigma_c(2520)^0$  mass  $m = 2517.5 \pm 1.4$  MeV

$m_{\Sigma_c(2520)^{++}} - m_{\Lambda_c^+} = 234.5 \pm 1.4$  MeV

$m_{\Sigma_c(2520)^+} - m_{\Lambda_c^+} = 231.0 \pm 2.3$  MeV

$m_{\Sigma_c(2520)^0} - m_{\Lambda_c^+} = 232.6 \pm 1.3$  MeV

$m_{\Sigma_c(2520)^{++}} - m_{\Sigma_c(2520)^0} = 1.9 \pm 1.7$  MeV


$\Sigma_c(2520)^{++}$  full width  $\Gamma = 18 \pm 5$  MeV

$\Sigma_c(2520)^+$  full width  $\Gamma < 17$  MeV, CL = 90%

$\Sigma_c(2520)^0$  full width  $\Gamma = 13 \pm 5$  MeV

$\Lambda_c^+ \pi$  is the only strong decay allowed to a  $\Sigma_c$  having this mass.

<b><math>\Sigma_c(2520)</math> DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100$ %	180



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2466.3 \pm 1.4$  MeV

Mean life  $\tau = (442 \pm 26) \times 10^{-15}$  s ( $S = 1.3$ )

$$c\tau = 132 \mu\text{m}$$

No absolute branching fractions have been measured. THE FOLLOWING ARE BRANCHING RATIOS RELATIVE TO  $\Xi^- \pi^+ \pi^+$ .

$\Xi_c^+ \text{ DECAY MODES}$	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level	$p$ (MeV/c)
$\Lambda K^- \pi^+ \pi^+$	[r] $0.58 \pm 0.18$		785
$\Lambda \bar{K}^*(892)^0 \pi^+$	[m,r] $< 0.29$	90%	603
$\Sigma(1385)^+ K^- \pi^+$	[m,r] $< 0.41$	90%	677
$\Sigma^+ K^- \pi^+$	[r] $1.18 \pm 0.31$		809
$\Sigma^+ \bar{K}^*(892)^0$	[m,r] $0.92 \pm 0.30$		654
$\Sigma^0 K^- \pi^+ \pi^+$	[r] $0.49 \pm 0.26$		734
$\Xi^0 \pi^+$	[r] $0.55 \pm 0.16$		876
$\Xi^- \pi^+ \pi^+$	[r] defined as 1		850
$\Xi(1530)^0 \pi^+$	[m,r] $< 0.2$	90%	749
$\Xi^0 \pi^+ \pi^0$	[r] $2.34 \pm 0.68$		855
$\Xi^0 \pi^+ \pi^+ \pi^-$	[r] $1.74 \pm 0.50$		817
$\Xi^0 e^+ \nu_e$	[r] $2.3 \begin{array}{l} +0.7 \\ -0.9 \end{array}$		883
$\rho K^- \pi^+$	[r] $0.21 \pm 0.03$		943
$\rho \bar{K}^*(892)^0$	[m,r] $0.12 \pm 0.02$		824

$\Xi_c^0$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2471.8 \pm 1.4$  MeV

$$m_{\Xi_c^0} - m_{\Xi_c^+} = 5.5 \pm 1.8$$
 MeV

$$\text{Mean life } \tau = (98^{+23}_{-15}) \times 10^{-15}$$
 s

$$c\tau = 29$$
  $\mu\text{m}$

### Decay asymmetry parameters

$$\Xi^- \pi^+ \quad \alpha = -0.6 \pm 0.4$$

$\Xi_c^0$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda \bar{K}^0$	seen	907
$\Lambda \bar{K}^0 \pi^+ \pi^-$	seen	788
$\Lambda K^- \pi^+ \pi^+ \pi^-$	seen	704
$\Xi^- \pi^+$	seen	876
$\Xi^- \pi^+ \pi^+ \pi^-$	seen	817
$p K^- \bar{K}^*(892)^0$	seen	408
$\Omega^- K^+$	seen	523
$\Xi^- e^+ \nu_e$	seen	883
$\Xi^- \ell^+ \text{anything}$	seen	—

$\Xi_c^{\prime +}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2574.1 \pm 3.3$  MeV

$$m_{\Xi_c^{\prime +}} - m_{\Xi_c^+} = 107.8 \pm 3.0$$
 MeV

The  $\Xi_c^{\prime +} - \Xi_c^+$  mass difference is too small for any strong decay to occur.

$\Xi_c^{\prime +}$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Xi_c^+ \gamma$	seen	106

$\Xi_c'^0$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2578.8 \pm 3.2$  MeV

$$m_{\Xi_c'^0} - m_{\Xi_c^0} = 107.0 \pm 2.9 \text{ MeV}$$

The  $\Xi_c'^0 - \Xi_c^0$  mass difference is too small for any strong decay to occur.

### $\Xi_c'^0$ DECAY MODES

Fraction ( $\Gamma_i/\Gamma$ )

$p$  (MeV/c)

$$\Xi_c^0 \gamma$$

seen

105

$\Xi_c(2645)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

$J^P$  has not been measured;  $\frac{3}{2}^+$  is the quark-model prediction.

$\Xi_c(2645)^+$  mass  $m = 2647.4 \pm 2.0$  MeV (S = 1.2)

$\Xi_c(2645)^0$  mass  $m = 2644.5 \pm 1.8$  MeV

$$m_{\Xi_c(2645)^+} - m_{\Xi_c^0} = 175.6 \pm 1.4 \text{ MeV} \quad (S = 1.7)$$

$$m_{\Xi_c(2645)^0} - m_{\Xi_c^+} = 178.2 \pm 1.1 \text{ MeV}$$

$\Xi_c(2645)^+$  full width  $\Gamma < 3.1$  MeV, CL = 90%

$\Xi_c(2645)^0$  full width  $\Gamma < 5.5$  MeV, CL = 90%

$\Xi_c \pi$  is the only strong decay allowed to a  $\Xi_c$  resonance having this mass.

### $\Xi_c(2645)$ DECAY MODES

Fraction ( $\Gamma_i/\Gamma$ )

$p$  (MeV/c)

$$\Xi_c^0 \pi^+$$

seen

103

$$\Xi_c^+ \pi^-$$

seen

107

## $\Xi_c(2790)$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

$J^P$  has not been measured;  $\frac{1}{2}^-$  is the quark-model prediction.

$\Xi_c(2790)^+$  mass =  $2790.0 \pm 3.5$  MeV

$\Xi_c(2790)^0$  mass =  $2790 \pm 4$  MeV

$m_{\Xi_c(2790)^+} - m_{\Xi_c^0} = 318.2 \pm 3.2$  MeV

$m_{\Xi_c(2790)^0} - m_{\Xi_c^+} = 324.0 \pm 3.3$  MeV

$\Xi_c(2790)^+$  width < 15 MeV, CL = 90%

$\Xi_c(2790)^0$  width < 12 MeV, CL = 90%

### $\Xi_c(2790)$ DECAY MODES

Fraction ( $\Gamma_i/\Gamma$ )

$p$  (MeV/c)

$\Xi_c' \pi$

seen

156

## $\Xi_c(2815)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

$J^P$  has not been measured;  $\frac{3}{2}^-$  is the quark-model prediction.

$\Xi_c(2815)^+$  mass  $m$  =  $2814.9 \pm 1.8$  MeV

$\Xi_c(2815)^0$  mass  $m$  =  $2819.0 \pm 2.5$  MeV

$m_{\Xi_c(2815)^+} - m_{\Xi_c^+} = 348.6 \pm 1.2$  MeV

$m_{\Xi_c(2815)^0} - m_{\Xi_c^0} = 347.2 \pm 2.1$  MeV

$\Xi_c(2815)^+$  full width  $\Gamma$  < 3.5 MeV, CL = 90%

$\Xi_c(2815)^0$  full width  $\Gamma$  < 6.5 MeV, CL = 90%

The  $\Xi_c \pi \pi$  modes are consistent with being entirely via  $\Xi_c(2645) \pi$ .

### $\Xi_c(2815)$ DECAY MODES

Fraction ( $\Gamma_i/\Gamma$ )

$p$  (MeV/c)

$\Xi_c^+ \pi^+ \pi^-$

seen

196

$\Xi_c^0 \pi^+ \pi^-$

seen

193



$$I(J^P) = 0(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2697.5 \pm 2.6$  MeV ( $S = 1.2$ )

Mean life  $\tau = (64 \pm 20) \times 10^{-15}$  s

$$c\tau = 19 \text{ } \mu\text{m}$$

No absolute branching fractions have been measured.

$\Omega_c^0$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Sigma^+ K^- K^- \pi^+$	seen	691
$\Xi^0 K^- \pi^+$	seen	902
$\Xi^- K^- \pi^+ \pi^+$	seen	832
$\Omega^- \pi^+$	seen	822
$\Omega^- \pi^+ \pi^0$	seen	798
$\Omega^- \pi^- \pi^+ \pi^+$	seen	754