

$\chi_{c1}(1P)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

 $\chi_{c1}(1P)$ MASS

<u>VALUE (MeV)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3510.51 ± 0.12 OUR AVERAGE					
3509.4 ± 0.9			BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3510.53 ± 0.04 ± 0.12	513		ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
3511.3 ± 0.4 ± 0.4	30		BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-X$
3512.3 ± 0.3 ± 4.0			¹ GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3507.4 ± 1.7		91	² LEMOIGNE	82 GOLI	190 $\pi^- \text{Be} \rightarrow \gamma 2\mu$
3510.4 ± 0.6			OREGLIA	82 CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
3510.1 ± 1.1		254	³ HIMEL	80 MRK2	$e^+e^- \rightarrow J/\psi 2\gamma$
3509 ± 11		21	BRANDELIK	79B DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
3507 ± 3			³ BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3505.0 ± 4 ± 4			^{3,4} TANENBAUM	78 MRK1	e^+e^-
3513 ± 7		367	³ BIDDICK	77 CNTR	$\psi(2S) \rightarrow \gamma X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
3500 ± 10		40	TANENBAUM	75 MRK1	Hadrons γ

¹ Using mass of $\psi(2S) = 3686.0$ MeV.² $J/\psi(1S)$ mass constrained to 3097 MeV.³ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.⁴ From a simultaneous fit to radiative and hadronic decay channels. **$\chi_{c1}(1P)$ WIDTH**

<u>VALUE (MeV)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.92 ± 0.13 OUR FIT					
0.88 ± 0.11 ± 0.08		513	ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<1.3		95	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-X$
<3.8		90	GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$

$\chi_{c1}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor
Hadronic decays		
Γ_1 $3(\pi^+\pi^-)$	$(6.3 \pm 1.4) \times 10^{-3}$	
Γ_2 $2(\pi^+\pi^-)$	$(5.6 \pm 2.6) \times 10^{-3}$	2.2
Γ_3 $\pi^+\pi^-K^+K^-$	$(4.9 \pm 1.2) \times 10^{-3}$	1.1
Γ_4 $\rho^0\pi^+\pi^-$	$(3.9 \pm 3.5) \times 10^{-3}$	
Γ_5 $K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	$(3.2 \pm 2.1) \times 10^{-3}$	
Γ_6 $K_S^0K^+\pi^-$	$(2.5 \pm 0.8) \times 10^{-3}$	
Γ_7 $\pi^+\pi^-p\bar{p}$	$(5.4 \pm 2.1) \times 10^{-4}$	
Γ_8 $K^+K^-K^+K^-$	$(4.2 \pm 1.9) \times 10^{-4}$	
Γ_9 $p\bar{p}$	$(7.2 \pm 1.3) \times 10^{-5}$	
Γ_{10} $\pi^+\pi^- + K^+K^-$	$< 2.1 \times 10^{-3}$	
Radiative decays		
Γ_{11} $\gamma J/\psi(1S)$	$(31.6 \pm 3.2) \%$	
Γ_{12} $\gamma\gamma$		

$\chi_{c1}(1P)$ PARTIAL WIDTHS

$\chi_{c1}(1P) \Gamma(i)\Gamma(\gamma J/\psi(1S))/\Gamma(\text{total})$

$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$	$\Gamma_9\Gamma_{11}/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
20.8 ± 2.3 OUR FIT			
21.3 ± 2.2 OUR AVERAGE			
$21.8 \pm 1.5 \pm 2.2$	⁵ ARMSTRONG 92	E760	$\bar{p}p \rightarrow e^+e^-\gamma$
$19.9^{+4.4}_{-4.0}$	⁵ BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-X$
⁵ Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.			

$\chi_{c1}(1P)$ BRANCHING RATIOS

HADRONIC DECAYS

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
6.3 ± 1.4 OUR AVERAGE			
$5.8 \pm 0.7 \pm 1.2$	⁶ BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c1}$
22 ± 8	⁷ TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$
$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
5.6 ± 2.6 OUR AVERAGE	Error includes scale factor of 2.2.		
$4.9 \pm 0.4 \pm 1.2$	⁶ BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c1}$
16 ± 5	⁷ TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$

$\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
49 ± 12 OUR AVERAGE Error includes scale factor of 1.1.			
45 ± 4 ± 11	⁶ BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$
90 ± 40	⁷ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
39 ± 35	⁷ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
32 ± 21	⁷ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

$\Gamma(K_S^0 K^+ \pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.46 ± 0.44 ± 0.65	⁶ BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$

$\Gamma(\pi^+ \pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.4 ± 2.1 OUR AVERAGE			
4.9 ± 1.3 ± 1.7	⁶ BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$
14 ± 9	⁷ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

$\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.42 ± 0.15 ± 0.12	⁶ BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$

$\Gamma(\rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
0.72 ± 0.13 OUR FIT	

$[\Gamma(\pi^+ \pi^-) + \Gamma(K^+ K^-)]/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<21		⁷ FELDMAN	77 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<38	90	⁷ BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma \chi_{c1}$

⁶ Using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087 \pm 0.008$.

⁷ Estimated using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

————— RADIATIVE DECAYS —————

$\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$

Γ_{11}/Γ

VALUE

DOCUMENT ID

0.316±0.032 OUR FIT

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

Γ_{12}/Γ

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0015 90 ⁸ YAMADA 77 DASP $e^+e^- \rightarrow 3\gamma$

⁸ Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

$\chi_{c1}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

$$B(\chi_{c1}(1P) \rightarrow p\bar{p}) \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}$$

VALUE (units 10^{-5})

DOCUMENT ID

TECN

COMMENT

2.0±0.5 OUR FIT

1.1±1.0

⁹ BAI

98I BES

$\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow \gamma\bar{p}p$

$$B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) \times B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))$$

VALUE (units 10^{-2})

DOCUMENT ID

TECN

COMMENT

2.66±0.15 OUR FIT

2.66±0.16 OUR AVERAGE

2.56±0.12±0.20

GAISER

86 CBAL

$\psi(2S) \rightarrow \gamma X$

2.78±0.30

¹⁰ OREGLIA

82 CBAL

$\psi(2S) \rightarrow \gamma\chi_{c1}$

2.2 ±0.5

¹¹ BRANDELIK

79B DASP

$\psi(2S) \rightarrow \gamma\chi_{c1}$

2.9 ±0.5

¹¹ BARTEL

78B CNTR

$\psi(2S) \rightarrow \gamma\chi_{c1}$

5.0 ±1.5

¹² BIDDICK

77 CNTR

$e^+e^- \rightarrow \gamma X$

2.8 ±0.9

¹⁰ WHITAKER

76 MRK1

e^+e^-

$$B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}$$

VALUE (units 10^{-2})

DOCUMENT ID

TECN

COMMENT

8.7±0.7 OUR FIT

8.5±2.1

¹³ HIMEL

80 MRK2

$\psi(2S) \rightarrow \gamma\chi_{c1}$

⁹ Calculated by us. The value for $B(\chi_{c1} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

¹⁰ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

¹¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

¹² Assumes isotropic gamma distribution.

¹³ The value for $B(\psi(2S) \rightarrow \gamma\chi_{c1}) \times B(\chi_{c1} \rightarrow \gamma J/\psi(1S))$ quoted in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.138 \pm 0.18$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

MULTIPOLE AMPLITUDES IN $\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$

$a_2 = M_2/\sqrt{E_1^2 + M_2^2}$ Magnetic quadrupole fractional transition amplitude

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.002^{+0.008}_{-0.017} OUR AVERAGE				
0.002 ± 0.032 ± 0.004	2090	AMBROGIANI 02	E835	$p\bar{p} \rightarrow \chi_{c1} \rightarrow J/\psi\gamma$
-0.002 ^{+0.008} _{-0.020}	921	OREGLIA 82	CBAL	$\psi(2S) \rightarrow \chi_{c1}\gamma \rightarrow J/\psi\gamma\gamma$

$\chi_{c1}(1P)$ REFERENCES

AMBROGIANI 02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
BAI 99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI 98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI 98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARMSTRONG 92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
Also 92B	PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BAGLIN 86B	PL B172 455	C. Baglin (LAPP, CERN, GENO, LYON, OSLO+)	
GAISER 86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
LEMOIGNE 82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
OREGLIA 82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
Also 82B	Private Comm.	M.J. Oreglia	(EFI)
HIMEL 80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
Also 82	Private Comm.	G. Trilling	(LBL, UCB)
BRANDELIK 79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL 78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM 78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
Also 82	Private Comm.	G. Trilling	(LBL, UCB)
BIDDICK 77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
FELDMAN 77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
YAMADA 77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)
WHITAKER 76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)
TANENBAUM 75	PRL 35 1323	W.M. Tanenbaum <i>et al.</i>	(LBL, SLAC)

OTHER RELATED PAPERS

BARBERIS 00G	PL B485 357	D. Barberis <i>et al.</i>	(Omega expt.)
BARATE 83	PL 121B 449	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, IND)
BRAUNSCH... 75B	PL 57B 407	W. Braunschweig <i>et al.</i>	(DASP Collab.)
SIMPSON 75	PRL 35 699	J.W. Simpson <i>et al.</i>	(STAN, PENN)