

$\phi(1680)$ $I^G(J^{PC}) = 0^-(1^{--})$ **$\phi(1680)$ MASS** **e^+e^- PRODUCTION**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1680±20 OUR ESTIMATE				
1681± 8 OUR AVERAGE				
1700±20		1 CLEGG	94 RVUE	$e^+e^- \rightarrow K^+K^-$, $K_S^0 K\pi$
1657±27	367	BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1680±10		2 BUON	82 DM1	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 1500		3 ACHASOV	98H RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$, $\omega\pi^+\pi^-$, K^+K^-
~ 1900		4 ACHASOV	98H RVUE	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1655±17		5 BISELLO	88B DM2	$e^+e^- \rightarrow K^+K^-$
1677±12		6 MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K\pi$

PHOTOPRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1726±22	BUSENITZ	89 TPS	$\gamma p \rightarrow K^+K^-X$
1760±20	ATKINSON	85C OMEG	20–70 $\gamma p \rightarrow K\bar{K}X$
1690±10	ASTON	81F OMEG	25–70 $\gamma p \rightarrow K^+K^-X$

1 Using BISELLO 88B and MANE 82 data.

2 From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_S^0 K_L^0$, $K_S^0 K^\pm \pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.

3 Using data from IVANOV 81, BARKOV 87, BISELLO 88B, DOLINSKY 91, and ANTONELLI 92.

4 Using the data from BISELLO 91C.

5 From global fit including ρ , ω , ϕ and $\rho(1700)$ assume mass 1570 MeV and width 510 MeV for ρ radial excitation.6 Fit to one channel only, neglecting interference with ω , $\rho(1700)$. **$\phi(1680)$ WIDTH** **e^+e^- PRODUCTION**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
150±50 OUR ESTIMATE				This is only an educated guess; the error given is larger than the error on the average of the published values.
• • • We do not use the following data for averages, fits, limits, etc. • • •				
300±60		7 CLEGG	94 RVUE	$e^+e^- \rightarrow K^+K^-$, $K_S^0 K\pi$
146±55	367	BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
207±45		8 BISELLO	88B DM2	$e^+e^- \rightarrow K^+K^-$
185±22		9 BUON	82 DM1	$e^+e^- \rightarrow$ hadrons
102±36		10 MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K\pi$

PHOTOPRODUCTION

VALUE (MeV)		DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
121±47	BUSENITZ	89 TPS	$\gamma p \rightarrow K^+ K^- X$	
80±40	ATKINSON	85C OMEG	20–70 $\gamma p \rightarrow K\bar{K}X$	
100±40	ASTON	81F OMEG	25–70 $\gamma p \rightarrow K^+ K^- X$	
7	Using BISELLO 88B and MANE 82 data.			
8	From global fit including ρ , ω , ϕ and $\rho(1700)$			
9	From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_S^0 K_L^0$, $K_S^0 K^\pm\pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.			
10	Fit to one channel only, neglecting interference with ω , $\rho(1700)$.			

$\phi(1680)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}^*(892) + c.c.$	dominant
Γ_2 $K_S^0 K\pi$	seen
Γ_3 $K\bar{K}$	seen
Γ_4 $e^+ e^-$	seen
Γ_5 $\omega\pi\pi$	not seen
Γ_6 $K^+ K^- \pi^0$	

$\phi(1680) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel (I) in e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(I)$ or the branching ratio $\Gamma(I)/\text{total}$.

$\Gamma(K\bar{K}^*(892)+c.c.) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_4/\Gamma$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.48±0.14	367	BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp$

$\phi(1680)$ BRANCHING RATIOS

$\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma(K_S^0 K\pi)$	Γ_1/Γ_2		
VALUE	DOCUMENT ID	TECN	COMMENT
dominant	MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp$

$\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892)+c.c.)$	Γ_3/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.07 ±0.01	BUON	82 DM1	e^+e^-

$\Gamma(\omega\pi\pi)/\Gamma(K\bar{K}^*(892)+\text{c.c.})$ Γ_5/Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.10	BUON	82 DM1	$e^+ e^-$

 $\phi(1680)$ REFERENCES

ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>
BUSENITZ	89	PR D40 1	J.K. Busenitz <i>et al.</i>
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>
		Translated from ZETFP 46 132.	
ATKINSON	85C	ZPHY C27 233	M. Atkinson <i>et al.</i>
BUON	82	PL 118B 221	J. Buon <i>et al.</i>
MANE	82	PL 112B 178	F. Mane <i>et al.</i>
ASTON	81F	PL 104B 231	D. Aston
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>

 Γ_5/Γ_1 **OTHER RELATED PAPERS**

ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ACHASOV	97F	PAN 60 2029	N.N. Achasov, A.A. Kozhevnikov	(NOVM)
		Translated from YAF 60 2212.		
ATKINSON	86C	ZPHY C30 541	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	84	NP B231 15	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	84B	NP B231 1	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	83C	NP B229 269	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORsay)
MANE	81	PL 99B 261	F. Mane <i>et al.</i>	(ORsay)
ASTON	80F	NP B174 269	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)