

# $\omega(1420)$

$I^G(J^{PC}) = 0^-(1^{--})$

## $\omega(1420)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>(1400–1450) OUR ESTIMATE</b>				

• • • We do not use the following data for averages, fits, limits, etc. • • •

1490 $\pm$ 50 $\pm$ 25	6.5k	1 ACHASOV	02E RVUE	$0.98\text{--}2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1450 $\pm$ 10		2 HENNER	02 RVUE	$1.2\text{--}2.0 e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$
1373 $\pm$ 70	177	3 AKHMETSHIN 00D	CMD2	$1.2\text{--}1.38 e^+ e^- \rightarrow \omega\pi^+\pi^-$
1370 $\pm$ 25	5095	ANISOVICH	00H SPEC	$0.0 p\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$
$1400^{+100}_{-200}$		4 ACHASOV	98H RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
$\sim 1400$		5 ACHASOV	98H RVUE	$e^+ e^- \rightarrow \omega\pi^+\pi^-$
$\sim 1460$		6 ACHASOV	98H RVUE	$e^+ e^- \rightarrow K^+ K^-$
1440 $\pm$ 70		7 CLEGG	94 RVUE	
1419 $\pm$ 31	315	8 ANTONELLI	92 DM2	$1.34\text{--}2.4 e^+ e^- \rightarrow \rho\pi$

<sup>1</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, and ACHASOV 02E data on the  $\pi^+ \pi^- \pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E.

<sup>2</sup> Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

<sup>3</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.

<sup>4</sup> Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.

<sup>5</sup> Using the data from ANTONELLI 92.

<sup>6</sup> Using the data from IVANOV 81 and BISELLO 88B.

<sup>7</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

<sup>8</sup> From a fit to two Breit-Wigner functions interfering between them and with the  $\omega, \phi$  tails with fixed  $(+, -, +)$  phases.

## $\omega(1420)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>(180–250) OUR ESTIMATE</b>				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1210^{+300}_{-200} \pm 170$	6.5k	9 ACHASOV	02E RVUE	$0.98\text{--}2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
199 $\pm$ 15		10 HENNER	02 RVUE	$1.2\text{--}2.0 e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$
188 $\pm$ 45	177	11 AKHMETSHIN 00D	CMD2	$1.2\text{--}1.38 e^+ e^- \rightarrow \omega\pi^+\pi^-$
$360^{+100}_{-60}$	5095	ANISOVICH	00H SPEC	$0.0 p\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$
240 $\pm$ 70		12 CLEGG	94 RVUE	
174 $\pm$ 59	315	13 ANTONELLI	92 DM2	$1.34\text{--}2.4 e^+ e^- \rightarrow \rho\pi$

- <sup>9</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, and ACHASOV 02E data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E.
- <sup>10</sup> Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.
- <sup>11</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.
- <sup>12</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
- <sup>13</sup> From a fit to two Breit-Wigner functions interfering between them and with the  $\omega,\phi$  tails with fixed  $(+,-,+)$  phases.

### $\omega(1420)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \rho\pi$	dominant
$\Gamma_2 \omega\pi\pi$	possibly seen
$\Gamma_3 b_1(1235)\pi$	seen
$\Gamma_4 e^+e^-$	

### $\omega(1420) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

$$\Gamma(\rho\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}^2 \quad \Gamma_1\Gamma_4/\Gamma^2$$

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.529 \pm 0.076 \pm 0.030$	15,16 ACHASOV	02E RVUE	$0.98-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0, \omega\pi^+\pi^-$
$0.625 \pm 0.160$	17,18 CLEGG	94 RVUE	
$0.466 \pm 0.178$	19,20 ANTONELLI	92 DM2	$1.34-2.4 e^+e^- \rightarrow \rho\pi$

$$\Gamma(\omega\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}^2 \quad \Gamma_2\Gamma_4/\Gamma^2$$

VALUE (units $10^{-8}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$0.68^{+2.04}_{-0.68} \pm 0.23$		15,16 ACHASOV	02E RVUE	$0.98-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0, \omega\pi^+\pi^-$
$1.3 \pm 1.3$	612	14 AKHMETSHIN 00D	CMD2	$1.2-2.4 e^+e^- \rightarrow \omega\pi^+\pi^-$

<sup>14</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.

<sup>15</sup> Calculated by us from the cross section at the peak.

<sup>16</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, and ACHASOV 02E data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E.

<sup>17</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

<sup>18</sup> From the partial and leptonic width given by the authors.

<sup>19</sup> From a fit to two Breit-Wigner functions interfering between them and with the  $\omega,\phi$  tails with fixed  $(+,-,+)$  phases.

<sup>20</sup> From the product of the leptonic width and partial branching ratio given by the authors.

**$\omega(1420)$  BRANCHING RATIOS** **$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$** 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
≈ 0.01	21 ACHASOV	02E RVUE	$0.98\text{--}2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0, \omega \pi^+ \pi^-$	
0.301 ± 0.029	22 HENNER	02 RVUE	$1.2\text{--}2.0 e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$	
possibly seen	AKHMETSHIN 00D	CMD2	$e^+ e^- \rightarrow \omega \pi^+ \pi^-$	

 **$\Gamma(\omega\pi\pi)/\Gamma(b_1(1235)\pi)$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma_3$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
0.60 ± 0.16	5095	ANISOVICH 00H	SPEC	$0.0 p\bar{p} \rightarrow \omega \pi^0 \pi^0 \pi^0$	

 **$\Gamma(\rho\pi)/\Gamma_{\text{total}}$** 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
≈ 0.99	21 ACHASOV	02E RVUE	$0.98\text{--}2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0, \omega \pi^+ \pi^-$	
0.699 ± 0.029	22 HENNER	02 RVUE	$1.2\text{--}2.0 e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$	

 **$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$** 

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
23 ± 1	22 HENNER	02 RVUE	$1.2\text{--}2.0 e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$	
21	From the combined fit of ANTONELLI 92, ACHASOV 01E, and ACHASOV 02E data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E.			
22	Assuming that the $\omega(1420)$ decays into $\rho \pi$ and $\omega \pi \pi$ only.			

 **$\omega(1420)$  REFERENCES**

ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>	
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ANISOVICH	00H	PL B485 341	A.V. Anisovich <i>et al.</i>	
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from ZETFP 46 132.		
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORsay)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)

## OTHER RELATED PAPERS

ACHASOV	02B	PAN 65 153 Translated from YAF 65 158.	N.N. Achasov, A.A. Kozhevnikov
CLOSE	02	PR D65 092003	F.E. Close, A. Donnachie, Yu.S. Kalashnikova
ACHASOV	00J	PR D62 117503	N.N. Achasov, A.A. Kozhevnikov
ABELE	99D	PL B468 178	A. Abele <i>et al.</i> (Crystal Barrel Collab.)
BELOZEROVA	98	PPN 29 63 Translated from FECAY 29 148.	T.S. Belozerova, V.K. Henner
ACHASOV	97F	PAN 60 2029 Translated from YAF 60 2212.	N.N. Achasov, A.A. Kozhevnikov (NOVM)
ATKINSON	87	ZPHY C34 157	M. Atkinson <i>et al.</i> (BONN, CERN, GLAS+)
ATKINSON	84	NP B231 15	M. Atkinson <i>et al.</i> (BONN, CERN, GLAS+)
ATKINSON	83B	PL 127B 132	M. Atkinson <i>et al.</i> (BONN, CERN, GLAS+)