

$$I(J^P) = \frac{1}{2}(0^-)$$

Quantum numbers not measured. Values shown are quark-model predictions.

See also the B^\pm/B^0 ADMIXTURE and $B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE sections.

B^\pm MASS

The fit uses m_{B^+} , $(m_{B^0} - m_{B^+})$, and m_{B^0} to determine m_{B^+} , m_{B^0} , and the mass difference.

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5279.0±0.5 OUR FIT				
5279.1±0.5 OUR AVERAGE				
5279.1±0.4 ±0.4	526	¹ CSORNA	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5279.1±1.7 ±1.4	147	ABE	96B CDF	$p\bar{p}$ at 1.8 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5278.8±0.54±2.0	362	ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5278.3±0.4 ±2.0		BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
5280.5±1.0 ±2.0		² ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5275.8±1.3 ±3.0	32	ALBRECHT	87C ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5278.2±1.8 ±3.0	12	³ ALBRECHT	87D ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5278.6±0.8 ±2.0		BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ CSORNA 00 uses fully reconstructed 526 $B^+ \rightarrow J/\psi(\prime) K^+$ events and invariant masses without beam constraint.

² ALBRECHT 90J assumes 10580 for $\Upsilon(4S)$ mass. Supersedes ALBRECHT 87C and ALBRECHT 87D.

³ Found using fully reconstructed decays with $J/\psi(1S)$. ALBRECHT 87D assume $m_{\Upsilon(4S)} = 10577$ MeV.

B^\pm MEAN LIFE

See $B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE section for data on B -hadron mean life averaged over species of bottom particles.

“OUR EVALUATION” is an average of the data listed below performed by the Heavy Flavor Averaging Group (HFAG) as described in our review “Production and Decay of b -flavored Hadrons” in the B^\pm Section of the Listings. The averaging procedure takes into account correlations between the measurements and asymmetric lifetime errors.

<u>VALUE (10^{-12} s)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.671±0.018 OUR NEW EVALUATION				$[(1.674 \pm 0.018) \times 10^{-12}$ s OUR 2002 EVALUATION]
Average is meaningless.				$[(1.669 \pm 0.017) \times 10^{-12}$ s OUR 2002 AVERAGE]
1.695±0.026±0.015		⁴ ABE	02H BELL	$e^+e^- \rightarrow \Upsilon(4S)$

1.636 ± 0.058 ± 0.025		⁵ ACOSTA	02C CDF	$p\bar{p}$ at 1.8 TeV	
1.673 ± 0.032 ± 0.023		⁴ AUBERT	01F BABR	$e^+e^- \rightarrow \Upsilon(4S)$	
1.648 ± 0.049 ± 0.035		⁶ BARATE	00R ALEP	$e^+e^- \rightarrow Z$	
1.643 ± 0.037 ± 0.025		⁷ ABBIENDI	99J OPAL	$e^+e^- \rightarrow Z$	
1.637 ± 0.058 ^{+0.045} _{-0.043}		⁶ ABE	98Q CDF	$p\bar{p}$ at 1.8 TeV	
1.66 ± 0.06 ± 0.03		⁷ ACCIARRI	98S L3	$e^+e^- \rightarrow Z$	
1.66 ± 0.06 ± 0.05		⁷ ABE	97J SLD	$e^+e^- \rightarrow Z$	
1.58 ^{+0.21} _{-0.18} ± 0.04 ± 0.03	94	⁵ BUSKULIC	96J ALEP	$e^+e^- \rightarrow Z$	
1.61 ± 0.16 ± 0.12		^{6,8} ABREU	95Q DLPH	$e^+e^- \rightarrow Z$	
1.72 ± 0.08 ± 0.06		⁹ ADAM	95 DLPH	$e^+e^- \rightarrow Z$	
1.52 ± 0.14 ± 0.09		⁶ AKERS	95T OPAL	$e^+e^- \rightarrow Z$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1.68 ± 0.07 ± 0.02		⁵ ABE	98B CDF	Repl. by ACOSTA 02C	
1.56 ± 0.13 ± 0.06		⁶ ABE	96C CDF	Repl. by ABE 98Q	
1.58 ± 0.09 ± 0.03		¹⁰ BUSKULIC	96J ALEP	$e^+e^- \rightarrow Z$	
1.58 ± 0.09 ± 0.04		⁶ BUSKULIC	96J ALEP	Repl. by BARATE 00R	
1.70 ± 0.09		¹¹ ADAM	95 DLPH	$e^+e^- \rightarrow Z$	
1.61 ± 0.16 ± 0.05	148	⁵ ABE	94D CDF	Repl. by ABE 98B	
1.30 ^{+0.33} _{-0.29} ± 0.16	92	⁶ ABREU	93D DLPH	Sup. by ABREU 95Q	
1.56 ± 0.19 ± 0.13	134	⁹ ABREU	93G DLPH	Sup. by ADAM 95	
1.51 ^{+0.30} _{-0.28} ± 0.12 ± 0.14	59	⁶ ACTON	93C OPAL	Sup. by AKERS 95T	
1.47 ^{+0.22} _{-0.19} ± 0.15 ± 0.14	77	⁶ BUSKULIC	93D ALEP	Sup. by BUSKULIC 96J	

⁴ Events are selected in which one B meson is fully reconstructed while the second B meson is reconstructed inclusively.

⁵ Measured mean life using fully reconstructed decays.

⁶ Data analyzed using $D/D^* \ell X$ event vertices.

⁷ Data analyzed using charge of secondary vertex.

⁸ ABREU 95Q assumes $B(B^0 \rightarrow D^{*-} \ell^+ \nu_\ell) = 3.2 \pm 1.7\%$.

⁹ Data analyzed using vertex-charge technique to tag B charge.

¹⁰ Combined result of $D/D^* \ell X$ analysis and fully reconstructed B analysis.

¹¹ Combined ABREU 95Q and ADAM 95 result.

B^+ DECAY MODES

B^- modes are charge conjugates of the modes below. Modes which do not identify the charge state of the B are listed in the B^\pm/B^0 ADMIXTURE section.

The branching fractions listed below assume 50% $B^0\bar{B}^0$ and 50% B^+B^- production at the $\Upsilon(4S)$. We have attempted to bring older measurements up to date by rescaling their assumed $\Upsilon(4S)$ production ratio to 50:50 and their assumed D , D_S , D^* , and ψ branching ratios to current values whenever this would affect our averages and best limits significantly.

Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Semileptonic and leptonic modes		
Γ_1 $\ell^+ \nu_\ell$ anything	[a] (10.2 \pm 0.9) %	
Γ_2 $\bar{D}^0 \ell^+ \nu_\ell$	[a] (2.15 \pm 0.22) %	
Γ_3 $\bar{D}^*(2007)^0 \ell^+ \nu_\ell$	[a] (6.5 \pm 0.5) %	
Γ_4 $\bar{D}_1(2420)^0 \ell^+ \nu_\ell$	(5.6 \pm 1.6) $\times 10^{-3}$	
Γ_5 $\bar{D}_2^*(2460)^0 \ell^+ \nu_\ell$	< 8 $\times 10^{-3}$	CL=90%
Γ_6 $\pi^0 e^+ \nu_e$	(9.0 \pm 2.8) $\times 10^{-5}$	
Γ_7 $\omega \ell^+ \nu_\ell$	[a] < 2.1 $\times 10^{-4}$	CL=90%
Γ_8 $\omega \mu^+ \nu_\mu$		
Γ_9 $\rho^0 \ell^+ \nu_\ell$	[a] (1.34 ^{+0.32} _{-0.35}) $\times 10^{-4}$	
Γ_{10} $e^+ \nu_e$	< 1.5 $\times 10^{-5}$	CL=90%
Γ_{11} $\mu^+ \nu_\mu$	< 2.1 $\times 10^{-5}$	CL=90%
Γ_{12} $\tau^+ \nu_\tau$	< 5.7 $\times 10^{-4}$	CL=90%
Γ_{13} $e^+ \nu_e \gamma$	< 2.0 $\times 10^{-4}$	CL=90%
Γ_{14} $\mu^+ \nu_\mu \gamma$	< 5.2 $\times 10^{-5}$	CL=90%
D, D^*, or D_S modes		
Γ_{15} $\bar{D}^0 \pi^+$	(4.98 \pm 0.29) $\times 10^{-3}$	
Γ_{16} $D_{CP(+1)} \pi^+$	[b]	
Γ_{17} $D_{CP(-1)} \pi^+$	[b]	
Γ_{18} $\bar{D}^0 \rho^+$	(1.34 \pm 0.18) %	
Γ_{19} $\bar{D}^0 K^+$	(3.7 \pm 0.6) $\times 10^{-4}$	S=1.1
Γ_{20} $D_{CP(+1)} K^+$	[b]	
Γ_{21} $D_{CP(-1)} K^+$	[b]	
Γ_{22} $\bar{D}^0 K^*(892)^+$	(6.1 \pm 2.3) $\times 10^{-4}$	
Γ_{23} $\bar{D}^0 K^+ \bar{K}^0$	(5.5 \pm 1.6) $\times 10^{-4}$	
Γ_{24} $\bar{D}^0 K^+ \bar{K}^*(892)^0$	(7.5 \pm 1.7) $\times 10^{-4}$	

Γ ₂₅	$\bar{D}^0 \pi^+ \pi^+ \pi^-$	(1.1 ± 0.4) %	
Γ ₂₆	$\bar{D}^0 \pi^+ \pi^+ \pi^-$ nonresonant	(5 ± 4) × 10 ⁻³	
Γ ₂₇	$\bar{D}^0 \pi^+ \rho^0$	(4.2 ± 3.0) × 10 ⁻³	
Γ ₂₈	$\bar{D}^0 a_1(1260)^+$	(5 ± 4) × 10 ⁻³	
Γ ₂₉	$\bar{D}^0 \omega \pi^+$	(4.1 ± 0.9) × 10 ⁻³	
Γ ₃₀	$D^{*}(2010)^- \pi^+ \pi^+$	(2.1 ± 0.6) × 10 ⁻³	
Γ ₃₁	$D^- \pi^+ \pi^+$	< 1.4 × 10 ⁻³	CL=90%
Γ ₃₂	$\bar{D}^{*}(2007)^0 \pi^+$	(4.6 ± 0.4) × 10 ⁻³	
Γ ₃₃	$\bar{D}^{*}(2007)^0 \omega \pi^+$	(4.5 ± 1.2) × 10 ⁻³	
Γ ₃₄	$\bar{D}^{*}(2007)^0 \rho^+$	(1.55 ± 0.31) %	
Γ ₃₅	$\bar{D}^{*}(2007)^0 K^+$	(3.6 ± 1.0) × 10 ⁻⁴	
Γ ₃₆	$\bar{D}^{*}(2007)^0 K^{*}(892)^+$	(7.2 ± 3.4) × 10 ⁻⁴	
Γ ₃₇	$\bar{D}^{*}(2007)^0 K^+ \bar{K}^0$	< 1.06 × 10 ⁻³	CL=90%
Γ ₃₈	$\bar{D}^{*}(2007)^0 K^+ K^{*}(892)^0$	(1.5 ± 0.4) × 10 ⁻³	
Γ ₃₉	$\bar{D}^{*}(2007)^0 \pi^+ \pi^+ \pi^-$	(9.4 ± 2.6) × 10 ⁻³	
Γ ₄₀	$\bar{D}^{*}(2007)^0 a_1(1260)^+$	(1.9 ± 0.5) %	
Γ ₄₁	$\bar{D}^{*}(2007)^0 \pi^- \pi^+ \pi^+ \pi^0$	(1.8 ± 0.4) %	
Γ ₄₂	$D^{*}(2010)^+ \pi^0$	< 1.7 × 10 ⁻⁴	CL=90%
Γ ₄₃	$\bar{D}^{*}(2010)^+ K^0$	< 9.5 × 10 ⁻⁵	CL=90%
Γ ₄₄	$D^{*}(2010)^- \pi^+ \pi^+ \pi^0$	(1.5 ± 0.7) %	
Γ ₄₅	$D^{*}(2010)^- \pi^+ \pi^+ \pi^+ \pi^-$	< 1 %	CL=90%
Γ ₄₆	$\bar{D}_1^{*}(2420)^0 \pi^+$	(1.5 ± 0.6) × 10 ⁻³	S=1.3
Γ ₄₇	$\bar{D}_1^{*}(2420)^0 \rho^+$	< 1.4 × 10 ⁻³	CL=90%
Γ ₄₈	$\bar{D}_2^{*}(2460)^0 \pi^+$	< 1.3 × 10 ⁻³	CL=90%
Γ ₄₉	$\bar{D}_2^{*}(2460)^0 \rho^+$	< 4.7 × 10 ⁻³	CL=90%
Γ ₅₀	$\bar{D}^0 D_s^+$	(1.3 ± 0.4) %	
Γ ₅₁	$\bar{D}^0 D_s^{*+}$	(9 ± 4) × 10 ⁻³	
Γ ₅₂	$\bar{D}^{*}(2007)^0 D_s^+$	(1.2 ± 0.5) %	
Γ ₅₃	$\bar{D}^{*}(2007)^0 D_s^{*+}$	(2.7 ± 1.0) %	
Γ ₅₄	$D_s^{(*)+} \bar{D}^{*0}$	(2.7 ± 1.2) %	
Γ ₅₅	$\bar{D}^{*}(2007)^0 D^{*}(2010)^+$	< 1.1 %	CL=90%
Γ ₅₆	$\bar{D}^0 D^{*}(2010)^+ + \bar{D}^{*}(2007)^0 D^+$	< 1.3 %	CL=90%
Γ ₅₇	$\bar{D}^0 D^+$	< 6.7 × 10 ⁻³	CL=90%
Γ ₅₈	$D_s^+ \pi^0$	< 2.0 × 10 ⁻⁴	CL=90%
Γ ₅₉	$D_s^{*+} \pi^0$	< 3.3 × 10 ⁻⁴	CL=90%
Γ ₆₀	$D_s^+ \eta$	< 5 × 10 ⁻⁴	CL=90%
Γ ₆₁	$D_s^{*+} \eta$	< 8 × 10 ⁻⁴	CL=90%
Γ ₆₂	$D_s^+ \rho^0$	< 4 × 10 ⁻⁴	CL=90%
Γ ₆₃	$D_s^{*+} \rho^0$	< 5 × 10 ⁻⁴	CL=90%
Γ ₆₄	$D_s^+ \omega$	< 5 × 10 ⁻⁴	CL=90%
Γ ₆₅	$D_s^{*+} \omega$	< 7 × 10 ⁻⁴	CL=90%

Γ_{66}	$D_s^+ a_1(1260)^0$	< 2.2	$\times 10^{-3}$	CL=90%
Γ_{67}	$D_s^{*+} a_1(1260)^0$	< 1.6	$\times 10^{-3}$	CL=90%
Γ_{68}	$D_s^+ \phi$	< 3.2	$\times 10^{-4}$	CL=90%
Γ_{69}	$D_s^{*+} \phi$	< 4	$\times 10^{-4}$	CL=90%
Γ_{70}	$D_s^+ \bar{K}^0$	< 1.1	$\times 10^{-3}$	CL=90%
Γ_{71}	$D_s^{*+} \bar{K}^0$	< 1.1	$\times 10^{-3}$	CL=90%
Γ_{72}	$D_s^+ \bar{K}^*(892)^0$	< 5	$\times 10^{-4}$	CL=90%
Γ_{73}	$D_s^{*+} \bar{K}^*(892)^0$	< 4	$\times 10^{-4}$	CL=90%
Γ_{74}	$D_s^- \pi^+ K^+$	< 8	$\times 10^{-4}$	CL=90%
Γ_{75}	$D_s^{*-} \pi^+ K^+$	< 1.2	$\times 10^{-3}$	CL=90%
Γ_{76}	$D_s^- \pi^+ K^*(892)^+$	< 6	$\times 10^{-3}$	CL=90%
Γ_{77}	$D_s^{*-} \pi^+ K^*(892)^+$	< 8	$\times 10^{-3}$	CL=90%

Charmonium modes

Γ_{78}	$\eta_c K^+$	(9.0 ± 2.7)	$\times 10^{-4}$	
Γ_{79}	$J/\psi(1S) K^+$	(1.00 ± 0.04)	$\times 10^{-3}$	
Γ_{80}	$J/\psi(1S) K^+ \pi^+ \pi^-$	(7.7 ± 2.0)	$\times 10^{-4}$	
Γ_{81}	$J/\psi(1S) K^*(892)^+$	(1.35 ± 0.10)	$\times 10^{-3}$	
Γ_{82}	$J/\psi(1S) K(1270)^+$	(1.8 ± 0.5)	$\times 10^{-3}$	
Γ_{83}	$J/\psi(1S) K(1400)^+$	< 5	$\times 10^{-4}$	CL=90%
Γ_{84}	$J/\psi(1S) \phi K^+$	$(8.8 \begin{smallmatrix} +3.7 \\ -3.3 \end{smallmatrix})$	$\times 10^{-5}$	
Γ_{85}	$J/\psi(1S) \pi^+$	(4.0 ± 0.5)	$\times 10^{-5}$	
Γ_{86}	$J/\psi(1S) \rho^+$	< 7.7	$\times 10^{-4}$	CL=90%
Γ_{87}	$J/\psi(1S) a_1(1260)^+$	< 1.2	$\times 10^{-3}$	CL=90%
Γ_{88}	$\psi(2S) K^+$	(6.8 ± 0.4)	$\times 10^{-4}$	
Γ_{89}	$\psi(2S) K^*(892)^+$	(9.2 ± 2.2)	$\times 10^{-4}$	
Γ_{90}	$\psi(2S) K^+ \pi^+ \pi^-$	(1.9 ± 1.2)	$\times 10^{-3}$	
Γ_{91}	$\chi_{c0}(1P) K^+$	$(6.0 \begin{smallmatrix} +2.4 \\ -2.1 \end{smallmatrix})$	$\times 10^{-4}$	
Γ_{92}	$\chi_{c1}(1P) K^+$	(6.8 ± 1.1)	$\times 10^{-4}$	
Γ_{93}	$\chi_{c1}(1P) K^*(892)^+$	< 2.1	$\times 10^{-3}$	CL=90%

K or K* modes

Γ_{94}	$K^0 \pi^+$	(1.87 ± 0.22)	$\times 10^{-5}$	
Γ_{95}	$K^+ \pi^0$	(1.17 ± 0.15)	$\times 10^{-5}$	
Γ_{96}	$\eta' K^+$	(7.5 ± 0.7)	$\times 10^{-5}$	
Γ_{97}	$\eta' K^*(892)^+$	< 3.5	$\times 10^{-5}$	CL=90%
Γ_{98}	ηK^+	< 6.9	$\times 10^{-6}$	CL=90%
Γ_{99}	$\eta K^*(892)^+$	$(2.6 \begin{smallmatrix} +1.0 \\ -0.9 \end{smallmatrix})$	$\times 10^{-5}$	
Γ_{100}	ωK^+	$(9.2 \begin{smallmatrix} +2.8 \\ -2.5 \end{smallmatrix})$	$\times 10^{-6}$	
Γ_{101}	$\omega K^*(892)^+$	< 8.7	$\times 10^{-5}$	CL=90%
Γ_{102}	$K^*(892)^0 \pi^+$	$(1.9 \begin{smallmatrix} +0.6 \\ -0.8 \end{smallmatrix})$	$\times 10^{-5}$	

Γ_{103}	$K^*(892)^+\pi^0$	< 3.1	$\times 10^{-5}$	CL=90%
Γ_{104}	$K^+\pi^-\pi^+$	(5.6 ± 1.0)	$\times 10^{-5}$	
Γ_{105}	$K^+\pi^-\pi^+$ nonresonant	< 2.8	$\times 10^{-5}$	CL=90%
Γ_{106}	$K^+f_0(980)$			
Γ_{107}	$K^+\rho^0$	< 1.2	$\times 10^{-5}$	CL=90%
Γ_{108}	$K_2^*(1430)^0\pi^+$	< 6.8	$\times 10^{-4}$	CL=90%
Γ_{109}	$K^-\pi^+\pi^+$	< 7.0	$\times 10^{-6}$	CL=90%
Γ_{110}	$K^-\pi^+\pi^+$ nonresonant	< 5.6	$\times 10^{-5}$	CL=90%
Γ_{111}	$K_1(1400)^0\pi^+$	< 2.6	$\times 10^{-3}$	CL=90%
Γ_{112}	$K^0\pi^+\pi^0$	< 6.6	$\times 10^{-5}$	CL=90%
Γ_{113}	$K^0\rho^+$	< 4.8	$\times 10^{-5}$	CL=90%
Γ_{114}	$K^*(892)^+\pi^+\pi^-$	< 1.1	$\times 10^{-3}$	CL=90%
Γ_{115}	$K^*(892)^+\rho^0$	< 7.4	$\times 10^{-5}$	CL=90%
Γ_{116}	$K^*(892)^+K^*(892)^0$	< 7.1	$\times 10^{-5}$	CL=90%
Γ_{117}	$K_1(1400)^+\rho^0$	< 7.8	$\times 10^{-4}$	CL=90%
Γ_{118}	$K_2^*(1430)^+\rho^0$	< 1.5	$\times 10^{-3}$	CL=90%
Γ_{119}	$K^+\bar{K}^0$	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{120}	$\bar{K}^0K^+\pi^0$	< 2.4	$\times 10^{-5}$	CL=90%
Γ_{121}	$K^+K^-\pi^+$	< 1.2	$\times 10^{-5}$	CL=90%
Γ_{122}	$K^+K^-\pi^+$ nonresonant	< 7.5	$\times 10^{-5}$	CL=90%
Γ_{123}	$K^+K^+\pi^-$	< 3.2	$\times 10^{-6}$	CL=90%
Γ_{124}	$K^+K^+\pi^-$ nonresonant	< 8.79	$\times 10^{-5}$	CL=90%
Γ_{125}	$K^+K^*(892)^0$	< 5.3	$\times 10^{-6}$	CL=90%
Γ_{126}	$K^+K^-K^+$	(3.5 ± 0.6)	$\times 10^{-5}$	
Γ_{127}	$K^+\phi$	$(7.9 \begin{smallmatrix} +2.0 \\ -1.8 \end{smallmatrix})$	$\times 10^{-6}$	S=1.6
Γ_{128}	$K^+K^-K^+$ nonresonant	< 3.8	$\times 10^{-5}$	CL=90%
Γ_{129}	$K^*(892)^+K^+K^-$	< 1.6	$\times 10^{-3}$	CL=90%
Γ_{130}	$K^*(892)^+\phi$	$(10 \begin{smallmatrix} +5 \\ -4 \end{smallmatrix})$	$\times 10^{-6}$	
Γ_{131}	$K_1(1400)^+\phi$	< 1.1	$\times 10^{-3}$	CL=90%
Γ_{132}	$K_2^*(1430)^+\phi$	< 3.4	$\times 10^{-3}$	CL=90%
Γ_{133}	$K^*(892)^+\gamma$	(3.8 ± 0.5)	$\times 10^{-5}$	
Γ_{134}	$K_1(1270)^+\gamma$	< 9.9	$\times 10^{-5}$	CL=90%
Γ_{135}	$K^+\pi^-\pi^+\gamma$	$(2.4 \begin{smallmatrix} +0.6 \\ -0.5 \end{smallmatrix})$	$\times 10^{-5}$	
Γ_{136}	$K^*(892)^0\pi^+\gamma$	$(2.0 \begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix})$	$\times 10^{-5}$	
Γ_{137}	$K^+\rho^0\gamma$	< 2.0	$\times 10^{-5}$	CL=90%
Γ_{138}	$K^+\pi^-\pi^+\gamma$ (NR)	< 9.2	$\times 10^{-6}$	CL=90%
Γ_{139}	$K_1(1400)^+\gamma$	< 5.0	$\times 10^{-5}$	CL=90%
Γ_{140}	$K_2^*(1430)^+\gamma$	< 1.4	$\times 10^{-3}$	CL=90%
Γ_{141}	$K^*(1680)^+\gamma$	< 1.9	$\times 10^{-3}$	CL=90%
Γ_{142}	$K_3^*(1780)^+\gamma$	< 5.5	$\times 10^{-3}$	CL=90%
Γ_{143}	$K_4^*(2045)^+\gamma$	< 9.9	$\times 10^{-3}$	CL=90%

Light unflavored meson modes

Γ_{144}	$\rho^+ \gamma$	< 1.3	$\times 10^{-5}$	CL=90%
Γ_{145}	$\pi^+ \pi^0$	(7.4 ± 2.4)	$\times 10^{-6}$	
Γ_{146}	$\pi^+ \pi^+ \pi^-$	< 1.3	$\times 10^{-4}$	CL=90%
Γ_{147}	$\rho^0 \pi^+$	(8.6 ± 2.0)	$\times 10^{-6}$	
Γ_{148}	$\pi^+ f_0(980)$	< 1.4	$\times 10^{-4}$	CL=90%
Γ_{149}	$\pi^+ f_2(1270)$	< 2.4	$\times 10^{-4}$	CL=90%
Γ_{150}	$\pi^+ \pi^- \pi^+$ nonresonant	< 4.1	$\times 10^{-5}$	CL=90%
Γ_{151}	$\pi^+ \pi^0 \pi^0$	< 8.9	$\times 10^{-4}$	CL=90%
Γ_{152}	$\rho^+ \pi^0$	< 4.3	$\times 10^{-5}$	CL=90%
Γ_{153}	$\pi^+ \pi^- \pi^+ \pi^0$	< 4.0	$\times 10^{-3}$	CL=90%
Γ_{154}	$\rho^+ \rho^0$	< 1.0	$\times 10^{-3}$	CL=90%
Γ_{155}	$a_1(1260)^+ \pi^0$	< 1.7	$\times 10^{-3}$	CL=90%
Γ_{156}	$a_1(1260)^0 \pi^+$	< 9.0	$\times 10^{-4}$	CL=90%
Γ_{157}	$\omega \pi^+$	$(6.4 \begin{smallmatrix} +1.8 \\ -1.6 \end{smallmatrix})$	$\times 10^{-6}$	S=1.3
Γ_{158}	$\omega \rho^+$	< 6.1	$\times 10^{-5}$	CL=90%
Γ_{159}	$\eta \pi^+$	< 5.7	$\times 10^{-6}$	CL=90%
Γ_{160}	$\eta' \pi^+$	< 7.0	$\times 10^{-6}$	CL=90%
Γ_{161}	$\eta' \rho^+$	< 3.3	$\times 10^{-5}$	CL=90%
Γ_{162}	$\eta \rho^+$	< 1.5	$\times 10^{-5}$	CL=90%
Γ_{163}	$\phi \pi^+$	< 1.4	$\times 10^{-6}$	CL=90%
Γ_{164}	$\phi \rho^+$	< 1.6	$\times 10^{-5}$	
Γ_{165}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^-$	< 8.6	$\times 10^{-4}$	CL=90%
Γ_{166}	$\rho^0 a_1(1260)^+$	< 6.2	$\times 10^{-4}$	CL=90%
Γ_{167}	$\rho^0 a_2(1320)^+$	< 7.2	$\times 10^{-4}$	CL=90%
Γ_{168}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$	< 6.3	$\times 10^{-3}$	CL=90%
Γ_{169}	$a_1(1260)^+ a_1(1260)^0$	< 1.3	%	CL=90%

Charged particle (h^\pm) modes

$$h^\pm = K^\pm \text{ or } \pi^\pm$$

Γ_{170}	$h^+ \pi^0$	$(1.6 \begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix})$	$\times 10^{-5}$	
Γ_{171}	ωh^+	$(1.38 \begin{smallmatrix} +0.27 \\ -0.24 \end{smallmatrix})$	$\times 10^{-5}$	
Γ_{172}	$h^+ X^0$ (Familon)	< 4.9	$\times 10^{-5}$	CL=90%

Baryon modes

Γ_{173}	$p \bar{p} \pi^+$	< 3.7	$\times 10^{-6}$	CL=90%
Γ_{174}	$p \bar{p} \pi^+$ nonresonant	< 5.3	$\times 10^{-5}$	CL=90%
Γ_{175}	$p \bar{p} \pi^+ \pi^+ \pi^-$	< 5.2	$\times 10^{-4}$	CL=90%
Γ_{176}	$p \bar{p} K^+$	$(4.3 \begin{smallmatrix} +1.2 \\ -1.0 \end{smallmatrix})$	$\times 10^{-6}$	
Γ_{177}	$p \bar{p} K^+$ nonresonant	< 8.9	$\times 10^{-5}$	CL=90%
Γ_{178}	$p \bar{\Lambda}$	< 2.2	$\times 10^{-6}$	CL=90%
Γ_{179}	$p \bar{\Lambda} \pi^+ \pi^-$	< 2.0	$\times 10^{-4}$	CL=90%

Γ_{180}	$\bar{\Delta}^0 p$		< 3.8	$\times 10^{-4}$	CL=90%
Γ_{181}	$\Delta^{++} \bar{p}$		< 1.5	$\times 10^{-4}$	CL=90%
Γ_{182}	$D^+ p \bar{p}$		< 1.5	$\times 10^{-5}$	CL=90%
Γ_{183}	$D^*(2010)^+ p \bar{p}$		< 1.5	$\times 10^{-5}$	CL=90%
Γ_{184}	$\Lambda_c^- p \pi^+$		(2.1 ± 0.7)	$\times 10^{-4}$	
Γ_{185}	$\Lambda_c^- p \pi^+ \pi^0$		(1.8 ± 0.6)	$\times 10^{-3}$	
Γ_{186}	$\Lambda_c^- p \pi^+ \pi^+ \pi^-$		(2.3 ± 0.7)	$\times 10^{-3}$	
Γ_{187}	$\Lambda_c^- p \pi^+ \pi^+ \pi^- \pi^0$		< 1.34	%	CL=90%
Γ_{188}	$\bar{\Sigma}_c(2455)^0 p$		< 8	$\times 10^{-5}$	CL=90%
Γ_{189}	$\bar{\Sigma}_c(2520)^0 p$		< 4.6	$\times 10^{-5}$	CL=90%
Γ_{190}	$\bar{\Sigma}_c(2455)^0 p \pi^0$		(4.4 ± 1.8)	$\times 10^{-4}$	
Γ_{191}	$\bar{\Sigma}_c(2455)^0 p \pi^- \pi^+$		(4.4 ± 1.7)	$\times 10^{-4}$	
Γ_{192}	$\Sigma_c(2455)^{-} p \pi^+ \pi^+$		(2.8 ± 1.2)	$\times 10^{-4}$	
Γ_{193}	$\Lambda_{c1}^- p \pi^+$		< 1.9	$\times 10^{-4}$	CL=90%

**Lepton Family number (LF) or Lepton number (L) violating modes, or
 $\Delta B = 1$ weak neutral current (B1) modes**

Γ_{194}	$\pi^+ e^+ e^-$	B1	< 3.9	$\times 10^{-3}$	CL=90%
Γ_{195}	$\pi^+ \mu^+ \mu^-$	B1	< 9.1	$\times 10^{-3}$	CL=90%
Γ_{196}	$K^+ e^+ e^-$	B1	< 9	$\times 10^{-7}$	CL=90%
Γ_{197}	$K^+ \mu^+ \mu^-$	B1	(10^{+5}_{-4})	$\times 10^{-7}$	
Γ_{198}	$K^+ \bar{\nu} \nu$	B1	< 2.4	$\times 10^{-4}$	CL=90%
Γ_{199}	$K^*(892)^+ e^+ e^-$	B1	< 8.9	$\times 10^{-6}$	CL=90%
Γ_{200}	$K^*(892)^+ \mu^+ \mu^-$	B1	< 3.9	$\times 10^{-6}$	CL=90%
Γ_{201}	$\pi^+ e^+ \mu^-$	LF	< 6.4	$\times 10^{-3}$	CL=90%
Γ_{202}	$\pi^+ e^- \mu^+$	LF	< 6.4	$\times 10^{-3}$	CL=90%
Γ_{203}	$K^+ e^+ \mu^-$	LF	< 8	$\times 10^{-7}$	CL=90%
Γ_{204}	$K^+ e^- \mu^+$	LF	< 6.4	$\times 10^{-3}$	CL=90%
Γ_{205}	$K^*(892)^+ e^\pm \mu^\mp$	LF	< 7.9	$\times 10^{-6}$	CL=90%
Γ_{206}	$\pi^- e^+ e^+$	L	< 1.6	$\times 10^{-6}$	CL=90%
Γ_{207}	$\pi^- \mu^+ \mu^+$	L	< 1.4	$\times 10^{-6}$	CL=90%
Γ_{208}	$\pi^- e^+ \mu^+$	L	< 1.3	$\times 10^{-6}$	CL=90%
Γ_{209}	$\rho^- e^+ e^+$	B1	< 2.6	$\times 10^{-6}$	CL=90%
Γ_{210}	$\rho^- \mu^+ \mu^+$	B1	< 5.0	$\times 10^{-6}$	CL=90%
Γ_{211}	$\rho^- e^+ \mu^+$	LF	< 3.3	$\times 10^{-6}$	CL=90%
Γ_{212}	$K^- e^+ e^+$	L	< 1.0	$\times 10^{-6}$	CL=90%
Γ_{213}	$K^- \mu^+ \mu^+$	L	< 1.8	$\times 10^{-6}$	CL=90%
Γ_{214}	$K^- e^+ \mu^+$	L	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{215}	$K^*(892)^- e^+ e^+$	B1	< 2.8	$\times 10^{-6}$	CL=90%
Γ_{216}	$K^*(892)^- \mu^+ \mu^+$	B1	< 8.3	$\times 10^{-6}$	CL=90%
Γ_{217}	$K^*(892)^- e^+ \mu^+$	LF	< 4.4	$\times 10^{-6}$	CL=90%

- [a] An ℓ indicates an e or a μ mode, not a sum over these modes.
 [b] An $CP(\pm 1)$ indicates the $CP=+1$ and $CP=-1$ eigenstates of the D^0 - \bar{D}^0 system.

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 9 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 1.1$ for 7 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$\times 85$	16
	$\times 79$

B^+ BRANCHING RATIOS

$\Gamma(\ell^+ \nu_\ell \text{ anything}) / \Gamma_{\text{total}}$ Γ_1 / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.1025 ± 0.0057 ± 0.0065	¹² ARTUSO	97 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.101 ± 0.018 ± 0.015	ATHANAS	94 CLE2	Sup. by ARTUSO 97
¹² ARTUSO 97 uses partial reconstruction of $B \rightarrow D^* \ell \nu_\ell$ and inclusive semileptonic branching ratio from BARISH 96B ($0.1049 \pm 0.0017 \pm 0.0043$).			

$\Gamma(\bar{D}^0 \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_2 / Γ

$\ell = e$ or μ , not sum over e and μ modes.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0215 ± 0.0022 OUR AVERAGE			

0.0221 ± 0.0013 ± 0.0019	¹³ BARTELT	99 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
--------------------------	-----------------------	---------	------------------------------------

0.016 ± 0.006 ± 0.003	¹⁴ FULTON	91 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	----------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0194 ± 0.0015 ± 0.0034	¹⁵ ATHANAS	97 CLE2	Repl. by BARTELT 99
--------------------------	-----------------------	---------	---------------------

¹³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹⁴ FULTON 91 assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\Upsilon(4S)$.

¹⁵ ATHANAS 97 uses missing energy and missing momentum to reconstruct neutrino.

$\Gamma(\bar{D}^*(2007)^0 \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_3 / Γ

$\ell = e$ or μ , not sum over e and μ modes.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.065 ± 0.005 OUR NEW AVERAGE		[0.053 ± 0.008 OUR 2002 AVERAGE]		

0.0650 ± 0.0020 ± 0.0043	¹⁶ ADAM	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
--------------------------	--------------------	---------	------------------------------------

0.066 ± 0.016 ± 0.015	¹⁷ ALBRECHT	92C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	------------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0650 ± 0.0020 ± 0.0043		18 BRIERE	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.0513 ± 0.0054 ± 0.0064	302	19 BARISH	95 CLE2	Repl. by ADAM 03
seen	398	20 SANGHERA	93 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.041 ± 0.008 ^{+0.008} / _{-0.009}		21 FULTON	91 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.070 ± 0.018 ± 0.014		22 ANTREASYAN 90B	CBAL	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁶ Simultaneous measurements of both $B^0 \rightarrow D^*(2010)^- \ell \nu$ and $B^+ \rightarrow \bar{D}^*(2007)^0 \ell \nu$.

¹⁷ ALBRECHT 92C reports $0.058 \pm 0.014 \pm 0.013$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^- \pi^+)$. Assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\Upsilon(4S)$.

¹⁸ The results are based on the same analysis and data sample reported in ADAM 03.

¹⁹ BARISH 95 use $B(D^0 \rightarrow K^- \pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*0} \rightarrow D^0 \pi^0) = (63.6 \pm 2.3 \pm 3.3)\%$.

²⁰ Combining $\bar{D}^{*0} \ell^+ \nu_\ell$ and $\bar{D}^{*-} \ell^+ \nu_\ell$ SANGHERA 93 test $V-A$ structure and fit the decay angular distributions to obtain $A_{FB} = 3/4 * (\Gamma^- - \Gamma^+) / \Gamma = 0.14 \pm 0.06 \pm 0.03$. Assuming a value of V_{cb} , they measure V , A_1 , and A_2 , the three form factors for the $D^* \ell \nu_\ell$ decay, where results are slightly dependent on model assumptions.

²¹ Assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\Upsilon(4S)$. Uncorrected for D and D^* branching ratio assumptions.

²² ANTREASYAN 90B is average over B and $\bar{D}^*(2010)$ charge states.

$\Gamma(\bar{D}_1(2420)^0 \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_4 / Γ

<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0056 ± 0.0013 ± 0.0009		23 ANASTASSOV 98	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

²³ ANASTASSOV 98 result is derived from the measurement of $B(B^+ \rightarrow \bar{D}_1^0 \ell^+ \nu_\ell) \times B(\bar{D}_1^0 \rightarrow D^{*+} \pi^-) = (0.373 \pm 0.085 \pm 0.052 \pm 0.024)\%$ by assuming $B(\bar{D}_1^0 \rightarrow D^{*+} \pi^-) = 67\%$, where the third error includes theoretical uncertainties.

$\Gamma(\bar{D}_2^*(2460)^0 \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_5 / Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 8 × 10⁻³	90	24 ANASTASSOV 98	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

²⁴ ANASTASSOV 98 result is derived from the measurement of $B(B^+ \rightarrow \bar{D}_2^{*0} \ell^+ \nu_\ell) \times B(\bar{D}_2^{*0} \rightarrow D^{*+} \pi^-) < 0.16\%$ at 90% CL by assuming $B(\bar{D}_2^{*0} \rightarrow D^{*+} \pi^-) = 20\%$.

$\Gamma(\pi^0 e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_6 / Γ

<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.9 ± 0.2 ± 0.2		25 ALEXANDER 96T	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<22	90	ANTREASYAN 90B	CBAL	$e^+e^- \rightarrow \Upsilon(4S)$
-----	----	----------------	------	-----------------------------------

²⁵ Derived based in the reported B^0 result by assuming isospin symmetry: $\Gamma(B^0 \rightarrow \pi^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \pi^0 \ell^+ \nu)$.

$\Gamma(\omega \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_7 / Γ
 $\ell = e \text{ or } \mu, \text{ not sum over } e \text{ and } \mu \text{ modes.}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-4}$	90	²⁶ BEAN	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

²⁶ BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\rho^0 \ell^+ \nu_\ell)$ and $\Gamma(\rho^- \ell^+ \nu_\ell)$ with this result, they obtain a limit $<(1.6-2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow \omega \ell^+ \nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.8-0.13$ at 90% CL is derived as well.

$\Gamma(\omega \mu^+ \nu_\mu) / \Gamma_{\text{total}}$ Γ_8 / Γ

VALUE	DOCUMENT ID	TECN
-------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ²⁷ ALBRECHT 91C ARG

²⁷ In ALBRECHT 91C, one event is fully reconstructed providing evidence for the $b \rightarrow u$ transition.

$\Gamma(\rho^0 \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_9 / Γ
 $\ell = e \text{ or } \mu, \text{ not sum over } e \text{ and } \mu \text{ modes.}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$1.34 \pm 0.15^{+0.28}_{-0.32}$		²⁸ BEHRENS	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
---------------------------------	--	-----------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.40 \pm 0.21^{+0.32}_{-0.33}$		²⁸ BEHRENS	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
---------------------------------	--	-----------------------	---------	------------------------------------

$1.2 \pm 0.2^{+0.3}_{-0.4}$		²⁸ ALEXANDER	96T CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------------	--	-------------------------	----------	------------------------------------

<2.1	90	²⁹ BEAN	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
--------	----	--------------------	----------	------------------------------------

²⁸ Derived based in the reported B^0 result by assuming isospin symmetry: $\Gamma(B^0 \rightarrow \rho^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \rho^0 \ell^+ \nu) \approx 2\Gamma(B^+ \rightarrow \omega \ell^+ \nu)$.

²⁹ BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\omega^0 \ell^+ \nu_\ell)$ and $\Gamma(\rho^- \ell^+ \nu_\ell)$ with this result, they obtain a limit $<(1.6-2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow \rho^0 \ell^+ \nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.8-0.13$ at 90% CL is derived as well.

$\Gamma(e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_{10} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-5}$	90	ARTUSO	95 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\mu^+ \nu_\mu) / \Gamma_{\text{total}}$ Γ_{11} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-5}$	90	ARTUSO	95 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\tau^+ \nu_\tau) / \Gamma_{\text{total}}$ Γ_{12} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.7 \times 10^{-4}$	90	³⁰ ACCIARRI	97F L3	$e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<8.3 \times 10^{-4}$	90	³¹ BARATE	01E ALEP	$e^+e^- \rightarrow Z$
$<8.4 \times 10^{-4}$	90	³² BROWDER	01 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$<1.04 \times 10^{-2}$	90	³³ ALBRECHT	95D ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<2.2 \times 10^{-3}$	90	ARTUSO	95 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$<1.8 \times 10^{-3}$	90	³⁴ BUSKULIC	95 ALEP	$e^+e^- \rightarrow Z$

³⁰ ACCIARRI 97F uses missing-energy technique and $f(b \rightarrow B^-) = (38.2 \pm 2.5)\%$.

³¹ The energy-flow and b -tagging algorithms were used.

³² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³³ ALBRECHT 95D use full reconstruction of one B decay as tag.

³⁴ BUSKULIC 95 uses same missing-energy technique as in $\bar{b} \rightarrow \tau^+ \nu_\tau X$, but analysis is restricted to endpoint region of missing-energy distribution.

$\Gamma(e^+ \nu_e \gamma)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-4}$	90	³⁵ BROWDER	97 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

³⁵ BROWDER 97 uses the hermiticity of the CLEO II detector to reconstruct the neutrino energy and momentum.

$\Gamma(\mu^+ \nu_\mu \gamma)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-5}$	90	³⁶ BROWDER	97 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

³⁶ BROWDER 97 uses the hermiticity of the CLEO II detector to reconstruct the neutrino energy and momentum.

$\Gamma(\bar{D}^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.00498 ± 0.00029 OUR NEW AVERAGE		[0.0053 ± 0.0005 OUR 2002 AVERAGE]		

0.00497 ± 0.00012 ± 0.00029 ^{37,38} AHMED 02B CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

0.0050 ± 0.0007 ± 0.0006 54 ³⁹ BORTOLETTO92 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

0.0054 $\begin{matrix} +0.0018 \\ -0.0015 \end{matrix}$ $\begin{matrix} +0.0012 \\ -0.0009 \end{matrix}$ 14 ⁴⁰ BEBEK 87 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0055 ± 0.0004 ± 0.0005 304 ⁴¹ ALAM 94 CLE2 Repl. by

0.0020 ± 0.0008 ± 0.0006 12 ³⁹ ALBRECHT 90J ARG $e^+e^- \rightarrow \Upsilon(4S)$

0.0019 ± 0.0010 ± 0.0006 7 ⁴² ALBRECHT 88K ARG $e^+e^- \rightarrow \Upsilon(4S)$

³⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³⁸ AHMED 02B reports an additional uncertainty on the branching ratios to account for 4.5% uncertainty on relative production of B^0 and B^+ , which is not included here.

³⁹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D .

⁴⁰ BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

⁴¹ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

⁴² ALBRECHT 88K assumes $B^0 \bar{B}^0 : B^+ B^-$ ratio is 45:55. Superseded by ALBRECHT 90J.

$\Gamma(\overline{D}^0 \rho^+)/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0134 ± 0.0018 OUR AVERAGE				
0.0135 ± 0.0012 ± 0.0015	212	⁴³ ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.013 ± 0.004 ± 0.004	19	⁴⁴ ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.021 ± 0.008 ± 0.009	10	⁴⁵ ALBRECHT	88K ARG	$e^+e^- \rightarrow \Upsilon(4S)$

⁴³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

⁴⁴ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D .

⁴⁵ ALBRECHT 88K assumes $B^0 \overline{B}^0 : B^+ B^-$ ratio is 45:55.

 $\Gamma(\overline{D}^0 K^+)/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.7 ± 0.6 OUR AVERAGE Error includes scale factor of 1.1.			
4.19 ± 0.57 ± 0.40	⁴⁶ ABE	01I BELL	$e^+e^- \rightarrow \Upsilon(4S)$
2.92 ± 0.80 ± 0.28	⁴⁷ ATHANAS	98 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

⁴⁶ ABE 01I reports $B(B^+ \rightarrow \overline{D}^0 K^+)/B(B^+ \rightarrow \overline{D}^0 \pi^+) = 0.079 \pm 0.009 \pm 0.006$. We multiply by our best value $B(B^+ \rightarrow \overline{D}^0 \pi^+) = (5.3 \pm 0.5) \times 10^{-3}$. Our first error is their experiment's error and the second error is systematic error from using our best value.

⁴⁷ ATHANAS 98 reports $[B(B^+ \rightarrow \overline{D}^0 K^+)]/[B(B^+ \rightarrow \overline{D}^0 \pi^+)] = 0.055 \pm 0.014 \pm 0.005$. We multiply by our best value $B(B^+ \rightarrow \overline{D}^0 \pi^+) = (5.3 \pm 0.5) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\overline{D}^0 K^+)/\Gamma(\overline{D}^0 \pi^+)$ Γ_{19}/Γ_{15}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.094 ± 0.009 ± 0.007	⁴⁸ ABE	03D BELL	$e^+e^- \rightarrow \Upsilon(4S)$

⁴⁸ Flavor specific D^0 meson is reconstructed via $D^0 \rightarrow K^- \pi^+$.

 $\Gamma(D_{CP(+1)} K^+)/\Gamma(D_{CP(+1)} \pi^+)$ Γ_{20}/Γ_{16}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.125 ± 0.036 ± 0.010	⁴⁹ ABE	03D BELL	$e^+e^- \rightarrow \Upsilon(4S)$

⁴⁹ $CP=+1$ eigenstate of $D^0 \overline{D}^0$ system is reconstructed via $K^+ K^-$ and $\pi^+ \pi^-$.

 $\Gamma(D_{CP(-1)} K^+)/\Gamma(D_{CP(-1)} \pi^+)$ Γ_{21}/Γ_{17}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.119 ± 0.028 ± 0.006	⁵⁰ ABE	03D BELL	$e^+e^- \rightarrow \Upsilon(4S)$

⁵⁰ $CP=-1$ eigenstate of $D^0 \overline{D}^0$ system is reconstructed via $K_S^0 \pi^0$, $K_S^0 \omega$, $K_S^0 \phi$, $K_S^0 \eta$, and $K_S^0 \eta'$.

 $\Gamma(\overline{D}^0 K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{22}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
(6.1 ± 1.6 ± 1.7) × 10⁻⁴	⁵¹ MAHAPATRA	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

⁵¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^0 K^+ \overline{K}^0)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$5.5 \pm 1.4 \pm 0.8$	52 DRUTSKOY 02	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
---	----------------	------	------------------------------------

⁵² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^0 K^+ \overline{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$7.5 \pm 1.3 \pm 1.1$	53 DRUTSKOY 02	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
---	----------------	------	------------------------------------

⁵³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{D}^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.0115 \pm 0.0029 \pm 0.0021$	54 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
--	-----------------	------	------------------------------------

⁵⁴ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\overline{D}^0 \pi^+ \pi^+ \pi^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.0051 \pm 0.0034 \pm 0.0023$	55 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
--	-----------------	------	------------------------------------

⁵⁵ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\overline{D}^0 \pi^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.0042 \pm 0.0023 \pm 0.0020$	56 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
--	-----------------	------	------------------------------------

⁵⁶ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\overline{D}^0 a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.0045 \pm 0.0019 \pm 0.0031$	57 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
--	-----------------	------	------------------------------------

⁵⁷ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\overline{D}^0 \omega \pi^+)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.0041 \pm 0.0007 \pm 0.0006$	58 ALEXANDER 01B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
--	------------------	------	------------------------------------

⁵⁸ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25_{-10}^{+10}$ MeV and width $547 \pm 86_{-45}^{+46}$ MeV.

$\Gamma(D^*(2010)^- \pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_{30}/Γ		
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
0.0021 ± 0.0006 OUR AVERAGE						
0.0019 ± 0.0007 ± 0.0003		14	⁵⁹ ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	
0.0026 ± 0.0014 ± 0.0007		11	⁶⁰ ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	
0.0024 ^{+0.0017 +0.0010} _{-0.0016 -0.0006}		3	⁶¹ BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.004	90		⁶² BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	
0.005 ± 0.002 ± 0.003		7	⁶³ ALBRECHT	87C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	

⁵⁹ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

⁶⁰ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D .

⁶¹ BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

⁶² BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$. The authors also find the product branching fraction into $D^{**} \pi$ followed by $D^{**} \rightarrow D^*(2010) \pi$ to be $0.0014^{+0.0008}_{-0.0006} \pm 0.0003$ where D^{**} represents all orbitally excited D mesons.

⁶³ ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

$\Gamma(D^- \pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_{31}/Γ		
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<0.0014						
	90		⁶⁴ ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.007	90		⁶⁵ BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	
0.0025 ^{+0.0041 +0.0024} _{-0.0023 -0.0008}		1	⁶⁶ BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

⁶⁴ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \rightarrow K^- \pi^+ \pi^+)$.

⁶⁵ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D . The product branching fraction into $D_0^*(2340) \pi$ followed by $D_0^*(2340) \rightarrow D \pi$ is < 0.005 at 90%CL and into $D_2^*(2460)$ followed by $D_2^*(2460) \rightarrow D \pi$ is < 0.004 at 90%CL.

⁶⁶ BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. $B(D^- \rightarrow K^+ \pi^- \pi^-) = (9.1 \pm 1.3 \pm 0.4)\%$ is assumed.

$$\Gamma(\bar{D}^*(2007)^0 \pi^+)/\Gamma_{\text{total}} \qquad \Gamma_{32}/\Gamma$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.0046 ± 0.0004 OUR AVERAGE

0.00434 ± 0.00047 ± 0.00018 ⁶⁷ BRANDENB... 98 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

0.0052 ± 0.0007 ± 0.0007 71 ⁶⁸ ALAM 94 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

0.0072 ± 0.0018 ± 0.0016 ⁶⁹ BORTOLETTO92 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

0.0040 ± 0.0014 ± 0.0012 9 ⁶⁹ ALBRECHT 90J ARG $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0027 ± 0.0044 ⁷⁰ BEBEK 87 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

⁶⁷ BRANDENBURG 98 assume equal production of B^+ and B^0 at $\Upsilon(4S)$ and use the D^* reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of $B(D^* \rightarrow D\pi)$.

⁶⁸ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0\pi^0)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

⁶⁹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

⁷⁰ This is a derived branching ratio, using the inclusive pion spectrum and other two-body B decays. BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$.

$$\Gamma(\bar{D}^*(2007)^0 \omega \pi^+)/\Gamma_{\text{total}} \qquad \Gamma_{33}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.0045 ± 0.0010 ± 0.0007 ⁷¹ ALEXANDER 01B CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

⁷¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega\pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25_{-5}^{+10}$ MeV and width $547 \pm 86_{-45}^{+46}$ MeV.

$$\Gamma(\bar{D}^*(2007)^0 \rho^+)/\Gamma_{\text{total}} \qquad \Gamma_{34}/\Gamma$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.0155 ± 0.0031 OUR AVERAGE

0.0168 ± 0.0021 ± 0.0028 86 ⁷² ALAM 94 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

0.010 ± 0.006 ± 0.004 7 ⁷³ ALBRECHT 90J ARG $e^+e^- \rightarrow \Upsilon(4S)$

⁷² ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0\pi^0)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$. The nonresonant $\pi^+\pi^0$ contribution under the ρ^+ is negligible.

⁷³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

$$\Gamma(\bar{D}^*(2007)^0 K^+)/\Gamma_{\text{total}} \qquad \Gamma_{35}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

(3.59 ± 0.97 ± 0.31) × 10⁻⁴ ⁷⁴ ABE 01i BELL $e^+e^- \rightarrow \Upsilon(4S)$

⁷⁴ ABE 01i reports $B(B^+ \rightarrow \bar{D}^*(2007)^0 K^+)/B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+) = 0.078 \pm 0.019 \pm 0.009$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+) = (4.6 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and the second error is systematic error from using our best value.

$\Gamma(\bar{D}^*(2007)^0 K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$(7.2 \pm 2.2 \pm 2.6) \times 10^{-4}$		⁷⁵ MAHAPATRA 02	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

⁷⁵ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and an unpolarized final state.

$\Gamma(\bar{D}^*(2007)^0 K^+ \bar{K}^0)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<10.6	90	⁷⁶ DRUTSKOY 02	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

⁷⁶ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 K^+ K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
15.3 ± 3.1 ± 2.9		⁷⁷ DRUTSKOY 02	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

⁷⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0094 ± 0.0020 ± 0.0017	48	^{78,79} ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

⁷⁸ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

⁷⁹ The three pion mass is required to be between 1.0 and 1.6 GeV consistent with an a_1 meson. (If this channel is dominated by a_1^+ , the branching ratio for $\bar{D}^{*0} a_1^+$ is twice that for $\bar{D}^{*0} \pi^+ \pi^+ \pi^-$.)

$\Gamma(\bar{D}^*(2007)^0 a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.0188 ± 0.0040 ± 0.0034		^{80,81} ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

⁸⁰ ALAM 94 value is twice their $\Gamma(\bar{D}^*(2007)^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ value based on their observation that the three pions are dominantly in the $a_1(1260)$ mass range 1.0 to 1.6 GeV.

⁸¹ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

$\Gamma(\bar{D}^*(2007)^0 \pi^- \pi^+ \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.0180 ± 0.0024 ± 0.0027		⁸² ALEXANDER 01B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

⁸² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25_{-5}^{+10}$ MeV and width $547 \pm 86_{-45}^{+46}$ MeV.

$\Gamma(D^*(2010)^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00017	90	⁸³ BRANDENB... 98	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

⁸³ BRANDENBURG 98 assume equal production of B^+ and B^0 at $\Upsilon(4S)$ and use the D^* partial reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of $B(D^* \rightarrow D \pi)$.

$\Gamma(\bar{D}^*(2010)^+ K^0)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.5 \times 10^{-5}$	90	⁸⁴ GRITSAN	01 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

⁸⁴ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.0152 \pm 0.0071 \pm 0.0001$	26	⁸⁵ ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.043 \pm 0.013 \pm 0.026$	24	⁸⁶ ALBRECHT	87C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------------	----	------------------------	---------	------------------------------------

⁸⁵ ALBRECHT 90J reports $0.018 \pm 0.007 \pm 0.005$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

⁸⁶ ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

$\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	90	⁸⁷ ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

⁸⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

$\Gamma(\bar{D}_1^*(2420)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0015 ± 0.0006 OUR AVERAGE		Error includes scale factor of 1.3.		
$0.0011 \pm 0.0005 \pm 0.0002$	8	⁸⁸ ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.0025 \pm 0.0007 \pm 0.0006$		⁸⁹ ALBRECHT	94D ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

⁸⁸ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and assuming $B(D_1(2420)^0 \rightarrow D^*(2010)^+ \pi^-) = 67\%$.

⁸⁹ ALBRECHT 94D assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ assuming $B(D_1(2420)^0 \rightarrow D^*(2010)^+ \pi^-) = 67\%$.

$\Gamma(\bar{D}_1^*(2420)^0 \rho^+)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0014	90	⁹⁰ ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

⁹⁰ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ assuming $B(D_1(2420)^0 \rightarrow D^*(2010)^+ \pi^-) = 67\%$.

$\Gamma(\overline{D}_2^*(2460)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{48}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0013	90	⁹¹ ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<0.0028	90	⁹² ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<0.0023	90	⁹³ ALBRECHT	94D ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

⁹¹ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \rightarrow K^- \pi^+ \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^+ \pi^-) = 30\%$.

⁹² ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \rightarrow K^- \pi^+ \pi^+)$, the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+ \pi^-) = 20\%$.

⁹³ ALBRECHT 94D assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+ \pi^-) = 30\%$.

$\Gamma(\overline{D}_2^*(2460)^0 \rho^+)/\Gamma_{\text{total}}$ Γ_{49}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0047	90	⁹⁴ ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<0.005	90	⁹⁵ ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

⁹⁴ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \rightarrow K^- \pi^+ \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^+ \pi^-) = 30\%$.

⁹⁵ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \rightarrow K^- \pi^+ \pi^+)$, the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+ \pi^-) = 20\%$.

$\Gamma(\overline{D}^0 D_s^+)/\Gamma_{\text{total}}$ Γ_{50}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.013 ± 0.004 OUR AVERAGE				
0.0122 ± 0.0032 $^{+0.0029}_{-0.0030}$		⁹⁶ GIBAUT	96 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.018 \pm 0.009 \pm 0.004$		⁹⁷ ALBRECHT	92G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.016 \pm 0.007 \pm 0.004$	5	⁹⁸ BORTOLETTO	90 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

⁹⁶ GIBAUT 96 reports $0.0126 \pm 0.0022 \pm 0.0025$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹⁷ ALBRECHT 92G reports $0.024 \pm 0.012 \pm 0.004$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 branching ratios, e.g., $B(D^0 \rightarrow K^- \pi^+) = 3.71 \pm 0.25\%$.

⁹⁸ BORTOLETTO 90 reports 0.029 ± 0.013 for $B(D_s^+ \rightarrow \phi \pi^+) = 0.02$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\overline{D}^0 D_s^{*+})/\Gamma_{\text{total}} \qquad \Gamma_{51}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.009 ± 0.004 OUR AVERAGE

0.0084 ± 0.0031 ^{+0.0020} _{-0.0021}	99 GIBAUT	96 CLE2	e ⁺ e ⁻ → $\Upsilon(4S)$
---	-----------	---------	--

0.012 ± 0.009 ± 0.003	100 ALBRECHT	92G ARG	e ⁺ e ⁻ → $\Upsilon(4S)$
-----------------------	--------------	---------	--

⁹⁹ GIBAUT 96 reports $0.0087 \pm 0.0027 \pm 0.0017$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁰⁰ ALBRECHT 92G reports $0.016 \pm 0.012 \pm 0.003$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 branching ratios, e.g., $B(D^0 \rightarrow K^- \pi^+) = 3.71 \pm 0.25\%$.

$$\Gamma(\overline{D}^{*}(2007)^0 D_s^+)/\Gamma_{\text{total}} \qquad \Gamma_{52}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.012 ± 0.005 OUR AVERAGE

0.014 ± 0.005 ± 0.003	101 GIBAUT	96 CLE2	e ⁺ e ⁻ → $\Upsilon(4S)$
-----------------------	------------	---------	--

0.010 ± 0.007 ± 0.002	102 ALBRECHT	92G ARG	e ⁺ e ⁻ → $\Upsilon(4S)$
-----------------------	--------------	---------	--

¹⁰¹ GIBAUT 96 reports $0.0140 \pm 0.0043 \pm 0.0035$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁰² ALBRECHT 92G reports $0.013 \pm 0.009 \pm 0.002$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 and $D^*(2007)^0$ branching ratios, e.g., $B(D^0 \rightarrow K^- \pi^+) = 3.71 \pm 0.25\%$ and $B(D^*(2007)^0 \rightarrow D^0 \pi^0) = 55 \pm 6\%$.

$$\Gamma(\overline{D}^{*}(2007)^0 D_s^{*+})/\Gamma_{\text{total}} \qquad \Gamma_{53}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.027 ± 0.010 OUR AVERAGE

0.030 ± 0.011 ± 0.007	103 GIBAUT	96 CLE2	e ⁺ e ⁻ → $\Upsilon(4S)$
-----------------------	------------	---------	--

0.023 ± 0.013 ± 0.006	104 ALBRECHT	92G ARG	e ⁺ e ⁻ → $\Upsilon(4S)$
-----------------------	--------------	---------	--

¹⁰³ GIBAUT 96 reports $0.0310 \pm 0.0088 \pm 0.0065$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁰⁴ ALBRECHT 92G reports $0.031 \pm 0.016 \pm 0.005$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 and $D^*(2007)^0$ branching ratios, e.g., $B(D^0 \rightarrow K^- \pi^+) = 3.71 \pm 0.25\%$ and $B(D^*(2007)^0 \rightarrow D^0 \pi^0) = 55 \pm 6\%$.

$$\Gamma(D_s^{(*)+} \overline{D}^{*0})/\Gamma_{\text{total}} \qquad \Gamma_{54}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

$(2.73 \pm 0.93 \pm 0.68) \times 10^{-2}$	105 AHMED	00B CLE2	e ⁺ e ⁻ → $\Upsilon(4S)$
---	-----------	----------	--

¹⁰⁵ AHMED 00B reports their experiment's uncertainties ($\pm 0.78 \pm 0.48 \pm 0.68$)%, where the first error is statistical, the second is systematic, and the third is the uncertainty in the $D_s \rightarrow \phi\pi$ branching fraction. We combine the first two in quadrature.

$\Gamma(\bar{D}^*(2007)^0 D^*(2010)^+)/\Gamma_{\text{total}}$ Γ_{55}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.011	90	BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

$[\Gamma(\bar{D}^0 D^*(2010)^+) + \Gamma(\bar{D}^*(2007)^0 D^+)]/\Gamma_{\text{total}}$ Γ_{56}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.013	90	BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

$\Gamma(\bar{D}^0 D^+)/\Gamma_{\text{total}}$ Γ_{57}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0067	90	BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

$\Gamma(D_s^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{58}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.00020	90	¹⁰⁶ ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁰⁶ ALEXANDER 93B reports $< 2.0 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$[\Gamma(D_s^+ \pi^0) + \Gamma(D_s^{*+} \pi^0)]/\Gamma_{\text{total}}$ $(\Gamma_{58} + \Gamma_{59})/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0007	90	¹⁰⁷ ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁰⁷ ALBRECHT 93E reports $< 0.9 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$\Gamma(D_s^{*+} \pi^0)/\Gamma_{\text{total}}$ Γ_{59}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.00033	90	¹⁰⁸ ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁰⁸ ALEXANDER 93B reports $< 3.2 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$\Gamma(D_s^+ \eta)/\Gamma_{\text{total}}$ Γ_{60}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0005	90	¹⁰⁹ ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁰⁹ ALEXANDER 93B reports $< 4.6 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$\Gamma(D_s^{*+} \eta)/\Gamma_{\text{total}}$ Γ_{61}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0008	90	¹¹⁰ ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹¹⁰ ALEXANDER 93B reports $< 7.5 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$\Gamma(D_s^+ \rho^0)/\Gamma_{\text{total}} \qquad \Gamma_{62}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0004	90	111 ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

111 ALEXANDER 93B reports $< 3.7 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$[\Gamma(D_s^+ \rho^0) + \Gamma(D_s^+ \bar{K}^*(892)^0)]/\Gamma_{\text{total}} \qquad (\Gamma_{62} + \Gamma_{72})/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0025	90	112 ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

112 ALBRECHT 93E reports $< 3.4 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$\Gamma(D_s^{*+} \rho^0)/\Gamma_{\text{total}} \qquad \Gamma_{63}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0005	90	113 ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

113 ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$[\Gamma(D_s^{*+} \rho^0) + \Gamma(D_s^{*+} \bar{K}^*(892)^0)]/\Gamma_{\text{total}} \qquad (\Gamma_{63} + \Gamma_{73})/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0015	90	114 ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

114 ALBRECHT 93E reports $< 2.0 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$\Gamma(D_s^+ \omega)/\Gamma_{\text{total}} \qquad \Gamma_{64}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0005	90	115 ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0025	90	116 ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	------------------	-----	------------------------------------

115 ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

116 ALBRECHT 93E reports $< 3.4 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$\Gamma(D_s^{*+} \omega)/\Gamma_{\text{total}} \qquad \Gamma_{65}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0007	90	117 ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0014	90	118 ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	------------------	-----	------------------------------------

117 ALEXANDER 93B reports $< 6.8 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

118 ALBRECHT 93E reports $< 1.9 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$\Gamma(D_s^+ a_1(1260)^0)/\Gamma_{\text{total}}$ Γ_{66}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0022	90	119 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

119 ALBRECHT 93E reports $< 3.0 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

 $\Gamma(D_s^{*+} a_1(1260)^0)/\Gamma_{\text{total}}$ Γ_{67}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0016	90	120 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

120 ALBRECHT 93E reports $< 2.2 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

 $\Gamma(D_s^+ \phi)/\Gamma_{\text{total}}$ Γ_{68}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.00032	90	121 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0013	90	122 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	--------------	---------	------------------------------------

121 ALEXANDER 93B reports $< 3.1 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

122 ALBRECHT 93E reports $< 1.7 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

 $\Gamma(D_s^{*+} \phi)/\Gamma_{\text{total}}$ Γ_{69}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0004	90	123 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0016	90	124 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	--------------	---------	------------------------------------

123 ALEXANDER 93B reports $< 4.2 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

124 ALBRECHT 93E reports $< 2.1 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

 $\Gamma(D_s^+ \bar{K}^0)/\Gamma_{\text{total}}$ Γ_{70}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0011	90	125 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0019	90	126 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	--------------	---------	------------------------------------

125 ALEXANDER 93B reports $< 10.3 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

126 ALBRECHT 93E reports $< 2.5 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$\Gamma(D_s^{*+} \bar{K}^0) / \Gamma_{\text{total}} \qquad \Gamma_{71} / \Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.0011	90	127 ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-------------------	----	-------------------	------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0023	90	128 ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	------------------	-----	------------------------------------

127 ALEXANDER 93B reports $< 10.9 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

128 ALBRECHT 93E reports $< 3.1 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$\Gamma(D_s^+ \bar{K}^*(892)^0) / \Gamma_{\text{total}} \qquad \Gamma_{72} / \Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.0005	90	129 ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-------------------	----	-------------------	------	------------------------------------

129 ALEXANDER 93B reports $< 4.4 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$\Gamma(D_s^{*+} \bar{K}^*(892)^0) / \Gamma_{\text{total}} \qquad \Gamma_{73} / \Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.0004	90	130 ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-------------------	----	-------------------	------	------------------------------------

130 ALEXANDER 93B reports $< 4.3 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$\Gamma(D_s^- \pi^+ K^+) / \Gamma_{\text{total}} \qquad \Gamma_{74} / \Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.0008	90	131 ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
-------------------	----	------------------	-----	------------------------------------

131 ALBRECHT 93E reports $< 1.1 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$\Gamma(D_s^{*-} \pi^+ K^+) / \Gamma_{\text{total}} \qquad \Gamma_{75} / \Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.0012	90	132 ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
-------------------	----	------------------	-----	------------------------------------

132 ALBRECHT 93E reports $< 1.6 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$$\Gamma(D_s^- \pi^+ K^*(892)^+) / \Gamma_{\text{total}} \qquad \Gamma_{76} / \Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

<0.006	90	133 ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
------------------	----	------------------	-----	------------------------------------

133 ALBRECHT 93E reports $< 8.6 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$\Gamma(D_s^{*-} \pi^+ K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.008	90	134 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹³⁴ ALBRECHT 93E reports $< 1.1 \times 10^{-2}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$.

$\Gamma(\eta_c K^+)/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.90 ± 0.27 OUR NEW AVERAGE	[(6.9 ^{+3.4} _{-3.0}) × 10 ⁻⁴ OUR 2002 AVERAGE]		

1.25 ± 0.14^{+0.39}_{-0.40} ¹³⁵ FANG 03 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

0.69^{+0.26}_{-0.21} ± 0.22 ¹³⁶ EDWARDS 01 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹³⁵ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹³⁶ EDWARDS 01 assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \rightarrow \gamma \eta_c)$ in those modes have been accounted for.

$\Gamma(J/\psi(1S) K^+)/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
10.0 ± 0.4 OUR FIT				

10.1 ± 0.4 OUR NEW AVERAGE [(10.1 ± 0.5) × 10⁻⁴ OUR 2002 AVERAGE]

10.1 ± 0.2 ± 0.7 ¹³⁷ ABE 03B BELL $e^+ e^- \rightarrow \Upsilon(4S)$

10.1 ± 0.3 ± 0.5 ¹³⁷ AUBERT 02 BABR $e^+ e^- \rightarrow \Upsilon(4S)$

10.2 ± 0.8 ± 0.7 ¹³⁷ JESSOP 97 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

9.3 ± 3.1 ± 0.2 ¹³⁸ BORTOLETTO92 CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

8.1 ± 3.5 ± 0.1 6 ¹³⁹ ALBRECHT 90J ARG $e^+ e^- \rightarrow \Upsilon(4S)$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

11.0 ± 1.5 ± 0.9 59 ¹³⁷ ALAM 94 CLE2 Repl. by JESSOP 97

22 ± 10 ± 2 BUSKULIC 92G ALEP $e^+ e^- \rightarrow Z$

7 ± 4 3 ¹⁴⁰ ALBRECHT 87D ARG $e^+ e^- \rightarrow \Upsilon(4S)$

10 ± 7 ± 2 3 ¹⁴¹ BEBEK 87 CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

9 ± 5 3 ¹⁴² ALAM 86 CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹³⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹³⁸ BORTOLETTO 92 reports $8 \pm 2 \pm 2$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹³⁹ ALBRECHT 90J reports $7 \pm 3 \pm 1$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹⁴⁰ ALBRECHT 87D assume $B^+ B^- / B^0 \bar{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.

¹⁴¹ BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

¹⁴² ALAM 86 assumes B^\pm / B^0 ratio is 60/40.

$\Gamma(J/\psi(1S)K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

0.77±0.20 OUR NEW AVERAGE $[(1.4 \pm 0.6) \times 10^{-3}$ OUR 2002 AVERAGE]

0.69±0.18±0.12			143 ACOSTA	02F CDF	$p\bar{p}$ 1.8 TeV
----------------	--	--	------------	---------	--------------------

1.40±0.82±0.02			144 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
----------------	--	--	------------------	------	-----------------------------------

1.40±0.91±0.02		6	145 ALBRECHT	87D ARG	$e^+e^- \rightarrow \Upsilon(4S)$
----------------	--	---	--------------	---------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.9		90	146 ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
------	--	----	--------------	---------	-----------------------------------

143 ACOSTA 02F uses as reference of $B(B \rightarrow J/\psi(1S)K^+) = (10.1 \pm 0.6) \times 10^{-4}$. The second error includes the systematic error and the uncertainties of the branching ratio.

144 BORTOLETTO 92 reports $1.2 \pm 0.6 \pm 0.4$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

145 ALBRECHT 87D reports 1.2 ± 0.8 for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. They actually report 0.0011 ± 0.0007 assuming $B^+B^-/B^0\bar{B}^0$ ratio is 55/45. We rescale to 50/50. Analysis explicitly removes $B^+ \rightarrow \psi(2S)K^+$.

146 ALBRECHT 90J reports < 1.6 for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593$. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{81}/Γ

For polarization information see the Listings at the end of the " B^0 Branching Ratios" section.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.35±0.10 OUR NEW AVERAGE $[(1.39 \pm 0.13) \times 10^{-3}$ OUR 2002 AVERAGE]

1.28±0.07±0.14			147 ABE	02N BELL	$e^+e^- \rightarrow \Upsilon(4S)$
----------------	--	--	---------	----------	-----------------------------------

1.37±0.09±0.11			147 AUBERT	02 BABR	$e^+e^- \rightarrow \Upsilon(4S)$
----------------	--	--	------------	---------	-----------------------------------

1.41±0.23±0.24			147 JESSOP	97 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
----------------	--	--	------------	---------	-----------------------------------

1.58±0.47±0.27			148 ABE	96H CDF	$p\bar{p}$ at 1.8 TeV
----------------	--	--	---------	---------	-----------------------

1.51±1.09±0.02			149 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
----------------	--	--	------------------	------	-----------------------------------

1.86±1.30±0.03		2	150 ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
----------------	--	---	--------------	---------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.78±0.51±0.23		13	147 ALAM	94 CLE2	Sup. by JESSOP 97
----------------	--	----	----------	---------	-------------------

147 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

148 ABE 96H assumes that $B(B^+ \rightarrow J/\psi K^+) = (1.02 \pm 0.14) \times 10^{-3}$.

149 BORTOLETTO 92 reports $1.3 \pm 0.9 \pm 0.3$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

150 ALBRECHT 90J reports $1.6 \pm 1.1 \pm 0.3$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)K^*(892)^+)/\Gamma(J/\psi(1S)K^+)$ Γ_{81}/Γ_{79}

VALUE	DOCUMENT ID	TECN	COMMENT
1.40 ± 0.11 OUR AVERAGE			
1.37 ± 0.10 ± 0.08	151 AUBERT	02 BABR	$e^+e^- \rightarrow \Upsilon(4S)$
1.45 ± 0.20 ± 0.17	152 JESSOP	97 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
1.92 ± 0.60 ± 0.17	ABE	96Q CDF	$p\bar{p}$

¹⁵¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹⁵² JESSOP 97 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The measurement is actually measured as an average over kaon charged and neutral states.

$\Gamma(J/\psi(1S)K(1270)^+)/\Gamma_{total}$ Γ_{82}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.80 ± 0.34 ± 0.39	153 ABE	01L BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁵³ Uses the PDG value of $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.00 \pm 0.10) \times 10^{-3}$.

$\Gamma(J/\psi(1S)K(1400)^+)/\Gamma(J/\psi(1S)K(1270)^+)$ Γ_{83}/Γ_{82}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.30	90	ABE	01L BELL	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(J/\psi(1S)\phi K^+)/\Gamma_{total}$ Γ_{84}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(8.8^{+3.5}_{-3.0} \pm 1.3) \times 10^{-5}$	154 ANASTASSOV 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁵⁴ ANASTASSOV 00 finds 10 events on a background of 0.5 ± 0.2 . Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$, a uniform Dalitz plot distribution, isotropic $J/\psi(1S)$ and ϕ decays, and $B(B^+ \rightarrow J/\psi(1S)\phi K^+) = B(B^0 \rightarrow J/\psi(1S)\phi K^0)$.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma_{total}$ Γ_{85}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(4.0 \pm 0.5) \times 10^{-5}$ OUR FIT			
$(3.8 \pm 0.6 \pm 0.3) \times 10^{-5}$	155 ABE	03B BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁵⁵ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma(J/\psi(1S)K^+)$ Γ_{85}/Γ_{79}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.040 ± 0.005 OUR FIT				
0.042 ± 0.007 OUR AVERAGE				
0.0391 ± 0.0078 ± 0.0019		AUBERT	02F BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.05 $^{+0.019}_{-0.017} \pm 0.001$		ABE	96R CDF	$p\bar{p}$ 1.8 TeV
0.052 ± 0.024		BISHAI	96 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.043 ± 0.023 5 ¹⁵⁶ ALEXANDER 95 CLE2 Sup. by BISHAI 96

¹⁵⁶ Assumes equal production of $B^+ B^-$ and $B^0 \bar{B}^0$ on $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\rho^+)/\Gamma_{total}$ Γ_{86}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 7.7 × 10⁻⁴	90	BISHAI	96 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(J/\psi(1S) a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-3}$	90	BISHAI	96 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\psi(2S) K^+)/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.8 ± 0.4 OUR NEW AVERAGE			[[$(6.6 \pm 0.6) \times 10^{-4}$ OUR 2002 AVERAGE]		
6.9 ± 0.6			157 ABE	03B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$6.4 \pm 0.5 \pm 0.8$			157 AUBERT	02 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$7.8 \pm 0.7 \pm 0.9$			157 RICHICHI	01 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$5.5 \pm 1.0 \pm 0.6$			158 ABE	98O CDF	$p\bar{p}$ 1.8 TeV
$18 \pm 8 \pm 4$		5	157 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.1 \pm 2.3 \pm 0.9$		7	157 ALAM	94 CLE2	Repl. by RICHICHI 01
< 5	90		157 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
22 ± 17		3	159 ALBRECHT	87D ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

157 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

158 ABE 98O reports $[B(B^+ \rightarrow \psi(2S) K^+)/B(B^+ \rightarrow J/\psi(1S) K^+)] = 0.558 \pm 0.082 \pm 0.056$. We multiply by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = (9.9 \pm 1.0) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

159 ALBRECHT 87D assume $B^+ B^- / B^0 \bar{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.

$\Gamma(\psi(2S) K^+)/\Gamma(J/\psi(1S) K^+)$ Γ_{88}/Γ_{79}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.64 \pm 0.06 \pm 0.07$	160 AUBERT	02 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

160 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S) K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$9.2 \pm 1.9 \pm 1.2$		161 RICHICHI	01 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<30	90	161 ALAM	94 CLE2	Repl. by RICHICHI 01
<35	90	161 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
<49	90	161 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

161 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S) K^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.0019 \pm 0.0011 \pm 0.0004$	3	162 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

162 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c0}(1P)K^+)/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$6.0^{+2.1}_{-1.8} \pm 1.1$		163 ABE	02B BELL	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------------	--	---------	----------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.8	90	164 EDWARDS	01 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
------	----	-------------	---------	-----------------------------------

163 ABE 02B measures the ratio of $B(B^+ \rightarrow \chi_c^0 K^+)/B(B^+ \rightarrow J/\psi(1S)K^+) = 0.60 + 0.21 - 0.18 \pm 0.05 \pm 0.08$, where the third error is due to the uncertainty in the $B(\chi_c^0 \rightarrow \pi^+ \pi^-)$, and uses $B(B^+ \rightarrow J/\psi(1S)K^+) = (10.0 \pm 1.0) \times 10^{-4}$ to obtain the result.

164 EDWARDS 01 assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \rightarrow \gamma \eta_c)$ in those modes have been accounted for.

$\Gamma(\chi_{c1}(1P)K^+)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.8 ± 1.1 OUR NEW AVERAGE		[(6.5 ± 1.1) × 10 ⁻⁴ OUR 2002 AVERAGE]		
----------------------------------	--	---	--	--

15.5 ± 5.4 ± 2.0		165 ACOSTA	02F CDF	$p\bar{p}$ 1.8 TeV
------------------	--	------------	---------	--------------------

6.5 ± 1.0 ± 0.6		166 AUBERT	02 BABR	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------	--	------------	---------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.7 ± 4.0 ± 0.9	6	167 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------	---	----------	---------	-----------------------------------

19 ± 13 ± 6		168 ALBRECHT	92E ARG	$e^+e^- \rightarrow \Upsilon(4S)$
-------------	--	--------------	---------	-----------------------------------

165 ACOSTA 02F uses as reference of $B(B \rightarrow J/\psi(1S)K^+) = (10.1 \pm 0.6) \times 10^{-4}$. The second error includes the systematic error and the uncertainties of the branching ratio.

166 AUBERT 02 reports $7.5 \pm 0.9 \pm 0.8$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$.

We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (31.6 \pm 2.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

167 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

168 ALBRECHT 92E assumes no $\chi_{c2}(1P)$ production and $B(\Upsilon(4S) \rightarrow B^+ B^-) = 50\%$.

$\Gamma(\chi_{c1}(1P)K^+)/\Gamma(J/\psi(1S)K^+)$ Γ_{92}/Γ_{79}

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.65 ± 0.09 OUR NEW AVERAGE	[0.65 ± 0.10 OUR 2002 AVERAGE]		
------------------------------------	--------------------------------	--	--

0.65 ± 0.07 ± 0.06	169 AUBERT	02 BABR	$e^+e^- \rightarrow \Upsilon(4S)$
---------------------------	------------	---------	-----------------------------------

169 AUBERT 02 reports $0.75 \pm 0.08 \pm 0.05$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$.

We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (31.6 \pm 2.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(1P)K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0021	90	170 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
-------------------	----	----------	---------	-----------------------------------

170 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^0\pi^+)/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE (units 10^{-5}) CL% DOCUMENT ID TECN COMMENT

1.87 ± 0.22 OUR NEW AVERAGE $[(1.73^{+0.27}_{-0.24}) \times 10^{-5}$ OUR 2002 AVERAGE]

1.94^{+0.31}_{-0.30} ± 0.16 171 CASEY 02 BELL e⁺e⁻ → $\Upsilon(4S)$

1.82^{+0.33}_{-0.30} ± 0.20 171 AUBERT 01E BABR e⁺e⁻ → $\Upsilon(4S)$

1.82^{+0.46}_{-0.40} ± 0.16 171 CRONIN-HEN..00 CLE2 e⁺e⁻ → $\Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.37^{+0.57}_{-0.48} ^{+0.19}_{-0.18} 171 ABE 01H BELL Repl. by CASEY 02

2.3 ^{+1.1}_{-1.0} ± 0.36 GODANG 98 CLE2 Repl. by CRONIN-HENNESSY 00

< 4.8 90 ASNER 96 CLE2 Repl. by GODANG 98

< 19 90 ALBRECHT 91B ARG e⁺e⁻ → $\Upsilon(4S)$

< 10 90 172 AVERY 89B CLEO e⁺e⁻ → $\Upsilon(4S)$

< 68 90 AVERY 87 CLEO e⁺e⁻ → $\Upsilon(4S)$

171 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

172 AVERY 89B reports $< 9 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^+\pi^0)/\Gamma_{\text{total}}$ Γ_{95}/Γ

VALUE (units 10^{-5}) CL% DOCUMENT ID TECN COMMENT

1.17 ± 0.15 OUR NEW AVERAGE $[(1.21 \pm 0.16) \times 10^{-5}$ OUR 2002 AVERAGE]

1.3 ^{+0.25}_{-0.24} ± 0.13 173 CASEY 02 BELL e⁺e⁻ → $\Upsilon(4S)$

1.08^{+0.21}_{-0.19} ± 0.10 173 AUBERT 01E BABR e⁺e⁻ → $\Upsilon(4S)$

1.16^{+0.30}_{-0.27} ^{+0.14}_{-0.13} 173 CRONIN-HEN..00 CLE2 e⁺e⁻ → $\Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.63^{+0.35}_{-0.33} ^{+0.16}_{-0.18} 173 ABE 01H BELL Repl. by CASEY 02

< 1.6 90 GODANG 98 CLE2 Repl. by CRONIN-HENNESSY 00

< 1.4 90 ASNER 96 CLE2 Repl. by GODANG 98

173 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+\pi^0)/\Gamma(K^0\pi^+)$ Γ_{95}/Γ_{94}

VALUE DOCUMENT ID TECN COMMENT

2.38^{+0.98}_{-1.10} ^{+0.39}_{-0.26} 174 ABE 01H BELL e⁺e⁻ → $\Upsilon(4S)$

174 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' K^+)/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

7.5 ± 0.7 OUR AVERAGE

$7.9^{+1.2}_{-1.1} \pm 0.9$	175 ABE	01M BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$7.0 \pm 0.8 \pm 0.5$	175 AUBERT	01G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$8.0^{+1.0}_{-0.9} \pm 0.7$	175 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.5^{+1.5}_{-1.4} \pm 0.9$	BEHRENS	98 CLE2	Repl. by RICHICHI 00
-----------------------------	---------	---------	----------------------

¹⁷⁵ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 3.5 × 10⁻⁵	90	¹⁷⁶ RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------------------	----	-------------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.3 × 10 ⁻⁴	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00
--------------------------	----	---------	---------	----------------------

¹⁷⁶ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K^+)/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 6.9 × 10⁻⁶	90	¹⁷⁷ RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------------------	----	-------------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.4 × 10 ⁻⁵	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00
--------------------------	----	---------	---------	----------------------

¹⁷⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

2.64^{+0.96}_{-0.82} ± 0.33		¹⁷⁸ RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
--	--	-------------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.0	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00
-------	----	---------	---------	----------------------

¹⁷⁸ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega K^+)/\Gamma_{\text{total}}$ Γ_{100}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

0.92^{+0.28}_{-0.25} ± 0.10				OUR NEW AVERAGE [(1.5 ^{+0.7} _{-0.6}) × 10 ⁻⁵ OUR 2000 AVERAGE]
--	--	--	--	---

0.92^{+0.26}_{-0.23} ± 0.10		¹⁷⁹ LU	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
--	--	-------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.4	90	¹⁷⁹ AUBERT	01G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-------	----	-----------------------	----------	------------------------------------

< 0.79	90	¹⁷⁹ JESSOP	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
--------	----	-----------------------	---------	------------------------------------

1.5 ^{+0.7} _{-0.6} ± 0.2		¹⁷⁹ BERGFELD	98 CLE2	Repl. by JESSOP 00
---	--	-------------------------	---------	--------------------

¹⁷⁹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.7 \times 10^{-5}$	90	180 BERGFELD	98	CLE2

¹⁸⁰ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{102}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$1.94^{+0.42+0.41}_{-0.39-0.71}$		181 GARMASH	02	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<11.9	90	182 ABE	00C	SLD $e^+ e^- \rightarrow Z$
< 1.6	90	183 JESSOP	00	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
<39	90	184 ADAM	96D	DLPH $e^+ e^- \rightarrow Z$
< 4.1	90	ASNER	96	CLE2 Repl. by JESSOP 00
<48	90	185 ABREU	95N	DLPH Sup. by ADAM 96D
<17	90	ALBRECHT	91B	ARG $e^+ e^- \rightarrow \Upsilon(4S)$
<15	90	186 AVERY	89B	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$
<26	90	AVERY	87	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹⁸¹ Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

¹⁸² ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

¹⁸³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹⁸⁴ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

¹⁸⁵ Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

¹⁸⁶ AVERY 89B reports $< 1.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{103}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.1 \times 10^{-5}$	90	187 JESSOP	00	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<9.9 \times 10^{-5}$	90	ASNER	96	CLE2 Repl. by JESSOP 00
-----------------------	----	-------	----	-------------------------

¹⁸⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ \pi^- \pi^+)/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(5.56 \pm 0.58 \pm 0.77) \times 10^{-5}$	188 GARMASH	02	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹⁸⁸ Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

$\Gamma(K^+ \pi^- \pi^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.8	90	BERGFELD	96B	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<33	90	189 ADAM	96D DLPH	$e^+e^- \rightarrow Z$
<40	90	190 ABREU	95N DLPH	Sup. by ADAM 96D
<33	90	ALBRECHT	91E ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<19	90	191 AVERY	89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

189 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

190 Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

191 AVERY 89B reports $< 1.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^+ f_0(980))/\Gamma_{\text{total}} \times B(f_0(980) \rightarrow \pi\pi)$ $\Gamma_{106}/\Gamma \times B$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$9.6^{+2.5+3.7}_{-2.3-1.7}$		192 GARMASH	02 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<80	90	193 AVERY	89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
-----	----	-----------	----------	-----------------------------------

192 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0\pi^+$ and $\bar{D}^0 \rightarrow K^+\pi^-$ with $B(B^+ \rightarrow \bar{D}^0\pi^+) \cdot B(\bar{D}^0 \rightarrow K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$. Only charged pions from the $f_0(980)$ are used.

193 AVERY 89B reports $< 7 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-5}$	90	194 GARMASH	02 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<8.6 \times 10^{-5}$	90	195 ABE	00C SLD	$e^+e^- \rightarrow Z$
$<1.7 \times 10^{-5}$	90	196 JESSOP	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$<1.2 \times 10^{-4}$	90	197 ADAM	96D DLPH	$e^+e^- \rightarrow Z$
$<1.9 \times 10^{-5}$	90	ASNER	96 CLE2	Repl. by JESSOP 00
$<1.9 \times 10^{-4}$	90	198 ABREU	95N DLPH	Sup. by ADAM 96D
$<1.8 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$<8 \times 10^{-5}$	90	199 AVERY	89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$<2.6 \times 10^{-4}$	90	AVERY	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

194 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0\pi^+$ and $\bar{D}^0 \rightarrow K^+\pi^-$ with $B(B^+ \rightarrow \bar{D}^0\pi^+) \cdot B(\bar{D}^0 \rightarrow K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

195 ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

196 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

197 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

198 Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

199 AVERY 89B reports $< 7 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_2^*(1430)^0\pi^+)/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.8 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_{109}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.0 \times 10^{-6}$	90	²⁰⁰ GARMASH	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

²⁰⁰ Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

$\Gamma(K^- \pi^+ \pi^+ \text{nonresonant})/\Gamma_{\text{total}}$ Γ_{110}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.6 \times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_1(1400)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{111}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.6 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^0 \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{112}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<66 \times 10^{-6}$	90	²⁰¹ ECKHART	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

²⁰¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^0 \rho^+)/\Gamma_{\text{total}}$ Γ_{113}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<4.8 \times 10^{-5}$	90	ASNER	96 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{114}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.1 \times 10^{-3}$	90	ALBRECHT	91E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{115}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.4 \times 10^{-5}$	90	²⁰² GODANG	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<9.0 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	----	----------	---------	------------------------------------

²⁰² Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 4.9×10^{-5} .

$\Gamma(K^*(892)^+ K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{116}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.1 \times 10^{-5}$	90	²⁰³ GODANG	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

²⁰³ Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 4.8×10^{-5} .

$\Gamma(K_1(1400)^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{117}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.8 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_2^*(1430)^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{118}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.5 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^+\bar{K}^0)/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.0 × 10⁻⁶ (CL = 90%)		[<2.4 × 10 ⁻⁶ (CL = 90%)		OUR 2002 BEST LIMIT]
<2.0 × 10⁻⁶	90	204 CASEY	02 BELL	e ⁺ e ⁻ → $\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<5.0 × 10 ⁻⁶	90	204 ABE	01H BELL	e ⁺ e ⁻ → $\Upsilon(4S)$
<2.4 × 10 ⁻⁶	90	204 AUBERT	01E BABR	e ⁺ e ⁻ → $\Upsilon(4S)$
<5.1 × 10 ⁻⁶	90	204 CRONIN-HEN..00	CLE2	e ⁺ e ⁻ → $\Upsilon(4S)$
<2.1 × 10 ⁻⁵	90	GODANG	98 CLE2	Repl. by CRONIN-HENNESSY 00

²⁰⁴ Assumes equal production of B⁺ and B⁰ at the $\Upsilon(4S)$.

$\Gamma(\bar{K}^0 K^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{120}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<24 × 10⁻⁶	90	205 ECKHART	02 CLE2	e ⁺ e ⁻ → $\Upsilon(4S)$

²⁰⁵ Assumes equal production of B⁺ and B⁰ at the $\Upsilon(4S)$.

$\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{121}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 × 10⁻⁵ (CL = 90%)		[<1.2 × 10 ⁻⁶ (CL = 90%)		OUR 2002 BEST LIMIT]
<1.2 × 10⁻⁵	90	206 GARMASH	02 BELL	e ⁺ e ⁻ → $\Upsilon(4S)$

²⁰⁶ Uses a reference decay mode B⁺ → $\bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with B(B⁺ → $\bar{D}^0 \pi^+$)·B($\bar{D}^0 \rightarrow K^+ \pi^-$) = (20.3 ± 2.0) × 10⁻⁵.

$\Gamma(K^+ K^- \pi^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.5 × 10⁻⁵	90	BERGFELD	96B CLE2	e ⁺ e ⁻ → $\Upsilon(4S)$

$\Gamma(K^+ K^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.2 × 10⁻⁶ (CL = 90%)		[<3.2 × 10 ⁻⁵ (CL = 90%)		OUR 2002 BEST LIMIT]
<3.2 × 10⁻⁶	90	207 GARMASH	02 BELL	e ⁺ e ⁻ → $\Upsilon(4S)$

²⁰⁷ Uses a reference decay mode B⁺ → $\bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with B(B⁺ → $\bar{D}^0 \pi^+$)·B($\bar{D}^0 \rightarrow K^+ \pi^-$) = (20.3 ± 2.0) × 10⁻⁵.

$\Gamma(K^+ K^+ \pi^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{124}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8.79 × 10⁻⁵	90	ABBIENDI	00B OPAL	e ⁺ e ⁻ → Z

$\Gamma(K^+ K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{125}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.3 × 10⁻⁶	90	208 JESSOP	00 CLE2	e ⁺ e ⁻ → $\Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.29 × 10 ⁻⁴	90	ABBIENDI	00B OPAL	e ⁺ e ⁻ → Z
<1.38 × 10 ⁻⁴	90	209 ABE	00C SLD	e ⁺ e ⁻ → Z

²⁰⁸ Assumes equal production of B⁺ and B⁰ at the $\Upsilon(4S)$.

²⁰⁹ ABE 00C assumes B(Z → b \bar{b})=(21.7 ± 0.1)% and the B fractions f_{B⁰}=f_{B⁺}=(39.7^{+1.8}_{-2.2})% and f_{B_s}=(10.5^{+1.8}_{-2.2})%.

$\Gamma(K^+ K^- K^+)/\Gamma_{\text{total}}$ Γ_{126}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$3.53 \pm 0.37 \pm 0.45$		210 GARMASH	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<20	90	211 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
<32	90	212 ABREU	95N DLPH	Sup. by ADAM 96D
<35	90	ALBRECHT	91E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

210 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

211 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

212 Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

$\Gamma(K^+ \phi)/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$7.9^{+2.0}_{-1.8}$ OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.		

$14.6^{+3.0}_{-2.8} \pm 2.0$		213 GARMASH	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$7.7^{+1.6}_{-1.4} \pm 0.8$		214 AUBERT	01D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$5.5^{+2.1}_{-1.8} \pm 0.6$		214 BRIERE	01 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<144	90	215 ABE	00C SLD	$e^+ e^- \rightarrow Z$
< 5	90	214 BERGFELD	98 CLE2	
<280	90	216 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
< 12	90	ASNER	96 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<440	90	217 ABREU	95N DLPH	Sup. by ADAM 96D
<180	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
< 90	90	218 AVERY	89B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
<210	90	AVERY	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

213 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.


214 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

215 ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

216 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

217 Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

218 AVERY 89B reports $< 8 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.



$$\Gamma(K^+\phi)/\Gamma_{\text{total}} \text{ (units } 10^{-6}\text{)}$$

$\Gamma(K^+K^-K^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<3.8	90	BERGFELD 96B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)^+K^+K^-)/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.6 × 10⁻³	90	ALBRECHT 91E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^*(892)^+\phi)/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
9.7^{+4.2}_{-3.4} ± 1.7	219	AUBERT 01D	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 22.5	90	219 BRIERE 01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 41	90	219 BERGFELD 98	CLE2	
< 70	90	ASNER 96	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 1300	90	ALBRECHT 91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

²¹⁹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K_1(1400)^+\phi)/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1 × 10⁻³	90	ALBRECHT 91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_2^*(1430)^+ \phi) / \Gamma_{\text{total}}$					Γ_{132} / Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$< 3.4 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	

$\Gamma(K^*(892)^+ \gamma) / \Gamma_{\text{total}}$					Γ_{133} / Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.8 ± 0.5 OUR AVERAGE					
$3.83 \pm 0.62 \pm 0.22$			220 AUBERT	02C BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$3.76^{+0.89}_{-0.83} \pm 0.28$			220 COAN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.7 \pm 3.1 \pm 1.1$		5	221 AMMAR	93 CLE2	Repl. by COAN 00
< 55	90		222 ALBRECHT	89G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
< 55	90		223 AVERY	89B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
< 180	90		AVERY	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

220 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

221 AMMAR 93 observed 4.1 ± 2.3 events above background.

222 Assumes the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$.

223 Assumes the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$.

$\Gamma(K^+ \pi^- \pi^+ \gamma) / \Gamma_{\text{total}}$					Γ_{135} / Γ
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$(2.4 \pm 0.5^{+0.4}_{-0.2}) \times 10^{-5}$		224,225 NISHIDA	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$	

224 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

225 $M_{K \pi \pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K^*(892)^0 \pi^+ \gamma) / \Gamma_{\text{total}}$					Γ_{136} / Γ
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$(2.0^{+0.7}_{-0.6} \pm 0.2) \times 10^{-5}$		226,227 NISHIDA	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$	

226 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

227 $M_{K \pi \pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K^+ \rho^0 \gamma) / \Gamma_{\text{total}}$					Γ_{137} / Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$< 2.0 \times 10^{-5}$	90	228,229 NISHIDA	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$	

228 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

229 $M_{K \pi \pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K^+ \pi^- \pi^+ \gamma \text{ (NR)}) / \Gamma_{\text{total}}$					Γ_{138} / Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$< 9.2 \times 10^{-6}$	90	230,231 NISHIDA	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$	

230 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

231 $M_{K \pi \pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K_1(1270)^+\gamma)/\Gamma_{\text{total}}$ Γ_{134}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<9.9 $\times 10^{-5}$ (CL = 90%) [<0.0073 (CL = 90%) OUR 2002 BEST LIMIT]

<9.9 $\times 10^{-5}$ 90 232 NISHIDA 02 BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0073 90 233 ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$

232 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

233 ALBRECHT 89G reports < 0.0066 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_1(1400)^+\gamma)/\Gamma_{\text{total}}$ Γ_{139}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<5.0 $\times 10^{-5}$ (CL = 90%) [<0.0022 (CL = 90%) OUR 2002 BEST LIMIT]

<5.0 $\times 10^{-5}$ 90 234 NISHIDA 02 BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0022 90 235 ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$

234 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

235 ALBRECHT 89G reports < 0.0020 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_2^*(1430)^+\gamma)/\Gamma_{\text{total}}$ Γ_{140}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0014 90 236 ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$

236 ALBRECHT 89G reports < 0.0013 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(1680)^+\gamma)/\Gamma_{\text{total}}$ Γ_{141}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0019 90 237 ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$

237 ALBRECHT 89G reports < 0.0017 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_3^*(1780)^+\gamma)/\Gamma_{\text{total}}$ Γ_{142}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0055 90 238 ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$

238 ALBRECHT 89G reports < 0.005 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_4^*(2045)^+\gamma)/\Gamma_{\text{total}}$ Γ_{143}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0099 90 239 ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$

239 ALBRECHT 89G reports < 0.0090 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\rho^+\gamma)/\Gamma_{\text{total}}$ Γ_{144}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.3 $\times 10^{-5}$ 90 240 COAN 00 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

240 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. No evidence for a nonresonant $K\pi\gamma$ contamination was seen; the central value assumes no contamination.

$\Gamma(\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{145}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$0.74^{+0.23}_{-0.22} \pm 0.09$		241 CASEY	02 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 1.34	90	241 ABE	01H BELL	$e^+e^- \rightarrow \Upsilon(4S)$
< 0.96	90	241 AUBERT	01E BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 1.27	90	241 CRONIN-HEN..00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 2.0	90	GODANG	98 CLE2	Repl. by CRONIN-HENNESSY 00
< 1.7	90	ASNER	96 CLE2	Repl. by GODANG 98
< 24	90	242 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<230	90	243 BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

241 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

242 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

243 BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$.

$\Gamma(\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{146}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.3 \times 10^{-4}$	90	244 ADAM	96D DLPH	$e^+e^- \rightarrow Z$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 2.2 \times 10^{-4}$	90	245 ABREU	95N DLPH	Sup. by ADAM 96D
$< 4.5 \times 10^{-4}$	90	246 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$< 1.9 \times 10^{-4}$	90	247 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

244 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

245 Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

246 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

247 BORTOLETTO 89 reports $< 1.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\rho^0\pi^+)/\Gamma_{\text{total}}$ Γ_{147}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.86 ± 0.20 OUR NEW AVERAGE			[[$(1.0 \pm 0.4) \times 10^{-5}$ OUR 2002 AVERAGE]		
$0.80^{+0.23}_{-0.20} \pm 0.07$			248 GORDON	02 BELL	$e^+e^- \rightarrow \Upsilon(rS)$
$1.04^{+0.33}_{-0.34} \pm 0.21$			248 JESSOP	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

< 8.3	90		249 ABE	00C SLD	$e^+e^- \rightarrow Z$
< 16	90		250 ADAM	96D DLPH	$e^+e^- \rightarrow Z$
< 4.3	90		ASNER	96 CLE2	Repl. by JESSOP 00
< 26	90		251 ABREU	95N DLPH	Sup. by ADAM 96D
< 15	90		252 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 17	90		253 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
< 23	90		253 BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
< 60	90	0	GILES	84 CLEO	Repl. by BEBEK 87

248 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

249 ABE 00C assumes $B(Z \rightarrow b\bar{b})=(21.7 \pm 0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$.

250 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

251 Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.

252 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

253 Papers assume the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$[\Gamma(K^*(892)^0\pi^+) + \Gamma(\rho^0\pi^+)]/\Gamma_{\text{total}}$ $(\Gamma_{102}+\Gamma_{147})/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$(17 \pm \frac{12}{8}) \times 10^{-5}$	90	254 ADAM	96D DLPH	$e^+e^- \rightarrow Z$

254 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

$\Gamma(\pi^+ f_0(980))/\Gamma_{\text{total}}$ Γ_{148}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-4}$	90	255 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

255 BORTOLETTO 89 reports $< 1.2 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+ f_2(1270))/\Gamma_{\text{total}}$ Γ_{149}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.4 \times 10^{-4}$	90	256 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

256 BORTOLETTO 89 reports $< 2.1 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+\pi^-\pi^+ \text{nonresonant})/\Gamma_{\text{total}}$ Γ_{150}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.1 \times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\pi^+\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.9 \times 10^{-4}$	90	257 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

257 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^+\pi^0)/\Gamma_{\text{total}}$ Γ_{152}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.3 \times 10^{-5}$	90	258 JESSOP	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<7.7 \times 10^{-5}$ 90 ASNER 96 CLE2 Repl. by JESSOP 00

$<5.5 \times 10^{-4}$ 90 259 ALBRECHT 90B ARG $e^+e^- \rightarrow \Upsilon(4S)$

258 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. Assumes no nonresonant contributions of $B^+ \rightarrow \pi^+\pi^0\pi^0$.

259 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\pi^+\pi^-\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{153}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.0 \times 10^{-3}$	90	260 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
260 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.				

$\Gamma(\rho^+\rho^0)/\Gamma_{\text{total}}$ Γ_{154}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-3}$	90	261 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
261 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.				

$\Gamma(a_1(1260)^+\pi^0)/\Gamma_{\text{total}}$ Γ_{155}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-3}$	90	262 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
262 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.				

$\Gamma(a_1(1260)^0\pi^+)/\Gamma_{\text{total}}$ Γ_{156}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.0 \times 10^{-4}$	90	263 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
263 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.				

$\Gamma(\omega\pi^+)/\Gamma_{\text{total}}$ Γ_{157}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$0.64^{+0.18}_{-0.16}$ OUR NEW AVERAGE Error includes scale factor of 1.3. See the ideogram below. $[(0.81^{+0.23}_{-0.20}) \times 10^{-5}]$ OUR 2002 AVERAGE Scale factor = 1.2]


$0.42^{+0.20}_{-0.18} \pm 0.05$	264 LU	02 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$0.66^{+0.21}_{-0.18} \pm 0.07$	264 AUBERT	01G BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.13^{+0.33}_{-0.29} \pm 0.14$	264 JESSOP	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.3	90	264 BERGFELD	98 CLE2	Repl. by JESSOP 00
< 40	90	265 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

264 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

265 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.



$$\Gamma(\omega\pi^+)/\Gamma_{\text{total}} \qquad \Gamma_{157}/\Gamma$$

$$\Gamma(\omega\rho^+)/\Gamma_{\text{total}} \qquad \Gamma_{158}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN
<6.1 × 10 ⁻⁵	90	266 BERGFELD	98 CLE2

266 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(\eta\pi^+)/\Gamma_{\text{total}} \qquad \Gamma_{159}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.7 × 10 ⁻⁶	90	267 RICHICHI	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5 × 10 ⁻⁵	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00
<7.0 × 10 ⁻⁴	90	268 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

267 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

268 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$$\Gamma(\eta'\pi^+)/\Gamma_{\text{total}} \qquad \Gamma_{160}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.0 × 10 ⁻⁶	90	269 ABE	01M BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.2 × 10 ⁻⁵	90	269 AUBERT	01G BABR	$e^+e^- \rightarrow \Upsilon(4S)$
<1.2 × 10 ⁻⁵	90	269 RICHICHI	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<3.1 × 10 ⁻⁵	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

269 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' \rho^+)/\Gamma_{\text{total}}$ Γ_{161}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.3 \times 10^{-5}$	90	270 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<4.7 \times 10^{-5}$	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

²⁷⁰ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta \rho^+)/\Gamma_{\text{total}}$ Γ_{162}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-5}$	90	271 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<3.2 \times 10^{-5}$	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

²⁷¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\phi \pi^+)/\Gamma_{\text{total}}$ Γ_{163}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-6}$	90	272 AUBERT	01D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1.53 \times 10^{-4}$	90	273 ABE	00C SLD	$e^+ e^- \rightarrow Z$
$<0.5 \times 10^{-5}$	90	272 BERGFELD	98 CLE2	

²⁷² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
²⁷³ ABE 00C assumes $B(Z \rightarrow b\bar{b})=(21.7 \pm 0.1)\%$ and the B fractions $f_{B^0}=f_{B^+}=(39.7^{+1.8}_{-2.2})\%$ and $f_{B_s}=(10.5^{+1.8}_{-2.2})\%$.

$\Gamma(\phi \rho^+)/\Gamma_{\text{total}}$ Γ_{164}/Γ

VALUE	DOCUMENT ID	TECN
$<1.6 \times 10^{-5}$	274 BERGFELD	98 CLE2

²⁷⁴ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^+ \pi^+ \pi^+ \pi^- \pi^-)/\Gamma_{\text{total}}$ Γ_{165}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.6 \times 10^{-4}$	90	275 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

²⁷⁵ ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\rho^0 a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{166}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.2 \times 10^{-4}$	90	276 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<6.0 \times 10^{-4}$	90	277 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
$<3.2 \times 10^{-3}$	90	276 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

²⁷⁶ BORTOLETTO 89 reports $< 5.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.
²⁷⁷ ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\rho^0 a_2(1320)^+)/\Gamma_{\text{total}}$ Γ_{167}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.2 \times 10^{-4}$	90	278 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.6 \times 10^{-3}$	90	279 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	----	-----------	---------	------------------------------------

278 BORTOLETTO 89 reports $< 6.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

279 BEBEK 87 reports $< 2.3 \times 10^{-3}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{168}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.3 \times 10^{-3}$	90	280 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

280 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^+ a_1(1260)^0)/\Gamma_{\text{total}}$ Γ_{169}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-2}$	90	281 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

281 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(h^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{170}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(1.6^{+0.6}_{-0.5} \pm 0.36) \times 10^{-5}$	GODANG	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\omega h^+)/\Gamma_{\text{total}}$ Γ_{171}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.38^{+0.27}_{-0.24}$ OUR NEW AVERAGE	[(1.4 ± 0.4) $\times 10^{-5}$ OUR 2002 AVERAGE]		

$1.34^{+0.33}_{-0.29} \pm 0.11$	282 LU	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
---------------------------------	--------	---------	------------------------------------

$1.43^{+0.36}_{-0.32} \pm 0.20$	282 JESSOP	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
---------------------------------	------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.5^{+0.8}_{-0.7} \pm 0.3$	282 BERGFELD	98 CLE2	Repl. by JESSOP 00
-----------------------------	--------------	---------	--------------------

282 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(h^+ X^0(\text{Familon}))/\Gamma_{\text{total}}$ Γ_{172}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.9 \times 10^{-5}$	90	283 AMMAR	01B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

283 AMMAR 01B searched for the two-body decay of the B meson to a massless neutral feebly-interacting particle X^0 such as the familon, the Nambu-Goldstone boson associated with a spontaneously broken global family symmetry.

$\Gamma(\rho\bar{p}\pi^+)/\Gamma_{\text{total}}$ Γ_{173}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 3.7	$\times 10^{-6}$ (CL = 90%)	[<1.6 $\times 10^{-4}$ (CL = 90%) OUR 2002 BEST LIMIT]		
< 3.7	$\times 10^{-6}$	90 ^{284,285} ABE	02K BELL	$e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••				
< 5.0	$\times 10^{-4}$	90 ²⁸⁶ ABREU	95N DLPH	Sup. by ADAM 96D
< 1.6	$\times 10^{-4}$	90 ²⁸⁷ BEBEK	89 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$(5.7 \pm 1.5 \pm 2.1) \times 10^{-4}$		288 ²⁸⁸ ALBRECHT	88F ARG	$e^+e^- \rightarrow \Upsilon(4S)$

²⁸⁴ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

²⁸⁵ Explicitly vetoes resonant production of $\rho\bar{p}$ from Charmonium states.

²⁸⁶ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

²⁸⁷ BEBEK 89 reports < 1.4×10^{-4} assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

²⁸⁸ ALBRECHT 88F reports $(5.2 \pm 1.4 \pm 1.9) \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\rho\bar{p}\pi^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{174}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 5.3 $\times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\rho\bar{p}\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{175}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 5.2 $\times 10^{-4}$	90	289 ²⁸⁹ ALBRECHT	88F ARG	$e^+e^- \rightarrow \Upsilon(4S)$

²⁸⁹ ALBRECHT 88F reports < 4.7×10^{-4} assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\rho\bar{p}K^+)/\Gamma_{\text{total}}$ Γ_{176}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(4.3^{+1.1}_{-0.9} \pm 0.5) \times 10^{-6}$	290,291 ^{290,291} ABE	02K BELL	$e^+e^- \rightarrow \Upsilon(4S)$

²⁹⁰ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

²⁹¹ Explicitly vetoes resonant production of $\rho\bar{p}$ from Charmonium states.

$\Gamma(\rho\bar{p}K^+ \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{177}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 8.9 $\times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\rho\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{178}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.2 $\times 10^{-6}$ (CL = 90%)		[<2.6 $\times 10^{-6}$ (CL = 90%) OUR 2002 BEST LIMIT]		
< 2.2 $\times 10^{-6}$	90	292 ²⁹² ABE	02O BELL	$e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••				
< 2.6 $\times 10^{-6}$	90	293 ²⁹³ COAN	99 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 6 $\times 10^{-5}$	90	294 ²⁹⁴ AVERY	89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
< 9.3 $\times 10^{-5}$	90	295 ²⁹⁵ ALBRECHT	88F ARG	$e^+e^- \rightarrow \Upsilon(4S)$

292 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

293 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

294 AVERY 89B reports $< 5 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

295 ALBRECHT 88F reports $< 8.5 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\rho \bar{\Lambda} \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{179} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.0 \times 10^{-4}$	90	296 ALBRECHT 88F	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

296 ALBRECHT 88F reports $< 1.8 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\bar{\Delta}^0 \rho) / \Gamma_{\text{total}}$ Γ_{180} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.8 \times 10^{-4}$	90	297 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

297 BORTOLETTO 89 reports $< 3.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\Delta^{++} \bar{p}) / \Gamma_{\text{total}}$ Γ_{181} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.5 \times 10^{-4}$	90	298 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

298 BORTOLETTO 89 reports $< 1.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(D^+ \rho \bar{p}) / \Gamma_{\text{total}}$ Γ_{182} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.5 \times 10^{-5}$	90	299 ABE 02W BELL		$e^+ e^- \rightarrow \Upsilon(4S)$

299 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^+ \rho \bar{p}) / \Gamma_{\text{total}}$ Γ_{183} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.5 \times 10^{-5}$	90	300 ABE 02W BELL		$e^+ e^- \rightarrow \Upsilon(4S)$

300 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\Lambda_c^- \rho \pi^+) / \Gamma_{\text{total}}$ Γ_{184} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

2.1 ± 0.7 OUR NEW AVERAGE [(6.2 ± 2.7) × 10⁻⁴ OUR 2002 AVERAGE]

2.4 ± 0.6 ± 0.6 301 DYTMAN 02 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

1.9 ± 0.5 ± 0.5 302 GABYSHEV 02 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.2 $^{+2.3}_{-2.0}$ ± 1.6 303 FU 97 CLE2 Repl. by DYTMAN 02

301 DYTMAN 02 reports $2.4^{+0.63}_{-0.62}$ for $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

302 GABYSHEV 02 reports $1.87^{+0.51}_{-0.49}$ for $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

303 FU 97 uses PDG 96 values of Λ_c branching fraction.

$\Gamma(\Lambda_c^- p \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{185}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
$1.81 \pm 0.29^{+0.52}_{-0.50}$		304,305 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.12 90 306 FU 97 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

³⁰⁴ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³⁰⁵ DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

³⁰⁶ FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\Lambda_c^- p \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{186}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
$2.25 \pm 0.25^{+0.63}_{-0.61}$		307,308 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.46 90 309 FU 97 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

³⁰⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³⁰⁸ DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

³⁰⁹ FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\Lambda_c^- p \pi^+ \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{187}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 1.34×10^{-2}		90 310 FU	97 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

³¹⁰ FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\bar{\Sigma}_c(2455)^0 p)/\Gamma_{\text{total}}$ Γ_{188}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.8×10^{-4}		90 311,312 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9.3×10^{-5} 90 311,313 GABYSHEV 02 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

³¹¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³¹² DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

³¹³ Uses the value for $\Lambda_c \rightarrow p K^- \pi^+$ branching ratio ($5.0 \pm 1.3\%$).

$\Gamma(\bar{\Sigma}_c(2520)^0 p)/\Gamma_{\text{total}}$ Γ_{189}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 4.6×10^{-5}		90 314,315 GABYSHEV	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

³¹⁴ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³¹⁵ Uses the value for $\Lambda_c \rightarrow p K^- \pi^+$ branching ratio ($5.0 \pm 1.3\%$).

$\Gamma(\overline{\Sigma}_c(2455)^0 p \pi^0)/\Gamma_{\text{total}}$ Γ_{190}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.4 \pm 1.4 \pm 1.1$	316,317 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

³¹⁶ DYTMAN 02 reports 4.4 ± 1.4 for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³¹⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\overline{\Sigma}_c(2455)^0 p \pi^- \pi^+)/\Gamma_{\text{total}}$ Γ_{191}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.4 \pm 1.3 \pm 1.1$	318,319 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

³¹⁸ DYTMAN 02 reports 4.4 ± 1.3 for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³¹⁹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\Sigma_c(2455)^{--} p \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_{192}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 1.0 \pm 0.7$	320,321 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

³²⁰ DYTMAN 02 reports 2.8 ± 1.0 for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³²¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\Lambda_c^- p \pi^+)/\Gamma_{\text{total}}$ Γ_{193}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.9 \times 10^{-4}$	90	^{322,323} DYTMAN	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

³²² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³²³ DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

$\Gamma(\pi^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{194}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.0039	90	³²⁴ WEIR	90B MRK2	$e^+ e^-$ 29 GeV

³²⁴ WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{195}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.0091	90	³²⁵ WEIR	90B MRK2	$e^+ e^-$ 29 GeV

³²⁵ WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{196}/Γ**

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9 \times 10^{-7}$ (CL = 90%)		[$< 1.4 \times 10^{-6}$ (CL = 90%)		OUR 2002 BEST LIMIT]
$< 0.9 \times 10^{-6}$	90	326 AUBERT	02L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 1.4 \times 10^{-6}$	90	326 ABE	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 2.4 \times 10^{-6}$	90	327 ANDERSON	01B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 9.9 \times 10^{-5}$	90	328 ALBRECHT	91E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 6.8 \times 10^{-3}$	90	329 WEIR	90B MRK2	$e^+ e^-$ 29 GeV
$< 6 \times 10^{-5}$	90	330 AVERY	89B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 2.5 \times 10^{-4}$	90	331 AVERY	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

326 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

327 The result is for di-lepton masses above 0.5 GeV.

328 ALBRECHT 91E reports $< 9.0 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

329 WEIR 90B assumes B^+ production cross section from LUND.

330 AVERY 89B reports $< 5 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

331 AVERY 87 reports $< 2.1 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{197}/Γ**

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$0.98^{+0.46+0.16}_{-0.36-0.16}$		332 ABE	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 1.2	90	332 AUBERT	02L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
< 3.68	90	333 ANDERSON	01B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
< 5.2	90	334 AFFOLDER	99B CDF	$p\bar{p}$ at 1.8 TeV
< 10	90	335 ABE	96L CDF	Repl. by AFFOLDER 99B
< 240	90	336 ALBRECHT	91E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
< 6400	90	337 WEIR	90B MRK2	$e^+ e^-$ 29 GeV
< 170	90	338 AVERY	89B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
< 380	90	339 AVERY	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

332 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

333 The result is for di-lepton masses above 0.5 GeV.

334 AFFOLDER 99B measured relative to $B^+ \rightarrow J/\psi(1S) K^+$.

335 ABE 96L measured relative to $B^+ \rightarrow J/\psi(1S) K^+$ using PDG 94 branching ratios.

336 ALBRECHT 91E reports $< 2.2 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

337 WEIR 90B assumes B^+ production cross section from LUND.

338 AVERY 89B reports $< 1.5 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

339 AVERY 87 reports $< 3.2 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K^+\bar{\nu})/\Gamma_{\text{total}}$ **Γ_{198}/Γ**

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.4 \times 10^{-4}$	90	340 BROWDER	01 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

³⁴⁰ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^+e^+e^-)/\Gamma_{\text{total}}$ **Γ_{199}/Γ**

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.9 \times 10^{-6}$	90	341 ABE	02 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

••• We do not use the following data for averages, fits, limits, etc. •••

$<9.5 \times 10^{-6}$	90	341 AUBERT	02L BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$<6.9 \times 10^{-4}$	90	342 ALBRECHT	91E ARG	$e^+e^- \rightarrow \Upsilon(4S)$

³⁴¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³⁴² ALBRECHT 91E reports $<6.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^+\mu^+\mu^-)/\Gamma_{\text{total}}$ **Γ_{200}/Γ**

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.9 \times 10^{-6}$	90	343 ABE	02 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

••• We do not use the following data for averages, fits, limits, etc. •••

$<17.0 \times 10^{-6}$	90	343 AUBERT	02L BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$<1.2 \times 10^{-3}$	90	344 ALBRECHT	91E ARG	$e^+e^- \rightarrow \Upsilon(4S)$

³⁴³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³⁴⁴ ALBRECHT 91E reports $<1.1 \times 10^{-3}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+e^+\mu^-)/\Gamma_{\text{total}}$ **Γ_{201}/Γ**

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	345 WEIR	90B MRK2	e^+e^- 29 GeV

³⁴⁵ WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^+e^-\mu^+)/\Gamma_{\text{total}}$ **Γ_{202}/Γ**

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	346 WEIR	90B MRK2	e^+e^- 29 GeV

³⁴⁶ WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+e^+\mu^-)/\Gamma_{\text{total}}$ **Γ_{203}/Γ**

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8 \times 10^{-7}$ (CL = 90%)				[<0.0064 (CL = 90%) OUR 2002 BEST LIMIT]

$<0.8 \times 10^{-6}$	90	347 AUBERT	02L BABR	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------	----	------------	----------	-----------------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

<0.0064	90	348 WEIR	90B MRK2	e^+e^- 29 GeV
-----------	----	----------	----------	-----------------

³⁴⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³⁴⁸ WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{204}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	349 WEIR	90B MRK2	$e^+ e^-$ 29 GeV

349 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^*(892)^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{205}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.9 × 10⁻⁶	90	350 AUBERT	02L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

350 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\pi^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{206}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.6 × 10⁻⁶ (CL = 90%)		[<0.0039 (CL = 90%) OUR 2002 BEST LIMIT]		

<1.6 × 10⁻⁶	90	351 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
----------------------------------	----	-------------	----------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0039	90	352 WEIR	90B MRK2	$e^+ e^-$ 29 GeV
---------	----	----------	----------	------------------

351 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

352 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{207}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.4 × 10⁻⁶ (CL = 90%)		[<0.0091 (CL = 90%) OUR 2002 BEST LIMIT]		

<1.4 × 10⁻⁶	90	353 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
----------------------------------	----	-------------	----------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0091	90	354 WEIR	90B MRK2	$e^+ e^-$ 29 GeV
---------	----	----------	----------	------------------

353 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

354 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{208}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 × 10⁻⁶ (CL = 90%)		[<0.0064 (CL = 90%) OUR 2002 BEST LIMIT]		

<1.3 × 10⁻⁶	90	355 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
----------------------------------	----	-------------	----------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0064	90	356 WEIR	90B MRK2	$e^+ e^-$ 29 GeV
---------	----	----------	----------	------------------

355 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

356 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\rho^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{209}/Γ

Test for $\Delta B = 1$ weak neutral current.

VALUE (units 10 ⁻⁶)	CL%	DOCUMENT ID	TECN	COMMENT
<2.6	90	357 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

357 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{210}/Γ

Test for $\Delta B = 1$ weak neutral current.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<5.0 90 358 EDWARDS 02B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

358 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\rho^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{211}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<3.3 90 359 EDWARDS 02B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

359 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{212}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.0 $\times 10^{-6}$ (CL = 90%) [<0.0039 (CL = 90%) OUR 2002 BEST LIMIT]

<1.0 $\times 10^{-6}$ 90 360 EDWARDS 02B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0039 90 361 WEIR 90B MRK2 $e^+ e^-$ 29 GeV

360 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

361 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{213}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.8 $\times 10^{-6}$ (CL = 90%) [<0.0091 (CL = 90%) OUR 2002 BEST LIMIT]

<1.8 $\times 10^{-6}$ 90 362 EDWARDS 02B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0091 90 363 WEIR 90B MRK2 $e^+ e^-$ 29 GeV

362 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

363 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{214}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.0 $\times 10^{-6}$ (CL = 90%) [<0.0064 (CL = 90%) OUR 2002 BEST LIMIT]

<2.0 $\times 10^{-6}$ 90 364 EDWARDS 02B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0064 90 365 WEIR 90B MRK2 $e^+ e^-$ 29 GeV

364 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

365 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^*(892)^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{215}/Γ

Test for $\Delta B = 1$ weak neutral current.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<2.8 90 366 EDWARDS 02B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

366 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{216}/Γ

Test for $\Delta B = 1$ weak neutral current.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<8.3	90	367 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

³⁶⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{217}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.4	90	368 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

³⁶⁸ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

CP VIOLATION

A_{CP} is defined as

$$\frac{B(B^- \rightarrow \bar{f}) - B(B^+ \rightarrow f)}{B(B^- \rightarrow \bar{f}) + B(B^+ \rightarrow f)},$$

the CP-violation charge asymmetry of inclusive B^- and B^+ decay.

$A_{CP}(B^+ \rightarrow J/\psi(1S)K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.007 ± 0.019 OUR NEW AVERAGE	[0.008 \pm 0.025 OUR 2002 AVERAGE]		
$-0.026 \pm 0.022 \pm 0.017$	ABE	03B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.003 \pm 0.030 \pm 0.004$	AUBERT	02F BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.018 \pm 0.043 \pm 0.004$	³⁶⁹ BONVICINI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

³⁶⁹ A +0.3% correction is applied due to a slightly higher reconstruction efficiency for the positive kaons.

$A_{CP}(B^+ \rightarrow J/\psi(1S)\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.01 ± 0.13 OUR NEW AVERAGE	[0.01 \pm 0.22 OUR 2002 AVERAGE]		
$-0.023 \pm 0.164 \pm 0.015$	ABE	03B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.01 \pm 0.22 \pm 0.01$	AUBERT	02F BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$A_{CP}(B^+ \rightarrow \psi(2S)K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.037 ± 0.025 OUR NEW AVERAGE	[0.02 \pm 0.09 OUR 2002 AVERAGE]		
$-0.042 \pm 0.020 \pm 0.017$	ABE	03B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.02 \pm 0.091 \pm 0.01$	³⁷⁰ BONVICINI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

³⁷⁰ A +0.3% correction is applied due to a slightly higher reconstruction efficiency for the positive kaons.

$A_{CP}(B^+ \rightarrow \bar{D}^0 K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.003 \pm 0.080 \pm 0.037$	³⁷¹ ABE	03D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

³⁷¹ Corresponds to 90% confidence range $-0.15 < A_{CP} < 0.16$.

$A_{CP}(B^+ \rightarrow D_{CP(+1)} K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.29 \pm 0.26 \pm 0.05$	372 ABE	03D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

372 Corresponds to 90% confidence range $-0.14 < A_{CP} < 0.73$.

$A_{CP}(B^+ \rightarrow D_{CP(-1)} K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.22 \pm 0.24 \pm 0.04$	373 ABE	03D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

373 Corresponds to 90% confidence range $-0.62 < A_{CP} < 0.18$.

$A_{CP}(B^+ \rightarrow \pi^+ \pi^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.30 \pm 0.30 \begin{smallmatrix} +0.06 \\ -0.04 \end{smallmatrix}$	374 CASEY	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

374 Corresponds to 90% confidence range $-0.23 < A_{CP} < +0.86$.

$A_{CP}(B^+ \rightarrow K^+ \pi^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.08 ± 0.11 OUR NEW AVERAGE $[-0.10 \pm 0.12$ OUR 2002 AVERAGE]			
$-0.02 \pm 0.19 \pm 0.02$	375 CASEY	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.00 \pm 0.18 \pm 0.04$	376 AUBERT	01E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-0.29 ± 0.23	377 CHEN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.059 \begin{smallmatrix} +0.222 +0.055 \\ -0.196 -0.017 \end{smallmatrix}$	378 ABE	01K BELL	Repl. by CASEY 02
---	---------	----------	-------------------

375 Corresponds to 90% confidence range $-0.35 < A_{CP} < +0.30$.

376 Corresponds to 90% confidence range $-0.30 < A_{CP} < +0.30$.

377 Corresponds to 90% confidence range $-0.67 < A_{CP} < 0.09$.

378 Corresponds to 90% confidence range $-0.40 < A_{CP} < 0.36$.

$A_{CP}(B^+ \rightarrow K_S^0 \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.19 ± 0.21 OUR NEW AVERAGE Error includes scale factor of 2.0. See the ideogram below. $[-0.05 \pm 0.14$ OUR 2002 AVERAGE]			

$0.46 \pm 0.15 \pm 0.02$	379 CASEY	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$-0.21 \pm 0.18 \pm 0.03$	380 AUBERT	01E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.18 ± 0.24	381 CHEN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •


$0.098 \begin{smallmatrix} +0.430 +0.020 \\ -0.343 -0.063 \end{smallmatrix}$	382 ABE	01K BELL	Repl. by CASEY 02
--	---------	----------	-------------------

379 Corresponds to 90% confidence range $+0.19 < A_{CP} < +0.72$.

380 Corresponds to 90% confidence range $-0.51 < A_{CP} < 0.09$.

381 Corresponds to 90% confidence range $-0.22 < A_{CP} < 0.56$.

382 Corresponds to 90% confidence range $-0.53 < A_{CP} < 0.82$.



$$A_{CP}(B^+ \rightarrow K_S^0 \pi^+)$$

$A_{CP}(B^+ \rightarrow K^+ \eta')$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.03 ± 0.05 OUR NEW AVERAGE	[-0.02 ± 0.07 OUR 2002 AVERAGE]		
-0.11 ± 0.11 ± 0.02	383 AUBERT	02E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-0.015 ± 0.070 ± 0.009	384 CHEN	02B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
0.03 ± 0.12	385 CHEN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.06 ± 0.15 ± 0.01	386 ABE	01M BELL	Repl. by CHEN 02B
383 Corresponds to 90% confidence range $-0.28 < A_{CP} < 0.07$.			
384 Corresponds to 90% confidence range $-0.13 < A_{CP} < 0.10$.			
385 Corresponds to 90% confidence range $-0.17 < A_{CP} < 0.23$.			
386 Corresponds to 90% confidence range $-0.20 < A_{CP} < 0.32$.			

$A_{CP}(B^+ \rightarrow \omega \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.21 ± 0.19 OUR AVERAGE			
-0.01 ^{+0.29} _{-0.31} ± 0.03	387 AUBERT	02E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-0.34 ± 0.25	388 CHEN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

387 Corresponds to 90% confidence range $-0.50 < A_{CP} < 0.46$.

388 Corresponds to 90% confidence range $-0.75 < A_{CP} < 0.07$.

$A_{CP}(B^+ \rightarrow \omega K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.21 ± 0.28 ± 0.03	389 LU	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

389 Corresponds to 90% confidence range $-0.70 < A_{CP} < +0.38$.

$A_{CP}(B^+ \rightarrow \phi K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.05 \pm 0.20 \pm 0.03$	³⁹⁰ AUBERT	02E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

³⁹⁰ Corresponds to 90% confidence range $-0.37 < A_{CP} < 0.28$. **$A_{CP}(B^+ \rightarrow \phi K^*(892)^+)$**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.43^{+0.36}_{-0.30} \pm 0.06$	³⁹¹ AUBERT	02E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

³⁹¹ Corresponds to 90% confidence range $-0.88 < A_{CP} < 0.18$. **B^\pm REFERENCES**

ABE	03B	PR D67 032003	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	03D	PRL 90 131803	K. Abe <i>et al.</i>	(BELLE Collab.)
ADAM	03	PR D67 032001	N.E. Adam <i>et al.</i>	(CLEO Collab.)
FANG	03	PRL 90 071801	F. Fang <i>et al.</i>	(BELLE Collab.)
ABE	02	PRL 88 021801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02B	PRL 88 031802	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02H	PRL 88 171801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02K	PRL 88 181803	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02N	PL B538 11	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02O	PR D65 091103R	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02W	PRL 89 151802	K. Abe <i>et al.</i>	(BELLE Collab.)
ACOSTA	02C	PR D65 092009	D. Acosta <i>et al.</i>	(CDF Collab.)
ACOSTA	02F	PR D66 052005	D. Acosta <i>et al.</i>	(CDF Collab.)
AHMED	02B	PR D66 031101R	S. Ahmed <i>et al.</i>	(CLEO Collab.)
AUBERT	02	PR D65 032001	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02C	PRL 88 101805	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02E	PR D65 051101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02F	PR D65 091101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02L	PRL 88 241801	B. Aubert <i>et al.</i>	(BaBar Collab.)
BRIERE	02	PRL 89 081803	R. Briere <i>et al.</i>	(CLEO Collab.)
CASEY	02	PR D66 092002	B.C.K. Casey <i>et al.</i>	(BELLE Collab.)
CHEN	02B	PL B546 196	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	02	PL B542 171	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
DYTMAN	02	PR D66 091101R	S.A. Dytman <i>et al.</i>	(CLEO Collab.)
ECKHART	02	PRL 89 251801	E. Eckhart <i>et al.</i>	(CLEO Collab.)
EDWARDS	02B	PR D65 111102R	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
GABYSHEV	02	PR D66 091102R	N. Gabyshev <i>et al.</i>	(BELLE Collab.)
GARMASH	02	PR D65 092005	A. Garmash <i>et al.</i>	(BELLE Collab.)
GODANG	02	PRL 88 021802	R. Godang <i>et al.</i>	(CLEO Collab.)
GORDON	02	PL B542 183	A. Gordon <i>et al.</i>	(BELLE Collab.)
LU	02	PRL 89 191801	R.-S. Lu <i>et al.</i>	(BELLE Collab.)
MAHAPATRA	02	PRL 88 101803	R. Mahapatra <i>et al.</i>	(CLEO Collab.)
NISHIDA	02	PRL 89 231801	S. Nishida <i>et al.</i>	(BELLE Collab.)
ABE	01H	PRL 87 101801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01I	PRL 87 111801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01K	PR D64 071101	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01L	PRL 87 161601	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01M	PL B517 309	K. Abe <i>et al.</i>	(BELLE Collab.)
ALEXANDER	01B	PR D64 092001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AMMAR	01B	PRL 87 271801	R. Ammar <i>et al.</i>	(CLEO Collab.)
ANDERSON	01B	PRL 87 181803	S. Anderson <i>et al.</i>	(CLEO Collab.)
AUBERT	01D	PRL 87 151801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01E	PRL 87 151802	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01F	PRL 87 201803	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01G	PRL 87 221802	B. Aubert <i>et al.</i>	(BaBar Collab.)
BARATE	01E	EPJ C19 213	R. Barate <i>et al.</i>	(ALEPH Collab.)
BRIERE	01	PRL 86 3718	R.A. Biere <i>et al.</i>	(CLEO Collab.)
BROWDER	01	PRL 86 2950	T.E. Browder <i>et al.</i>	(CLEO Collab.)
EDWARDS	01	PRL 86 30	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
GRITSAN	01	PR D64 077501	A. Gritsan <i>et al.</i>	(CLEO Collab.)
RICHICHI	01	PR D63 031103R	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ABBIENDI	00B	PL B476 233	G. Abbiendi <i>et al.</i>	(OPAL Collab.)

ABE	00C	PR D62 071101R	K. Abe <i>et al.</i>	(SLD Collab.)
AHMED	00B	PR D62 112003	S. Ahmed <i>et al.</i>	(CLEO Collab.)
ANASTASSOV	00	PRL 84 1393	A. Anastassov <i>et al.</i>	(CLEO Collab.)
BARATE	00R	PL B492 275	R. Barate <i>et al.</i>	(ALEPH Collab.)
BEHRENS	00	PR D61 052001	B.H. Behrens <i>et al.</i>	(CLEO Collab.)
BONVICINI	00	PRL 84 5940	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
CHEN	00	PRL 85 525	S. Chen <i>et al.</i>	(CLEO Collab.)
COAN	00	PRL 84 5283	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN...	00	PRL 85 515	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
CSORNA	00	PR D61 111101	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
JESSOP	00	PRL 85 2881	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
RICHICHI	00	PRL 85 520	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ABBIENDI	99J	EPJ C12 609	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AFFOLDER	99B	PRL 83 3378	T. Affolder <i>et al.</i>	(CDF Collab.)
BARTELT	99	PRL 82 3746	J. Bartelt <i>et al.</i>	(CLEO Collab.)
COAN	99	PR D59 111101	T.E. Coan <i>et al.</i>	(CLEO Collab.)
ABE	98B	PR D57 5382	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98O	PR D58 072001	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98Q	PR D58 092002	F. Abe <i>et al.</i>	(CDF Collab.)
ACCIARRI	98S	PL B438 417	M. Acciarri <i>et al.</i>	(L3 Collab.)
ANASTASSOV	98	PRL 80 4127	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ATHANAS	98	PRL 80 5493	M. Athanas <i>et al.</i>	(CLEO Collab.)
BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
BEHRENS	98	PRL 80 3710	B.H. Behrens <i>et al.</i>	(CLEO Collab.)
BERGFELD	98	PRL 81 272	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BRANDENB...	98	PRL 80 2762	G. Brandenbrug <i>et al.</i>	(CLEO Collab.)
GODANG	98	PRL 80 3456	R. Godang <i>et al.</i>	(CLEO Collab.)
ABE	97J	PRL 79 590	K. Abe <i>et al.</i>	(SLD Collab.)
ACCIARRI	97F	PL B396 327	M. Acciarri <i>et al.</i>	(L3 Collab.)
ARTUSO	97	PL B399 321	M. Artuso <i>et al.</i>	(CLEO Collab.)
ATHANAS	97	PRL 79 2208	M. Athanas <i>et al.</i>	(CLEO Collab.)
BROWDER	97	PR D56 11	T. Browder <i>et al.</i>	(CLEO Collab.)
FU	97	PRL 79 3125	X. Fu <i>et al.</i>	(CLEO Collab.)
JESSOP	97	PRL 79 4533	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
ABE	96B	PR D53 3496	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96C	PRL 76 4462	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96H	PRL 76 2015	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96L	PRL 76 4675	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96Q	PR D54 6596	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96R	PRL 77 5176	F. Abe <i>et al.</i>	(CDF Collab.)
ADAM	96D	ZPHY C72 207	W. Adam <i>et al.</i>	(DELPHI Collab.)
ALEXANDER	96T	PRL 77 5000	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ASNER	96	PR D53 1039	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BERGFELD	96B	PRL 77 4503	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BISHAI	96	PL B369 186	M. Bishai <i>et al.</i>	(CLEO Collab.)
BUSKULIC	96J	ZPHY C71 31	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
GIBAUT	96	PR D53 4734	D. Gibaut <i>et al.</i>	(CLEO Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	
ABREU	95N	PL B357 255	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95Q	ZPHY C68 13	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ADAM	95	ZPHY C68 363	W. Adam <i>et al.</i>	(DELPHI Collab.)
AKERS	95T	ZPHY C67 379	R. Akers <i>et al.</i>	(OPAL Collab.)
ALBRECHT	95D	PL B353 554	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	95	PL B341 435	J. Alexander <i>et al.</i>	(CLEO Collab.)
Also	95C	PL B347 469 (erratum)	J. Alexander <i>et al.</i>	(CLEO Collab.)
ARTUSO	95	PRL 75 785	M. Artuso <i>et al.</i>	(CLEO Collab.)
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	95	PL B343 444	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABE	94D	PRL 72 3456	F. Abe <i>et al.</i>	(CDF Collab.)
ALAM	94	PR D50 43	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	94D	PL B335 526	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ATHANAS	94	PRL 73 3503	M. Athanas <i>et al.</i>	(CLEO Collab.)
Also	95	PRL 74 3090 (erratum)	M. Athanas <i>et al.</i>	(CLEO Collab.)
PDG	94	PR D50 1173	L. Montanet <i>et al.</i>	(CERN, LBL, BOST+)
STONE	94	HEPSY 93-11	S. Stone	
Published in B Decays, 2nd Edition, World Scientific, Singapore				
ABREU	93D	ZPHY C57 181	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	93G	PL B312 253	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	93C	PL B307 247	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ALBRECHT	93E	ZPHY C60 11	H. Albrecht <i>et al.</i>	(ARGUS Collab.)

ALEXANDER	93B	PL B319 365	J. Alexander <i>et al.</i>	(CLEO Collab.)
AMMAR	93	PRL 71 674	R. Ammar <i>et al.</i>	(CLEO Collab.)
BEAN	93B	PRL 70 2681	A. Bean <i>et al.</i>	(CLEO Collab.)
BUSKULIC	93D	PL B307 194	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
Also	94H	PL B325 537 (errata)	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
SANGHERA	93	PR D47 791	S. Sanghera <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92C	PL B275 195	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92E	PL B277 209	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92G	ZPHY C54 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
BUSKULIC	92G	PL B295 396	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ALBRECHT	91B	PL B254 288	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91C	PL B255 297	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91E	PL B262 148	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BERKELMAN	91	ARNPS 41 1	K. Berkelman, S. Stone	(CORN, SYRA)
"Decays of <i>B</i> Mesons"				
FULTON	91	PR D43 651	R. Fulton <i>et al.</i>	(CLEO Collab.)
ALBRECHT	90B	PL B241 278	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	90J	ZPHY C48 543	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANTREASYAN	90B	ZPHY C48 553	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
BORTOLETTO	90	PRL 64 2117	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
Also	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
WEIR	90B	PR D41 1384	A.J. Weir <i>et al.</i>	(Mark II Collab.)
ALBRECHT	89G	PL B229 304	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AVERY	89B	PL B223 470	P. Avery <i>et al.</i>	(CLEO Collab.)
BEBEK	89	PRL 62 8	C. Bebek <i>et al.</i>	(CLEO Collab.)
BORTOLETTO	89	PRL 62 2436	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
ALBRECHT	88F	PL B209 119	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88K	PL B215 424	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87C	PL B185 218	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87D	PL B199 451	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AVERY	87	PL B183 429	P. Avery <i>et al.</i>	(CLEO Collab.)
BEBEK	87	PR D36 1289	C. Bebek <i>et al.</i>	(CLEO Collab.)
ALAM	86	PR D34 3279	M.S. Alam <i>et al.</i>	(CLEO Collab.)
PDG	86	PL 170B	M. Aguilar-Benitez <i>et al.</i>	(CERN, CIT+)
GILES	84	PR D30 2279	R. Giles <i>et al.</i>	(CLEO Collab.)