

Extra Dimensions

For explanation of terms used and discussion of significant model dependence of following limits, see the "Extra Dimensions Review." Limits are expressed in conventions of Giudice, Rattazzi, and Wells as explained in the Review. Footnotes describe originally quoted limit. n indicates the number of extra dimensions.

Limits not encoded here are summarized in the "Extra Dimensions Review."

A REVIEW GOES HERE – Check our WWW List of Reviews

Mass Limits on M_H

This section includes limits on mass scale of coefficient of dim-8 operators from KK graviton exchange in models with large extra dimensions. Ambiguity of UV divergent summation is represented by unknown λ . Quoted limits on M_H arbitrarily assume $|\lambda|=1$. Different papers use slightly different definitions of the mass scale; two popular conventions, M_H and Λ_T , are discussed in the above Review on "Extra Dimensions."

<u>VALUE (TeV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>0.84	95	1 ACHARD	02D L3	$e^+e^- \rightarrow \gamma\gamma$ ($\lambda=+1$)
>0.99	95	1 ACHARD	02D L3	$e^+e^- \rightarrow \gamma\gamma$ ($\lambda=-1$)
>1.2	95	2 ABBOTT	01 D0	$p\bar{p} \rightarrow e^+e^-, \gamma\gamma, \lambda=+1$
>1.1	95	2 ABBOTT	01 D0	$p\bar{p} \rightarrow e^+e^-, \gamma\gamma, \lambda=-1$
>0.60	95	3 ABBIENDI	00R OPAL	$e^+e^- \rightarrow \mu^+\mu, \lambda=+1$
>0.63	95	3 ABBIENDI	00R OPAL	$e^+e^- \rightarrow \mu^+\mu, \lambda=-1$
>0.63	95	3 ABBIENDI	00R OPAL	$e^+e^- \rightarrow \tau^+\tau^-, \lambda=+1$
>0.50	95	3 ABBIENDI	00R OPAL	$e^+e^- \rightarrow \tau^+\tau^-, \lambda=-1$
>0.68	95	3 ABBIENDI	00R OPAL	$e^+e^- \rightarrow \mu^+\mu, \tau^+\tau^-, \lambda=+1$
>0.61	95	3 ABBIENDI	00R OPAL	$e^+e^- \rightarrow \mu^+\mu, \tau^+\tau^-, \lambda=-1$
		4 ABREU	00A DLPH	
>0.559	95	5 ABREU	00S DLPH	$e^+e^- \rightarrow \mu^+\mu^-, \lambda=-1$
>0.649	95	5 ABREU	00S DLPH	$e^+e^- \rightarrow \mu^+\mu^-, \lambda=+1$
>0.450	95	5 ABREU	00S DLPH	$e^+e^- \rightarrow \tau^+\tau^-, \lambda=-1$
>0.564	95	5 ABREU	00S DLPH	$e^+e^- \rightarrow \tau^+\tau^-, \lambda=+1$
>0.542	95	5 ABREU	00S DLPH	$e^+e^- \rightarrow \mu^+\mu^-, \tau^+\tau^-, \lambda=-1$
>0.680	95	5 ABREU	00S DLPH	$e^+e^- \rightarrow \mu^+\mu^-, \tau^+\tau^-, \lambda=+1$
>0.48	95	6 ADLOFF	00 H1	$ep \rightarrow eX, \lambda=+1$
>0.72	95	6 ADLOFF	00 H1	$ep \rightarrow eX, \lambda=-1$
> 15–28	99.7	7 CHANG	00B RVUE	Electroweak
>0.98	95	8 CHEUNG	00 RVUE	$e^+e^- \rightarrow \gamma\gamma$
> 0.29–0.38	95	9 GRAESSER	00 RVUE	$(g-2)_\mu$
> 0.50–1.1	95	10 HAN	00 RVUE	Electroweak

>2.0	95	11	MATHEWS	00	RVUE	$\bar{p}p \rightarrow jj$ ($\lambda=\pm 1$)	
>1.0	95	12	MELE	00	RVUE	$e^+e^- \rightarrow VV$, $\lambda=+1$	
>1.1	95	12	MELE	00	RVUE	$e^+e^- \rightarrow VV$, $\lambda=-1$	
		13	ABBIENDI	99P	OPAL		
		14	ACCIARRI	99M	L3		
		15	ACCIARRI	99S	L3		
>1.412	95	16	BOURILKOV	99		$e^+e^- \rightarrow e^+e^-$, $\lambda=+1$	
>1.077	95	16	BOURILKOV	99		$e^+e^- \rightarrow e^+e^-$, $\lambda=-1$	
		1	ACHARD	02		search for s -channel graviton exchange effects in $e^+e^- \rightarrow \gamma\gamma$ at $E_{\text{cm}} = 192\text{--}209$ GeV.	
		2	ABBOTT	01		search for variations in differential cross sections to e^+e^- and $\gamma\gamma$ final states at the Tevatron. Bounds scale as $\lambda^{1/4}$.	
		3	ABBIENDI	00R		uses e^+e^- collisions at $\sqrt{s} = 189$ GeV. Bound is on M_H , where $\Lambda_T^4 = \frac{\pi}{2} M_H^4 / \lambda $.	
		4	ABREU	00A		search for s -channel graviton exchange effects in $e^+e^- \rightarrow \gamma\gamma$ at $E_{\text{cm}} = 189\text{--}202$ GeV. The limits $M_S > 713$ (691) GeV for $\lambda=+1$ (-1) are obtained, where M_S is a string mass scale.	
		5	ABREU	00S		uses e^+e^- collisions at $\sqrt{s} = 183$ and 189 GeV. Bound is on M_H , where $\Lambda_T^4 = \frac{\pi}{2} M_H^4 / \lambda $.	
		6	ADLOFF	00		uses ep collisions at $\sqrt{s} = 300$ GeV. Bound is on M_H , where $\Lambda_T^4 = \frac{\pi}{2} M_H^4 / \lambda $.	
		7	CHANG	00B		derive 3σ limit on M_H of (28,19,15) TeV for $n=(2,4,6)$ respectively assuming the presence of a torsional coupling in the gravitational action. Highly model dependent.	
		8	CHEUNG	00		obtains limits from anomalous diphoton production at OPAL due to graviton exchange. Original limit for $n=4$. However, unknown UV theory renders n dependence unreliable. Original paper works in HLZ convention. Limit scales as $\lambda^{1/4}$.	
		9	GRAESSER	00		obtains a bound from graviton contributions to $g-2$ of the muon through loops of 0.29 TeV for $n=2$ and 0.38 TeV for $n=4,6$. Limits scale as $\lambda^{1/2}$. However calculational scheme not well-defined without specification of high-scale theory. See the "Extra Dimensions Review."	
		10	HAN	00		calculates corrections to gauge boson self-energies from KK graviton loops and constrain them using S and T . Bounds on M_H range from 0.5 TeV ($n=6$) to 1.1 TeV ($n=2$); see text. Limits have strong dependence, λ^{n+2} , on unknown λ coefficient.	
		11	MATHEWS	00		search for evidence of graviton exchange in CDF and $D\bar{D}$ dijet production data. See their Table 2 for slightly stronger n -dependent bounds. Limits expressed in terms of $\tilde{M}_S^4 = m_H^4/8$.	
		12	MELE	00		obtains bound from KK graviton contributions to $e^+e^- \rightarrow VV$ ($V=\gamma, W, Z$) at LEP. Authors use Hewett conventions. Limits scale as $\lambda^{1/4}$.	
		13	ABBIENDI	99P		search for s -channel graviton exchange effects in $e^+e^- \rightarrow \gamma\gamma$ at $E_{\text{cm}} = 189$ GeV. The limits $G_+ > 660$ GeV and $G_- > 634$ GeV are obtained from combined $E_{\text{cm}} = 183$ and 189 GeV data, where G_{\pm} is a scale related to the fundamental gravity scale.	
		14	ACCIARRI	99M		search for the reaction $e^+e^- \rightarrow \gamma G$ and s -channel graviton exchange effects in $e^+e^- \rightarrow \gamma\gamma, W^+W^-, ZZ, e^+e^-, \mu^+\mu^-, \tau^+\tau^-, q\bar{q}$ at $E_{\text{cm}} = 183$ GeV. Limits on the gravity scale are listed in their Tables 1 and 2.	
		15	ACCIARRI	99S		search for the reaction $e^+e^- \rightarrow ZG$ and s -channel graviton exchange effects in $e^+e^- \rightarrow \gamma\gamma, W^+W^-, ZZ, e^+e^-, \mu^+\mu^-, \tau^+\tau^-, q\bar{q}$ at $E_{\text{cm}} = 189$ GeV. Limits on the gravity scale are listed in their Tables 1 and 2.	
		16	BOURILKOV	99		performs global analysis of LEP data on e^+e^- collisions at $\sqrt{s} = 183$ and 189 GeV. Bound is on Λ_T .	

Limits on R from On-Shell Production of Gravitons

This section include limits on on-shell (OS) production of gravitons, usually in astrophysical systems. Bounds quoted are on R the assumed common radius of the flat extra dimensions. n is the number of extra dimensions. Strong model dependence on light KK spectrum as explained in the "Extra Dimensions Review."

VALUE (nm)	CL%	DOCUMENT ID	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
< 9.6	($n=2$)	17 HANNESTAD 02	Astrophysics
< 0.04	($n=3$)	17 HANNESTAD 02	Astrophysics
< 0.07	($n=2$)	18 HANNESTAD 02	Astrophysics
< 0.001	($n=3$)	18 HANNESTAD 02	Astrophysics
< 660	($n=2$) 95	19 HANHART 01	Supernova
< 0.8	($n=3$) 95	19 HANHART 01	Supernova
< 90	($n=2$) 95	20 HANNESTAD 01	Astrophysics
< 0.19	($n=3$) 95	20 HANNESTAD 01	Astrophysics

¹⁷ HANNESTAD 02 obtain a limit on R from emission of KK gravitons in supernovae which subsequently decay to $\gamma\gamma$, e^+e^- , and $\nu\bar{\nu}$, contaminating the diffuse γ -ray background.

¹⁸ HANNESTAD 02 obtain a limit on R from bounds on the heating of neutron stars by the decay of the halo of KK gravitons produced in SN explosions and which surround neutron stars.

¹⁹ HANHART 01 obtain bounds on R from limits on graviton cooling of supernova SN 1987a using numerical simulations of proto-neutron star neutrino emission.

²⁰ HANNESTAD 01 obtain a limit on R from emission of KK gravitons in supernovae which subsequently decay to $\gamma\gamma$, e^+e^- , and $\nu\bar{\nu}$, contaminating the diffuse γ -ray background.

Limits on M_D from On-Shell Production of Gravitons

This section include limits on on-shell (OS) production of gravitons, usually in astrophysical systems. Bounds quoted are on M_D , which is related to the fundamental scale \bar{M}_D , introduced in Eq. (3) of the above note on "Extra Dimensions" by $M_D = (2\pi)^n/(2+n)\bar{M}_D$. Strong model dependence on light KK spectrum as explained in the "Extra Dimensions Review."

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$\gtrsim 1-2$		21 ANCHORDOQ.02B	RVUE	Cosmic Rays
$\gtrsim 1000$		22 FAIRBAIRN 01		Cosmology
>1.10	95	23 ABREU 00Z	DLPH	$e^+e^- \rightarrow \gamma G$
$\gtrsim 3-4$		24 CASSISI 00		Red giants
		25 ACCIARRI 99M	L3	
		26 ACCIARRI 99R	L3	
		27 ACCIARRI 99S	L3	

²¹ ANCHORDOQUI 02B derive bound on M_D from non-observation of black hole production in high-energy cosmic rays. Bound is stronger for larger n , but depends sensitively on threshold for black hole production.

²² FAIRBAIRN 01 obtains bound on the fundamental scale M_f , defined by $4\pi G = (1/(RM_f))^n/M_f^2$, from overproduction of KK modes in the early universe.

²³ ABREU 00Z uses e^+e^- collisions at $\sqrt{s} = 189$ GeV. Bound is $n=2$, and is equivalent to bound on common radius of $R < 0.4$ nm. For $n=4$ and 6, they obtain > 0.68 and 0.51 TeV, respectively. Systematic errors reduce limits by 9%, 3%, and 3% respectively.

- ²⁴ CASSISI 00 obtains bound from red-giant cooling, assuming $n=2$. Bound is on the scale M_S defined by $R^n = M_{Pl}^2 / (M_S^{n+2} \Omega_n)$ where Ω_n is the volume of the n -dimensional sphere with unit radius.
- ²⁵ ACCIARRI 99M search for the reaction $e^+ e^- \rightarrow \gamma G$ and s -channel graviton exchange effects in $e^+ e^- \rightarrow \gamma\gamma, W^+ W^-, ZZ, e^+ e^-, \mu^+ \mu^-, \tau^+ \tau^-, q\bar{q}$ at $E_{cm}=183$ GeV. Limits on the gravity scale are listed in their Tables 1 and 2.
- ²⁶ ACCIARRI 99R search for the reaction $e^+ e^- \rightarrow \gamma G$ at $E_{cm}=189$ GeV. Limits on the gravity scale are listed in their Table 4.
- ²⁷ ACCIARRI 99S search for the reaction $e^+ e^- \rightarrow ZG$ and s -channel graviton exchange effects in $e^+ e^- \rightarrow \gamma\gamma, W^+ W^-, ZZ, e^+ e^-, \mu^+ \mu^-, \tau^+ \tau^-, q\bar{q}$ at $E_{cm}=189$ GeV. Limits on the gravity scale are listed in their Tables 1 and 2.

Limits on Mass of Radion

This section includes limits on mass of radion, usually in context of Randall-Sundrum models. See the "Extra Dimension Review" for discussion of model dependence.

VALUE (GeV)	DOCUMENT ID	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●		
$\gtrsim 35$	28 MAHANTA 00	$Z \rightarrow$ radion $\ell\bar{\ell}$
>120	29 MAHANTA 00B	$p\bar{p} \rightarrow$ radion $\rightarrow \gamma\gamma$
²⁸ MAHANTA 00 obtain bound on radion mass in the RS model. Bound is from Higgs boson search at LEP I.		
²⁹ MAHANTA 00B uses $p\bar{p}$ collisions at $\sqrt{s}=1.8$ TeV; production via gluon-gluon fusion. Authors assume a radion vacuum expectation value of 1 TeV.		

Limits on $1/R = M_c$

This section includes limits on $1/R = M_c$, the compactification scale in models with TeV extra dimensions, due to exchange of Standard Model KK excitations. See the "Extra Dimension Review" for discussion of model dependence.

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>3.3	95	30 CORNET 00	RVUE	Electroweak
$> 3.3-3.8$	95	31 RIZZO 00	RVUE	Electroweak
³⁰ CORNET 00 translates a bound on the coefficient of the 4-fermion operator $(\bar{\ell}\gamma_\mu\tau^a\ell)(\bar{\ell}\gamma^\mu\tau^a\ell)$ derived by Hagiwara and Matsumoto into a limit on the mass scale of KK W bosons.				
³¹ RIZZO 00 obtains limits from global electroweak fits in models with a Higgs in the bulk (3.8 TeV) or on the standard brane (3.3 TeV).				

Limits on M_S

This section includes limits on the cutoff scale or string scale as indicated (TeV).

VALUE (TeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
>0.49	32 ACCIARRI 00P L3		$e^+ e^- \rightarrow e^+ e^-$
³² ACCIARRI 00P uses $e^+ e^-$ collisions at $\sqrt{s}=183$ and 189 GeV. Bound on string scale M_S from massive string modes. M_S is defined in hep-ph/0001166 by $M_S(1/\pi)^{1/8}\alpha^{-1/4} = M$ where $(4\pi G)^{-1} = M^{n+2}R^n$.			

REFERENCES FOR Extra Dimensions

ACHARD	02	PL B524 65	P. Achard <i>et al.</i>	(L3 Collab.)
ACHARD	02D	PL B531 28	P. Achard <i>et al.</i>	(L3 Collab.)
ANCHORDOQ...	02B	PR D66 103002	L. Anchordoqui <i>et al.</i>	
HANNESTAD	02	PRL 88 071301	S. Hannestad, G. Raffelt	
ABBOTT	01	PRL 86 1156	B. Abbott <i>et al.</i>	(D0 Collab.)
FAIRBAIRN	01	PL B508 335	M. Fairbairn	
HANHART	01	PL B509 1	C. Hanhart <i>et al.</i>	
HANNESTAD	01	PRL 87 051301	S. Hannestad, G. Raffelt	
ABBIENDI	00R	EPJ C13 553	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	00A	PL B491 67	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00S	PL B485 45	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00Z	EPJ C17 53	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	00P	PL B489 81	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADLOFF	00	PL B479 358	C. Adloff <i>et al.</i>	(H1 Collab.)
CASSISI	00	PL B481 323	S. Cassisi <i>et al.</i>	
CHANG	00B	PRL 85 3765	L.N. Chang <i>et al.</i>	
CHEUNG	00	PR D61 015005	K. Cheung	
CORNET	00	PR D61 037701	F. Cornet, M. Relano, J. Rico	
GRAESSER	00	PR D61 074019	M.L. Graesser	
HAN	00	PR D62 125018	T. Han, D. Marfatia, R.-J. Zhang	
MAHANTA	00	PL B480 176	U. Mahanta, S. Rakshit	
MAHANTA	00B	PL B483 196	U. Mahanta, A. Datta	
MATHEWS	00	JHEP 0007 008	P. Mathews, S. Raychaudhuri, K. Sridhar	
MELE	00	PR D61 117901	S. Mele, E. Sanchez	
RIZZO	00	PR D61 016007	T.G. Rizzo, J.D. Wells	
ABBIENDI	99P	PL B465 303	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ACCIARRI	99M	PL B464 135	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	99R	PL B470 268	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	99S	PL B470 281	M. Acciarri <i>et al.</i>	(L3 Collab.)
BOURILKOV	99	JHEP 08 006	D. Bourilkov	
