

$f_0(1710)$

$I^G(J^{PC}) = 0^+(0^{++})$

THE $f_0(1710)$

Updated April 2002 by M. Doser (CERN).

The $f_0(1710)$ is seen in the radiative decay $J/\psi(1S) \rightarrow \gamma f_0(1710)$; therefore $C = +1$. It decays into 2η and $K_S^0 K_S^0$, which implies $I^G J^{PC} = 0^+(even)^{++}$. The spin of the $f_0(1710)$ has been controversial, but evidence for spin 0 has accumulated recently in all production modes.

An analysis of radiative $J/\psi(1S)$ decays at BES into $\pi^+\pi^-\pi^+\pi^-$ (BAI 00) clearly favors spin 0. Combined amplitude analyses of the K^+K^- , K_SK_S , and $\pi^+\pi^-$ systems produced in $J/\psi(1S)$ radiative decay by MARK III (CHEN 91 and more recently DUNWOODIE 97) find a large spin-0 component, as well as reproducing known parameters of the $f_2(1270)$ and $f'_2(1525)$. In addition, a recent reanalysis (BUGG 95) of the 4π channel from MARK III, allowing both $\rho\rho$ and two $\pi\pi$ S waves, also finds a 0^{++} assignment for the $f_0(1710)$. Earlier analyses of this final state (BISSELLO 89B, BALTRUSAITIS 86B) found only pseudoscalar activity in the $f_0(1710)$ region, but considered only the process $J/\psi(1S) \rightarrow \gamma\rho\rho$. Similarly, earlier analyses of the K^+K^- system based on less statistics (BALTRUSAITIS 87, BAI 96) found a spin of 2 for the $f_0(1710)$.

A similar situation is present in central production, with earlier analyses favoring spin 2 over spin 0 (ARMSTRONG 89D). More recent analyses with greater statistics [BARBERIS 99 (K^+K^- , K_SK_S), BARBERIS 99B ($\pi^+\pi^-$), and FRENCH 99 (K^+K^-)], however, clearly indicate spin 0, and exclude spin 2. Generally, analyses preferring spin 2 concentrate on angular distributions in the $f_J(1710)$ region, and do not include possible interferences or distortion due to the nearby $f'_2(1525)$.

The $f_0(1710)$ is also observed in $K\bar{K}$ (FALVARD 88) in $J/\psi(1S) \rightarrow \omega K\bar{K}$ and $J/\psi(1S) \rightarrow \phi K\bar{K}$, but with no spin-parity analysis, as well as in $\eta\eta$ in radiative $J/\psi(1S)$ decays (EDWARDS 82). It is also clearly seen in 300-GeV/c pp central production in both K^+K^- and $K_S^0K_S^0$ (ARMSTRONG 89D). Mass and width are determined via a fit to non-interfering Breit-Wigners over a polynomial background, which leads to large systematic errors for the width. ARMSTRONG 93C also sees a broad peak in $\eta\eta$ at 1747 MeV, which may be the $f_0(1710)$.

This resonance is not observed in the hypercharge-exchange reactions $K^-p \rightarrow K_S^0K_S^0\Lambda$ (ASTON 88D) and $K^-p \rightarrow K_S^0K_S^0Y^*$ (BOLONKIN 86); these non-observations are explained by a spin of 0 (LINDENBAUM 92). It is not observed in $\bar{p}p$ interactions, neither via its $\pi\pi$ nor its $\eta\eta$ decay (AMSLER 02). A possible observation in $\gamma\gamma$ collisions leading to K_SK_S (BRACCINI 99, but no spin determination), and a non-observation in $\gamma\gamma \rightarrow \pi^+\pi^-$ (BARATE 00E), are consistent with a large $\bar{s}s$ component.

References

References may be found at the end of the $f_0(1710)$ Listing.

$f_0(1710)$ MASS

VALUE (MeV) 1714 \pm 5	EVTS OUR AVERAGE	DOCUMENT ID	TECN	COMMENT
1740 \pm 4	$+^{10}_{-25}$	1 BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
1740 $+^{30}_{-25}$		1 BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
1698 \pm 18		2 BARBERIS	00E	$450 \text{ } pp \rightarrow p_f \eta\eta p_s$
1710 \pm 12	\pm 11	3 BARBERIS	99D OMEG	$450 \text{ } pp \rightarrow K^+K^-$, $\pi^+\pi^-$
1710 \pm 25		4 FRENCH	99	$300 \text{ } pp \rightarrow p_f(K^+K^-)p_s$
1707 \pm 10		5 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+K^-$, $K_S^0K_S^0$
1698 \pm 15		5 AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1720 \pm 10	\pm 10	6 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+K^-$
1742 \pm 15		5 WILLIAMS	84 MPSF	$200 \pi^- N \rightarrow 2K_S^0 X$
1670 \pm 50		BLOOM	83 CBAL	$J/\psi \rightarrow \gamma 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1726 ± 7	74	⁷ CHEKANOV	04	ZEUS	$e p \rightarrow K_S^0 K_S^0 X$	
$1732+15$		⁸ ANISOVICH	03	RVUE		
1682 ± 16		TIKHOMIROV	03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$	
1670 ± 26	3651	^{1,9} NICHITIU	02	OBLX		
1767 ± 14	221	¹⁰ ACCIARRI	01H L3		$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{cm} = 91, 183-209 \text{ GeV}$	
1770 ± 12		^{11,12} ANISOVICH	99B	SPEC	$0.6-1.2 p\bar{p} \rightarrow \eta\eta\pi^0$	
1730 ± 15		¹ BARBERIS	99	OMEG	$p p \rightarrow p_s p_f K^+ K^-$	
1750 ± 20		¹ BARBERIS	99B	OMEG	$p p \rightarrow p_s p_f \pi^+ \pi^-$	
1750 ± 30		¹³ ANISOVICH	98B	RVUE	Compilation	
1720 ± 39		BAI	98H	BES	$J/\psi \rightarrow \gamma\pi^0\pi^0$	
1775 ± 1.5	57	¹⁴ BARKOV	98		$\pi^- p \rightarrow K_S^0 K_S^0 n$	
1690 ± 11		¹⁵ ABREU	96C	DLPH	$Z^0 \rightarrow K^+ K^- + X$	
1696 ± 5	^{+ 9} -34	⁶ BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$	
1781 ± 8	⁺¹⁰ -31	¹ BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$	
1768 ± 14		BALOSHIN	95	SPEC	$40 \pi^- C \rightarrow K_S^0 K_S^0 X$	
1750 ± 15		¹⁶ BUGG	95	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$	
1620 ± 16		⁶ BUGG	95	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$	
1748 ± 10		⁵ ARMSTRONG	93C	E760	$\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$	
~ 1750		BREAKSTONE	93	SFM	$p p \rightarrow p p \pi^+ \pi^- \pi^+ \pi^-$	
1744 ± 15		¹⁷ ALDE	92D	GAM2	$38 \pi^- p \rightarrow \eta\eta n$	
1713 ± 10		¹⁸ ARMSTRONG	89D	OMEG	$300 p p \rightarrow p p K^+ K^-$	
1706 ± 10		¹⁸ ARMSTRONG	89D	OMEG	$300 p p \rightarrow p p K_S^0 K_S^0$	
1700 ± 15		⁶ BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$	
1720 ± 60		¹ BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$	
1638 ± 10		¹⁹ FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$	
1690 ± 4		²⁰ FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$	
1755 ± 8		²¹ ALDE	86C	GAM2	$38 \pi^- p \rightarrow n2\eta$	
1730^{+2}_{-10}		²² LONGACRE	86	RVUE	$22 \pi^- p \rightarrow n2K_S^0$	
1650 ± 50		BURKE	82	MRK2	$J/\psi \rightarrow \gamma 2\rho$	
1640 ± 50		^{23,24} EDWARDS	82D	CBAL	$J/\psi \rightarrow \gamma 2\eta$	
1730 ± 10	± 20	²⁵ ETKIN	82C	MPS	$23 \pi^- p \rightarrow n2K_S^0$	

¹ $J^P = 0^+$.

²T-matrix pole.

³Supersedes BARBERIS 99 and BARBERIS 99B.

⁴ $J^P = 0^+$, supersedes by ARMSTRONG 89D.

⁵No $J^P C$ determination.

⁶ $J^P = 2^+$.

⁷Systematic errors not estimated.

⁸ K-matrix pole, assuming $J^P = 0^+$, from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

⁹ Decaying to $f_0(1370)\pi\pi$.

¹⁰ Spin 2 dominant, isospin not determined, could also be $I=1$.

¹¹ $J^P = 0^+$.

¹² Not seen by AMSLER 02.

¹³ T-matrix pole, assuming $J^P = 0^+$

¹⁴ No J^{PC} determination.

¹⁵ No J^{PC} determination, width not determined.

¹⁶ From a fit to the 0^+ partial wave.

¹⁷ ALDE 92D combines all the GAMS-2000 data.

¹⁸ $J^P = 2^+$, superseded by FRENCH 99.

¹⁹ From an analysis ignoring interference with $f'_2(1525)$.

²⁰ From an analysis including interference with $f'_2(1525)$.

²¹ Superseded by ALDE 92D.

²² Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

²³ $J^P = 2^+$ preferred.

²⁴ From fit neglecting nearby $f'_2(1525)$. Replaced by BLOOM 83.

²⁵ Superseded by LONGACRE 86.

$f_0(1710)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
140 \pm 10	OUR AVERAGE		Error includes scale factor of 1.2.		
166	\pm 5	$+15$	26 BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
120	\pm 40	50	26 BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
120	\pm 26		27 BARBERIS	00E	450 $p\bar{p} \rightarrow p_f \eta \eta p_s$
126	\pm 16	± 18	28 BARBERIS	99D OMEG	450 $p\bar{p} \rightarrow K^+ K^-$, $\pi^+ \pi^-$
105	\pm 34		29 FRENCH	99	300 $p\bar{p} \rightarrow p_f(K^+ K^-)p_s$
166.4 \pm 33.2			30 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$, $K_S^0 K_S^0$
136	\pm 28		30 AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
130	\pm 20		31 BALTRUSAIT..	87 MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
57	\pm 38		5 WILLIAMS	84 MPSF	200 $\pi^- N \rightarrow 2K_S^0 X$
160	\pm 80		BLOOM	83 CBAL	$J/\psi \rightarrow \gamma 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

38	\pm 20	74	32 CHEKANOV	04 ZEUS	$e\bar{p} \rightarrow K_S^0 K_S^0 X$
144	\pm 30		33,34 ANISOVICH	03 RVUE	
320	\pm 50		34,35 ANISOVICH	03 RVUE	
102	\pm 26		TIKHOMIROV	03 SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$

267	\pm 44	3651	^{26,36}	NICHITIU	02	OBLX	
187	\pm 60	221	³⁷	ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0,$	
						$E_{\text{cm}}^{\text{ee}} = 91,$	
						183–209 GeV	
220	\pm 40	38,39	ANISOVICH	99B	SPEC	$0.6\text{--}1.2 p\bar{p} \rightarrow \eta\eta\pi^0$	
100	\pm 25	26	BARBERIS	99	OMEG	$p p \rightarrow p_s p_f K^+ K^-$	
160	\pm 30	26	BARBERIS	99B	OMEG	$450 p p \rightarrow p_s p_f \pi^+ \pi^-$	
250	\pm 140	40	ANISOVICH	98B	RVUE	Compilation	
30	\pm 7	57	41	BARKOV	98	$\pi^- p \rightarrow K_S^0 K_S^0 n$	
103	\pm 18	⁺³⁰ ₋₁₁	31	BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
85	\pm 24	⁺²² ₋₁₉	26	BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
56	\pm 19			BALOSHIN	95	SPEC	$40 \pi^- C \rightarrow K_S^0 K_S^0 X$
160	\pm 40			42	BUGG	95	MRK3
160	\pm 60					$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$	
160	\pm 20			31	BUGG	95	MRK3
264	\pm 25					$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$	
200	to 300			30	ARMSTRONG	93C	E760
						$\bar{p}p \rightarrow \pi^0 \eta\eta \rightarrow 6\gamma$	
						BREAKSTONE	93 SFM
						$pp \rightarrow pp\pi^+\pi^-\pi^+\pi^-$	
< 80		90		43	ALDE	92D	GAM2
181	\pm 30					$38 \pi^- p \rightarrow \eta\eta N^*$	
104	\pm 30			44	ARMSTRONG	89D	OMEG
30	\pm 20					$300 pp \rightarrow ppK^+K^-$	
350	\pm 150			44	ARMSTRONG	89D	OMEG
148	\pm 17					$300 pp \rightarrow ppK_S^0 K_S^0$	
184	\pm 6			31	BOLONKIN	88	SPEC
122	\pm 74					$40 \pi^- p \rightarrow K_S^0 K_S^0 n$	
200	\pm 100			26	BOLONKIN	88	SPEC
220	\pm 100					$40 \pi^- p \rightarrow K_S^0 K_S^0 n$	
200.0	\pm 156.0			45	FALVARD	88	DM2
	\pm 9.0					$J/\psi \rightarrow \phi K^+ K^-$	
				46	FALVARD	88	DM2
						$J/\psi \rightarrow \phi K^+ K^-$	
				47	LONGACRE	86	RVUE
						$22 \pi^- p \rightarrow n2K_S^0$	
					BURKE	82	MRK2
						$J/\psi \rightarrow \gamma 2\rho$	
				48,49	EDWARDS	82D	CBAL
						$J/\psi \rightarrow \gamma 2\eta$	
				50	ETKIN	82B	MPS
						$23 \pi^- p \rightarrow n2K_S^0$	

26 $J^P = 0^+$.

27 T-matrix pole.

28 Supersedes BARBERIS 99 and BARBERIS 99B.

29 $J^P = 0^+$, supersedes by ARMSTRONG 89D.

30 No $J^P C$ determination.

31 $J^P = 2^+$.

32 Systematic errors not estimated.

33 (Solution I)

- ³⁴ K-matrix pole, assuming $J^P = 0^+$, from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.
- ³⁵ (Solution I)
- ³⁶ Decaying to $f_0(1370)\pi\pi$.
- ³⁷ Spin 2 dominant, isospin not determined, could also be $I=1$.
- ³⁸ $J^P = 0^+$.
- ³⁹ Not seen by AMSLER 02.
- ⁴⁰ T-matrix pole, assuming $J^P = 0^+$
- ⁴¹ No $J^P C$ determination.
- ⁴² From a fit to the 0^+ partial wave.
- ⁴³ ALDE 92D combines all the GAMS-2000 data.
- ⁴⁴ $J^P = 2^+$, (0^+ excluded).
- ⁴⁵ From an analysis ignoring interference with $f'_2(1525)$.
- ⁴⁶ From an analysis including interference with $f'_2(1525)$.
- ⁴⁷ Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.
- ⁴⁸ $J^P = 2^+$ preferred.
- ⁴⁹ From fit neglecting nearby $f'_2(1525)$. Replaced by BLOOM 83.
- ⁵⁰ From an amplitude analysis of the $K_S^0 K_S^0$ system, superseded by LONGACRE 86.

$f_0(1710)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	seen
Γ_2 $\eta\eta$	seen
Γ_3 $\pi\pi$	seen
Γ_4 $\gamma\gamma$	

$f_0(1710) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_4/\Gamma$
VALUE (eV)	CL \%
<110	
95	52 BEHREND 89C CELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
$49 \pm 11 \pm 13$	53 ACCIARRI 01H L3 $\gamma\gamma \rightarrow K_S^0 K_S^0$, $E_{\text{cm}}^{ee} = 91, 183-209$ GeV
<480	95 ALBRECHT 90G ARG $\gamma\gamma \rightarrow K^+ K^-$
<280	95 52 ALTHOFF 85B TASS $\gamma\gamma \rightarrow K\bar{K}\pi$

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_4/\Gamma$
VALUE (keV)	CL \%
<0.82	95 51 BARATE 00E ALEP $\gamma\gamma \rightarrow \pi^+ \pi^-$

⁵¹ Assuming spin 0.

⁵² Assuming helicity 2.

⁵³ Spin 2 dominant, isospin not determined, could also be $I=1$.

$f_0(1710)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_1/Γ
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$0.38^{+0.09}_{-0.19}$	54,55 LONGACRE	86 MPS	$22 \pi^- p \rightarrow n2K_S^0$	

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	Γ_2/Γ
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.18^{+0.03}_{-0.13}$	54,55 LONGACRE	86 RVUE	

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
not seen	AMSLER	02 CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta\eta, \pi^0 \pi^0 \pi^0$	
$0.039^{+0.002}_{-0.024}$	54,55 LONGACRE	86 RVUE		

$\Gamma(\pi\pi)/\Gamma(K\bar{K})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ_1
$0.2 \pm 0.024 \pm 0.036$	BARBERIS	99D OMEG	$450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$5.8^{+9.1}_{-5.5}$	56 ANISOVICH	02D SPEC	Combined fit	
0.39 ± 0.14	ARMSTRONG 91	OMEG	$300 pp \rightarrow pp\pi\pi, ppK\bar{K}$	

$\Gamma(\eta\eta)/\Gamma(K\bar{K})$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2/Γ_1
0.48 ± 0.15		BARBERIS	00E	$450 pp \rightarrow p_f \eta\eta p_s$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$0.46^{+0.70}_{-0.38}$		56 ANISOVICH	02D SPEC	Combined fit	
<0.02	90	57 PROKOSHKIN 91	GA24	$300 \pi^- p \rightarrow \pi^- p\eta\eta$	

⁵⁴ From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2.
⁵⁵ Fit with constrained inelasticity.
⁵⁶ From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n$, $\eta\eta n$, $\eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.
⁵⁷ Combining results of GAM4 with those of ARMSTRONG 89D.

$f_0(1710)$ REFERENCES

CHEKANOV	04	PL B578 33	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>	
		Translated from YAF 65 1583.		
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARATE	00E	PL B472 189	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>	
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)
FRENCH	99	PL B460 213	B. French <i>et al.</i>	(WA76 Collab.)
ANISOVICH	98B	UFN 41 419	V.V. Anisovich <i>et al.</i>	
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARKOV	98	JEPTL 68 764	B.P. Barkov <i>et al.</i>	
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)
		Translated from YAF 58 50.		
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BREAKSTONE	93	ZPHY C58 251	A.M. Breakstone <i>et al.</i>	(IOWA, CERN, DORT+)
ALDE	92D	PL B284 457	D.M. Alde <i>et al.</i>	(GAM2 Collab.)
Also	91	SJNP 54 451	D.M. Alde <i>et al.</i>	(GAM2 Collab.)
		Translated from YAF 54 745.		
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2, GAM4 Collab.)
		Translated from DANS 316 900.		
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LAPO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LAPO, CLER, FRAS+)
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
ALDE	86C	PL B182 105	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)
WILLIAMS	84	PR D30 877	E.G.H. Williams <i>et al.</i>	(VAND, NDAM, TUFTS+)
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
ETKIN	82C	PR D25 2446	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)

OTHER RELATED PAPERS

ANISOVICH	03B	PAN 66 741	V.V. Anisovich, V.A. Nikonorov, A.V. Sarantsev	
		Translated from YAF 66 772.		
AMSLER	02B	PL B541 22	C. Amsler	
JIN	02	PR D66 057505	H. Jin, X. Zhang	
KLEEFELD	02	PR D66 034007	F. Kleefeld <i>et al.</i>	
RUPP	02	PR D65 078501	G. Rupp, E. van Beveren, M.D. Scadron	
SHAKIN	02	PR D65 078502	C.M. Shakin, H. Wang	
TESHIMA	02	JPG 28 1391	T. Teshima, I. Kitamura, N. Morisita	
VOLKOV	02	PAN 65 1657	M.K. Volkov, V.L. Yudichev	
		Translated from YAF 65 1701.		
LI	01B	EPJ C19 529	D.-M. Li, H. Yu, Q.-X. Shen	
VOLKOV	01	PAN 64 2006	M.K. Volkov, V.L. Yudichev	
		Translated from YAF 64 2091.		
ANISOVICH	99H	PL B467 289	A.V. Anisovich, V.V. Anisovich	
GRYGOREV	99	PAN 62 470	V.K. Grygorev <i>et al.</i>	
		Translated from YAF 62 513.		

PROKOSHKIN	99	PAN 62 356	Yu.D. Prokoshkin
		Translated from YAF 62	396.
ANISOVICH	97	PL B395 123	A.V. Anisovich, A.V. Sarantsev (PNPI)
LINDENBAUM	92	PL B274 492	S.J. Lindenbaum, R.S. Longacre (BNL)
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i> (DM2 Collab.)
ASTON	88D	NP B301 525	D. Aston <i>et al.</i> (SLAC, NAGO, CINC, INUS)
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i> (Axial Field Spec. Collab.)
ARMSTRONG	86B	PL 167B 133	T.A. Armstrong <i>et al.</i> (ATHU, BARI, BIRM+)
BALTRUSAIT...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i> (Mark III Collab.)
ALTHOFF	83	PL 121B 216	M. Althoff <i>et al.</i> (TASSO Collab.)
BARNETT	83B	PL 120B 455	B. Barnett <i>et al.</i> (JHU)
BARNES	82B	NP B198 380	T. Barnes, F.E. Close, S. Monaghan (RHEL, OXFTP)
TANIMOTO	82	PL 116B 198	M. Tanimoto (BIEL)
