

$b\bar{b}$ MESONS

$\Upsilon(1S)$

$I^G(J^{PC}) = 0^-(1^{--})$

Mass $m = 9460.30 \pm 0.26$ MeV ($S = 3.3$)

Full width $\Gamma = 53.0 \pm 1.5$ keV

$\Gamma_{ee} = 1.314 \pm 0.029$ keV

$\Upsilon(1S)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\tau^+ \tau^-$	$(2.67^{+0.14}_{-0.16})\%$		4384
$e^+ e^-$	$(2.38 \pm 0.11)\%$		4730
$\mu^+ \mu^-$	$(2.48 \pm 0.06)\%$		4729
Hadronic decays			
$\eta'(958)$ anything	$(2.8 \pm 0.4)\%$		—
$J/\psi(1S)$ anything	$(1.1 \pm 0.4) \times 10^{-3}$		4223
$\rho\pi$	$< 2 \times 10^{-4}$	90%	4697
$\pi^+ \pi^-$	$< 5 \times 10^{-4}$	90%	4728
$K^+ K^-$	$< 5 \times 10^{-4}$	90%	4704
$p\bar{p}$	$< 5 \times 10^{-4}$	90%	4636
$\pi^0 \pi^+ \pi^-$	$< 1.84 \times 10^{-5}$	90%	4725
Radiative decays			
$\gamma \pi^+ \pi^-$	$(6.3 \pm 1.8) \times 10^{-5}$		4728
$\gamma \pi^0 \pi^0$	$(1.7 \pm 0.7) \times 10^{-5}$		4728
$\gamma 2h^+ 2h^-$	$(7.0 \pm 1.5) \times 10^{-4}$		4720
$\gamma 3h^+ 3h^-$	$(5.4 \pm 2.0) \times 10^{-4}$		4703
$\gamma 4h^+ 4h^-$	$(7.4 \pm 3.5) \times 10^{-4}$		4679
$\gamma \pi^+ \pi^- K^+ K^-$	$(2.9 \pm 0.9) \times 10^{-4}$		4686
$\gamma 2\pi^+ 2\pi^-$	$(2.5 \pm 0.9) \times 10^{-4}$		4720
$\gamma 3\pi^+ 3\pi^-$	$(2.5 \pm 1.2) \times 10^{-4}$		4703
$\gamma 2\pi^+ 2\pi^- K^+ K^-$	$(2.4 \pm 1.2) \times 10^{-4}$		4658
$\gamma \pi^+ \pi^- p\bar{p}$	$(1.5 \pm 0.6) \times 10^{-4}$		4604
$\gamma 2\pi^+ 2\pi^- p\bar{p}$	$(4 \pm 6) \times 10^{-5}$		4563
$\gamma 2K^+ 2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$		4601
$\gamma \eta'(958)$	$< 1.6 \times 10^{-5}$	90%	4682
$\gamma \eta$	$< 2.1 \times 10^{-5}$	90%	4714
$\gamma f'_2(1525)$	$< 1.4 \times 10^{-4}$	90%	4607
$\gamma f_2(1270)$	$(8 \pm 4) \times 10^{-5}$		4644
$\gamma \eta(1405)$	$< 8.2 \times 10^{-5}$	90%	4625
$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$< 2.6 \times 10^{-4}$	90%	4576

$\gamma f_0(2200) \rightarrow \gamma K^+ K^-$	< 2	$\times 10^{-4}$	90%	4475
$\gamma f_J(2220) \rightarrow \gamma K^+ K^-$	< 1.5	$\times 10^{-5}$	90%	4469
$\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-$	< 1.2	$\times 10^{-5}$	90%	—
$\gamma f_J(2220) \rightarrow \gamma p\bar{p}$	< 1.6	$\times 10^{-5}$	90%	—
$\gamma \eta(2225) \rightarrow \gamma \phi\phi$	< 3	$\times 10^{-3}$	90%	4469
γX ($X =$ pseudoscalar with $m < 7.2$ GeV)	< 3	$\times 10^{-5}$	90%	—
$\gamma X\bar{X}$ ($X\bar{X} =$ vectors with $m < 3.1$ GeV)	< 1	$\times 10^{-3}$	90%	—

$\chi_{b0}(1P)$ [a]

$I^G(J^{PC}) = 0^+(0^{++})$
 J needs confirmation.

Mass $m = 9859.9 \pm 1.0$ MeV

$\chi_{b0}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\gamma \gamma(1S)$	< 6 %	90%	391

$\chi_{b1}(1P)$ [a]

$I^G(J^{PC}) = 0^+(1^{++})$
 J needs confirmation.

Mass $m = 9892.7 \pm 0.6$ MeV (S = 1.1)

$\chi_{b1}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\gamma \gamma(1S)$	(35 \pm 8) %	423

$\chi_{b2}(1P)$ [a]

$I^G(J^{PC}) = 0^+(2^{++})$
 J needs confirmation.

Mass $m = 9912.6 \pm 0.5$ MeV (S = 1.1)

$\chi_{b2}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\gamma \gamma(1S)$	(22 \pm 4) %	442

$\Upsilon(2S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 10.02326 \pm 0.00031$ GeV

Full width $\Gamma = 43 \pm 6$ keV

$\Gamma_{ee} = 0.576 \pm 0.024$ keV

$\Upsilon(2S)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\gamma(1S)\pi^+\pi^-$	(18.8 \pm 0.6) %		475
$\gamma(1S)\pi^0\pi^0$	(9.0 \pm 0.8) %		480
$\tau^+\tau^-$	(1.7 \pm 1.6) %		4686
$\mu^+\mu^-$	(1.31 \pm 0.21) %		5011
e^+e^-	(1.34 \pm 0.20) %		5012
$\gamma(1S)\pi^0$	< 1.1 $\times 10^{-3}$	90%	531
$\gamma(1S)\eta$	< 2 $\times 10^{-3}$	90%	126
$J/\psi(1S)$ anything	< 6 $\times 10^{-3}$	90%	4533

Radiative decays

$\gamma\chi_{b1}(1P)$	(6.8 \pm 0.7) %	130
$\gamma\chi_{b2}(1P)$	(7.0 \pm 0.6) %	110
$\gamma\chi_{b0}(1P)$	(3.8 \pm 0.6) %	162
$\gamma f_0(1710)$	< 5.9 $\times 10^{-4}$	90%
$\gamma f'_2(1525)$	< 5.3 $\times 10^{-4}$	90%
$\gamma f_2(1270)$	< 2.41 $\times 10^{-4}$	90%

$\chi_{b0}(2P)$ ^[a]

$$I^G(J^{PC}) = 0^+(0^{++})$$

J needs confirmation.

Mass $m = 10.2321 \pm 0.0006$ GeV

$\chi_{b0}(2P)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\gamma\Upsilon(2S)$	(4.6 \pm 2.1) %	207
$\gamma\Upsilon(1S)$	(9 \pm 6) $\times 10^{-3}$	743

$\chi_{b1}(2P)$ ^[a]

$$I^G(J^{PC}) = 0^+(1^{++})$$

J needs confirmation.

Mass $m = 10.2552 \pm 0.0005$ GeV

$$m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)} = 23.5 \pm 1.0 \text{ MeV}$$

$\chi_{b1}(2P)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor	p (MeV/c)
$\gamma\Upsilon(2S)$	(21 \pm 4) %	1.5	229
$\gamma\Upsilon(1S)$	(8.5 \pm 1.3) %	1.3	764

$\chi_{b2}(2P)$ [a]

$J^G(J^{PC}) = 0^+(2^{++})$
J needs confirmation.

Mass $m = 10.2685 \pm 0.0004$ GeV

$m_{\chi_{b2}(2P)} - m_{\chi_{b1}(2P)} = 13.5 \pm 0.6$ MeV

$\chi_{b2}(2P)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\gamma \Upsilon(2S)$	(16.2 \pm 2.4) %	242
$\gamma \Upsilon(1S)$	(7.1 \pm 1.0) %	776

$\Upsilon(3S)$

$J^G(J^{PC}) = 0^-(1^{--})$

Mass $m = 10.3552 \pm 0.0005$ GeV

Full width $\Gamma = 26.3 \pm 3.4$ keV

$\Upsilon(3S)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\Upsilon(2S)$ anything	(10.6 \pm 0.8) %		296
$\Upsilon(2S)\pi^+\pi^-$	(2.8 \pm 0.6) %	S=2.2	177
$\Upsilon(2S)\pi^0\pi^0$	(2.00 \pm 0.32) %		190
$\Upsilon(2S)\gamma\gamma$	(5.0 \pm 0.7) %		327
$\Upsilon(1S)\pi^+\pi^-$	(4.48 \pm 0.21) %		813
$\Upsilon(1S)\pi^0\pi^0$	(2.06 \pm 0.28) %		816
$\Upsilon(1S)\eta$	< 2.2 $\times 10^{-3}$	CL=90%	677
$\mu^+\mu^-$	(1.81 \pm 0.17) %		5177
e^+e^-	seen		5178

Radiative decays

$\gamma \chi_{b2}(2P)$	(11.4 \pm 0.8) %	S=1.3	86
$\gamma \chi_{b1}(2P)$	(11.3 \pm 0.6) %		100
$\gamma \chi_{b0}(2P)$	(5.4 \pm 0.6) %	S=1.1	122

$\Upsilon(4S)$
or **$\Upsilon(10580)$**

$J^G(J^{PC}) = 0^-(1^{--})$

Mass $m = 10.5800 \pm 0.0035$ GeV

Full width $\Gamma = 20 \pm 4$ MeV

$\Gamma_{ee} = 0.248 \pm 0.031$ keV (S = 1.3)

$\Upsilon(4S)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$B\bar{B}$	> 96 %	95%	335
non- $B\bar{B}$	< 4 %	95%	-
e^+e^-	(2.8 \pm 0.7) $\times 10^{-5}$		5290

$J/\psi(1S)$ anything	< 1.9	$\times 10^{-4}$	95%	—
D^{*+} anything + c.c.	< 7.4	%	90%	5099
ϕ anything	< 2.3	$\times 10^{-3}$	90%	5240
$\Upsilon(1S)$ anything	< 4	$\times 10^{-3}$	90%	1053
$\Upsilon(1S)\pi^+\pi^-$	< 1.2	$\times 10^{-4}$	90%	1027
$\Upsilon(2S)\pi^+\pi^-$	< 3.9	$\times 10^{-4}$	90%	469

$\Upsilon(10860)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 10.865 \pm 0.008$ GeV ($S = 1.1$)

Full width $\Gamma = 110 \pm 13$ MeV

$\Gamma_{ee} = 0.31 \pm 0.07$ keV ($S = 1.3$)

$\Upsilon(10860)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$e^+ e^-$

$(2.8 \pm 0.7) \times 10^{-6}$

5432

$\Upsilon(11020)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 11.019 \pm 0.008$ GeV

Full width $\Gamma = 79 \pm 16$ MeV

$\Gamma_{ee} = 0.130 \pm 0.030$ keV

$\Upsilon(11020)$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$e^+ e^-$

$(1.6 \pm 0.5) \times 10^{-6}$

5510

NOTES

[a] Spectroscopic labeling for these states is theoretical, pending experimental information.