

**$\Delta(1910)$   $P_{31}$**  $I(J^P) = \frac{3}{2}(\frac{1}{2}^+)$  Status: \*\*\*\*

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

 **$\Delta(1910)$  BREIT-WIGNER MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1870 to 1920 (<math>\approx 1910</math>) OUR ESTIMATE</b>			
1882 $\pm 10$	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
1910 $\pm 40$	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1888 $\pm 20$	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1995 $\pm 12$	VRANA 00	DPWA	Multichannel
2152	ARNDT 95	DPWA	$\pi N \rightarrow N\pi$
1960.1 $\pm 21.0$	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
2121.4 $^{+13.0}_{-14.3}$	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
1921	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1899	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1790	<sup>2</sup> LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

 **$\Delta(1910)$  BREIT-WIGNER WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>190 to 270 (<math>\approx 250</math>) OUR ESTIMATE</b>			
239 $\pm 25$	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$
225 $\pm 50$	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
280 $\pm 50$	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
713 $\pm 465$	VRANA 00	DPWA	Multichannel
760	ARNDT 95	DPWA	$\pi N \rightarrow N\pi$
152.9 $\pm 60.0$	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
172.2 $\pm 37.0$	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
351	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
230	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
170	<sup>2</sup> LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

 **$\Delta(1910)$  POLE POSITION****REAL PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1830 to 1880 (<math>\approx 1855</math>) OUR ESTIMATE</b>			
1874	<sup>3</sup> HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
1880 $\pm 30$	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1880	VRANA	00	DPWA	Multichannel
1810	ARNDT	95	DPWA	$\pi N \rightarrow N\pi$
1950	ARNDT	91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1792 or 1801	<sup>2</sup> LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$

## -2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>200 to 500 (<math>\approx 350</math>) OUR ESTIMATE</b>			
283	<sup>3</sup> HOEHLER	93	SPED $\pi N \rightarrow \pi N$
200±40	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
496	VRANA	00	DPWA Multichannel
494	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
398	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
172 or 165	<sup>2</sup> LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$

## $\Delta(1910)$ ELASTIC POLE RESIDUE

### MODULUS $|r|$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
38	HOEHLER	93	SPED $\pi N \rightarrow \pi N$
20±4	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
53	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
37	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

### PHASE $\theta$

VALUE (°)	DOCUMENT ID	TECN	COMMENT
– 90±30	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
– 176	ARNDT	95	DPWA $\pi N \rightarrow N\pi$
– 91	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90

## $\Delta(1910)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $N\pi$	15–30 %
$\Gamma_2$ $\Sigma K$	
$\Gamma_3$ $N\pi\pi$	
$\Gamma_4$ $\Delta\pi$	
$\Gamma_5$ $\Delta(1232)\pi$ , P-wave	

$\Gamma_6$	$N\rho$		
$\Gamma_7$	$N\rho, S=3/2, P\text{-wave}$		
$\Gamma_8$	$N(1440)\pi$		
$\Gamma_9$	$N(1440)\pi, P\text{-wave}$		
$\Gamma_{10}$	$N\gamma$	0.0–0.2 %	
$\Gamma_{11}$	$N\gamma, \text{ helicity}=1/2$	0.0–0.2 %	

## $\Delta(1910)$ BRANCHING RATIOS

### $\Gamma(N\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
<b>0.15 to 0.3 OUR ESTIMATE</b>				
0.23±0.08	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$	
0.19±0.03	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.24±0.06	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.29±0.21	VRANA 00	DPWA	Multichannel	
0.26	ARNDT 95	DPWA	$\pi N \rightarrow N\pi$	
0.17	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	
0.40	<sup>1</sup> CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	

### $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1910) \rightarrow \Sigma K$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
< 0.03	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.019	LIVANOS 80	DPWA	$\pi p \rightarrow \Sigma K$	
0.082 to 0.184	<sup>4</sup> DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$	

Note: Signs of couplings from  $\pi N \rightarrow N\pi\pi$  analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the  $\Delta(1620) S_{31}$  coupling to  $\Delta(1232)\pi$ .

### $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1910) \rightarrow \Delta(1232)\pi, P\text{-wave}$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
+0.06	<sup>2</sup> LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	

### $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1910) \rightarrow N\rho, S=3/2, P\text{-wave}$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_7)^{1/2}/\Gamma$
+0.29	<sup>2</sup> LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

+0.17	<sup>5</sup> NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$
-------	----------------------------	------	-----------------------------

### $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1910) \rightarrow N(1440)\pi, P\text{-wave}$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_9)^{1/2}/\Gamma$
-0.39±0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N & N\pi\pi$	

$\Gamma(N(1440)\pi)/\Gamma_{\text{total}}$	$\Gamma_8/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
0.56 ± 0.07	VRANA	00	DPWA Multichannel

## $\Delta(1910)$ PHOTON DECAY AMPLITUDES

### $\Delta(1910) \rightarrow N\gamma$ , helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV $^{-1/2}$ )	DOCUMENT ID	TECN	COMMENT
<b>+0.003±0.014 OUR ESTIMATE</b>			
-0.002 ± 0.008	ARNDT	96	IPWA $\gamma N \rightarrow \pi N$
0.014 ± 0.030	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.025 ± 0.011	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.012 ± 0.005	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.031 ± 0.004	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.005 ± 0.030	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.032 ± 0.003	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.035 ± 0.021	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

## $\Delta(1910)$ FOOTNOTES

- <sup>1</sup> CHEW 80 reports four resonances in the  $P_{31}$  wave — see also the  $\Delta(1750)$ . Problems with this analysis are discussed in section 2.1.11 of HOEHLER 83.  
<sup>2</sup> LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to  $\pi N \rightarrow N\pi\pi$  data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.  
<sup>3</sup> See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of  $N$  and  $\Delta$  resonances as determined from Argand diagrams of  $\pi N$  elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.  
<sup>4</sup> The range given for DEANS 75 is from the four best solutions.  
<sup>5</sup> Evidence for this coupling is weak; see NOVOSELLER 78. This coupling assumes the mass is near 1820 MeV.

## $\Delta(1910)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

VRANA	00	PRPL 328 181	T.P. Vrana, S.A. Dytman,, T.-S.H. Lee	(PITT+)
ARNDT	96	PR C53 430	R.A. Arndt, I.I. Strakovsky, R.L. Workman	(VPI)
ARNDT	95	PR C52 2120	R.A. Arndt <i>et al.</i>	(VPI, BRCO)
HOEHLER	93	$\pi N$ Newsletter 9 1	G. Hohler	(KARL)
LI	93	PR C47 2759	Z.J. Li <i>et al.</i>	(VPI)
MANLEY	92	PR D45 4002	D.M. Manley, E.M. Saleski	(KENT) IJP
Also	84	PR D30 904	D.M. Manley <i>et al.</i>	(VPI)
ARNDT	91	PR D43 2131	R.A. Arndt <i>et al.</i>	(VPI, TELE) IJP
CANDLIN	84	NP B238 477	D.J. Candlin <i>et al.</i>	(EDIN, RAL, LOWC)
CRAWFORD	83	NP B211 1	R.L. Crawford, W.T. Morton	(GLAS)
HOEHLER	83	Landolt-Bornstein 1/9B2	G. Hohler	(KARLT)
PDG	82	PL 111B	M. Roos <i>et al.</i>	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	N. Awaji, R. Kajikawa	(NAGO)
Also	82	NP B197 365	K. Fujii <i>et al.</i>	(NAGO)
ARAI	80	Toronto Conf. 93	I. Arai	(INUS)
Also	82	NP B194 251	I. Arai, H. Fujii	(INUS)

CHEW	80	Toronto Conf. 123	D.M. Chew	(LBL) IJP
CRAWFORD	80	Toronto Conf. 107	R.L. Crawford	(GLAS)
CUTKOSKY	80	Toronto Conf. 19	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
Also	79	PR D20 2839	R.E. Cutkosky <i>et al.</i>	(CMU, LBL) IJP
LIVANOS	80	Toronto Conf. 35	P. Livanos <i>et al.</i>	(SACL) IJP
HOEHLER	79	PDAT 12-1	G. Hohler <i>et al.</i>	(KARLT) IJP
Also	80	Toronto Conf. 3	R. Koch	(KARLT) IJP
BARBOUR	78	NP B141 253	I.M. Barbour, R.L. Crawford, N.H. Parsons	(GLAS)
NOVOSELLER	78	NP B137 509	D.E. Novoseller	(CIT) IJP
Also	78B	NP B137 445	D.E. Novoseller	(CIT) IJP
LONGACRE	77	NP B122 493	R.S. Longacre, J. Dolbeau	(SACL) IJP
Also	76	NP B108 365	J. Dolbeau <i>et al.</i>	(SACL) IJP
DEANS	75	NP B96 90	S.R. Deans <i>et al.</i>	(SFLA, ALAH) IJP