

$\Xi(1950)$ $I(J^P) = \frac{1}{2}(??)$ Status: ***

We list here everything reported between 1875 and 2000 MeV. The accumulated evidence for a Ξ near 1950 MeV seems strong enough to include a $\Xi(1950)$ in the main Baryon Table, but not much can be said about its properties. In fact, there may be more than one Ξ near this mass.

 $\Xi(1950)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1950±15 OUR ESTIMATE				
1955± 6		ADAMOVICH	99B WA89	Σ^- nucleus, 345 GeV
1944± 9	129	BIAGI	87 SPEC	$\Xi^- \text{Be} \rightarrow (\Xi^- \pi^+) \pi^- X$
1963± 5±2	63	BIAGI	87C SPEC	$\Xi^- \text{Be} \rightarrow (\Lambda \bar{K}^0) X$
1937± 7	150	BIAGI	81 SPEC	SPS hyperon beam
1961±18	139	BRIEFEL	77 HBC	2.87 $K^- p \rightarrow \Xi^- \pi^+ X$
1936±22	44	BRIEFEL	77 HBC	2.87 $K^- p \rightarrow \Xi^0 \pi^- X$
1964±10	56	BRIEFEL	77 HBC	$\Xi(1530)\pi$
1900±12		DIBIANCA	75 DBC	$\Xi \pi$
1952±11	25	ROSS	73C	$(\Xi\pi)^-$
1956± 6	29	BADIER	72 HBC	$\Xi \pi, \Xi \pi \pi, Y K$
1955±14	21	GOLDWASSER	70 HBC	$\Xi \pi$
1894±18	66	DAUBER	69 HBC	$\Xi \pi$
1930±20	27	ALITTI	68 HBC	$\Xi^- \pi^+$
1933±16	35	BADIER	65 HBC	$\Xi^- \pi^+$

 $\Xi(1950)$ WIDTH

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
60±20 OUR ESTIMATE				
68±22		ADAMOVICH	99B WA89	Σ^- nucleus, 345 GeV
100±31	129	BIAGI	87 SPEC	$\Xi^- \text{Be} \rightarrow (\Xi^- \pi^+) \pi^- X$
25±15±1.2	63	BIAGI	87C SPEC	$\Xi^- \text{Be} \rightarrow (\Lambda \bar{K}^0) X$
60± 8	150	BIAGI	81 SPEC	SPS hyperon beam
159±57	139	BRIEFEL	77 HBC	2.87 $K^- p \rightarrow \Xi^- \pi^+ X$
87±26	44	BRIEFEL	77 HBC	2.87 $K^- p \rightarrow \Xi^0 \pi^- X$
60±39	56	BRIEFEL	77 HBC	$\Xi(1530)\pi$
63±78		DIBIANCA	75 DBC	$\Xi \pi$
38±10		ROSS	73C	$(\Xi\pi)^-$
35±11	29	BADIER	72 HBC	$\Xi \pi, \Xi \pi \pi, Y K$
56±26	21	GOLDWASSER	70 HBC	$\Xi \pi$
98±23	66	DAUBER	69 HBC	$\Xi \pi$
80±40	27	ALITTI	68 HBC	$\Xi^- \pi^+$
140±35	35	BADIER	65 HBC	$\Xi^- \pi^+$

$\Xi(1950)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Lambda \bar{K}$	seen
Γ_2 $\Sigma \bar{K}$	possibly seen
Γ_3 $\Xi \pi$	seen
Γ_4 $\Xi(1530) \pi$	
Γ_5 $\Xi \pi \pi$ (not $\Xi(1530) \pi$)	

$\Xi(1950)$ BRANCHING RATIOS

$\Gamma(\Sigma \bar{K})/\Gamma(\Lambda \bar{K})$						Γ_2/Γ_1
<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2.3	90	0	BIAGI	87C SPEC	Ξ^- Be 116 GeV	
$\Gamma(\Sigma \bar{K})/\Gamma_{\text{total}}$						Γ_2/Γ
<u>VALUE</u>		<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
possibly seen		17	HASSALL	81 HBC	$K^- p$ 6.5 GeV/c	
$\Gamma(\Xi \pi)/\Gamma(\Xi(1530) \pi)$						Γ_3/Γ_4
<u>VALUE</u>			<u>DOCUMENT ID</u>	<u>TECN</u>		
$2.8^{+0.7}_{-0.6}$			APSELL	70 HBC		
$\Gamma(\Xi \pi \pi \text{ (not } \Xi(1530) \pi))/\Gamma(\Xi(1530) \pi)$						Γ_5/Γ_4
<u>VALUE</u>			<u>DOCUMENT ID</u>	<u>TECN</u>		
0.0 ± 0.3			APSELL	70 HBC		

$\Xi(1950)$ REFERENCES

ADAMOVICH	99B	EPJ C11 271	M.I. Adamovich <i>et al.</i>	(CERN WA89 Collab.)
BIAGI	87	ZPHY C34 15	S.F. Biagi <i>et al.</i>	(BRIS, CERN, GEVA+)
BIAGI	87C	ZPHY C34 175	S.F. Biagi <i>et al.</i>	(BRIS, CERN, GEVA+)
BIAGI	81	ZPHY C9 305	S.F. Biagi <i>et al.</i>	(BRIS, CAVE, GEVA+)
HASSALL	81	NP B189 397	J.K. Hassall <i>et al.</i>	(CAVE, MSU)
BRIEFEL	77	PR D16 2706	E. Briefel <i>et al.</i>	(BRAN, UMD, SYRA+)
	Also	70 Duke Conf. 317	E. Briefel <i>et al.</i>	(BRAN, UMD, SYRA+)
Hyperon Resonances, 1970				
DIBIANCA	75	NP B98 137	F.A. Dibianca, R.J. Endorf	(CMU)
ROSS	73C	Purdue Conf. 345	R.T. Ross, J.L. Lloyd, D. Radojicic	(OXF)
BADIER	72	NP B37 429	J. Badier <i>et al.</i>	(EPOL)
APSELL	70	PRL 24 777	S.P. Apsell <i>et al.</i>	(BRAN, UMD, SYRA+)
GOLDWASSER	70	PR D1 1960	E.L. Goldwasser, P.F. Schultz	(ILL)
DAUBER	69	PR 179 1262	P.M. Dauber <i>et al.</i>	(LRL)
ALITTI	68	PRL 21 1119	J. Alitti <i>et al.</i>	(BNL, SYRA)
BADIER	65	PL 16 171	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)