

$B^\pm$ 

$$I(J^P) = \frac{1}{2}(0^-)$$

Quantum numbers not measured. Values shown are quark-model predictions.

See also the  $B^\pm/B^0$  ADMIXTURE and  $B^\pm/B_s^0/B_s^0/b$ -baryon ADMIXTURE sections.

## $B^\pm$ MASS

The fit uses  $m_{B^+}$ ,  $(m_{B^0} - m_{B^+})$ , and  $m_{B^0}$  to determine  $m_{B^+}$ ,  $m_{B^0}$ , and the mass difference.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5279.0±0.5 OUR FIT</b>				
<b>5279.1±0.5 OUR AVERAGE</b>				
5279.1±0.4 ±0.4	526	1 CSORNA	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
5279.1±1.7 ±1.4	147	ABE	96B CDF	$p\bar{p}$ at 1.8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5278.8±0.54±2.0	362	ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
5278.3±0.4 ±2.0		BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
5280.5±1.0 ±2.0		2 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$
5275.8±1.3 ±3.0	32	ALBRECHT	87C ARG	$e^+ e^- \rightarrow \gamma(4S)$
5278.2±1.8 ±3.0	12	3 ALBRECHT	87D ARG	$e^+ e^- \rightarrow \gamma(4S)$
5278.6±0.8 ±2.0		BEBEK	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

<sup>1</sup> CSORNA 00 uses fully reconstructed  $526 B^+ \rightarrow J/\psi(') K^+$  events and invariant masses without beam constraint.

<sup>2</sup> ALBRECHT 90J assumes 10580 for  $\gamma(4S)$  mass. Supersedes ALBRECHT 87C and ALBRECHT 87D.

<sup>3</sup> Found using fully reconstructed decays with  $J/\psi(1S)$ . ALBRECHT 87D assume  $m\gamma(4S) = 10577$  MeV.

## $B^\pm$ MEAN LIFE

See  $B^\pm/B_s^0/b$ -baryon ADMIXTURE section for data on  $B$ -hadron mean life averaged over species of bottom particles.

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account corrections between the measurements and asymmetric lifetime errors.

VALUE ( $10^{-12}$ s)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.638±0.011 OUR EVALUATION</b>				
1.635±0.011±0.011		4 ABE	05B BELL	$e^+ e^- \rightarrow \gamma(4S)$
1.624±0.014±0.018		5 ABDALLAH	04E DLPH	$e^+ e^- \rightarrow Z$
1.636±0.058±0.025		6 ACOSTA	02C CDF	$p\bar{p}$ at 1.8 TeV
1.673±0.032±0.023		7 AUBERT	01F BABR	$e^+ e^- \rightarrow \gamma(4S)$
1.648±0.049±0.035		8 BARATE	00R ALEP	$e^+ e^- \rightarrow Z$
1.643±0.037±0.025		9 ABBIENDI	99J OPAL	$e^+ e^- \rightarrow Z$

$1.637 \pm 0.058^{+0.045}_{-0.043}$	<sup>8</sup> ABE	98Q CDF	$p\bar{p}$ at 1.8 TeV
$1.66 \pm 0.06 \pm 0.03$	<sup>9</sup> ACCIARRI	98S L3	$e^+ e^- \rightarrow Z$
$1.66 \pm 0.06 \pm 0.05$	<sup>9</sup> ABE	97J SLD	$e^+ e^- \rightarrow Z$
$1.58^{+0.21}_{-0.18} \pm 0.04$	94	<sup>6</sup> BUSKULIC	$e^+ e^- \rightarrow Z$
$1.61 \pm 0.16 \pm 0.12$		<sup>8,10</sup> ABREU	$e^+ e^- \rightarrow Z$
$1.72 \pm 0.08 \pm 0.06$		<sup>11</sup> ADAM	$e^+ e^- \rightarrow Z$
$1.52 \pm 0.14 \pm 0.09$		<sup>8</sup> AKERS	$e^+ e^- \rightarrow Z$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
$1.695 \pm 0.026 \pm 0.015$	<sup>7</sup> ABE	02H BELL	Repl. by ABE 05B
$1.68 \pm 0.07 \pm 0.02$	<sup>6</sup> ABE	98B CDF	Repl. by ACOSTA 02C
$1.56 \pm 0.13 \pm 0.06$	<sup>8</sup> ABE	96C CDF	Repl. by ABE 98Q
$1.58 \pm 0.09 \pm 0.03$		<sup>12</sup> BUSKULIC	$e^+ e^- \rightarrow Z$
$1.58 \pm 0.09 \pm 0.04$		<sup>8</sup> BUSKULIC	96J ALEP Repl. by BARATE 00R
$1.70 \pm 0.09$		<sup>13</sup> ADAM	$e^+ e^- \rightarrow Z$
$1.61 \pm 0.16 \pm 0.05$	148	<sup>6</sup> ABE	94D CDF Repl. by ABE 98B
$1.30^{+0.33}_{-0.29} \pm 0.16$	92	<sup>8</sup> ABREU	93D DLPN Sup. by ABREU 95Q
$1.56 \pm 0.19 \pm 0.13$	134	<sup>11</sup> ABREU	93G DLPN Sup. by ADAM 95
$1.51^{+0.30}_{-0.28} \pm 0.12_{-0.14}$	59	<sup>8</sup> ACTON	93C OPAL Sup. by AKERS 95T
$1.47^{+0.22}_{-0.19} \pm 0.15_{-0.14}$	77	<sup>8</sup> BUSKULIC	93D ALEP Sup. by BUSKULIC 96J

<sup>4</sup> Measurement performed using a combined fit of  $CP$ -violation, mixing and lifetimes.

<sup>5</sup> Measurement performed using an inclusive reconstruction and  $B$  flavor identification technique.

<sup>6</sup> Measured mean life using fully reconstructed decays.

<sup>7</sup> Events are selected in which one  $B$  meson is fully reconstructed while the second  $B$  meson is reconstructed inclusively.

<sup>8</sup> Data analyzed using  $D/D^*\ell X$  event vertices.

<sup>9</sup> Data analyzed using charge of secondary vertex.

<sup>10</sup> ABREU 95Q assumes  $B(B^0 \rightarrow D^{**-} \ell^+ \nu_\ell) = 3.2 \pm 1.7\%$ .

<sup>11</sup> Data analyzed using vertex-charge technique to tag  $B$  charge.

<sup>12</sup> Combined result of  $D/D^*\ell X$  analysis and fully reconstructed  $B$  analysis.

<sup>13</sup> Combined ABREU 95Q and ADAM 95 result.

## $B^+$ DECAY MODES

$B^-$  modes are charge conjugates of the modes below. Modes which do not identify the charge state of the  $B$  are listed in the  $B^\pm/B^0$  ADMIXTURE section.

The branching fractions listed below assume 50%  $B^0\bar{B}^0$  and 50%  $B^+B^-$  production at the  $\Upsilon(4S)$ . We have attempted to bring older measurements up to date by rescaling their assumed  $\Upsilon(4S)$  production ratio to 50:50 and their assumed  $D$ ,  $D_s$ ,  $D^*$ , and  $\psi$  branching ratios to current values whenever this would affect our averages and best limits significantly.

Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.

For inclusive branching fractions, e.g.,  $B \rightarrow D^\pm$  anything, the values usually are multiplicities, not branching fractions. They can be greater than one.

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
<b>Semileptonic and leptonic modes</b>		
$\Gamma_1 \ell^+ \nu_\ell$ anything	[a] ( 10.2 ± 0.9 ) %	
$\Gamma_2 \bar{D}^0 \ell^+ \nu_\ell$	[a] ( 2.15 ± 0.22 ) %	
$\Gamma_3 \bar{D}^*(2007)^0 \ell^+ \nu_\ell$	[a] ( 6.5 ± 0.5 ) %	
$\Gamma_4 \bar{D}_1(2420)^0 \ell^+ \nu_\ell$	( 5.6 ± 1.6 ) × 10 <sup>-3</sup>	
$\Gamma_5 \bar{D}_2^*(2460)^0 \ell^+ \nu_\ell$	< 8 × 10 <sup>-3</sup>	CL=90%
$\Gamma_6 \pi^0 e^+ \nu_e$	( 9.0 ± 2.8 ) × 10 <sup>-5</sup>	
$\Gamma_7 \eta \ell^+ \nu_\ell$	( 8 ± 4 ) × 10 <sup>-5</sup>	
$\Gamma_8 \omega \ell^+ \nu_\ell$	[a] ( 1.3 ± 0.6 ) × 10 <sup>-4</sup>	
$\Gamma_9 \omega \mu^+ \nu_\mu$		
$\Gamma_{10} \rho^0 \ell^+ \nu_\ell$	[a] ( 1.34 <sup>+ 0.32</sup> <sub>- 0.35</sub> ) × 10 <sup>-4</sup>	
$\Gamma_{11} p\bar{p} e^+ \nu_e$	< 5.2 × 10 <sup>-3</sup>	CL=90%
$\Gamma_{12} e^+ \nu_e$	< 1.5 × 10 <sup>-5</sup>	CL=90%
$\Gamma_{13} \mu^+ \nu_\mu$	< 6.6 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{14} \tau^+ \nu_\tau$	< 5.7 × 10 <sup>-4</sup>	CL=90%
$\Gamma_{15} e^+ \nu_e \gamma$	< 2.0 × 10 <sup>-4</sup>	CL=90%
$\Gamma_{16} \mu^+ \nu_\mu \gamma$	< 5.2 × 10 <sup>-5</sup>	CL=90%
<b>Inclusive modes</b>		
$\Gamma_{17} D^0 X$	( 9.8 ± 1.1 ) %	
$\Gamma_{18} \bar{D}^0 X$	( 79 ± 5 ) %	
$\Gamma_{19} D^+ X$	( 3.8 ± 1.0 ) %	
$\Gamma_{20} D^- X$	( 9.8 ± 1.8 ) %	
$\Gamma_{21} D_s^+ X$	( 14 <sup>+ 5</sup> <sub>- 4</sub> ) %	
$\Gamma_{22} D_s^- X$	< 2.2 %	CL=90%
$\Gamma_{23} \Lambda_c^+ X$	( 2.9 <sup>+ 1.4</sup> <sub>- 1.1</sub> ) %	
$\Gamma_{24} \bar{\Lambda}_c^- X$	( 3.5 <sup>+ 1.5</sup> <sub>- 1.2</sub> ) %	
$\Gamma_{25} \bar{c} X$	( 98 ± 6 ) %	
$\Gamma_{26} c X$	( 33 <sup>+ 6</sup> <sub>- 4</sub> ) %	
$\Gamma_{27} \bar{c} c X$	( 131 <sup>+ 10</sup> <sub>- 8</sub> ) %	

**$D$ ,  $D^*$ , or  $D_s$  modes**

$\Gamma_{28}$	$\overline{D}^0 \pi^+$	$( -4.91 \pm 0.21 ) \times 10^{-3}$
$\Gamma_{29}$	$D_{CP(+1)} \pi^+$	[b]
$\Gamma_{30}$	$D_{CP(-1)} \pi^+$	[b]
$\Gamma_{31}$	$\overline{D}^0 \rho^+$	$( -1.34 \pm 0.18 ) \%$
$\Gamma_{32}$	$\overline{D}^0 K^+$	$( -3.7 \pm 0.6 ) \times 10^{-4}$ S=1.1
$\Gamma_{33}$	$D_{CP(+1)} K^+$	[b]
$\Gamma_{34}$	$D_{CP(-1)} K^+$	[b]
$\Gamma_{35}$	$[K^- \pi^+]_D K^+ + [K^+ \pi^-]_D K^-$	[c]
$\Gamma_{36}$	$[K^+ \pi^-]_D K^+ + [K^- \pi^+]_D K^-$	[c]
$\Gamma_{37}$	$\overline{D}^0 K^*(892)^+$	$( -6.3 \pm 0.8 ) \times 10^{-4}$
$\Gamma_{38}$	$\overline{D}^0 K^+ \overline{K}^0$	$( -5.5 \pm 1.6 ) \times 10^{-4}$
$\Gamma_{39}$	$\overline{D}^0 K^+ \overline{K}^*(892)^0$	$( -7.5 \pm 1.7 ) \times 10^{-4}$
$\Gamma_{40}$	$\overline{D}^0 \pi^+ \pi^+ \pi^-$	$( -1.1 \pm 0.4 ) \%$
$\Gamma_{41}$	$\overline{D}^0 \pi^+ \pi^+ \pi^-$ nonresonant	$( -5 \pm 4 ) \times 10^{-3}$
$\Gamma_{42}$	$\overline{D}^0 \pi^+ \rho^0$	$( -4.2 \pm 3.0 ) \times 10^{-3}$
$\Gamma_{43}$	$\overline{D}^0 a_1(1260)^+$	$( -5 \pm 4 ) \times 10^{-3}$
$\Gamma_{44}$	$\overline{D}^0 \omega \pi^+$	$( -4.1 \pm 0.9 ) \times 10^{-3}$
$\Gamma_{45}$	$D^*(2010)^- \pi^+ \pi^+$	$( -1.35 \pm 0.22 ) \times 10^{-3}$
$\Gamma_{46}$	$D^- \pi^+ \pi^+$	$( -1.02 \pm 0.16 ) \times 10^{-3}$
$\Gamma_{47}$	$\overline{D}_2^*(2462)^0 \pi^+$ $\times B(\overline{D}_2^*(2462)^0 \rightarrow D^- \pi^+)$	$( -3.4 \pm 0.8 ) \times 10^{-4}$
$\Gamma_{48}$	$\overline{D}_0^*(2308)^0 \pi^+$ $\times B(\overline{D}_0^*(2308)^0 \rightarrow D^- \pi^+)$	$( -6.1 \pm 1.9 ) \times 10^{-4}$
$\Gamma_{49}$	$\overline{D}_1(2421)^0 \pi^+$ $\times B(\overline{D}_1(2421)^0 \rightarrow D^{*-} \pi^+)$	$( -6.8 \pm 1.5 ) \times 10^{-4}$
$\Gamma_{50}$	$\overline{D}_2^*(2462)^0 \pi^+$ $\times B(\overline{D}_2^*(2462)^0 \rightarrow D^{*-} \pi^+)$	$( -1.8 \pm 0.5 ) \times 10^{-4}$
$\Gamma_{51}$	$\overline{D}'_1(2427)^0 \pi^+$ $\times B(\overline{D}'_1(2427)^0 \rightarrow D^{*-} \pi^+)$	$( -5.0 \pm 1.2 ) \times 10^{-4}$
$\Gamma_{52}$	$\overline{D}^*(2007)^0 \pi^+$	$( -4.6 \pm 0.4 ) \times 10^{-3}$
$\Gamma_{53}$	$\overline{D}^*(2007)^0 \omega \pi^+$	$( -4.5 \pm 1.2 ) \times 10^{-3}$
$\Gamma_{54}$	$\overline{D}^*(2007)^0 \rho^+$	$( -9.8 \pm 1.7 ) \times 10^{-3}$
$\Gamma_{55}$	$\overline{D}^*(2007)^0 K^+$	$( -3.6 \pm 1.0 ) \times 10^{-4}$
$\Gamma_{56}$	$\overline{D}^*(2007)^0 K^* \overline{(892)}^+$	$( -8.1 \pm 1.4 ) \times 10^{-4}$
$\Gamma_{57}$	$\overline{D}^*(2007)^0 K^+ \overline{K}^0$	$< 1.06 \times 10^{-3}$ CL=90%
$\Gamma_{58}$	$\overline{D}^*(2007)^0 K^+ K^*(892)^0$	$( -1.5 \pm 0.4 ) \times 10^{-3}$
$\Gamma_{59}$	$\overline{D}^*(2007)^0 \pi^+ \pi^+ \pi^-$	$( -1.03 \pm 0.12 ) \%$
$\Gamma_{60}$	$\overline{D}^*(2007)^0 a_1(1260)^+$	$( -1.9 \pm 0.5 ) \%$
$\Gamma_{61}$	$\overline{D}^*(2007)^0 \pi^- \pi^+ \pi^+ \pi^0$	$( -1.8 \pm 0.4 ) \%$

$\Gamma_{62}$	$\overline{D}^{*0} 3\pi^+ 2\pi^-$	$( 5.7 \pm 1.2 ) \times 10^{-3}$	
$\Gamma_{63}$	$D^*(2010)^+ \pi^0$	$< 1.7 \times 10^{-4}$	CL=90%
$\Gamma_{64}$	$\overline{D}^*(2010)^+ K^0$	$< 9.5 \times 10^{-5}$	CL=90%
$\Gamma_{65}$	$D^*(2010)^- \pi^+ \pi^+ \pi^0$	$( 1.5 \pm 0.7 ) \%$	
$\Gamma_{66}$	$D^*(2010)^- \pi^+ \pi^+ \pi^+ \pi^-$	$( 2.6 \pm 0.4 ) \times 10^{-3}$	
$\Gamma_{67}$	$\overline{D}_1^*(2420)^0 \pi^+$	$( 1.5 \pm 0.6 ) \times 10^{-3}$	S=1.3
$\Gamma_{68}$	$\overline{D}_1^*(2420)^0 \rho^+$	$< 1.4 \times 10^{-3}$	CL=90%
$\Gamma_{69}$	$\overline{D}_2^*(2460)^0 \pi^+$	$< 1.3 \times 10^{-3}$	CL=90%
$\Gamma_{70}$	$\overline{D}_2^*(2460)^0 \rho^+$	$< 4.7 \times 10^{-3}$	CL=90%
$\Gamma_{71}$	$\overline{D}^0 D_s^+$	$( 1.3 \pm 0.4 ) \%$	
$\Gamma_{72}$	$D_{sJ}(2317)^+ \overline{D}^0 \times$ $B(D_{sJ}(2317)^+ \rightarrow D_s^+ \pi^0)$	$( 9.0 \pm 3.3 ) \times 10^{-4}$	
$\Gamma_{73}$	$D_{sJ}(2317)^+ \overline{D}^0 \times$ $B(D_{sJ}(2317)^+ \rightarrow D_s^{*+} \gamma)$	$< 7.6 \times 10^{-4}$	CL=90%
$\Gamma_{74}$	$D_{sJ}(2317)^+ \overline{D}^*(2010)^0 \times$ $B(D_{sJ}(2317)^+ \rightarrow D_s^+ \pi^0)$	$( 9 \pm 7 ) \times 10^{-4}$	
$\Gamma_{75}$	$D_{sJ}(2457)^+ \overline{D}^0 \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0)$	$( 1.7 \pm 0.7 ) \times 10^{-3}$	S=1.1
$\Gamma_{76}$	$D_{sJ}(2457)^+ \overline{D}^0 \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)$	$( 5.8 \pm 1.9 ) \times 10^{-4}$	
$\Gamma_{77}$	$D_{sJ}(2457)^+ \overline{D}^0 \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^+ \pi^-)$	$< 2.2 \times 10^{-4}$	CL=90%
$\Gamma_{78}$	$D_{sJ}(2457)^+ \overline{D}^0 \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^0)$	$< 2.7 \times 10^{-4}$	CL=90%
$\Gamma_{79}$	$D_{sJ}(2457)^+ \overline{D}^0 \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \gamma)$	$< 9.8 \times 10^{-4}$	CL=90%
$\Gamma_{80}$	$D_{sJ}(2457)^+ \overline{D}^*(2010)^0 \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0)$	$( 7.6 \pm 3.6 ) \times 10^{-3}$	
$\Gamma_{81}$	$D_{sJ}(2457)^+ \overline{D}^*(2010)^0 \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)$	$( 1.4 \pm 0.7 ) \times 10^{-3}$	
$\Gamma_{82}$	$\overline{D}^0 D_{sJ}(2536)^+ \times$ $B(D_{sJ}(2536)^+ \rightarrow D^*(2007)^0 K^+)$	$< 2 \times 10^{-4}$	CL=90%
$\Gamma_{83}$	$\overline{D}^*(2007)^0 D_{sJ}(2536)^+ \times$ $B(D_{sJ}(2536)^+ \rightarrow D^*(2007)^0 K^+)$	$< 7 \times 10^{-4}$	CL=90%
$\Gamma_{84}$	$\overline{D}^0 D_{sJ}(2573)^+ \times$ $B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)$	$< 2 \times 10^{-4}$	CL=90%
$\Gamma_{85}$	$\overline{D}^*(2007)^0 D_{sJ}(2573)^+ \times$ $B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)$	$< 5 \times 10^{-4}$	CL=90%

$\Gamma_{86}$	$\overline{D}^0 D_s^{*+}$	$(9 \pm 4) \times 10^{-3}$	
$\Gamma_{87}$	$\overline{D}^*(2007)^0 D_s^+$	$(1.2 \pm 0.5) \%$	
$\Gamma_{88}$	$\overline{D}^*(2007)^0 D_s^{*+}$	$(2.7 \pm 1.0) \%$	
$\Gamma_{89}$	$D_s^{(*)+} \overline{D}^{**0}$	$(2.7 \pm 1.2) \%$	
$\Gamma_{90}$	$\overline{D}^*(2007)^0 D^*(2010)^+$	$< 1.1 \%$	CL=90%
$\Gamma_{91}$	$\overline{D}^0 D^*(2010)^+ +$ $\overline{D}^*(2007)^0 D^+$	$< 1.3 \%$	CL=90%
$\Gamma_{92}$	$\overline{D}^0 D^+$	$< 6.7 \times 10^{-3}$	CL=90%
$\Gamma_{93}$	$\overline{D}^0 D^+ K^0$	$< 2.8 \times 10^{-3}$	CL=90%
$\Gamma_{94}$	$\overline{D}^*(2007)^0 D^+ K^0$	$< 6.1 \times 10^{-3}$	CL=90%
$\Gamma_{95}$	$\overline{D}^0 \overline{D}^*(2010)^+ K^0$	$(5.2 \pm 1.2) \times 10^{-3}$	
$\Gamma_{96}$	$\overline{D}^*(2007)^0 D^*(2010)^+ K^0$	$(7.8 \pm 2.6) \times 10^{-3}$	
$\Gamma_{97}$	$\overline{D}^0 D^0 K^+$	$(1.37 \pm 0.32) \times 10^{-3}$	S=1.5
$\Gamma_{98}$	$\overline{D}^*(2010)^0 D^0 K^+$	$< 3.8 \times 10^{-3}$	CL=90%
$\Gamma_{99}$	$\overline{D}^0 D^*(2007)^0 K^+$	$(4.7 \pm 1.0) \times 10^{-3}$	
$\Gamma_{100}$	$\overline{D}^*(2007)^0 D^*(2007)^0 K^+$	$(5.3 \pm 1.6) \times 10^{-3}$	
$\Gamma_{101}$	$D^- D^+ K^+$	$< 4 \times 10^{-4}$	CL=90%
$\Gamma_{102}$	$D^- D^*(2010)^+ K^+$	$< 7 \times 10^{-4}$	CL=90%
$\Gamma_{103}$	$D^*(2010)^- D^+ K^+$	$(1.5 \pm 0.4) \times 10^{-3}$	
$\Gamma_{104}$	$D^*(2010)^- D^*(2010)^+ K^+$	$< 1.8 \times 10^{-3}$	CL=90%
$\Gamma_{105}$	$(\overline{D} + \overline{D}^*)(D + D^*) K$	$(3.5 \pm 0.6) \%$	
$\Gamma_{106}$	$D_s^+ \pi^0$	$< 2.0 \times 10^{-4}$	CL=90%
$\Gamma_{107}$	$D_s^{*+} \pi^0$	$< 3.3 \times 10^{-4}$	CL=90%
$\Gamma_{108}$	$D_s^+ \eta$	$< 5 \times 10^{-4}$	CL=90%
$\Gamma_{109}$	$D_s^{*+} \eta$	$< 8 \times 10^{-4}$	CL=90%
$\Gamma_{110}$	$D_s^+ \rho^0$	$< 4 \times 10^{-4}$	CL=90%
$\Gamma_{111}$	$D_s^{*+} \rho^0$	$< 5 \times 10^{-4}$	CL=90%
$\Gamma_{112}$	$D_s^+ \omega$	$< 5 \times 10^{-4}$	CL=90%
$\Gamma_{113}$	$D_s^{*+} \omega$	$< 7 \times 10^{-4}$	CL=90%
$\Gamma_{114}$	$D_s^+ a_1(1260)^0$	$< 2.2 \times 10^{-3}$	CL=90%
$\Gamma_{115}$	$D_s^{*+} a_1(1260)^0$	$< 1.6 \times 10^{-3}$	CL=90%
$\Gamma_{116}$	$D_s^+ \phi$	$< 3.2 \times 10^{-4}$	CL=90%
$\Gamma_{117}$	$D_s^{*+} \phi$	$< 4 \times 10^{-4}$	CL=90%
$\Gamma_{118}$	$D_s^+ \overline{K}^0$	$< 1.1 \times 10^{-3}$	CL=90%
$\Gamma_{119}$	$D_s^{*+} \overline{K}^0$	$< 1.1 \times 10^{-3}$	CL=90%
$\Gamma_{120}$	$D_s^+ \overline{K}^*(892)^0$	$< 5 \times 10^{-4}$	CL=90%
$\Gamma_{121}$	$D_s^{*+} \overline{K}^*(892)^0$	$< 4 \times 10^{-4}$	CL=90%
$\Gamma_{122}$	$D_s^- \pi^+ K^+$	$< 8 \times 10^{-4}$	CL=90%
$\Gamma_{123}$	$D_s^{*-} \pi^+ K^+$	$< 1.2 \times 10^{-3}$	CL=90%
$\Gamma_{124}$	$D_s^- \pi^+ K^*(892)^+$	$< 6 \times 10^{-3}$	CL=90%
$\Gamma_{125}$	$D_s^{*-} \pi^+ K^*(892)^+$	$< 8 \times 10^{-3}$	CL=90%

**Charmonium modes**

$\Gamma_{126}$	$\eta_c K^+$	$(1.03 \pm 0.21) \times 10^{-3}$	
$\Gamma_{127}$	$J/\psi(1S) K^+$	$(9.9 \pm 0.4) \times 10^{-4}$	
$\Gamma_{128}$	$J/\psi(1S) K^+ \pi^+ \pi^-$	$(7.7 \pm 2.0) \times 10^{-4}$	
$\Gamma_{129}$	$X(3872) K^+$	seen	
$\Gamma_{130}$	$X(3872) K^+$ $\times B(X(3872) \rightarrow D^0 \bar{D}^0)$	$< 6.0 \times 10^{-5}$	CL=90%
$\Gamma_{131}$	$X(3872) K^+$ $\times B(X(3872) \rightarrow D^+ D^-)$	$< 4.0 \times 10^{-5}$	CL=90%
$\Gamma_{132}$	$X(3872) K^+$ $\times B(X(3872) \rightarrow D^0 \bar{D}^0 \pi^0)$	$< 6.0 \times 10^{-5}$	CL=90%
$\Gamma_{133}$	$X(3872) K^+$ $\times B(X(3872) \rightarrow J/\psi(1S) \eta)$	$< 7.7 \times 10^{-6}$	CL=90%
$\Gamma_{134}$	$J/\psi(1S) K^*(892)^+$	$(1.35 \pm 0.10) \times 10^{-3}$	
$\Gamma_{135}$	$J/\psi(1S) K(1270)^+$	$(1.8 \pm 0.5) \times 10^{-3}$	
$\Gamma_{136}$	$J/\psi(1S) K(1400)^+$	$< 5 \times 10^{-4}$	CL=90%
$\Gamma_{137}$	$J/\psi(1S) \eta K^+$	$(1.08 \pm 0.33) \times 10^{-4}$	
$\Gamma_{138}$	$J/\psi(1S) \phi K^+$	$(5.2 \pm 1.7) \times 10^{-5}$	S=1.2
$\Gamma_{139}$	$J/\psi(1S) \pi^+$	$(4.8 \pm 0.6) \times 10^{-5}$	S=1.5
$\Gamma_{140}$	$J/\psi(1S) \rho^+$	$< 7.7 \times 10^{-4}$	CL=90%
$\Gamma_{141}$	$J/\psi(1S) a_1(1260)^+$	$< 1.2 \times 10^{-3}$	CL=90%
$\Gamma_{142}$	$J/\psi(1S) p \bar{\Lambda}$	$(1.2 \pm 0.9) \times 10^{-5}$	
$\Gamma_{143}$	$\psi(2S) K^+$	$(6.8 \pm 0.4) \times 10^{-4}$	
$\Gamma_{144}$	$\psi(2S) K^*(892)^+$	$(9.2 \pm 2.2) \times 10^{-4}$	
$\Gamma_{145}$	$\psi(2S) K^+ \pi^+ \pi^-$	$(1.9 \pm 1.2) \times 10^{-3}$	
$\Gamma_{146}$	$\psi(3770) K^+$	$(4.8 \pm 1.3) \times 10^{-4}$	
$\Gamma_{147}$	$\psi(3770) K^+$ $\times B(\psi(3770) \rightarrow D^0 \bar{D}^0)$	$(3.4 \pm 0.9) \times 10^{-4}$	
$\Gamma_{148}$	$\psi(3770) K^+$ $\times B(\psi(3770) \rightarrow D^+ D^- K^+)$	$(1.4 \pm 0.8) \times 10^{-4}$	
$\Gamma_{149}$	$\chi_{c0}(1P) K^+$	$(3.0 \pm 0.6) \times 10^{-4}$	S=1.1
$\Gamma_{150}$	$\chi_{c1}(1P) K^+$	$(6.8 \pm 1.1) \times 10^{-4}$	
$\Gamma_{151}$	$\chi_{c1}(1P) K^*(892)^+$	$< 2.1 \times 10^{-3}$	CL=90%

**K or  $K^*$  modes**

$\Gamma_{152}$	$K^0 \pi^+$	$(2.18 \pm 0.14) \times 10^{-5}$	
$\Gamma_{153}$	$K^+ \pi^0$	$(1.25 \pm 0.10) \times 10^{-5}$	
$\Gamma_{154}$	$\eta' K^+$	$(7.8 \pm 0.5) \times 10^{-5}$	
$\Gamma_{155}$	$\eta' K^*(892)^+$	$< 1.4 \times 10^{-5}$	CL=90%
$\Gamma_{156}$	$\eta K^+$	$(3.3 \pm 0.8) \times 10^{-6}$	
$\Gamma_{157}$	$\eta K^*(892)^+$	$(2.6 \pm 0.4) \times 10^{-5}$	
$\Gamma_{158}$	$\omega K^+$	$(5.1 \pm 0.7) \times 10^{-6}$	
$\Gamma_{159}$	$\omega K^*(892)^+$	$< 8.7 \times 10^{-5}$	CL=90%
$\Gamma_{160}$	$a_0^+ K^0$	$< 3.9 \times 10^{-6}$	CL=90%

$\Gamma_{161}$	$a_0^0 K^+$	$< 2.5 \times 10^{-6}$	CL=90%
$\Gamma_{162}$	$K^*(892)^0 \pi^+$	$(1.60^{+0.22}_{-0.39}) \times 10^{-5}$	
$\Gamma_{163}$	$K^*(892)^+ \pi^0$	$< 3.1 \times 10^{-5}$	CL=90%
$\Gamma_{164}$	$K^+ \pi^- \pi^+$	$(5.7 \pm 0.4) \times 10^{-5}$	
$\Gamma_{165}$	$K^+ \pi^- \pi^+ \text{nonresonant}$	$< 1.7 \times 10^{-5}$	CL=90%
$\Gamma_{166}$	$K^+ f_0(980) \times B(f_0 \rightarrow \pi^+ \pi^-)$	$(9.3 \pm 2.1) \times 10^{-6}$	
$\Gamma_{167}$	$K^+ \rho^0$	$< 6.2 \times 10^{-6}$	CL=90%
$\Gamma_{168}$	$K_2^*(1430)^0 \pi^+$	$< 6.8 \times 10^{-4}$	CL=90%
$\Gamma_{169}$	$K^- \pi^+ \pi^+$	$< 1.8 \times 10^{-6}$	CL=90%
$\Gamma_{170}$	$K^- \pi^+ \pi^+ \text{nonresonant}$	$< 5.6 \times 10^{-5}$	CL=90%
$\Gamma_{171}$	$K_1(1400)^0 \pi^+$	$< 2.6 \times 10^{-3}$	CL=90%
$\Gamma_{172}$	$K^0 \pi^+ \pi^0$	$< 6.6 \times 10^{-5}$	CL=90%
$\Gamma_{173}$	$K^0 \rho^+$	$< 4.8 \times 10^{-5}$	CL=90%
$\Gamma_{174}$	$K^*(892)^+ \pi^+ \pi^-$	$< 1.1 \times 10^{-3}$	CL=90%
$\Gamma_{175}$	$K^*(892)^+ \rho^0$	$(1.1 \pm 0.4) \times 10^{-5}$	
$\Gamma_{176}$	$K^*(892)^+ K^*(892)^0$	$< 7.1 \times 10^{-5}$	CL=90%
$\Gamma_{177}$	$K_1(1400)^+ \rho^0$	$< 7.8 \times 10^{-4}$	CL=90%
$\Gamma_{178}$	$K_2^*(1430)^+ \rho^0$	$< 1.5 \times 10^{-3}$	CL=90%
$\Gamma_{179}$	$K^+ \bar{K}^0$	$< 2.5 \times 10^{-6}$	CL=90%
$\Gamma_{180}$	$\bar{K}^0 K^+ \pi^0$	$< 2.4 \times 10^{-5}$	CL=90%
$\Gamma_{181}$	$K^+ K_S^0 K_S^0$	$(1.15 \pm 0.13) \times 10^{-5}$	
$\Gamma_{182}$	$K_S^0 K_S^0 \pi^+$	$< 3.2 \times 10^{-6}$	CL=90%
$\Gamma_{183}$	$K^+ K^- \pi^+$	$< 6.3 \times 10^{-6}$	CL=90%
$\Gamma_{184}$	$K^+ K^- \pi^+ \text{nonresonant}$	$< 7.5 \times 10^{-5}$	CL=90%
$\Gamma_{185}$	$K^+ K^+ \pi^-$	$< 1.3 \times 10^{-6}$	CL=90%
$\Gamma_{186}$	$K^+ K^+ \pi^- \text{nonresonant}$	$< 8.79 \times 10^{-5}$	CL=90%
$\Gamma_{187}$	$K^+ K^*(892)^0$	$< 5.3 \times 10^{-6}$	CL=90%
$\Gamma_{188}$	$K^+ f_J(2220)$		
$\Gamma_{189}$	$K^+ K^- K^+$	$(3.08 \pm 0.21) \times 10^{-5}$	
$\Gamma_{190}$	$K^+ \phi$	$(9.3 \pm 1.0) \times 10^{-6}$	S=1.3
$\Gamma_{191}$	$K^+ K^- K^+ \text{nonresonant}$	$< 3.8 \times 10^{-5}$	CL=90%
$\Gamma_{192}$	$K^*(892)^+ K^+ K^-$	$< 1.6 \times 10^{-3}$	CL=90%
$\Gamma_{193}$	$K^*(892)^+ \phi$	$(9.6 \pm 3.0) \times 10^{-6}$	S=1.9
$\Gamma_{194}$	$K_1(1400)^+ \phi$	$< 1.1 \times 10^{-3}$	CL=90%
$\Gamma_{195}$	$K_2^*(1430)^+ \phi$	$< 3.4 \times 10^{-3}$	CL=90%
$\Gamma_{196}$	$K^+ \phi \phi$	$(2.6^{+1.1}_{-0.9}) \times 10^{-6}$	
$\Gamma_{197}$	$K^*(892)^+ \gamma$	$(4.03 \pm 0.26) \times 10^{-5}$	
$\Gamma_{198}$	$K_1(1270)^+ \gamma$	$< 9.9 \times 10^{-5}$	CL=90%
$\Gamma_{199}$	$\phi K^+ \gamma$	$(3.4 \pm 1.0) \times 10^{-6}$	
$\Gamma_{200}$	$K^+ \pi^- \pi^+ \gamma$	$(2.4^{+0.6}_{-0.5}) \times 10^{-5}$	
$\Gamma_{201}$	$K^*(892)^0 \pi^+ \gamma$	$(2.0^{+0.7}_{-0.6}) \times 10^{-5}$	

$\Gamma_{202}$	$K^+ \rho^0 \gamma$	<	2.0	$\times 10^{-5}$	CL=90%
$\Gamma_{203}$	$K^+ \pi^- \pi^+ \gamma$ nonresonant	<	9.2	$\times 10^{-6}$	CL=90%
$\Gamma_{204}$	$K_1(1400)^+ \gamma$	<	5.0	$\times 10^{-5}$	CL=90%
$\Gamma_{205}$	$K_2^*(1430)^+ \gamma$	(	1.4 $\pm$ 0.4	) $\times 10^{-5}$	
$\Gamma_{206}$	$K^*(1680)^+ \gamma$	<	1.9	$\times 10^{-3}$	CL=90%
$\Gamma_{207}$	$K_3^*(1780)^+ \gamma$	<	5.5	$\times 10^{-3}$	CL=90%
$\Gamma_{208}$	$K_4^*(2045)^+ \gamma$	<	9.9	$\times 10^{-3}$	CL=90%

### Light unflavored meson modes

$\Gamma_{209}$	$\rho^+ \gamma$	<	1.8	$\times 10^{-6}$	CL=90%
$\Gamma_{210}$	$\pi^+ \pi^0$	(	5.1 $\pm$ 0.8 0.9	) $\times 10^{-6}$	
$\Gamma_{211}$	$\pi^+ \pi^+ \pi^-$	(	1.1 $\pm$ 0.4	) $\times 10^{-5}$	
$\Gamma_{212}$	$\rho^0 \pi^+$	(	9.2 $\pm$ 1.1	) $\times 10^{-6}$	
$\Gamma_{213}$	$\pi^+ f_0(980)$	<	1.4	$\times 10^{-4}$	CL=90%
$\Gamma_{214}$	$\pi^+ f_2(1270)$	<	2.4	$\times 10^{-4}$	CL=90%
$\Gamma_{215}$	$\pi^+ \pi^- \pi^+$ nonresonant	<	4.1	$\times 10^{-5}$	CL=90%
$\Gamma_{216}$	$\pi^+ \pi^0 \pi^0$	<	8.9	$\times 10^{-4}$	CL=90%
$\Gamma_{217}$	$\rho^+ \pi^0$	(	1.09 $\pm$ 0.27	) $\times 10^{-5}$	
$\Gamma_{218}$	$\pi^+ \pi^- \pi^+ \pi^0$	<	4.0	$\times 10^{-3}$	CL=90%
$\Gamma_{219}$	$\rho^+ \rho^0$	(	2.6 $\pm$ 0.6	) $\times 10^{-5}$	
$\Gamma_{220}$	$a_1(1260)^+ \pi^0$	<	1.7	$\times 10^{-3}$	CL=90%
$\Gamma_{221}$	$a_1(1260)^0 \pi^+$	<	9.0	$\times 10^{-4}$	CL=90%
$\Gamma_{222}$	$\omega \pi^+$	(	5.9 $\pm$ 1.0	) $\times 10^{-6}$	S=1.2
$\Gamma_{223}$	$\omega \rho^+$	<	6.1	$\times 10^{-5}$	CL=90%
$\Gamma_{224}$	$\eta \pi^+$	(	4.8 $\pm$ 1.3 1.1	) $\times 10^{-6}$	S=1.4
$\Gamma_{225}$	$\eta' \pi^+$	<	4.5	$\times 10^{-6}$	CL=90%
$\Gamma_{226}$	$\eta' \rho^+$	<	2.2	$\times 10^{-5}$	CL=90%
$\Gamma_{227}$	$\eta \rho^+$	<	1.4	$\times 10^{-5}$	CL=90%
$\Gamma_{228}$	$\phi \pi^+$	<	4.1	$\times 10^{-7}$	CL=90%
$\Gamma_{229}$	$\phi \rho^+$	<	1.6	$\times 10^{-5}$	
$\Gamma_{230}$	$a_0^0 \pi^+$	<	5.8	$\times 10^{-6}$	CL=90%
$\Gamma_{231}$	$\pi^+ \pi^+ \pi^+ \pi^- \pi^-$	<	8.6	$\times 10^{-4}$	CL=90%
$\Gamma_{232}$	$\rho^0 a_1(1260)^+$	<	6.2	$\times 10^{-4}$	CL=90%
$\Gamma_{233}$	$\rho^0 a_2(1320)^+$	<	7.2	$\times 10^{-4}$	CL=90%
$\Gamma_{234}$	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$	<	6.3	$\times 10^{-3}$	CL=90%
$\Gamma_{235}$	$a_1(1260)^+ a_1(1260)^0$	<	1.3	%	CL=90%

### Charged particle ( $h^\pm$ ) modes

$$h^\pm = K^\pm \text{ or } \pi^\pm$$

$\Gamma_{236}$	$h^+ \pi^0$	(	1.6 $\pm$ 0.7 0.6	) $\times 10^{-5}$	
$\Gamma_{237}$	$\omega h^+$	(	1.38 $\pm$ 0.27 0.24	) $\times 10^{-5}$	
$\Gamma_{238}$	$h^+ X^0$ (Familon)	<	4.9	$\times 10^{-5}$	CL=90%

### Baryon modes

$\Gamma_{239}$	$p\bar{p}\pi^+$	$( -3.1 \pm 0.8 ) \times 10^{-6}$		
$\Gamma_{240}$	$p\bar{p}\pi^+$ nonresonant	$< 5.3 \times 10^{-5}$	CL=90%	
$\Gamma_{241}$	$p\bar{p}\pi^+\pi^+\pi^-$	$< 5.2 \times 10^{-4}$	CL=90%	
$\Gamma_{242}$	$p\bar{p}K^+$	$( -5.7 \pm 0.9 ) \times 10^{-6}$		
$\Gamma_{243}$	$p\bar{p}K^+$ nonresonant	$< 8.9 \times 10^{-5}$	CL=90%	
$\Gamma_{244}$	$p\bar{p}K^*(892)^+$	$( 1.03^{+0.38}_{-0.33} ) \times 10^{-5}$		
$\Gamma_{245}$	$p\bar{\Lambda}$	$< 1.5 \times 10^{-6}$	CL=90%	
$\Gamma_{246}$	$p\bar{\Lambda}\gamma$			
$\Gamma_{247}$	$p\bar{\Sigma}\gamma$			
$\Gamma_{248}$	$p\bar{\Lambda}\pi^+\pi^-$	$< 2.0 \times 10^{-4}$	CL=90%	
$\Gamma_{249}$	$\Lambda\bar{\Lambda}\pi^+$	$( 2.9^{+1.0}_{-0.8} ) \times 10^{-6}$		
$\Gamma_{250}$	$\Lambda\bar{\Lambda}K^+$	$< 2.8 \times 10^{-6}$	CL=90%	
$\Gamma_{251}$	$\bar{\Delta}^0 p$	$< 3.8 \times 10^{-4}$	CL=90%	
$\Gamma_{252}$	$\Delta^{++}\bar{p}$	$< 1.5 \times 10^{-4}$	CL=90%	
$\Gamma_{253}$	$D^+ p\bar{p}$	$< 1.5 \times 10^{-5}$	CL=90%	
$\Gamma_{254}$	$D^*(2010)^+ p\bar{p}$	$< 1.5 \times 10^{-5}$	CL=90%	
$\Gamma_{255}$	$\bar{\Lambda}_c^- p\pi^+$	$( 2.1 \pm 0.7 ) \times 10^{-4}$		
$\Gamma_{256}$	$\bar{\Lambda}_c^- p\pi^+\pi^0$	$( 1.8 \pm 0.6 ) \times 10^{-3}$		
$\Gamma_{257}$	$\bar{\Lambda}_c^- p\pi^+\pi^+\pi^-$	$( 2.3 \pm 0.7 ) \times 10^{-3}$		
$\Gamma_{258}$	$\bar{\Lambda}_c^- p\pi^+\pi^+\pi^-\pi^0$	$< 1.34 \%$	CL=90%	
$\Gamma_{259}$	$\bar{\Sigma}_c(2455)^0 p$	$< 8 \times 10^{-5}$	CL=90%	
$\Gamma_{260}$	$\bar{\Sigma}_c(2520)^0 p$	$< 4.6 \times 10^{-5}$	CL=90%	
$\Gamma_{261}$	$\bar{\Sigma}_c(2455)^0 p\pi^0$	$( 4.4 \pm 1.8 ) \times 10^{-4}$		
$\Gamma_{262}$	$\bar{\Sigma}_c(2455)^0 p\pi^-\pi^+$	$( 4.4 \pm 1.7 ) \times 10^{-4}$		
$\Gamma_{263}$	$\bar{\Sigma}_c(2455)^{--} p\pi^+\pi^+$	$( 2.8 \pm 1.2 ) \times 10^{-4}$		
$\Gamma_{264}$	$\bar{\Lambda}_c(2593)^-/\bar{\Lambda}_c(2625)^- p\pi^+$	$< 1.9 \times 10^{-4}$	CL=90%	

### Lepton Family number ( $LF$ ) or Lepton number ( $L$ ) violating modes, or $\Delta B = 1$ weak neutral current ( $B1$ ) modes

$\Gamma_{265}$	$\pi^+ e^+ e^-$	$B1$	$< 3.9 \times 10^{-3}$	CL=90%
$\Gamma_{266}$	$\pi^+ \mu^+ \mu^-$	$B1$	$< 9.1 \times 10^{-3}$	CL=90%
$\Gamma_{267}$	$K^+ e^+ e^-$	$B1$	$( 8.0^{+2.2}_{-1.9} ) \times 10^{-7}$	S=1.4
$\Gamma_{268}$	$K^+ \mu^+ \mu^-$	$B1$	$( 3.4^{+1.9}_{-1.4} ) \times 10^{-7}$	S=1.7
$\Gamma_{269}$	$K^+ \ell^+ \ell^-$	$B1$	$[a] ( 5.3 \pm 1.1 ) \times 10^{-7}$	
$\Gamma_{270}$	$K^+ \bar{\nu}_\nu$	$B1$	$< 2.4 \times 10^{-4}$	CL=90%
$\Gamma_{271}$	$K^*(892)^+ e^+ e^-$	$B1$	$< 4.6 \times 10^{-6}$	CL=90%
$\Gamma_{272}$	$K^*(892)^+ \mu^+ \mu^-$	$B1$	$< 2.2 \times 10^{-6}$	CL=90%
$\Gamma_{273}$	$K^*(892)^+ \ell^+ \ell^-$	$B1$	$[a] < 2.2 \times 10^{-6}$	CL=90%
$\Gamma_{274}$	$\pi^+ e^+ \mu^-$	$LF$	$< 6.4 \times 10^{-3}$	CL=90%
$\Gamma_{275}$	$\pi^+ e^- \mu^+$	$LF$	$< 6.4 \times 10^{-3}$	CL=90%

$\Gamma_{276}$	$K^+ e^+ \mu^-$	$LF$	<	8	$\times 10^{-7}$	CL=90%
$\Gamma_{277}$	$K^+ e^- \mu^+$	$LF$	<	6.4	$\times 10^{-3}$	CL=90%
$\Gamma_{278}$	$K^*(892)^+ e^\pm \mu^\mp$	$LF$	<	7.9	$\times 10^{-6}$	CL=90%
$\Gamma_{279}$	$\pi^- e^+ e^+$	$L$	<	1.6	$\times 10^{-6}$	CL=90%
$\Gamma_{280}$	$\pi^- \mu^+ \mu^+$	$L$	<	1.4	$\times 10^{-6}$	CL=90%
$\Gamma_{281}$	$\pi^- e^+ \mu^+$	$L$	<	1.3	$\times 10^{-6}$	CL=90%
$\Gamma_{282}$	$\rho^- e^+ e^+$	$L$	<	2.6	$\times 10^{-6}$	CL=90%
$\Gamma_{283}$	$\rho^- \mu^+ \mu^+$	$L$	<	5.0	$\times 10^{-6}$	CL=90%
$\Gamma_{284}$	$\rho^- e^+ \mu^+$	$LF$	<	3.3	$\times 10^{-6}$	CL=90%
$\Gamma_{285}$	$K^- e^+ e^+$	$L$	<	1.0	$\times 10^{-6}$	CL=90%
$\Gamma_{286}$	$K^- \mu^+ \mu^+$	$L$	<	1.8	$\times 10^{-6}$	CL=90%
$\Gamma_{287}$	$K^- e^+ \mu^+$	$L$	<	2.0	$\times 10^{-6}$	CL=90%
$\Gamma_{288}$	$K^*(892)^- e^+ e^+$	$L$	<	2.8	$\times 10^{-6}$	CL=90%
$\Gamma_{289}$	$K^*(892)^- \mu^+ \mu^+$	$L$	<	8.3	$\times 10^{-6}$	CL=90%
$\Gamma_{290}$	$K^*(892)^- e^+ \mu^+$	$LF$	<	4.4	$\times 10^{-6}$	CL=90%

[a] An  $\ell$  indicates an  $e$  or a  $\mu$  mode, not a sum over these modes.

[b] An  $CP(\pm 1)$  indicates the  $CP=+1$  and  $CP=-1$  eigenstates of the  $D^0$ - $\overline{D}^0$  system.

[c]  $D$  denotes  $D^0$  or  $\overline{D}^0$ .

## CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 9 measurements and one constraint to determine 3 parameters. The overall fit has a  $\chi^2 = 4.1$  for 7 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$$\begin{array}{c|cc} x_{139} & & 30 \\ \hline & x_{127} & \end{array}$$

## $B^+$ BRANCHING RATIOS

$\Gamma(\ell^+ \nu_\ell \text{anything}) / \Gamma_{\text{total}}$	$\Gamma_1 / \Gamma$
VALUE	DOCUMENT ID

**0.1025 ± 0.0057 ± 0.0065** 14 ARTUSO 97 CLE2  $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.101 ± 0.018 ± 0.015 ATHANAS 94 CLE2 Sup. by ARTUSO 97

<sup>14</sup> ARTUSO 97 uses partial reconstruction of  $B \rightarrow D^* \ell \nu_\ell$  and inclusive semileptonic branching ratio from BARISH 96B ( $0.1049 \pm 0.0017 \pm 0.0043$ ).

$\Gamma(\overline{D}^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$ 

$\ell = e$  or  $\mu$ , not sum over  $e$  and  $\mu$  modes.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0215 ± 0.0022 OUR AVERAGE</b>			
0.0221 ± 0.0013 ± 0.0019	15 BARTEL	99 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.016 ± 0.006 ± 0.003	16 FULTON	91 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0194 ± 0.0015 ± 0.0034	17 ATHANAS	97 CLE2	Repl. by BARTEL 99
15 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ .			
16 FULTON 91 assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\gamma(4S)$ .			
17 ATHANAS 97 uses missing energy and missing momentum to reconstruct neutrino.			

 $\Gamma(\overline{D}^*(2007)^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$ 

$\ell = e$  or  $\mu$ , not sum over  $e$  and  $\mu$  modes.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.065 ± 0.005 OUR AVERAGE</b>				
0.0650 ± 0.0020 ± 0.0043	18 ADAM	03 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
0.066 ± 0.016 ± 0.015	19 ALBRECHT	92C ARG	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0650 ± 0.0020 ± 0.0043	20 BRIERE	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
0.0513 ± 0.0054 ± 0.0064	302 BARISH	95 CLE2	Repl. by ADAM 03	
seen	398	22 SANGHERA	93 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.041 ± 0.008 + 0.008 - 0.009	23 FULTON	91 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
0.070 ± 0.018 ± 0.014	24 ANTREASYAN	90B CBAL	$e^+ e^- \rightarrow \gamma(4S)$	
18 Simultaneous measurements of both $B^0 \rightarrow D^*(2010)^- \ell \nu$ and $B^+ \rightarrow \overline{D}(2007)^0 \ell \nu$ .				
19 ALBRECHT 92C reports $0.058 \pm 0.014 \pm 0.013$ . We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^- \pi^+)$ . Assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\gamma(4S)$ .				
20 The results are based on the same analysis and data sample reported in ADAM 03.				
21 BARISH 95 use $B(D^0 \rightarrow K^- \pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*0} \rightarrow D^0 \pi^0) = (63.6 \pm 2.3 \pm 3.3)\%$ .				
22 Combining $\overline{D}^{*0} \ell^+ \nu_\ell$ and $\overline{D}^{*-} \ell^+ \nu_\ell$ SANGHERA 93 test $V-A$ structure and fit the decay angular distributions to obtain $A_{FB} = 3/4 * (\Gamma^- - \Gamma^+)/\Gamma = 0.14 \pm 0.06 \pm 0.03$ . Assuming a value of $V_{cb}$ , they measure $V$ , $A_1$ , and $A_2$ , the three form factors for the $D^* \ell \nu_\ell$ decay, where results are slightly dependent on model assumptions.				
23 Assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\gamma(4S)$ . Uncorrected for $D$ and $D^*$ branching ratio assumptions.				
24 ANTREASYAN 90B is average over $B$ and $\overline{D}^*(2010)$ charge states.				

 $\Gamma(\overline{D}_1(2420)^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0056 ± 0.0013 ± 0.0009</b>			
25 ANASTASSOV	98	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
25 ANASTASSOV 98 result is derived from the measurement of $B(B^+ \rightarrow \overline{D}_1^0 \ell^+ \nu_\ell) \times B(\overline{D}_1^0 \rightarrow D^{*+} \pi^-) = (0.373 \pm 0.085 \pm 0.052 \pm 0.024)\%$ by assuming $B(\overline{D}_1^0 \rightarrow D^{*+} \pi^-) = 67\%$ , where the third error includes theoretical uncertainties.			

$\Gamma(\overline{D}_2^*(2460)^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$	$\Gamma_5/\Gamma$
$\text{VALUE}$ $\overline{\text{CL}\%}$	$\text{DOCUMENT ID}$ $\text{TECN}$ $\text{COMMENT}$
$<8 \times 10^{-3}$ 90	26 ANASTASSOV 98 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

26 ANASTASSOV 98 result is derived from the measurement of  $B(B^+ \rightarrow \overline{D}_2^{*0} \ell^+ \nu_\ell) \times B(\overline{D}_2^{*0} \rightarrow D^{*+} \pi^-) < 0.16\%$  at 90% CL by assuming  $B(\overline{D}_2^{*0} \rightarrow D^{*+} \pi^-) = 20\%$ .

$\Gamma(\pi^0 e^+ \nu_e)/\Gamma_{\text{total}}$	$\Gamma_6/\Gamma$
$\text{VALUE (units } 10^{-4})$ $\text{CL\%}$	$\text{DOCUMENT ID}$ $\text{TECN}$ $\text{COMMENT}$
$0.9 \pm 0.2 \pm 0.2$	27 ALEXANDER 96T CLE2 $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$<22$ 90	ANTREASYAN 90B CBAL $e^+ e^- \rightarrow \gamma(4S)$
27 Derived based in the reported $B^0$ result by assuming isospin symmetry: $\Gamma(B^0 \rightarrow \pi^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \pi^0 \ell^+ \nu)$ .	

$\Gamma(\eta \ell^+ \nu_\ell)/\Gamma_{\text{total}}$	$\Gamma_7/\Gamma$
$\text{VALUE (units } 10^{-4})$	$\text{DOCUMENT ID}$ $\text{TECN}$ $\text{COMMENT}$
$0.84 \pm 0.31 \pm 0.18$	28 ATHAR 03 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

28 ATHAR 03 reports systematic errors  $0.16 \pm 0.09$ , which are experimental systematic and systematic due to model dependence. We combine these in quadrature.

$\Gamma(\omega \ell^+ \nu_\ell)/\Gamma_{\text{total}}$	$\Gamma_8/\Gamma$
$\ell = e \text{ or } \mu, \text{ not sum over } e \text{ and } \mu \text{ modes.}$	
$\text{VALUE (units } 10^{-4})$ $\text{CL\%}$	$\text{DOCUMENT ID}$ $\text{TECN}$ $\text{COMMENT}$
$1.3 \pm 0.4 \pm 0.4$	29 SCHWANDA 04 BELL $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$<2.1$ 90	30 BEAN 93B CLE2 $e^+ e^- \rightarrow \gamma(4S)$
29 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ .	
30 BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\rho^0 \ell^+ \nu_\ell)$ and $\Gamma(\rho^- \ell^+ \nu_\ell)$ with this result, they obtain a limit $<(1.6-2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow \omega \ell^+ \nu_\ell$ . The range corresponds to the ISGW, WSB, and KS models. An upper limit on $ V_{ub}/V_{cb}  < 0.8-0.13$ at 90% CL is derived as well.	

$\Gamma(\omega \mu^+ \nu_\mu)/\Gamma_{\text{total}}$	$\Gamma_9/\Gamma$
$\text{VALUE}$	$\text{DOCUMENT ID}$ $\text{TECN}$
• • • We do not use the following data for averages, fits, limits, etc. • • •	

seen 31 ALBRECHT 91C ARG

31 In ALBRECHT 91C, one event is fully reconstructed providing evidence for the  $b \rightarrow u$  transition.

$\Gamma(\rho^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$	$\Gamma_{10}/\Gamma$
$\ell = e \text{ or } \mu, \text{ not sum over } e \text{ and } \mu \text{ modes.}$	
$\text{VALUE (units } 10^{-4})$ $\text{CL\%}$	$\text{DOCUMENT ID}$ $\text{TECN}$ $\text{COMMENT}$
$1.34 \pm 0.15 \pm 0.28$	32 BEHRENS 00 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.40 \pm 0.21$	$+0.32$	$-0.33$	<sup>32</sup> BEHRENS	00	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$1.2 \pm 0.2$	$+0.3$	$-0.4$	<sup>32</sup> ALEXANDER	96T	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$<2.1$	90		<sup>33</sup> BEAN	93B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
						<sup>32</sup> Derived based in the reported $B^0$ result by assuming isospin symmetry: $\Gamma(B^0 \rightarrow \rho^- \ell^+ \nu_\ell) = 2\Gamma(B^+ \rightarrow \rho^0 \ell^+ \nu_\ell) \approx 2\Gamma(B^+ \rightarrow \omega \ell^+ \nu_\ell)$ .
						<sup>33</sup> BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\omega^0 \ell^+ \nu_\ell)$ and $\Gamma(\rho^- \ell^+ \nu_\ell)$ with this result, they obtain a limit $<(1.6\text{--}2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow \rho^0 \ell^+ \nu_\ell$ . The range corresponds to the ISGW, WSB, and KS models. An upper limit on $ V_{ub}/V_{cb}  < 0.8\text{--}0.13$ at 90% CL is derived as well.

### $\Gamma(p\bar{p}e^+\nu_e)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{11}/\Gamma$
$<5.2 \times 10^{-3}$	90	<sup>34</sup> ADAM	03B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

<sup>34</sup> Based on phase-space model; if V-A model is used, the 90% CL upper limit becomes  $< 1.2 \times 10^{-3}$ .

### $\Gamma(e^+\nu_e)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{12}/\Gamma$
$<1.5 \times 10^{-5}$	90	ARTUSO	95	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

### $\Gamma(\mu^+\nu_\mu)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{13}/\Gamma$
$<6.6 \times 10^{-6}$	90	AUBERT	040	BABR	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.1 \times 10^{-5}$	90	ARTUSO	95	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
-----------------------	----	--------	----	------	----------------------------------

### $\Gamma(\tau^+\nu_\tau)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{14}/\Gamma$
$<5.7 \times 10^{-4}$	90	<sup>35</sup> ACCIARRI	97F	L3	$e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<8.3 \times 10^{-4}$	90	<sup>36</sup> BARATE	01E	ALEP	$e^+ e^- \rightarrow Z$
$<8.4 \times 10^{-4}$	90	<sup>37</sup> BROWDER	01	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$<1.04 \times 10^{-2}$	90	<sup>38</sup> ALBRECHT	95D	ARG	$e^+ e^- \rightarrow \gamma(4S)$
$<2.2 \times 10^{-3}$	90	ARTUSO	95	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$<1.8 \times 10^{-3}$	90	<sup>39</sup> BUSKULIC	95	ALEP	$e^+ e^- \rightarrow Z$

<sup>35</sup> ACCIARRI 97F uses missing-energy technique and  $f(b \rightarrow B^-) = (38.2 \pm 2.5)\%$ .

<sup>36</sup> The energy-flow and  $b$ -tagging algorithms were used.

<sup>37</sup> Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

<sup>38</sup> ALBRECHT 95D use full reconstruction of one  $B$  decay as tag.

<sup>39</sup> BUSKULIC 95 uses same missing-energy technique as in  $\bar{B} \rightarrow \tau^+\nu_\tau X$ , but analysis is restricted to endpoint region of missing-energy distribution.

$\Gamma(e^+\nu_e\gamma)/\Gamma_{\text{total}}$				$\Gamma_{15}/\Gamma$
VALUE	CL %	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-4}$	90	40	BROWDER 97	CLE2 $e^+e^- \rightarrow \gamma(4S)$

40 BROWDER 97 uses the hermiticity of the CLEO II detector to reconstruct the neutrino energy and momentum.

$\Gamma(\mu^+\nu_\mu\gamma)/\Gamma_{\text{total}}$				$\Gamma_{16}/\Gamma$
VALUE	CL %	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-5}$	90	41	BROWDER 97	CLE2 $e^+e^- \rightarrow \gamma(4S)$

41 BROWDER 97 uses the hermiticity of the CLEO II detector to reconstruct the neutrino energy and momentum.

$\Gamma(D^0 X)/\Gamma_{\text{total}}$				$\Gamma_{17}/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.098 \pm 0.009 \pm 0.006$	42 AUBERT,BE	04B BABR	$e^+e^- \rightarrow \gamma(4S)$	

42 Events are selected by completely reconstructing one  $B$  and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\bar{D}^0 X)/\Gamma_{\text{total}}$				$\Gamma_{18}/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.793 \pm 0.025^{+0.045}_{-0.044}$	43 AUBERT,BE	04B BABR	$e^+e^- \rightarrow \gamma(4S)$	

43 Events are selected by completely reconstructing one  $B$  and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D^0 X)/[\Gamma(D^0 X) + \Gamma(\bar{D}^0 X)]$				$\Gamma_{17}/(\Gamma_{17}+\Gamma_{18})$
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.110 \pm 0.010 \pm 0.003$	AUBERT,BE	04B BABR	$e^+e^- \rightarrow \gamma(4S)$	

$\Gamma(D^+ X)/\Gamma_{\text{total}}$				$\Gamma_{19}/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.038 \pm 0.009 \pm 0.005$	44 AUBERT,BE	04B BABR	$e^+e^- \rightarrow \gamma(4S)$	

44 Events are selected by completely reconstructing one  $B$  and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D^- X)/\Gamma_{\text{total}}$				$\Gamma_{20}/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.098 \pm 0.012 \pm 0.014$	45 AUBERT,BE	04B BABR	$e^+e^- \rightarrow \gamma(4S)$	

45 Events are selected by completely reconstructing one  $B$  and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(D^+ X)/[\Gamma(D^+ X) + \Gamma(D^- X)]$				$\Gamma_{19}/(\Gamma_{19}+\Gamma_{20})$
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.278 \pm 0.052 \pm 0.009$	AUBERT,BE	04B BABR	$e^+e^- \rightarrow \gamma(4S)$	

$\Gamma(D_s^+ X)/\Gamma_{\text{total}}$ VALUE**0.143±0.016<sup>+0.051</sup><sub>-0.034</sub>**DOCUMENT ID46 AUBERT,BE 04B BABR  $e^+ e^- \rightarrow \gamma(4S)$  $\Gamma_{21}/\Gamma$ 

|

46 Events are selected by completely reconstructing one  $B$  and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(D_s^- X)/\Gamma_{\text{total}}$ VALUE**<0.022**CL%

90

DOCUMENT ID47 AUBERT,BE 04B BABR  $e^+ e^- \rightarrow \gamma(4S)$  $\Gamma_{22}/\Gamma$ 

|

47 Events are selected by completely reconstructing one  $B$  and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(D_s^+ X)/[\Gamma(D_s^+ X) + \Gamma(D_s^- X)]$ VALUE**0.966±0.039<sup>±0.012</sup>**DOCUMENT IDAUBERT,BE 04B BABR  $e^+ e^- \rightarrow \gamma(4S)$  $\Gamma_{21}/(\Gamma_{21}+\Gamma_{22})$ 

|

 $\Gamma(D_s^- X)/[\Gamma(D_s^+ X) + \Gamma(D_s^- X)]$ VALUE**<0.126**CL%

90

DOCUMENT IDAUBERT,BE 04B BABR  $e^+ e^- \rightarrow \gamma(4S)$  $\Gamma_{22}/(\Gamma_{21}+\Gamma_{22})$ 

|

 $\Gamma(\Lambda_c^+ X)/\Gamma_{\text{total}}$ VALUE**0.029±0.008<sup>+0.011</sup><sub>-0.007</sub>**DOCUMENT ID48 AUBERT,BE 04B BABR  $e^+ e^- \rightarrow \gamma(4S)$  $\Gamma_{23}/\Gamma$ 

|

48 Events are selected by completely reconstructing one  $B$  and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(\bar{\Lambda}_c^- X)/\Gamma_{\text{total}}$ VALUE**0.035±0.008<sup>+0.013</sup><sub>-0.009</sub>**DOCUMENT ID49 AUBERT,BE 04B BABR  $e^+ e^- \rightarrow \gamma(4S)$  $\Gamma_{24}/\Gamma$ 

|

49 Events are selected by completely reconstructing one  $B$  and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(\Lambda_c^+ X)/[\Gamma(\Lambda_c^+ X) + \Gamma(\bar{\Lambda}_c^- X)]$ VALUE**0.452±0.090<sup>±0.003</sup>**DOCUMENT IDAUBERT,BE 04B BABR  $e^+ e^- \rightarrow \gamma(4S)$  $\Gamma_{23}/(\Gamma_{23}+\Gamma_{24})$ 

|

 $\Gamma(\bar{c} X)/\Gamma_{\text{total}}$ VALUE**0.983±0.030<sup>+0.054</sup><sub>-0.051</sub>**DOCUMENT ID50 AUBERT,BE 04B BABR  $e^+ e^- \rightarrow \gamma(4S)$  $\Gamma_{25}/\Gamma$ 

|

50 Events are selected by completely reconstructing one  $B$  and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(cX)/\Gamma_{\text{total}}$	$\Gamma_{26}/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
$0.330 \pm 0.022^{+0.055}_{-0.037}$	51 AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$

51 Events are selected by completely reconstructing one  $B$  and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\bar{c}cX)/\Gamma_{\text{total}}$	$\Gamma_{27}/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
$1.313 \pm 0.037^{+0.088}_{-0.075}$	52 AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$

52 Events are selected by completely reconstructing one  $B$  and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\bar{D}^0\pi^+)/\Gamma_{\text{total}}$	$\Gamma_{28}/\Gamma$			
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT

#### **4.91 $\pm$ 0.21 OUR AVERAGE**

4.84 $\pm$ 0.27 $\pm$ 0.11	53 AUBERT,B	04P BABR	$e^+ e^- \rightarrow \gamma(4S)$
4.97 $\pm$ 0.12 $\pm$ 0.29	54,55 AHMED	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
5.0 $\pm$ 0.7 $\pm$ 0.6	54 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
5.4 $\pm$ 1.8 $\pm$ 1.2	14 BEBEK	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.5 $\pm$ 0.4 $\pm$ 0.5	304 ALAM	94 CLE2	Repl. by AHMED 02B
2.0 $\pm$ 0.8 $\pm$ 0.6	12 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$
1.9 $\pm$ 1.0 $\pm$ 0.6	7 ALBRECHT	88K ARG	$e^+ e^- \rightarrow \gamma(4S)$

53 AUBERT,B 04P reports  $[B(B^+ \rightarrow \bar{D}^0\pi^+) \times B(D^0 \rightarrow K^-\pi^+)] = (1.846 \pm 0.032 \pm 0.097) \times 10^{-4}$ . We divide by our best value  $B(D^0 \rightarrow K^-\pi^+) = (3.81 \pm 0.09) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

54 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

55 AHMED 02B reports an additional uncertainty on the branching ratios to account for 4.5% uncertainty on relative production of  $B^0$  and  $B^+$ , which is not included here.

56 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$  and uses the Mark III branching fractions for the  $D$ .

57 BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

58 ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$  and use the CLEO II absolute  $B(D^0 \rightarrow K^-\pi^+)$  and the PDG 1992  $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$  and  $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$ .

59 ALBRECHT 88K assumes  $B^0\bar{B}^0:B^+B^-$  ratio is 45:55. Superseded by ALBRECHT 90J.

$\Gamma(\bar{D}^0\rho^+)/\Gamma_{\text{total}}$	$\Gamma_{31}/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0134 <math>\pm</math> 0.0018 OUR AVERAGE</b>				
0.0135 $\pm$ 0.0012 $\pm$ 0.0015	212 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
0.013 $\pm$ 0.004 $\pm$ 0.004	19 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.021 $\pm$ 0.008 $\pm$ 0.009	10 ALBRECHT	88K ARG	$e^+ e^- \rightarrow \gamma(4S)$	

<sup>60</sup> ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the CLEO II absolute  $B(D^0 \rightarrow K^- \pi^+)$  and the PDG 1992  $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$  and  $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$ .

<sup>61</sup> Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and uses the Mark III branching fractions for the  $D$ .

<sup>62</sup> ALBRECHT 88K assumes  $B^0 \bar{B}^0 : B^+ B^-$  ratio is 45:55.

### $\Gamma(\bar{D}^0 K^+)/\Gamma_{\text{total}}$

$\Gamma_{32}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.7 ± 0.6 OUR AVERAGE</b>	Error includes scale factor of 1.1.		
4.19 ± 0.57 ± 0.40	63 ABE 01I	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
2.92 ± 0.80 ± 0.28	64 ATHANAS 98	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<sup>63</sup> ABE 01I reports $B(B^+ \rightarrow \bar{D}^0 K^+)/B(B^+ \rightarrow \bar{D}^0 \pi^+) = 0.079 \pm 0.009 \pm 0.006$ . We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 \pi^+) = (5.3 \pm 0.5) \times 10^{-3}$ . Our first error is their experiment's error and the second error is systematic error from using our best value.			
<sup>64</sup> ATHANAS 98 reports $[B(B^+ \rightarrow \bar{D}^0 K^+)]/[B(B^+ \rightarrow \bar{D}^0 \pi^+)] = 0.055 \pm 0.014 \pm 0.005$ . We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 \pi^+) = (5.3 \pm 0.5) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.			

### $\Gamma(\bar{D}^0 K^+)/\Gamma(\bar{D}^0 \pi^+)$

$\Gamma_{32}/\Gamma_{28}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0830 ± 0.0035 OUR AVERAGE</b>			
0.0831 ± 0.0035 ± 0.002	AUBERT 04N	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.099 <sup>+0.014</sup> <sub>-0.012</sub> <sup>+0.007</sup> <sub>-0.006</sub>	BORNHEIM 03	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.077 ± 0.005 ± 0.006	65 SWAIN 03	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.094 ± 0.009 ± 0.007	65 ABE 03D	BELL	Repl. by SWAIN 03
65 Flavor specific $D^0$ meson is reconstructed via $D^0 \rightarrow K^- \pi^+$ .			

### $\Gamma(D_{CP(+1)} K^+)/\Gamma(D_{CP(+1)} \pi^+)$

$\Gamma_{33}/\Gamma_{29}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.090 ± 0.013 OUR AVERAGE</b>			
0.088 ± 0.016 ± 0.005	66 AUBERT 04N	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.093 ± 0.018 ± 0.008	66 SWAIN 03	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.125 ± 0.036 ± 0.010	66 ABE 03D	BELL	Repl. by SWAIN 03
66 $CP=+1$ eigenstate of $D^0 \bar{D}^0$ system is reconstructed via $K^+ K^-$ and $\pi^+ \pi^-$ .			

### $\Gamma(D_{CP(-1)} K^+)/\Gamma(D_{CP(-1)} \pi^+)$

$\Gamma_{34}/\Gamma_{30}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.108 ± 0.019 ± 0.007</b>	67 SWAIN 03	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.119 ± 0.028 ± 0.006	67 ABE 03D	BELL	Repl. by SWAIN 03
67 $CP=-1$ eigenstate of $D^0 \bar{D}^0$ system is reconstructed via $K_S^0 \pi^0$ , $K_S^0 \omega$ , $K_S^0 \phi$ , $K_S^0 \eta$ , and $K_S^0 \eta'$ .			

$$\frac{\Gamma([K^-\pi^+]_D K^+ + [K^+\pi^-]_D K^-)/\Gamma([K^+\pi^-]_D K^+ + [K^-\pi^+]_D K^-)}{\Gamma_{35}/\Gamma_{36}}$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<0.026	90	68 AUBERT,B	04L BABR

68 AUBERT,B 04L extract a constraint on the magnitude of the ratio of amplitudes  $|A(B^+ \rightarrow D^0 K^+)/A(B^+ \rightarrow \bar{D}^0 K^+)| < 0.22$  at 90% CL.

$$\frac{\Gamma(\bar{D}^0 K^*(892)^+)/\Gamma_{\text{total}}}{\Gamma_{37}/\Gamma}$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>6.3±0.8 OUR AVERAGE</b>			

6.3±0.7±0.5	69 AUBERT	04Q BABR	
6.1±1.6±1.7	69 MAHAPATRA	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

69 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$$\frac{\Gamma(\bar{D}^0 K^+ \bar{K}^0)/\Gamma_{\text{total}}}{\Gamma_{38}/\Gamma}$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.5±1.4±0.8</b>	70 DRUTSKOY	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$

70 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$$\frac{\Gamma(\bar{D}^0 K^+ \bar{K}^*(892)^0)/\Gamma_{\text{total}}}{\Gamma_{39}/\Gamma}$$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>7.5±1.3±1.1</b>	71 DRUTSKOY	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$

71 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$$\frac{\Gamma(\bar{D}^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}}{\Gamma_{40}/\Gamma}$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0115±0.0029±0.0021</b>	72 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

72 BORTOLETTO 92 assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$  and uses Mark III branching fractions for the  $D$ .

$$\frac{\Gamma(\bar{D}^0 \pi^+ \pi^+ \pi^- \text{ nonresonant})/\Gamma_{\text{total}}}{\Gamma_{41}/\Gamma}$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0051±0.0034±0.0023</b>	73 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

73 BORTOLETTO 92 assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$  and uses Mark III branching fractions for the  $D$ .

$$\frac{\Gamma(\bar{D}^0 \pi^+ \rho^0)/\Gamma_{\text{total}}}{\Gamma_{42}/\Gamma}$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0042±0.0023±0.0020</b>	74 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

74 BORTOLETTO 92 assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$  and uses Mark III branching fractions for the  $D$ .

$$\frac{\Gamma(\bar{D}^0 a_1(1260)^+)/\Gamma_{\text{total}}}{\Gamma_{43}/\Gamma}$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0045±0.0019±0.0031</b>	75 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

75 BORTOLETTO 92 assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$  and uses Mark III branching fractions for the  $D$ .

$\Gamma(\overline{D}^0 \omega \pi^+)/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{44}/\Gamma$
<b>0.0041 ± 0.0007 ± 0.0006</b>	76 ALEXANDER	01B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
76 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ . The signal is consistent with all observed $\omega \pi^+$ having proceeded through the $\rho'^+$ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.				

 $\Gamma(D^*(2010)^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{45}/\Gamma$
<b>0.00135 ± 0.00022 OUR AVERAGE</b>						
0.00125 ± 0.00008 ± 0.00022			77 ABE	04D BELL	$e^+ e^- \rightarrow \gamma(4S)$	
0.0019 ± 0.0007 ± 0.0003	14		78 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
0.0026 ± 0.0014 ± 0.0007	11		79 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$	
0.0024 +0.0017 -0.0016 +0.0010 -0.0006	3		80 BEBEK	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.004	90		81 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
0.005 ± 0.002 ± 0.003	7		82 ALBRECHT	87C ARG	$e^+ e^- \rightarrow \gamma(4S)$	

77 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .78 ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$  and use the CLEO II  $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$  and absolute  $B(D^0 \rightarrow K^- \pi^+)$  and the PDG 1992  $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$  and  $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$ .79 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$  and uses the Mark III branching fractions for the  $D$ .

80 BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

81 BORTOLETTO 92 assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$  and uses Mark III branching fractions for the  $D$  and  $D^*(2010)$ . The authors also find the product branching fraction into  $D^{**} \pi$  followed by  $D^{**} \rightarrow D^*(2010) \pi$  to be  $0.0014^{+0.0008}_{-0.0006} \pm 0.0003$  where  $D^{**}$  represents all orbitally excited  $D$  mesons.82 ALBRECHT 87C use PDG 86 branching ratios for  $D$  and  $D^*(2010)$  and assume  $B(\gamma(4S) \rightarrow B^+ B^-) = 55\%$  and  $B(\gamma(4S) \rightarrow B^0 \bar{B}^0) = 45\%$ . Superseded by ALBRECHT 90J. $\Gamma(D^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{46}/\Gamma$
<b>1.02 ± 0.04 ± 0.15</b>			83 ABE	04D BELL	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.4	90	84 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
<7	90	85 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
2.5 +4.1 -2.3 +2.4 -0.8	1	86 BEBEK	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

<sup>83</sup> Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

<sup>84</sup> ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the Mark III  $B(D^+ \rightarrow K^-\pi^+\pi^+)$ .

<sup>85</sup> BORTOLETTO 92 assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and uses Mark III branching fractions for the  $D$ . The product branching fraction into  $D_0^*(2340)\pi$  followed by  $D_0^*(2340) \rightarrow D\pi$  is  $< 0.005$  at 90%CL and into  $D_2^*(2460)$  followed by  $D_2^*(2460) \rightarrow D\pi$  is  $< 0.004$  at 90%CL.

<sup>86</sup> BEBEK 87 assume the  $\Upsilon(4S)$  decays 43% to  $B^0\bar{B}^0$ .  $B(D^- \rightarrow K^+\pi^-\pi^-) = (9.1 \pm 1.3 \pm 0.4)\%$  is assumed.

$$\Gamma(\bar{D}_2^*(2462)^0\pi^+ \times B(\bar{D}_2^*(2462)^0 \rightarrow D^-\pi^+))/\Gamma_{\text{total}} \quad \Gamma_{47}/\Gamma$$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.4±0.3±0.72</b>	87 ABE	04D BELL	$e^+e^- \rightarrow \Upsilon(4S)$

<sup>87</sup> Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$$\Gamma(\bar{D}_0^*(2308)^0\pi^+ \times B(\bar{D}_0^*(2308)^0 \rightarrow D^-\pi^+))/\Gamma_{\text{total}} \quad \Gamma_{48}/\Gamma$$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>6.1±0.6±1.8</b>	88 ABE	04D BELL	$e^+e^- \rightarrow \Upsilon(4S)$

<sup>88</sup> Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$$\Gamma(\bar{D}_1(2421)^0\pi^+ \times B(\bar{D}_1(2421)^0 \rightarrow D^{*-}\pi^+))/\Gamma_{\text{total}} \quad \Gamma_{49}/\Gamma$$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>6.8±0.7±1.3</b>	89 ABE	04D BELL	$e^+e^- \rightarrow \Upsilon(4S)$

<sup>89</sup> Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$$\Gamma(\bar{D}_2^*(2462)^0\pi^+ \times B(\bar{D}_2^*(2462)^0 \rightarrow D^{*-}\pi^+))/\Gamma_{\text{total}} \quad \Gamma_{50}/\Gamma$$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.8±0.3±0.4</b>	90 ABE	04D BELL	$e^+e^- \rightarrow \Upsilon(4S)$

<sup>90</sup> Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$$\Gamma(\bar{D}'_1(2427)^0\pi^+ \times B(\bar{D}'_1(2427)^0 \rightarrow D^{*-}\pi^+))/\Gamma_{\text{total}} \quad \Gamma_{51}/\Gamma$$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>5.0±0.4±1.1</b>	91 ABE	04D BELL	$e^+e^- \rightarrow \Upsilon(4S)$

<sup>91</sup> Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$$\Gamma(\bar{D}^*(2007)^0\pi^+)/\Gamma_{\text{total}} \quad \Gamma_{52}/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0046 ± 0.0004 OUR AVERAGE</b>				
0.00434 ± 0.00047 ± 0.00018		92 BRANDENB...	98 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.0052 ± 0.0007 ± 0.0007	71	93 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.0072 ± 0.0018 ± 0.0016		94 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.0040 ± 0.0014 ± 0.0012	9	94 ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
0.0027 ± 0.0044		95 BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

92 BRANDENBURG 98 assume equal production of  $B^+$  and  $B^0$  at  $\Upsilon(4S)$  and use the  $D^*$  reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of  $B(D^* \rightarrow D\pi)$ .

93 ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the CLEO II  $B(D^*(2007)^0 \rightarrow D^0\pi^0)$  and absolute  $B(D^0 \rightarrow K^-\pi^+)$  and the PDG 1992  $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$  and  $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$ .

94 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and uses Mark III branching fractions for the  $D$  and  $D^*(2010)$ .

95 This is a derived branching ratio, using the inclusive pion spectrum and other two-body  $B$  decays. BEBEK 87 assume the  $\Upsilon(4S)$  decays 43% to  $B^0\bar{B}^0$ .

### $\Gamma(\bar{D}^*(2007)^0\omega\pi^+)/\Gamma_{\text{total}}$

### $\Gamma_{53}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0045±0.0010±0.0007</b>	96 ALEXANDER	01B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

96 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . The signal is consistent with all observed  $\omega\pi^+$  having proceeded through the  $\rho^+$  resonance at mass  $1349 \pm 25^{+10}_{-5}$  MeV and width  $547 \pm 86^{+46}_{-45}$  MeV.

### $\Gamma(\bar{D}^*(2007)^0\rho^+)/\Gamma_{\text{total}}$

### $\Gamma_{54}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0098±0.0017 OUR AVERAGE</b>				
0.0098±0.0006±0.0017		97 CSORNA	03 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.010 ± 0.006 ± 0.004	7	98 ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0168±0.0021±0.0028	86	99 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

97 Assumes equal production of  $B^0$  and  $B^+$  at the  $\Upsilon(4S)$  resonance. The second error combines the systematic and theoretical uncertainties in quadrature. CSORNA 03 includes data used in ALAM 94. A full angular fit to three complex helicity amplitudes is performed.

98 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and uses Mark III branching fractions for the  $D$  and  $D^*(2010)$ .

99 ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the CLEO II  $B(D^*(2007)^0 \rightarrow D^0\pi^0)$  and absolute  $B(D^0 \rightarrow K^-\pi^+)$  and the PDG 1992  $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$  and  $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$ . The nonresonant  $\pi^+\pi^0$  contribution under the  $\rho^+$  is negligible.

### $\Gamma(\bar{D}^*(2007)^0K^+)/\Gamma_{\text{total}}$

### $\Gamma_{55}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.59±0.97±0.31</b>	100 ABE	01I BELL	$e^+e^- \rightarrow \Upsilon(4S)$

100 ABE 01I reports  $B(B^+ \rightarrow \bar{D}^*(2007)^0 K^+)/B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+) = 0.078 \pm 0.019 \pm 0.009$ . We multiply by our best value  $B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+) = (4.6 \pm 0.4) \times 10^{-3}$ . Our first error is their experiment's error and the second error is systematic error from using our best value.

$\Gamma(\bar{D}^*(2007)^0 K^*(892)^+)/\Gamma_{\text{total}}$   $\Gamma_{56}/\Gamma$ 

VALUE (units $10^{-4}$ )		DOCUMENT ID	TECN	COMMENT
<b>8.1±1.4 OUR AVERAGE</b>				
8.3±1.1±1.0	101	AUBERT	04K BABR	$e^+ e^- \rightarrow \gamma(4S)$
7.2±2.2±2.6	102	MAHAPATRA	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
101 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ .				
102 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ and an unpolarized final state.				

 $\Gamma(\bar{D}^*(2007)^0 K^+ \bar{K}^0)/\Gamma_{\text{total}}$   $\Gamma_{57}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<10.6	90	103 DRUTSKOY	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$
103 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ .				

 $\Gamma(\bar{D}^*(2007)^0 K^+ K^*(892)^0)/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>15.3±3.1±2.9</b>	104 DRUTSKOY	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$
104 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ .			

 $\Gamma(\bar{D}^*(2007)^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{59}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.03 ±0.12 OUR AVERAGE</b>				
1.055±0.047±0.129	105 MAJUMDER	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$	
0.94 ±0.20 ±0.17	48,106,107 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
105 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ .				
106 ALAM 94 assume equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$ .				
107 The three pion mass is required to be between 1.0 and 1.6 GeV consistent with an $a_1$ meson. (If this channel is dominated by $a_1^+$ , the branching ratio for $\bar{D}^{*0} a_1^+$ is twice that for $\bar{D}^{*0} \pi^+ \pi^+ \pi^-$ .)				

 $\Gamma(\bar{D}^*(2007)^0 a_1(1260)^+)/\Gamma_{\text{total}}$   $\Gamma_{60}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0188±0.0040±0.0034</b>	108,109 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
108 ALAM 94 value is twice their $\Gamma(\bar{D}^*(2007)^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ value based on their observation that the three pions are dominantly in the $a_1(1260)$ mass range 1.0 to 1.6 GeV.			
109 ALAM 94 assume equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$ .			

 $\Gamma(\bar{D}^*(2007)^0 \pi^- \pi^+ \pi^+ \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{61}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0180±0.0024±0.0027</b>	110 ALEXANDER	01B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
110 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ . The signal is consistent with all observed $\omega \pi^+$ having proceeded through the $\rho'^+$ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.			

$\Gamma(\overline{D}^{*0} 3\pi^+ 2\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{62}/\Gamma$ 

<u>VALUE</u> (units $10^{-3}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.67±0.91±0.85</b>	111 MAJUMDER 04	BELL	$e^+ e^- \rightarrow \gamma(4S)$

111 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . $\Gamma(D^*(2010)^+ \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.00017</b>	90	112 BRANDENB... 98	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

112 BRANDENBURG 98 assume equal production of  $B^+$  and  $B^0$  at  $\gamma(4S)$  and use the  $D^*$  partial reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of  $B(D^* \rightarrow D\pi)$ . $\Gamma(\overline{D}^*(2010)^+ K^0)/\Gamma_{\text{total}}$   $\Gamma_{64}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;9.5 × 10<sup>-5</sup></b>	90	113 GRITSAN 01	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

113 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . $\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0152±0.0071±0.0001</b>	26	114 ALBRECHT 90J	ARG	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.043 ± 0.013 ± 0.026      24      115 ALBRECHT 87C ARG       $e^+ e^- \rightarrow \gamma(4S)$ 114 ALBRECHT 90J reports  $0.018 \pm 0.007 \pm 0.005$  for  $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$ . We rescale to our best value  $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$  and uses Mark III branching fractions for the  $D$ .115 ALBRECHT 87C use PDG 86 branching ratios for  $D$  and  $D^*(2010)$  and assume  $B(\gamma(4S) \rightarrow B^+ B^-) = 55\%$  and  $B(\gamma(4S) \rightarrow B^0 \overline{B}^0) = 45\%$ . Superseded by ALBRECHT 90J. $\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$ 

<u>VALUE</u> (units $10^{-3}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.56±0.26±0.33</b>	90	116 MAJUMDER 04	BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10      90      117 ALBRECHT 90J ARG       $e^+ e^- \rightarrow \gamma(4S)$ 116 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .117 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$  and uses Mark III branching fractions for the  $D$  and  $D^*(2010)$ . $\Gamma(\overline{D}_1^*(2420)^0 \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0015±0.0006 OUR AVERAGE</b>		Error includes scale factor of 1.3.		

0.0011±0.0005±0.0002      8      118 ALAM 94 CLE2       $e^+ e^- \rightarrow \gamma(4S)$ 0.0025±0.0007±0.0006      119 ALBRECHT 94D ARG       $e^+ e^- \rightarrow \gamma(4S)$

- 118 ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the CLEO II  $B(D^*(2010)^+ \rightarrow D^0\pi^+)$  and absolute  $B(D^0 \rightarrow K^-\pi^+)$  and the PDG 1992  $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$  and assuming  $B(D_1(2420)^0 \rightarrow D^*(2010)^+\pi^-) = 67\%$ .  
 119 ALBRECHT 94D assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the CLEO II  $B(D^*(2010)^+ \rightarrow D^0\pi^+)$  assuming  $B(D_1(2420)^0 \rightarrow D^*(2010)^+\pi^-) = 67\%$ .

$\Gamma(\bar{D}_1^*(2420)^0\rho^+)/\Gamma_{\text{total}}$	$\Gamma_{68}/\Gamma$			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0014	90	120 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

120 ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the CLEO II  $B(D^*(2010)^+ \rightarrow D^0\pi^+)$  assuming  $B(D_1(2420)^0 \rightarrow D^*(2010)^+\pi^-) = 67\%$ .

$\Gamma(\bar{D}_2^*(2460)^0\pi^+)/\Gamma_{\text{total}}$	$\Gamma_{69}/\Gamma$			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0013	90	121 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.0028	90	122 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<0.0023	90	123 ALBRECHT	94D ARG	$e^+e^- \rightarrow \Upsilon(4S)$

121 ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the Mark III  $B(D^+ \rightarrow K^-\pi^+\pi^+)$  and  $B(D_2^*(2460)^0 \rightarrow D^+\pi^-) = 30\%$ .

122 ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the Mark III  $B(D^+ \rightarrow K^-\pi^+\pi^+)$ , the CLEO II  $B(D^*(2010)^+ \rightarrow D^0\pi^+)$  and  $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+\pi^-) = 20\%$ .

123 ALBRECHT 94D assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the CLEO II  $B(D^*(2010)^+ \rightarrow D^0\pi^+)$  and  $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+\pi^-) = 30\%$ .

$\Gamma(\bar{D}_2^*(2460)^0\rho^+)/\Gamma_{\text{total}}$	$\Gamma_{70}/\Gamma$			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0047	90	124 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<0.005	90	125 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

124 ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the Mark III  $B(D^+ \rightarrow K^-\pi^+\pi^+)$  and  $B(D_2^*(2460)^0 \rightarrow D^+\pi^-) = 30\%$ .

125 ALAM 94 assume equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$  and use the Mark III  $B(D^+ \rightarrow K^-\pi^+\pi^+)$ , the CLEO II  $B(D^*(2010)^+ \rightarrow D^0\pi^+)$  and  $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+\pi^-) = 20\%$ .

$\Gamma(\bar{D}^0 D_s^+)/\Gamma_{\text{total}}$	$\Gamma_{71}/\Gamma$			
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.013 <math>\pm</math> 0.004 OUR AVERAGE</b>				
0.0122 $\pm$ 0.0032 $^{+0.0029}_{-0.0030}$	126 GIBAUT	96 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$	
0.018 $\pm$ 0.009 $\pm$ 0.004	127 ALBRECHT	92G ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
0.016 $\pm$ 0.007 $\pm$ 0.004	5 BORTOLETTO90	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

- 126 GIBAUT 96 reports  $0.0126 \pm 0.0022 \pm 0.0025$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$ . We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- 127 ALBRECHT 92G reports  $0.024 \pm 0.012 \pm 0.004$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990  $D^0$  branching ratios, e.g.,  $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$ .
- 128 BORTOLETTO 90 reports  $0.029 \pm 0.013$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.02$ . We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_{sJ}(2317)^+ \bar{D}^0 \times B(D_{sJ}(2317)^+ \rightarrow D_s^+ \pi^0)) / \Gamma_{\text{total}} \quad \Gamma_{72}/\Gamma$$

<u>VALUE</u> (units $10^{-3}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.90<sup>+0.33</sup><sub>-0.30</sub> OUR AVERAGE</b>			
1.00 <sup>+0.43</sup> <sub>-0.26</sub> $\pm 0.24$	129,130 AUBERT,B	04S BABR	$e^+e^- \rightarrow \gamma(4S)$
0.81 <sup>+0.33</sup> <sub>-0.30</sub> $\pm 0.19$	129,131 KROKOVNY	03B BELL	$e^+e^- \rightarrow \gamma(4S)$
129 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ .			
130 AUBERT,B 04S reports $(1.0 \pm 0.3 \pm 0.4) \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.			
131 KROKOVNY 03B reports $(0.81 \pm 0.30 \pm 0.27) \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.			

$$\Gamma(D_{sJ}(2317)^+ \bar{D}^0 \times B(D_{sJ}(2317)^+ \rightarrow D_s^{*+} \gamma)) / \Gamma_{\text{total}} \quad \Gamma_{73}/\Gamma$$

<u>VALUE</u> (units $10^{-3}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.76	90	132 KROKOVNY	03B BELL	$e^+e^- \rightarrow \gamma(4S)$
132 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ .				

$$\Gamma(D_{sJ}(2317)^+ \bar{D}^*(2010)^0 \times B(D_{sJ}(2317)^+ \rightarrow D_s^+ \pi^0)) / \Gamma_{\text{total}} \quad \Gamma_{74}/\Gamma$$

<u>VALUE</u> (units $10^{-3}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.9<sup>+0.6</sup><sub>-0.3</sub></b>	133 AUBERT,B	04S BABR	$e^+e^- \rightarrow \gamma(4S)$

133 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$$\Gamma(D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0)) / \Gamma_{\text{total}} \quad \Gamma_{75}/\Gamma$$

<u>VALUE</u> (units $10^{-3}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.7<sup>+0.7</sup><sub>-0.7</sub> OUR AVERAGE</b>	Error includes scale factor of 1.1.		
2.7 <sup>+1.0</sup> <sub>-0.8</sub> <sup>+0.6</sup> <sub>-0.7</sub>	134,135 AUBERT,B	04S BABR	$e^+e^- \rightarrow \gamma(4S)$
1.2 <sup>+0.6</sup> <sub>-0.5</sub> <sup>+0.3</sup>	134,136 KROKOVNY	03B BELL	$e^+e^- \rightarrow \gamma(4S)$

134 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

135 AUBERT,B 04S reports  $(2.7 \pm 0.7^{+1.0}_{-0.8}) \times 10^{-3}$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$ .

We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

136 KROKOVNY 03B reports  $(1.19^{+0.61}_{-0.49} \pm 0.36) \times 10^{-3}$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$ .

We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)) / \Gamma_{\text{total}} \quad \Gamma_{76}/\Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

### 0.58±0.19 OUR AVERAGE

$0.60^{+0.24}_{-0.17} {}^{+0.14}_{-0.15}$  137,138 AUBERT,B 04S BABR  $e^+ e^- \rightarrow \gamma(4S)$

$0.56^{+0.19}_{-0.18} {}^{+0.13}_{-0.14}$  137,139 KROKOVNY 03B BELL  $e^+ e^- \rightarrow \gamma(4S)$

137 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

138 AUBERT,B 04S reports  $(0.6 \pm 0.2^{+0.2}_{-0.1}) \times 10^{-3}$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$ .

We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

139 KROKOVNY 03B reports  $(0.56^{+0.16}_{-0.15} \pm 0.17) \times 10^{-3}$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$ .

We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{77}/\Gamma$$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.22	90	140	KROKOVNY 03B BELL	$e^+ e^- \rightarrow \gamma(4S)$

140 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$$\Gamma(D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^0)) / \Gamma_{\text{total}} \quad \Gamma_{78}/\Gamma$$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.27	90	141	KROKOVNY 03B BELL	$e^+ e^- \rightarrow \gamma(4S)$

141 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$$\Gamma(D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \gamma)) / \Gamma_{\text{total}} \quad \Gamma_{79}/\Gamma$$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.98	90	142	KROKOVNY 03B BELL	$e^+ e^- \rightarrow \gamma(4S)$

142 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$$\Gamma(D_{sJ}(2457)^+ \bar{D}^*(2010)^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0)) / \Gamma_{\text{total}} \quad \Gamma_{80}/\Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
$7.6 \pm 1.7^{+3.2}_{-2.4}$	143	AUBERT,B 04S BABR	$e^+ e^- \rightarrow \gamma(4S)$

143 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$$\Gamma(D_{sJ}(2457)^+ \bar{D}^*(2010)^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)) / \Gamma_{\text{total}} \quad \Gamma_{81}/\Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>1.4 \pm 0.4^{+0.6}_{-0.4}</math></b>	144 AUBERT,B	04S BABR	$e^+ e^- \rightarrow \gamma(4S)$

144 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$$\Gamma(\bar{D}^0 D_{sJ}(2536)^+ \times B(D_{sJ}(2536)^+ \rightarrow D^*(2007)^0 K^+)) / \Gamma_{\text{total}} \quad \Gamma_{82}/\Gamma$$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2</b>	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

$$\Gamma(\bar{D}^*(2007)^0 D_{sJ}(2536)^+ \times B(D_{sJ}(2536)^+ \rightarrow D^*(2007)^0 K^+)) / \Gamma_{\text{total}} \quad \Gamma_{83}/\Gamma$$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;7</b>	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

$$\Gamma(\bar{D}^0 D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)) / \Gamma_{\text{total}} \quad \Gamma_{84}/\Gamma$$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2</b>	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

$$\Gamma(\bar{D}^*(2007)^0 D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)) / \Gamma_{\text{total}} \quad \Gamma_{85}/\Gamma$$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

$$\Gamma(\bar{D}^0 D_s^{*+}) / \Gamma_{\text{total}} \quad \Gamma_{86}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.009 ± 0.004 OUR AVERAGE</b>			

0.0084 ± 0.0031 $^{+0.0020}_{-0.0021}$	145 GIBAUT	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.012 ± 0.009 ± 0.003	146 ALBRECHT	92G ARG	$e^+ e^- \rightarrow \gamma(4S)$

145 GIBAUT 96 reports  $0.0087 \pm 0.0027 \pm 0.0017$  for  $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$ . We rescale to our best value  $B(D_s^+ \rightarrow \phi \pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.  
 146 ALBRECHT 92G reports  $0.016 \pm 0.012 \pm 0.003$  for  $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$ . We rescale to our best value  $B(D_s^+ \rightarrow \phi \pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.  
 Assumes PDG 1990  $D^0$  branching ratios, e.g.,  $B(D^0 \rightarrow K^- \pi^+) = 3.71 \pm 0.25\%$ .

$$\Gamma(\bar{D}^*(2007)^0 D_s^+)/\Gamma_{\text{total}} \quad \Gamma_{87}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.012 ± 0.005 OUR AVERAGE</b>			

0.014 ± 0.005 ± 0.003	147 GIBAUT	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.010 ± 0.007 ± 0.002	148 ALBRECHT	92G ARG	$e^+ e^- \rightarrow \gamma(4S)$

147 GIBAUT 96 reports  $0.0140 \pm 0.0043 \pm 0.0035$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$ . We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

148 ALBRECHT 92G reports  $0.013 \pm 0.009 \pm 0.002$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990  $D^0$  and  $D^*(2007)^0$  branching ratios, e.g.,  $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$  and  $B(D^*(2007)^0 \rightarrow D^0\pi^0) = 55 \pm 6\%$ .

$\Gamma(\bar{D}^*(2007)^0 D_s^{*+})/\Gamma_{\text{total}}$	$\Gamma_{88}/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.027 ± 0.010 OUR AVERAGE</b>			
$0.030 \pm 0.011 \pm 0.007$	149 GIBAUT	96 CLE2	$e^+e^- \rightarrow \gamma(4S)$
$0.023 \pm 0.013 \pm 0.006$	150 ALBRECHT	92G ARG	$e^+e^- \rightarrow \gamma(4S)$
149 GIBAUT 96 reports $0.0310 \pm 0.0088 \pm 0.0065$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.			
150 ALBRECHT 92G reports $0.031 \pm 0.016 \pm 0.005$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 $D^0$ and $D^*(2007)^0$ branching ratios, e.g., $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$ and $B(D^*(2007)^0 \rightarrow D^0\pi^0) = 55 \pm 6\%$ .			

$\Gamma(D_s^{(*)+}\bar{D}^{**0})/\Gamma_{\text{total}}$	$\Gamma_{89}/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>(2.73 ± 0.93 ± 0.68) × 10<sup>-2</sup></b>			
151 AHMED	00B CLE2	$e^+e^- \rightarrow \gamma(4S)$	
151 AHMED 00B reports their experiment's uncertainties ( $\pm 0.78 \pm 0.48 \pm 0.68\%$ ), where the first error is statistical, the second is systematic, and the third is the uncertainty in the $D_s \rightarrow \phi\pi$ branching fraction. We combine the first two in quadrature.			

$\Gamma(\bar{D}^*(2007)^0 D^*(2010)^+)/\Gamma_{\text{total}}$	$\Gamma_{90}/\Gamma$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.011	90	BARATE	98Q ALEP	$e^+e^- \rightarrow Z$

$[\Gamma(\bar{D}^0 D^*(2010)^+) + \Gamma(\bar{D}^*(2007)^0 D^+)]/\Gamma_{\text{total}}$	$\Gamma_{91}/\Gamma$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.013	90	BARATE	98Q ALEP	$e^+e^- \rightarrow Z$

$\Gamma(\bar{D}^0 D^+)/\Gamma_{\text{total}}$	$\Gamma_{92}/\Gamma$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0067	90	BARATE	98Q ALEP	$e^+e^- \rightarrow Z$

$\Gamma(\bar{D}^0 D^+ K^0)/\Gamma_{\text{total}}$	$\Gamma_{93}/\Gamma$			
VALUE (units 10 <sup>-3</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<2.8	90	152 AUBERT	03X BABR	$e^+e^- \rightarrow \gamma(4S)$

152 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(\bar{D}^*(2007)^0 D^+ K^0)/\Gamma_{\text{total}}$   $\Gamma_{94}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;6.1</b>	90	153 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

153 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma(\bar{D}^0 \bar{D}^*(2010)^+ K^0)/\Gamma_{\text{total}}$   $\Gamma_{95}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>5.2^{+1.0}_{-0.9} \pm 0.7</math></b>	154 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

154 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma(\bar{D}^*(2007)^0 D^*(2010)^+ K^0)/\Gamma_{\text{total}}$   $\Gamma_{96}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>7.8^{+2.3}_{-2.1} \pm 1.4</math></b>	155 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

155 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma(\bar{D}^0 D^0 K^+)/\Gamma_{\text{total}}$   $\Gamma_{97}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.37 \pm 0.32</math> OUR AVERAGE</b>	Error includes scale factor of 1.5.		

$1.17 \pm 0.21 \pm 0.15$	156 CHISTOV	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$
$1.9 \pm 0.3 \pm 0.3$	156 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

156 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma(\bar{D}^*(2010)^0 D^0 K^+)/\Gamma_{\text{total}}$   $\Gamma_{98}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;3.8</b>	90	157 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

157 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma(\bar{D}^0 D^*(2007)^0 K^+)/\Gamma_{\text{total}}$   $\Gamma_{99}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>4.7 \pm 0.7 \pm 0.7</math></b>	158 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

158 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma(\bar{D}^*(2007)^0 D^*(2007)^0 K^+)/\Gamma_{\text{total}}$   $\Gamma_{100}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>5.3^{+1.1}_{-1.0} \pm 1.2</math></b>	159 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

159 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma(D^- D^+ K^+)/\Gamma_{\text{total}}$   $\Gamma_{101}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.4</b>	90	160 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.90$	90	160 CHISTOV	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$
---------	----	-------------	---------	----------------------------------

160 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(D^- D^*(2010)^+ K^+)/\Gamma_{\text{total}}$   $\Gamma_{102}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	161 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

161 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(D^*(2010)^- D^+ K^+)/\Gamma_{\text{total}}$   $\Gamma_{103}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.5±0.3±0.2</b>	162 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

162 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(D^*(2010)^- D^*(2010)^+ K^+)/\Gamma_{\text{total}}$   $\Gamma_{104}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	163 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

163 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma((\bar{D}+\bar{D}^*)(D+D^*)K)/\Gamma_{\text{total}}$   $\Gamma_{105}/\Gamma$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.5±0.3±0.5</b>	164 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

164 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(D_s^+ \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{106}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00020	90	165 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

165 ALEXANDER 93B reports  $< 2.0 \times 10^{-4}$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .

$[\Gamma(D_s^+ \pi^0) + \Gamma(D_s^{*+} \pi^0)]/\Gamma_{\text{total}}$   $(\Gamma_{106} + \Gamma_{107})/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0007	90	166 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$

166 ALBRECHT 93E reports  $< 0.9 \times 10^{-3}$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .

$\Gamma(D_s^{*+} \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{107}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00033	90	167 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

167 ALEXANDER 93B reports  $< 3.2 \times 10^{-4}$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .

$\Gamma(D_s^+ \eta)/\Gamma_{\text{total}}$   $\Gamma_{108}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0005	90	168 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

168 ALEXANDER 93B reports  $< 4.6 \times 10^{-4}$  for  $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value  $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .

$\Gamma(D_s^{*+} \eta)/\Gamma_{\text{total}}$				$\Gamma_{109}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0008	90	169 ALEXANDER 93B CLE2	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$	
169 ALEXANDER 93B reports $< 7.5 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.036$ .				

$\Gamma(D_s^{*+} \rho^0)/\Gamma_{\text{total}}$				$\Gamma_{110}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0004	90	170 ALEXANDER 93B CLE2	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$	
170 ALEXANDER 93B reports $< 3.7 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.036$ .				

$[\Gamma(D_s^{*+} \rho^0) + \Gamma(D_s^{*+} \bar{K}^*(892)^0)]/\Gamma_{\text{total}}$				$(\Gamma_{110} + \Gamma_{120})/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0025	90	171 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$	
171 ALBRECHT 93E reports $< 3.4 \times 10^{-3}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.036$ .				

$\Gamma(D_s^{*+} \rho^0)/\Gamma_{\text{total}}$				$\Gamma_{111}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0005	90	172 ALEXANDER 93B CLE2	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$	
172 ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.036$ .				

$[\Gamma(D_s^{*+} \rho^0) + \Gamma(D_s^{*+} \bar{K}^*(892)^0)]/\Gamma_{\text{total}}$				$(\Gamma_{111} + \Gamma_{121})/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0015	90	173 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$	
173 ALBRECHT 93E reports $< 2.0 \times 10^{-3}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.036$ .				

$\Gamma(D_s^{*+} \omega)/\Gamma_{\text{total}}$				$\Gamma_{112}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0005	90	174 ALEXANDER 93B CLE2	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0025	90	175 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$	
174 ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.036$ .				
175 ALBRECHT 93E reports $< 3.4 \times 10^{-3}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.036$ .				

$\Gamma(D_s^{*+} \omega)/\Gamma_{\text{total}}$					$\Gamma_{113}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0007	90	176 ALEXANDER 93B CLE2	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0014	90	177 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
176 ALEXANDER 93B reports $< 6.8 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.036$ .					
177 ALBRECHT 93E reports $< 1.9 \times 10^{-3}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.036$ .					

$\Gamma(D_s^+ a_1(1260)^0)/\Gamma_{\text{total}}$					$\Gamma_{114}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0022	90	178 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
178 ALBRECHT 93E reports $< 3.0 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					

$\Gamma(D_s^{*+} a_1(1260)^0)/\Gamma_{\text{total}}$					$\Gamma_{115}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0016	90	179 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
179 ALBRECHT 93E reports $< 2.2 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					

$\Gamma(D_s^+ \phi)/\Gamma_{\text{total}}$					$\Gamma_{116}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.00032	90	180 ALEXANDER 93B CLE2	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0013	90	181 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
180 ALEXANDER 93B reports $< 3.1 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					
181 ALBRECHT 93E reports $< 1.7 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					

$\Gamma(D_s^{*+} \phi)/\Gamma_{\text{total}}$					$\Gamma_{117}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0004	90	182 ALEXANDER 93B CLE2	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0016	90	183 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
182 ALEXANDER 93B reports $< 4.2 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.036$ .					
183 ALBRECHT 93E reports $< 2.1 \times 10^{-3}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.036$ .					

$\Gamma(D_s^+ \bar{K}^0)/\Gamma_{\text{total}}$					$\Gamma_{118}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0011	90	184 ALEXANDER 93B CLE2	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0019	90	185 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
184 ALEXANDER 93B reports $< 10.3 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					
185 ALBRECHT 93E reports $< 2.5 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					

$\Gamma(D_s^{*+} \bar{K}^0)/\Gamma_{\text{total}}$					$\Gamma_{119}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0011	90	186 ALEXANDER 93B CLE2	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0023	90	187 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
186 ALEXANDER 93B reports $< 10.9 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					
187 ALBRECHT 93E reports $< 3.1 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					

$\Gamma(D_s^+ \bar{K}^*(892)^0)/\Gamma_{\text{total}}$					$\Gamma_{120}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0005	90	188 ALEXANDER 93B CLE2	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
188 ALEXANDER 93B reports $< 4.4 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					

$\Gamma(D_s^{*+} \bar{K}^*(892)^0)/\Gamma_{\text{total}}$					$\Gamma_{121}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0004	90	189 ALEXANDER 93B CLE2	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
189 ALEXANDER 93B reports $< 4.3 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					

$\Gamma(D_s^- \pi^+ K^+)/\Gamma_{\text{total}}$					$\Gamma_{122}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0008	90	190 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
190 ALBRECHT 93E reports $< 1.1 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					

$\Gamma(D_s^{*-} \pi^+ K^+)/\Gamma_{\text{total}}$					$\Gamma_{123}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0012	90	191 ALBRECHT 93E ARG	e <sup>+</sup> e <sup>-</sup> → $\gamma(4S)$		
191 ALBRECHT 93E reports $< 1.6 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .					

$\Gamma(D_s^- \pi^+ K^*(892)^+)/\Gamma_{\text{total}}$				$\Gamma_{124}/\Gamma$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.006</b>	90	192 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$
192 ALBRECHT 93E reports $< 8.6 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .				

$\Gamma(D_s^{*-} \pi^+ K^*(892)^+)/\Gamma_{\text{total}}$				$\Gamma_{125}/\Gamma$
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.008</b>	90	193 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$
193 ALBRECHT 93E reports $< 1.1 \times 10^{-2}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$ . We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$ .				

$\Gamma(\eta_c K^+)/\Gamma_{\text{total}}$				$\Gamma_{126}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b><math>1.03 \pm 0.21</math> OUR AVERAGE</b>				
1.3 $\pm 0.2$ $\pm 0.4$	194 AUBERT,B	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$1.25 \pm 0.14$ $^{+0.39}_{-0.40}$	195 FANG	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$	
$0.69^{+0.26}_{-0.21} \pm 0.22$	196 EDWARDS	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
194 AUBERT,B 04B reports $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c(1S) \rightarrow K\bar{K}\pi)] = (0.074 \pm 0.005 \pm 0.007) \times 10^{-3}$ . We divide by our best value $B(\eta_c(1S) \rightarrow K\bar{K}\pi) = (5.7 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				
195 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ .				
196 EDWARDS 01 assumes equal production of $B^0$ and $B^+$ at the $\gamma(4S)$ . The correlated uncertainties (28.3)% from $B(J/\psi(1S) \rightarrow \gamma\eta_c)$ in those modes have been accounted for.				

$\Gamma(\eta_c K^+)/\Gamma(J/\psi(1S) K^+)$				$\Gamma_{126}/\Gamma_{127}$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b><math>1.33 \pm 0.10 \pm 0.43</math></b>	197 AUBERT,B	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$	
197 Uses BABAR measurement of $B(B^+ \rightarrow J/\psi K^+) = (10.1 \pm 0.3 \pm 0.5) \times 10^{-4}$ .				

$\Gamma(J/\psi(1S) K^+)/\Gamma_{\text{total}}$				$\Gamma_{127}/\Gamma$
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>9.9 \pm 0.4</math> OUR FIT</b>				
<b><math>10.1 \pm 0.4</math> OUR AVERAGE</b>				
10.1 $\pm 0.2 \pm 0.7$	198 ABE	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$	
10.1 $\pm 0.3 \pm 0.5$	198 AUBERT	02 BABR	$e^+ e^- \rightarrow \gamma(4S)$	
10.2 $\pm 0.8 \pm 0.7$	198 JESSOP	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
9.3 $\pm 3.1 \pm 0.2$	199 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
8.1 $\pm 3.5 \pm 0.1$	6 200 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
11.0 $\pm 1.5 \pm 0.9$	59 198 ALAM	94 CLE2	Repl. by JESSOP 97	
22 $\pm 10 \pm 2$		BUSKULIC	92G ALEP	$e^+ e^- \rightarrow Z$
7 $\pm 4$	3 201 ALBRECHT	87D ARG	$e^+ e^- \rightarrow \gamma(4S)$	
10 $\pm 7 \pm 2$	3 202 BEBEK	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
9 $\pm 5$	3 203 ALAM	86 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	

- 198 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .  
 199 BORTOLETTO 92 reports  $(8 \pm 2 \pm 2) \times 10^{-4}$  for  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$ . We rescale to our best value  $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .  
 200 ALBRECHT 90J reports  $(7 \pm 3 \pm 1) \times 10^{-4}$  for  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$ . We rescale to our best value  $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .  
 201 ALBRECHT 87D assume  $B^+ B^- / B^0 \bar{B}^0$  ratio is 55/45. Superseded by ALBRECHT 90J.  
 202 BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.  
 203 ALAM 86 assumes  $B^\pm / B^0$  ratio is 60/40.

 $\Gamma(J/\psi(1S) K^+ \pi^+ \pi^-) / \Gamma_{\text{total}}$  $\Gamma_{128} / \Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.77 ± 0.20 OUR AVERAGE</b>					
0.69 ± 0.18 ± 0.12			204 ACOSTA	02F CDF	$p\bar{p}$ 1.8 TeV
1.40 ± 0.82 ± 0.02			205 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
1.40 ± 0.91 ± 0.02	6	206 ALBRECHT	87D ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.9	90	207 ALBRECHT	90J ARG		$e^+ e^- \rightarrow \Upsilon(4S)$

- 204 ACOSTA 02F uses as reference of  $B(B \rightarrow J/\psi(1S) K^+) = (10.1 \pm 0.6) \times 10^{-4}$ . The second error includes the systematic error and the uncertainties of the branching ratio.  
 205 BORTOLETTO 92 reports  $(1.2 \pm 0.6 \pm 0.4) \times 10^{-3}$  for  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$ . We rescale to our best value  $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .  
 206 ALBRECHT 87D reports  $(1.2 \pm 0.8) \times 10^{-3}$  for  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$ . We rescale to our best value  $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. They actually report  $0.0011 \pm 0.0007$  assuming  $B^+ B^- / B^0 \bar{B}^0$  ratio is 55/45. We rescale to 50/50. Analysis explicitly removes  $B^+ \rightarrow \psi(2S) K^+$ .  
 207 ALBRECHT 90J reports  $< 1.6 \times 10^{-3}$  for  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069$ . We rescale to our best value  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593$ . Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

 $\Gamma(X(3872) K^+) / \Gamma_{\text{total}}$  $\Gamma_{129} / \Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	208 CHOI	03 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
208 CHOI 03 reports $B(B^+ \rightarrow X(3872) K^+) \times B(X(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) / B(B^+ \rightarrow \psi(2S) K^+) \times B(\psi(2S) \rightarrow \pi^+ \pi^0 J/\psi(1S)) = 0.063 \pm 0.012(\text{stat}) \pm 0.007(\text{syst})$ where the $X(3872)$ is a new meson state with a mass $3872.0 \pm 0.6 \pm 0.5$ MeV and a full width $< 2.3$ MeV at 90%CL.			

$\Gamma(X(3872)K^+ \times B(X(3872) \rightarrow D^0\bar{D}^0))/\Gamma_{\text{total}}$   $\Gamma_{130}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.0 \times 10^{-5}$	90	209 CHISTOV	04 BELL	$e^+e^- \rightarrow \gamma(4S)$

209 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . $\Gamma(X(3872)K^+ \times B(X(3872) \rightarrow D^+D^-))/\Gamma_{\text{total}}$   $\Gamma_{131}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.0 \times 10^{-5}$	90	210 CHISTOV	04 BELL	$e^+e^- \rightarrow \gamma(4S)$

210 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . $\Gamma(X(3872)K^+ \times B(X(3872) \rightarrow D^0\bar{D}^0\pi^0))/\Gamma_{\text{total}}$   $\Gamma_{132}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.0 \times 10^{-5}$	90	211 CHISTOV	04 BELL	$e^+e^- \rightarrow \gamma(4S)$

211 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . $\Gamma(X(3872)K^+ \times B(X(3872) \rightarrow J/\psi(1S)\eta))/\Gamma_{\text{total}}$   $\Gamma_{133}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.7 \times 10^{-6}$	90	212 AUBERT	04Y BABR	$e^+e^- \rightarrow \gamma(4S)$

212 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . $\Gamma(J/\psi(1S)K^*(892)^+)/\Gamma_{\text{total}}$   $\Gamma_{134}/\Gamma$ For polarization information see the Listings at the end of the " $B^0$  Branching Ratios" section.

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.35±0.10 OUR AVERAGE</b>				
1.28±0.07±0.14		213 ABE	02N BELL	$e^+e^- \rightarrow \gamma(4S)$
1.37±0.09±0.11		213 AUBERT	02 BABR	$e^+e^- \rightarrow \gamma(4S)$
1.41±0.23±0.24		213 JESSOP	97 CLE2	$e^+e^- \rightarrow \gamma(4S)$
1.58±0.47±0.27		214 ABE	96H CDF	$p\bar{p}$ at 1.8 TeV
1.51±1.09±0.02		215 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
1.86±1.30±0.03	2	216 ALBRECHT	90J ARG	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.78±0.51±0.23      13      213 ALAM      94 CLE2 Sup. by JESSOP 97

213 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .214 ABE 96H assumes that  $B(B^+ \rightarrow J/\psi K^+) = (1.02 \pm 0.14) \times 10^{-3}$ .215 BORTOLETTO 92 reports  $(1.3 \pm 0.9 \pm 0.3) \times 10^{-3}$  for  $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$ . We rescale to our best value  $B(J/\psi(1S) \rightarrow e^+e^-) = (5.93 \pm 0.10) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .216 ALBRECHT 90J reports  $(1.6 \pm 1.1 \pm 0.3) \times 10^{-3}$  for  $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$ . We rescale to our best value  $B(J/\psi(1S) \rightarrow e^+e^-) = (5.93 \pm 0.10) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

### $\Gamma(J/\psi(1S)K^*(892)^+)/\Gamma(J/\psi(1S)K^+)$ $\Gamma_{134}/\Gamma_{127}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.40±0.11 OUR AVERAGE</b>			
1.37±0.10±0.08	217 AUBERT	02 BABR	$e^+ e^- \rightarrow \gamma(4S)$
1.45±0.20±0.17	218 JESSOP	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
1.92±0.60±0.17	ABE	96Q CDF	$p\bar{p}$

217 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

218 JESSOP 97 assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . The measurement is actually measured as an average over kaon charged and neutral states.

### $\Gamma(J/\psi(1S)K(1270)^+)/\Gamma_{\text{total}}$ $\Gamma_{135}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.80±0.34±0.39</b>	219 ABE	01L BELL	$e^+ e^- \rightarrow \gamma(4S)$

219 Uses the PDG value of  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.00 \pm 0.10) \times 10^{-3}$ .

### $\Gamma(J/\psi(1S)K(1400)^+)/\Gamma(J/\psi(1S)K(1270)^+)$ $\Gamma_{136}/\Gamma_{135}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.30	90	ABE	01L BELL	$e^+ e^- \rightarrow \gamma(4S)$

### $\Gamma(J/\psi(1S)\eta K^+)/\Gamma_{\text{total}}$ $\Gamma_{137}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>10.8±2.3±2.4</b>	220 AUBERT	04Y BABR	$e^+ e^- \rightarrow \gamma(4S)$

220 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

### $\Gamma(J/\psi(1S)\phi K^+)/\Gamma_{\text{total}}$ $\Gamma_{138}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>(5.2±1.7) × 10<sup>-5</sup> OUR AVERAGE</b> Error includes scale factor of 1.2.			
(4.4±1.4±0.5) × 10 <sup>-5</sup>	221 AUBERT	03O BABR	$e^+ e^- \rightarrow \gamma(4S)$
(8.8 <sup>+3.5</sup> <sub>-3.0</sub> ± 1.3) × 10 <sup>-5</sup>	222 ANASTASSOV 00	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

221 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

222 ANASTASSOV 00 finds 10 events on a background of  $0.5 \pm 0.2$ . Assumes equal production of  $B^0$  and  $B^+$  at the  $\gamma(4S)$ , a uniform Dalitz plot distribution, isotropic  $J/\psi(1S)$  and  $\phi$  decays, and  $B(B^+ \rightarrow J/\psi(1S)\phi K^+) = B(B^0 \rightarrow J/\psi(1S)\phi K^0)$ .

### $\Gamma(J/\psi(1S)\pi^+)/\Gamma_{\text{total}}$ $\Gamma_{139}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>(4.8±0.6) × 10<sup>-5</sup> OUR FIT</b> Error includes scale factor of 1.5.			
<b>(3.8±0.6±0.3) × 10<sup>-5</sup></b>	223 ABE	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$

223 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(J/\psi(1S)\pi^+)/\Gamma(J/\psi(1S)K^+)$   $\Gamma_{139}/\Gamma_{127}$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.049 ± 0.005 OUR FIT</b>		Error includes scale factor of 1.4.		
<b>0.053 ± 0.004 OUR AVERAGE</b>				
0.0537 ± 0.0045 ± 0.0011		AUBERT	04P BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.05 ± 0.019 ± 0.001		ABE	96R CDF	$p\bar{p} 1.8 \text{ TeV}$
0.052 ± 0.024		BISHAI	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0391 ± 0.0078 ± 0.0019		AUBERT	02F BABR	Repl. by AUBERT 04P
0.043 ± 0.023	5	224 ALEXANDER	95 CLE2	Sup. by BISHAI 96
224 Assumes equal production of $B^+ B^-$ and $B^0 \bar{B}^0$ on $\gamma(4S)$ .				

 $\Gamma(J/\psi(1S)\rho^+)/\Gamma_{\text{total}}$   $\Gamma_{140}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;7.7 × 10⁻⁴</b>	90	BISHAI	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(J/\psi(1S)a_1(1260)^+)/\Gamma_{\text{total}}$   $\Gamma_{141}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;1.2 × 10⁻³</b>	90	BISHAI	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(J/\psi(1S)p\bar{\Lambda})/\Gamma_{\text{total}}$   $\Gamma_{142}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>(12 ± 9) × 10⁻⁶</b>		225 AUBERT	03K BABR	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 4.1 × 10⁻⁵	90	ZANG	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$
225 Assumes equal production of $B^+$ and $B^0$ at the $\gamma(4S)$ .				

 $\Gamma(\psi(2S)K^+)/\Gamma_{\text{total}}$   $\Gamma_{143}/\Gamma$ 

<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>6.8 ± 0.4 OUR AVERAGE</b>					
6.9 ± 0.6		226 ABE	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$	
6.4 ± 0.5 ± 0.8		226 AUBERT	02 BABR	$e^+ e^- \rightarrow \gamma(4S)$	
7.8 ± 0.7 ± 0.9		226 RICHICHI	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
5.5 ± 1.0 ± 0.6		227 ABE	980 CDF	$p\bar{p} 1.8 \text{ TeV}$	
18 ± 8 ± 4	5	226 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.1 ± 2.3 ± 0.9	7	226 ALAM	94 CLE2	Repl. by RICHICHI 01	
< 5	90	226 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
22 ± 17	3	228 ALBRECHT	87D ARG	$e^+ e^- \rightarrow \gamma(4S)$	

226 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

227 ABE 980 reports  $[B(B^+ \rightarrow \psi(2S)K^+)/[B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.558 \pm 0.082 \pm 0.056$ . We multiply by our best value  $B(B^+ \rightarrow J/\psi(1S)K^+) = (9.9 \pm 1.0) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

228 ALBRECHT 87D assume  $B^+ B^- / B^0 \bar{B}^0$  ratio is 55/45. Superseded by ALBRECHT 90J.

### $\Gamma(\psi(2S)K^+)/\Gamma(J/\psi(1S)K^+)$

$\Gamma_{143}/\Gamma_{127}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.64 ± 0.06 ± 0.07</b>	229 AUBERT	02 BABR	$e^+ e^- \rightarrow \gamma(4S)$

229 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

### $\Gamma(\psi(2S)K^*(892)^+)/\Gamma_{\text{total}}$

$\Gamma_{144}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>9.2 ± 1.9 ± 1.2</b>		230 RICHICHI	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
<30	90	230 ALAM	94 CLE2	Repl. by RICHICHI 01
<35	90	230 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
<49	90	230 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$

230 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

### $\Gamma(\psi(2S)K^+\pi^+\pi^-)/\Gamma_{\text{total}}$

$\Gamma_{145}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0019 ± 0.0011 ± 0.0004</b>	3	231 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$

231 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

### $\Gamma(\psi(3770)K^+)/\Gamma_{\text{total}}$

$\Gamma_{146}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.48 ± 0.11 ± 0.07</b>	232 CHISTOV	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$

232 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

### $\Gamma(\psi(3770)K^+ \times B(\psi(3770) \rightarrow D^0 \bar{D}^0))/\Gamma_{\text{total}}$

$\Gamma_{147}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.34 ± 0.08 ± 0.05</b>	233 CHISTOV	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$

233 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

### $\Gamma(\psi(3770)K^+ \times B(\psi(3770) \rightarrow D^+ D^- K^+))/\Gamma_{\text{total}}$

$\Gamma_{148}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.14 ± 0.08 ± 0.02</b>	234 CHISTOV	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$

234 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(\chi_{c0}(1P)K^+)/\Gamma_{\text{total}}$   $\Gamma_{149}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>3.0±0.6 OUR AVERAGE</b>				Error includes scale factor of 1.1.
2.7±0.7		235 AUBERT	04T BABR	$e^+e^- \rightarrow \gamma(4S)$
3.0±0.8±0.3		236 AUBERT,B	04P BABR	$e^+e^- \rightarrow \gamma(4S)$
6.0 <sup>+2.1</sup> <sub>-1.8</sub> ±1.1		237 ABE	02B BELL	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.8 90 238 EDWARDS 01 CLE2  $e^+e^- \rightarrow \gamma(4S)$

235 The measurement performed using decay channels  $\chi_c^0 \rightarrow \pi^+\pi^-$  and  $\chi_c^0 \rightarrow K^+K^-$ . The ratio of the branching ratios for these channels is found to be consistent with world average.

236 AUBERT 04P reports  $B(B^+ \rightarrow \chi_c^0 K^+) \times B(\chi_c^0 \rightarrow \pi^+\pi^-) = (1.5 \pm 0.4 \pm 0.1) \times 10^{-6}$  and used PDG value of  $B(\chi_c^0 \rightarrow \pi\pi) = (7.4 \pm 0.8) \times 10^{-3}$  and Clebsh-Gordan coefficient to compute  $B(B^\pm \rightarrow \chi_c^0 K^\pm)$ .

237 ABE 02B measures the ratio of  $B(B^+ \rightarrow \chi_c^0 K^+)/B(B^+ \rightarrow J/\psi(1S)K^+) = 0.60 + 0.21 - 0.18 \pm 0.05 \pm 0.08$ , where the third error is due to the uncertainty in the  $B(\chi_c^0 \rightarrow \pi^+\pi^-)$ , and uses  $B(B^+ \rightarrow J/\psi(1S)K^+) = (10.0 \pm 1.0) \times 10^{-4}$  to obtain the result.

238 EDWARDS 01 assumes equal production of  $B^0$  and  $B^+$  at the  $\gamma(4S)$ . The correlated uncertainties (28.3)% from  $B(J/\psi(1S) \rightarrow \gamma\eta_c)$  in those modes have been accounted for.

$\Gamma(\chi_{c1}(1P)K^+)/\Gamma_{\text{total}}$   $\Gamma_{150}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.8±1.1 OUR AVERAGE</b>				
15.5±5.4±2.0		239 ACOSTA	02F CDF	$p\bar{p} 1.8 \text{ TeV}$
6.5±1.0±0.5		240 AUBERT	02 BABR	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.7±4.0±0.9 6 241 ALAM 94 CLE2  $e^+e^- \rightarrow \gamma(4S)$

19±13±6 242 ALBRECHT 92E ARG  $e^+e^- \rightarrow \gamma(4S)$

239 ACOSTA 02F uses as reference of  $B(B \rightarrow J/\psi(1S)K^+) = (10.1 \pm 0.6) \times 10^{-4}$ . The second error includes the systematic error and the uncertainties of the branching ratio.

240 AUBERT 02 reports  $(7.5 \pm 0.9 \pm 0.8) \times 10^{-4}$  for  $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$ . We rescale to our best value  $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (31.6 \pm 2.3) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

241 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

242 ALBRECHT 92E assumes no  $\chi_{c2}(1P)$  production and  $B(\gamma(4S) \rightarrow B^+ B^-) = 50\%$ .

$\Gamma(\chi_{c1}(1P)K^+)/\Gamma(J/\psi(1S)K^+)$   $\Gamma_{150}/\Gamma_{127}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.65±0.07±0.05</b>	243 AUBERT	02 BABR	$e^+e^- \rightarrow \gamma(4S)$

243 AUBERT 02 reports  $0.75 \pm 0.08 \pm 0.05$  for  $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$ .

We rescale to our best value  $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (31.6 \pm 2.3) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(\chi_{c1}(1P) K^*(892)^+)/\Gamma_{\text{total}}$   $\Gamma_{151}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0021	90	244 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

244 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$\Gamma(K^0 \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{152}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>2.18±0.14 OUR AVERAGE</b>				
2.23±0.17±0.11	245 AUBERT	04M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$	
2.20±0.19±0.11	245 CHAO	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$	
1.88 <sup>+0.37</sup> <sub>-0.33</sub> <sup>+0.21</sup> <sub>-0.18</sub>	245 BORNHEIM	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.94 <sup>+0.31</sup> <sub>-0.30</sub> <sup>+0.16</sup>	245 CASEY	02 BELL	Repl. by CHAO 04
1.37 <sup>+0.57</sup> <sub>-0.48</sub> <sup>+0.19</sup> <sub>-0.18</sub>	245 ABE	01H BELL	Repl. by CASEY 02
1.82 <sup>+0.33</sup> <sub>-0.30</sub> <sup>+0.20</sup>	245 AUBERT	01E BABR	Repl. by AUBERT 04M
1.82 <sup>+0.46</sup> <sub>-0.40</sub> <sup>+0.16</sup>	245 CRONIN-HEN..00	CLE2	Repl. by BORNHEIM 03
2.3 <sup>+1.1</sup> <sub>-1.0</sub> <sup>±0.36</sup>	GODANG	98 CLE2	Repl. by CRONIN-HENNESSY 00
< 4.8	90 ASNER	96 CLE2	Repl. by GODANG 98
<19	90 ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
<10	90 Avery	89B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
<68	90 Avery	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

245 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

246 Avery 89B reports  $< 9 \times 10^{-5}$  assuming the  $\Upsilon(4S)$  decays 43% to  $B^0 \bar{B}^0$ . We rescale to 50%.

$\Gamma(K^+ \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{153}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>1.25±0.10 OUR AVERAGE</b>				
1.20±0.13 <sup>+0.13</sup> <sub>-0.09</sub>	247 CHAO	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$	
1.28 <sup>+0.12</sup> <sub>-0.11</sub> <sup>±0.10</sup>	247 AUBERT	03L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$	
1.29 <sup>+0.24</sup> <sub>-0.22</sub> <sup>+0.12</sup> <sub>-0.11</sub>	247 BORNHEIM	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.3 <sup>+0.25</sup> <sub>-0.24</sub> <sup>±0.13</sup>	247 CASEY	02 BELL	Repl. by CHAO 04
1.63 <sup>+0.35</sup> <sub>-0.33</sub> <sup>+0.16</sup> <sub>-0.18</sub>	247 ABE	01H BELL	Repl. by CASEY 02
1.08 <sup>+0.21</sup> <sub>-0.19</sub> <sup>±0.10</sup>	247 AUBERT	01E BABR	Repl. by AUBERT 03L
1.16 <sup>+0.30</sup> <sub>-0.27</sub> <sup>+0.14</sup> <sub>-0.13</sub>	247 CRONIN-HEN..00	CLE2	Repl. by BORNHEIM 03
<1.6	90 GODANG	98 CLE2	Repl. by CRONIN-HENNESSY 00
<1.4	90 ASNER	96 CLE2	Repl. by GODANG 98

247 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$\Gamma(K^+\pi^0)/\Gamma(K^0\pi^+)$ 

VALUE	CL%
<b>2.38 <math>\pm 0.98 \pm 0.39</math></b>	
<b><math>-1.10 \pm 0.26</math></b>	

 $\Gamma_{153}/\Gamma_{152}$ 

DOCUMENT ID	TECN	COMMENT
ABE	01H BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

 $\Gamma(\eta' K^+)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%
<b>7.8 <math>\pm 0.5</math> OUR AVERAGE</b>	

DOCUMENT ID	TECN	COMMENT
248 AUBERT	03W BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
248 ABE	01M BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
248 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>		
248 AUBERT	01G BABR	Repl. by AUBERT 03W
6.5 $\pm 1.5$	98 CLE2	BEHRENS Repl. by RICHICHI 00

248 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . $\Gamma(\eta' K^*(892)^+)/\Gamma_{\text{total}}$ 

VALUE	CL%
<b>&lt;1.4 <math>\times 10^{-5}</math></b>	90

DOCUMENT ID	TECN	COMMENT
249 AUBERT,B	04D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

**• • • We do not use the following data for averages, fits, limits, etc. • • •**

DOCUMENT ID	TECN	COMMENT
249 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
BEHRENS	98 CLE2	Repl. by RICHICHI 00

249 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . $\Gamma(\eta K^+)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-6}$ )	CL%
<b>3.3 <math>\pm 0.8</math> OUR AVERAGE</b>	

DOCUMENT ID	TECN	COMMENT
250 AUBERT	04H BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
250 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

**• • • We do not use the following data for averages, fits, limits, etc. • • •**

DOCUMENT ID	TECN	COMMENT
BEHRENS	98 CLE2	Repl. by RICHICHI 00

250 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . $\Gamma(\eta K^*(892)^+)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%
<b>2.6 <math>\pm 0.4</math> OUR AVERAGE</b>	

DOCUMENT ID	TECN	COMMENT
251 AUBERT,B	04D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
251 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

**• • • We do not use the following data for averages, fits, limits, etc. • • •**

DOCUMENT ID	TECN	COMMENT
BEHRENS	98 CLE2	Repl. by RICHICHI 00

251 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$\Gamma(\omega K^+)/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-5}$ )	<u>CL%</u>
<b><math>0.51 \pm 0.07</math> OUR AVERAGE</b>	
$0.48 \pm 0.08 \pm 0.04$	
$0.65^{+0.13}_{-0.12} \pm 0.06$	
$0.32^{+0.24}_{-0.19} \pm 0.08$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.92^{+0.26}_{-0.23} \pm 0.10$	252 LU	02 BELL	Repl. by WANG 04A
$<0.4$	90	252 AUBERT	$01G$ BABR $e^+ e^- \rightarrow \gamma(4S)$
$1.5^{+0.7}_{-0.6} \pm 0.2$		252 BERGFELD	98 CLE2 Repl. by JESSOP 00

252 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma(\omega K^*(892)^+)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>
$<8.7 \times 10^{-5}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>
253 BERGFELD	98 CLE2

253 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma(a_0^0 K^+)/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>
$<2.5$	90

<u>DOCUMENT ID</u>	<u>TECN</u>
254 AUBERT,BE	04 BABR

254 Assumes equal production of charged and neutral  $B$  mesons from  $\gamma(4S)$  decays.

 $\Gamma(a_0^+ K^0)/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>
$<3.9$	90

<u>DOCUMENT ID</u>	<u>TECN</u>
255 AUBERT,BE	04 BABR

255 Assumes equal production of charged and neutral  $B$  mesons from  $\gamma(4S)$  decays.

 $\Gamma(K^*(892)^0 \pi^+)/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-5}$ )	<u>CL%</u>
<b><math>1.60^{+0.22}_{-0.39}</math> OUR AVERAGE</b>	

$1.55 \pm 0.18^{+0.15}_{-0.40}$	256,257 AUBERT,B	04P BABR
$1.94^{+0.42}_{-0.39}^{+0.41}_{-0.71}$	258 GARMASH	02 BELL

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<11.9$	90	259 ABE	00C SLD	$e^+ e^- \rightarrow Z$
$< 1.6$	90	260 JESSOP	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$<39$	90	261 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
$< 4.1$	90	ASNER	96 CLE2	Repl. by JESSOP 00
$<48$	90	262 ABREU	95N DLPH	Sup. by ADAM 96D
$<17$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \gamma(4S)$
$<15$	90	263 AVERY	89B CLEO	$e^+ e^- \rightarrow \gamma(4S)$
$<26$	90	AVERY	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma_{158}/\Gamma$ 

$0.48 \pm 0.08 \pm 0.04$	252 AUBERT	04H BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.65^{+0.13}_{-0.12} \pm 0.06$	252 WANG	04A BELL	$e^+ e^- \rightarrow \gamma(4S)$
$0.32^{+0.24}_{-0.19} \pm 0.08$	252 JESSOP	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.92^{+0.26}_{-0.23} \pm 0.10$	252 LU	02 BELL	Repl. by WANG 04A
$<0.4$	90	252 AUBERT	$01G$ BABR $e^+ e^- \rightarrow \gamma(4S)$
$1.5^{+0.7}_{-0.6} \pm 0.2$		252 BERGFELD	98 CLE2 Repl. by JESSOP 00

252 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma_{159}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>
$<8.7 \times 10^{-5}$	90

<u>DOCUMENT ID</u>	<u>TECN</u>
253 BERGFELD	98 CLE2

253 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma_{161}/\Gamma$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>
$<2.5$	90

<u>DOCUMENT ID</u>	<u>TECN</u>
254 AUBERT,BE	04 BABR

254 Assumes equal production of charged and neutral  $B$  mesons from  $\gamma(4S)$  decays.

 $\Gamma_{160}/\Gamma$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>
$<3.9$	90

<u>DOCUMENT ID</u>	<u>TECN</u>
255 AUBERT,BE	04 BABR

255 Assumes equal production of charged and neutral  $B$  mesons from  $\gamma(4S)$  decays.

 $\Gamma_{162}/\Gamma$ 

<u>VALUE</u> (units $10^{-5}$ )	<u>CL%</u>
<b><math>1.60^{+0.22}_{-0.39}</math> OUR AVERAGE</b>	

$1.55 \pm 0.18^{+0.15}_{-0.40}$	256,257 AUBERT,B	04P BABR
$1.94^{+0.42}_{-0.39}^{+0.41}_{-0.71}$	258 GARMASH	02 BELL

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<11.9$	90	259 ABE	00C SLD	$e^+ e^- \rightarrow Z$
$< 1.6$	90	260 JESSOP	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$<39$	90	261 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
$< 4.1$	90	ASNER	96 CLE2	Repl. by JESSOP 00
$<48$	90	262 ABREU	95N DLPH	Sup. by ADAM 96D
$<17$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \gamma(4S)$
$<15$	90	263 AVERY	89B CLEO	$e^+ e^- \rightarrow \gamma(4S)$
$<26$	90	AVERY	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

- 256 AUBERT 04P also report a branching ratio for  $B^+ \rightarrow$  "higher  $K^*$  resonances"  $\pi^+$ ,  $K^* \rightarrow K^+ \pi^-$ ,  $(25.1 \pm 2.0^{+11.0}_{-5.7}) \times 10^{-6}$ . |
- 257 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . |
- 258 Uses a reference decay mode  $B^+ \rightarrow \bar{D}^0 \pi^+$  and  $\bar{D}^0 \rightarrow K^+ \pi^-$  with  $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$ . |
- 259 ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$ . |
- 260 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . |
- 261 ADAM 96D assumes  $f_{B^0} = f_{B^+} = 0.39$  and  $f_{B_s} = 0.12$ . |
- 262 Assumes a  $B^0$ ,  $B^-$  production fraction of 0.39 and a  $B_s$  production fraction of 0.12. |
- 263 Avery 89B reports  $< 1.3 \times 10^{-4}$  assuming the  $\Upsilon(4S)$  decays 43% to  $B^0 \bar{B}^0$ . We rescale to 50%. |

### $\Gamma(K^*(892)^+ \pi^0)/\Gamma_{\text{total}}$ $\Gamma_{163}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.1 \times 10^{-5}$	90	264 JESSOP	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$< 9.9 \times 10^{-5}$	90	ASNER	96 CLE2	Repl. by JESSOP 00
264 Assumes equal production of $B^+$ and $B^0$ at the $\Upsilon(4S)$ .				

### $\Gamma(K^+ \pi^- \pi^+)/\Gamma_{\text{total}}$ $\Gamma_{164}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>(5.7 ± 0.4 ) × 10<sup>-5</sup> OUR AVERAGE</b>			
$(5.36 \pm 0.31 \pm 0.51) \times 10^{-5}$	265 GARMASH	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$(5.91 \pm 0.38 \pm 0.32) \times 10^{-5}$	266 AUBERT	03M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$(5.56 \pm 0.58 \pm 0.77) \times 10^{-5}$	267 GARMASH	02 BELL	Repl. by GARMASH 04
265 Assumes equal production of $B^+$ and $B^0$ at the $\Upsilon(4S)$ .			
266 Assumes equal production of $B^0$ and $B^+$ at the $\Upsilon(4S)$ ; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.			
267 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$ .			

### $\Gamma(K^+ \pi^- \pi^+ \text{nonresonant})/\Gamma_{\text{total}}$ $\Gamma_{165}/\Gamma$

VALUE (units 10 <sup>-5</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.7$	90	268 AUBERT,B	04P BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$< 33$	90	269 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
$< 2.8$	90	BERGFELD	96B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 40$	90	270 ABREU	95N DLPH	Sup. by ADAM 96D
$< 33$	90	ALBRECHT	91E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 19$	90	271 Avery	89B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
268 Assumes equal production of $B^+$ and $B^0$ at the $\Upsilon(4S)$ .				
269 ADAM 96D assumes $f_{B^0} = f_{B^+} = 0.39$ and $f_{B_s} = 0.12$ .				
270 Assumes a $B^0$ , $B^-$ production fraction of 0.39 and a $B_s$ production fraction of 0.12.				
271 Avery 89B reports $< 1.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$ . We rescale to 50%.				

$\Gamma(K^+ f_0(980) \times B(f_0 \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$   $\Gamma_{166}/\Gamma$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b><math>9.3 \pm 2.1</math> OUR AVERAGE</b>					
$9.2 \pm 1.2^{+2.1}_{-2.6}$	272	AUBERT,B	04P BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$9.6^{+2.5+3.7}_{-2.3-1.7}$	273	GARMASH	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<80$	90	274 Avery	89B CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
272 AUBERT,B 04P also reports $B(B^+ \rightarrow \text{"higher } f^0 \text{ resonances"} \pi^+, f(980)^0 \rightarrow \pi^+ \pi^-) = (3.2 \pm 1.2^{+6.0}_{-2.9}) \times 10^{-6}$ .					
273 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$ . Only charged pions from the $f_0(980)$ are used.					
274 Avery 89B reports $< 7 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$ . We rescale to 50%.					

 $\Gamma(K^+ \rho^0)/\Gamma_{\text{total}}$   $\Gamma_{167}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<6.2 \times 10^{-6}$	90	275 AUBERT,B	04P BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<1.2 \times 10^{-5}$	90	276 GARMASH	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$	
$<8.6 \times 10^{-5}$	90	277 ABE	00C SLD	$e^+ e^- \rightarrow Z$	
$<1.7 \times 10^{-5}$	90	278 JESSOP	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
$<1.2 \times 10^{-4}$	90	279 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$	
$<1.9 \times 10^{-5}$	90	ASNER	96 CLE2	Repl. by JESSOP 00	
$<1.9 \times 10^{-4}$	90	280 ABREU	95N DLPH	Sup. by ADAM 96D	
$<1.8 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \gamma(4S)$	
$<8 \times 10^{-5}$	90	281 Avery	89B CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
$<2.6 \times 10^{-4}$	90	AVERY	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	

275 AUBERT 04P reports a central value of  $(3.9 \pm 1.2^{+1.3}_{-3.5}) \times 10^{-6}$  for this branching ratio.

276 Uses a reference decay mode  $B^+ \rightarrow \bar{D}^0 \pi^+$  and  $\bar{D}^0 \rightarrow K^+ \pi^-$  with  $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$ .

277 ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$ .

278 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

279 ADAM 96D assumes  $f_{B^0} = f_{B^-} = 0.39$  and  $f_{B_s} = 0.12$ .

280 Assumes a  $B^0$ ,  $B^-$  production fraction of 0.39 and a  $B_s$  production fraction of 0.12.

281 Avery 89B reports  $< 7 \times 10^{-5}$  assuming the  $\gamma(4S)$  decays 43% to  $B^0 \bar{B}^0$ . We rescale to 50%.

 $\Gamma(K_2^*(1430)^0 \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{168}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<6.8 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \gamma(4S)$	

$\Gamma(K^-\pi^+\pi^+)/\Gamma_{\text{total}}$  $\Gamma_{169}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	90	282 AUBERT	03M BABR	$e^+e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<4.5 \times 10^{-6}$	90	283 GARMASH	04 BELL	$e^+e^- \rightarrow \gamma(4S)$
$<7.0 \times 10^{-6}$	90	284 GARMASH	02 BELL	$e^+e^- \rightarrow \gamma(4S)$

282 Assumes equal production of  $B^0$  and  $B^+$  at the  $\gamma(4S)$ ; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

283 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

284 Uses a reference decay mode  $B^+ \rightarrow \bar{D}^0\pi^+$  and  $\bar{D}^0 \rightarrow K^+\pi^-$  with  $B(B^+ \rightarrow \bar{D}^0\pi^+) \cdot B(\bar{D}^0 \rightarrow K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$ .

 $\Gamma(K^-\pi^+\pi^+\text{nonresonant})/\Gamma_{\text{total}}$  $\Gamma_{170}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.6 \times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+e^- \rightarrow \gamma(4S)$

 $\Gamma(K_1(1400)^0\pi^+)/\Gamma_{\text{total}}$  $\Gamma_{171}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.6 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \gamma(4S)$

 $\Gamma(K^0\pi^+\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{172}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<66 \times 10^{-6}$	90	285 ECKHART	02 CLE2	$e^+e^- \rightarrow \gamma(4S)$

285 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

 $\Gamma(K^0\rho^+)/\Gamma_{\text{total}}$  $\Gamma_{173}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.8 \times 10^{-5}$	90	ASNER	96 CLE2	$e^+e^- \rightarrow \gamma(4S)$

 $\Gamma(K^*(892)^+\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{174}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-3}$	90	ALBRECHT	91E ARG	$e^+e^- \rightarrow \gamma(4S)$

 $\Gamma(K^*(892)^+\rho^0)/\Gamma_{\text{total}}$  $\Gamma_{175}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$10.6^{+3.0}_{-2.6} \pm 2.4$		286 AUBERT	03V BABR	$e^+e^- \rightarrow \gamma(4S)$

$\bullet \bullet \bullet$  We do not use the following data for averages, fits, limits, etc.  $\bullet \bullet \bullet$

$< 74$	90	287 GODANG	02 CLE2	$e^+e^- \rightarrow \gamma(4S)$
$<900$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \gamma(4S)$

286 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

287 Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to  $4.9 \times 10^{-5}$ .

 $\Gamma(K^*(892)^+K^*(892)^0)/\Gamma_{\text{total}}$  $\Gamma_{176}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.1 \times 10^{-5}$	90	288 GODANG	02 CLE2	$e^+e^- \rightarrow \gamma(4S)$

288 Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to  $4.8 \times 10^{-5}$ .

$\Gamma(K_1(1400)^+\rho^0)/\Gamma_{\text{total}}$

VALUE	CL%
$<7.8 \times 10^{-4}$	90

$\Gamma_{177}/\Gamma$

DOCUMENT ID	TECN	COMMENT
ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_2^*(1430)^+\rho^0)/\Gamma_{\text{total}}$

VALUE	CL%
$<1.5 \times 10^{-3}$	90

$\Gamma_{178}/\Gamma$

DOCUMENT ID	TECN	COMMENT
ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^+\bar{K}^0)/\Gamma_{\text{total}}$

VALUE	CL%
$<2.5 \times 10^{-6}$	90

$\Gamma_{179}/\Gamma$

DOCUMENT ID	TECN	COMMENT
289 AUBERT	04M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.3 \times 10^{-6}$

289 CHAO 04 BELL  $e^+ e^- \rightarrow \Upsilon(4S)$

$<3.3 \times 10^{-6}$

289 BORNHEIM 03 CLE2  $e^+ e^- \rightarrow \Upsilon(4S)$

$<2.0 \times 10^{-6}$

289 CASEY 02 BELL Repl. by CHAO 04

$<5.0 \times 10^{-6}$

289 ABE 01H BELL  $e^+ e^- \rightarrow \Upsilon(4S)$

$<2.4 \times 10^{-6}$

289 AUBERT 01E BABR  $e^+ e^- \rightarrow \Upsilon(4S)$

$<5.1 \times 10^{-6}$

289 CRONIN-HEN..00 CLE2  $e^+ e^- \rightarrow \Upsilon(4S)$

$<2.1 \times 10^{-5}$

289 GODANG 98 CLE2 Repl. by CRONIN-

HENNESSY 00

289 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$\Gamma(\bar{K}^0 K^+ \pi^0)/\Gamma_{\text{total}}$

VALUE	CL%
$<24 \times 10^{-6}$	90

$\Gamma_{180}/\Gamma$

DOCUMENT ID	TECN	COMMENT
290 ECKHART	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

290 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$\Gamma(K^+ K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units $10^{-6}$ )	CL%
<b>11.5 ± 1.3 OUR AVERAGE</b>	

$\Gamma_{181}/\Gamma$

DOCUMENT ID	TECN	COMMENT
291 AUBERT,B	04V BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

DOCUMENT ID	TECN	COMMENT
291 GARMASH	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

291 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$\Gamma(K_S^0 K_S^0 \pi^+)/\Gamma_{\text{total}}$

VALUE (units $10^{-6}$ )	CL%
$<3.2$	90

$\Gamma_{182}/\Gamma$

DOCUMENT ID	TECN	COMMENT
292 GARMASH	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

292 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$

VALUE	CL%
$< 6.3 \times 10^{-6}$	90

$\Gamma_{183}/\Gamma$

DOCUMENT ID	TECN	COMMENT
293 AUBERT	03M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<13 \times 10^{-6}$

294 GARMASH 04 BELL  $e^+ e^- \rightarrow \Upsilon(4S)$

$< 1.2 \times 10^{-5}$

295 GARMASH 02 BELL  $e^+ e^- \rightarrow \Upsilon(4S)$

- 293 Assumes equal production of  $B^0$  and  $B^+$  at the  $\Upsilon(4S)$ ; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.  
 294 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .  
 295 Uses a reference decay mode  $B^+ \rightarrow \bar{D}^0\pi^+$  and  $\bar{D}^0 \rightarrow K^+\pi^-$  with  $B(B^+ \rightarrow \bar{D}^0\pi^+) \cdot B(\bar{D}^0 \rightarrow K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$ .

$\Gamma(K^+K^-\pi^+ \text{ nonresonant})/\Gamma_{\text{total}}$   $\Gamma_{184}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.5 \times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^+K^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{185}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-6}$	90	296 AUBERT	03M BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<2.4 \times 10^{-6}$	90	297 GARMASH	04 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$<3.2 \times 10^{-6}$	90	298 GARMASH	02 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

- 296 Assumes equal production of  $B^0$  and  $B^+$  at the  $\Upsilon(4S)$ ; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

- 297 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .  
 298 Uses a reference decay mode  $B^+ \rightarrow \bar{D}^0\pi^+$  and  $\bar{D}^0 \rightarrow K^+\pi^-$  with  $B(B^+ \rightarrow \bar{D}^0\pi^+) \cdot B(\bar{D}^0 \rightarrow K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$ .

$\Gamma(K^+K^+\pi^- \text{ nonresonant})/\Gamma_{\text{total}}$   $\Gamma_{186}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.79 \times 10^{-5}$	90	ABBIENDI	00B OPAL	$e^+e^- \rightarrow Z$

$\Gamma(K^+K^*(892)^0)/\Gamma_{\text{total}}$   $\Gamma_{187}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.3 \times 10^{-6}$	90	299 JESSOP	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<1.29 \times 10^{-4}$	90	ABBIENDI	00B OPAL	$e^+e^- \rightarrow Z$
$<1.38 \times 10^{-4}$	90	300 ABE	00C SLD	$e^+e^- \rightarrow Z$

- 299 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

- 300 ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$ .

$\Gamma(K^+f_J(2220))/\Gamma_{\text{total}}$   $\Gamma_{188}/\Gamma$

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
not seen	301 HUANG	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
301 No evidence is found for such decay and set a limit on $B(B^+ \rightarrow f_J(2220)) \times B(f_J(2220) \rightarrow \phi\phi) < 1.2 \times 10^{-6}$ at 90%CL where the $f_J(2220)$ is a possible glueball state.			

**$\Gamma(K^+ K^- K^+)/\Gamma_{\text{total}}$**   **$\Gamma_{189}/\Gamma$**

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>3.08 \pm 0.21</math> OUR AVERAGE</b>				
3.28 $\pm 0.18 \pm 0.28$		302 GARMASH	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$
2.96 $\pm 0.21 \pm 0.16$		303 AUBERT	03M BABR	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.53 $\pm 0.37 \pm 0.45$		304 GARMASH	02 BELL	Repl. by GARMASH 04
<20	90	305 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
<32	90	306 ABREU	95N DLPH	Sup. by ADAM 96D
<35	90	ALBRECHT	91E ARG	$e^+ e^- \rightarrow \gamma(4S)$

302 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

303 Assumes equal production of  $B^0$  and  $B^+$  at the  $\gamma(4S)$ ; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

304 Uses a reference decay mode  $B^+ \rightarrow \bar{D}^0 \pi^+$  and  $\bar{D}^0 \rightarrow K^+ \pi^-$  with  $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$ .

305 ADAM 96D assumes  $f_{B^0} = f_{B^-} = 0.39$  and  $f_{B_s} = 0.12$ .

306 Assumes a  $B^0$ ,  $B^-$  production fraction of 0.39 and a  $B_s$  production fraction of 0.12.

**$\Gamma(K^+ \phi)/\Gamma_{\text{total}}$**   **$\Gamma_{190}/\Gamma$**

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>9.3 \pm 1.0</math> OUR AVERAGE</b>				
Error includes scale factor of 1.3. See the ideogram below.				
10.0 $\pm 0.9$	$\pm 0.5$	307 AUBERT	04A BABR	$e^+ e^- \rightarrow \gamma(4S)$
9.4 $\pm 1.1$	$\pm 0.7$	308 CHEN	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
5.5 $\pm 2.1$	$\pm 0.6$	308 BRIERE	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

14.6  $\pm 3.0$

$\pm 2.0$

309 GARMASH

02 BELL

Repl. by CHEN 03B

7.7  $\pm 1.6$

$\pm 0.8$

308 AUBERT

01D BABR

$e^+ e^- \rightarrow \gamma(4S)$

<144

90

310 ABE

00C SLD

$e^+ e^- \rightarrow Z$

< 5

90

308 BERGFELD

98 CLE2

<280

90

311 ADAM

96D DLPH

$e^+ e^- \rightarrow Z$

< 12

90

ASNER

96 CLE2

$e^+ e^- \rightarrow \gamma(4S)$

<440

90

312 ABREU

95N DLPH

Sup. by ADAM 96D

<180

90

ALBRECHT

91B ARG

$e^+ e^- \rightarrow \gamma(4S)$

< 90

90

313 AVERY

89B CLEO

$e^+ e^- \rightarrow \gamma(4S)$

<210

90

AVERY

87 CLEO

$e^+ e^- \rightarrow \gamma(4S)$

307 Assumes equal production of  $B^+$  and  $B^0$  at  $\gamma(4S)$ .

308 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .


309 Uses a reference decay mode  $B^+ \rightarrow \bar{D}^0 \pi^+$  and  $\bar{D}^0 \rightarrow K^+ \pi^-$  with  $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$ .

310 ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7 \pm 1.8)\%$  and  $f_{B_s} = (10.5 \pm 1.8)\%$ .

311 ADAM 96D assumes  $f_{B^0} = f_{B^-} = 0.39$  and  $f_{B_s} = 0.12$ .

312 Assumes a  $B^0$ ,  $B^-$  production fraction of 0.39 and a  $B_s$  production fraction of 0.12.

313 Avery 89B reports  $< 8 \times 10^{-5}$  assuming the  $\Upsilon(4S)$  decays 43% to  $B^0 \bar{B}^0$ . We rescale to 50%.



### $\Gamma(K^+ K^- K^+ \text{nonresonant})/\Gamma_{\text{total}}$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.8$	90	BERGFELD	96B	$e^+ e^- \rightarrow \Upsilon(4S)$

### $\Gamma(K^*(892)^+ K^+ K^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-3}$	90	ALBRECHT	91E	$e^+ e^- \rightarrow \Upsilon(4S)$

### $\Gamma(K^*(892)^+ \phi)/\Gamma_{\text{total}}$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>9.6 \pm 3.0</math> OUR AVERAGE</b>				Error includes scale factor of 1.9.
$12.7^{+2.2}_{-2.0} \pm 1.1$		314 AUBERT	03V BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$6.7^{+2.1}_{-1.9} \pm 0.7$		314 CHEN	03B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.7^{+4.2}_{-3.4} \pm 1.7$	314 AUBERT	01D BABR	Repl. by AUBERT 03V
$< 22.5$	90	314 BRIERE	01 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
$< 41$	90	314 BERGFELD	98 CLE2
$< 70$	90	ASNER	96 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
$< 1300$	90	ALBRECHT	91B ARG $e^+ e^- \rightarrow \Upsilon(4S)$

314 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$\Gamma(K_1(1400)^+\phi)/\Gamma_{\text{total}}$		$\Gamma_{194}/\Gamma$		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_2^*(1430)^+\phi)/\Gamma_{\text{total}}$		$\Gamma_{195}/\Gamma$		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.4 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^+\phi\phi)/\Gamma_{\text{total}}$		$\Gamma_{196}/\Gamma$		
VALUE (units $10^{-6}$ )		DOCUMENT ID	TECN	COMMENT
$2.6^{+1.1}_{-0.9} \pm 0.3$		315 HUANG	03 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

315 Assumes equal production of  $B^0$  and  $B^+$  at the  $\Upsilon(4S)$  and for a  $\phi\phi$  invariant mass below  $2.85 \text{ GeV}/c^2$ .

$\Gamma(K^*(892)^+\gamma)/\Gamma_{\text{total}}$		$\Gamma_{197}/\Gamma$		
VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>4.03 \pm 0.26 \text{ OUR AVERAGE}</math></b>				
$3.87 \pm 0.28 \pm 0.26$		316 AUBERT,BE	04A BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$4.25 \pm 0.31 \pm 0.24$		317 NAKAO	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$3.76^{+0.89}_{-0.83} \pm 0.28$		317 COAN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.83 \pm 0.62 \pm 0.22$	317 AUBERT	02C BABR	Repl. by AUBERT,BE 04A
$5.7 \pm 3.1 \pm 1.1$	318 AMMAR	93 CLE2	Repl. by COAN 00
$< 55$	90	319 ALBRECHT	89G ARG
$< 55$	90	319 AVERY	89B CLEO
$< 180$	90	AVERY	87 CLEO

316 Uses the production ratio of charged and neutral B from  $\Upsilon(4S)$  decays  $R^{+/0} = 1.006 \pm 0.048$ .

317 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

318 AMMAR 93 observed  $4.1 \pm 2.3$  events above background.

319 Assumes the  $\Upsilon(4S)$  decays 43% to  $B^0\bar{B}^0$ .

$\Gamma(K_1(1270)^+\gamma)/\Gamma_{\text{total}}$		$\Gamma_{198}/\Gamma$		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.9 \times 10^{-5}$	90	320 NISHIDA	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.0073$	90	321 ALBRECHT	89G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------	----	--------------	---------	------------------------------------

320 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

321 ALBRECHT 89G reports  $< 0.0066$  assuming the  $\Upsilon(4S)$  decays 45% to  $B^0\bar{B}^0$ . We rescale to 50%.

$\Gamma(\phi K^+\gamma)/\Gamma_{\text{total}}$		$\Gamma_{199}/\Gamma$		
VALUE (units $10^{-6}$ )		DOCUMENT ID	TECN	COMMENT
$3.4 \pm 0.9 \pm 0.4$		322 DRUTSKOY	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

322 Assumes equal production of  $B^+$  and  $B^0$  at  $\Upsilon(4S)$ .

$\Gamma(K^+\pi^-\pi^+\gamma)/\Gamma_{\text{total}}$	$\Gamma_{200}/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$(2.4 \pm 0.5 \pm 0.4) \times 10^{-5}$	323,324 NISHIDA	02 BELL	$e^+e^- \rightarrow \gamma(4S)$

323 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .324  $M_{K\pi\pi} < 2.4 \text{ GeV}/c^2$ .

$\Gamma(K^*(892)^0\pi^+\gamma)/\Gamma_{\text{total}}$	$\Gamma_{201}/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$(2.0 \pm 0.7 \pm 0.2) \times 10^{-5}$	325,326 NISHIDA	02 BELL	$e^+e^- \rightarrow \gamma(4S)$

325 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .326  $M_{K\pi\pi} < 2.4 \text{ GeV}/c^2$ .

$\Gamma(K^+\rho^0\gamma)/\Gamma_{\text{total}}$	$\Gamma_{202}/\Gamma$			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.0 \times 10^{-5}$	90	327,328 NISHIDA	02 BELL	$e^+e^- \rightarrow \gamma(4S)$

327 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .328  $M_{K\pi\pi} < 2.4 \text{ GeV}/c^2$ .

$\Gamma(K^+\pi^-\pi^+\gamma \text{ nonresonant})/\Gamma_{\text{total}}$	$\Gamma_{203}/\Gamma$			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9.2 \times 10^{-6}$	90	329,330 NISHIDA	02 BELL	$e^+e^- \rightarrow \gamma(4S)$

329 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .330  $M_{K\pi\pi} < 2.4 \text{ GeV}/c^2$ .

$\Gamma(K_1(1400)^+\gamma)/\Gamma_{\text{total}}$	$\Gamma_{204}/\Gamma$			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.0 \times 10^{-5}$	90	331 NISHIDA	02 BELL	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0022 90 332 ALBRECHT 89G ARG  $e^+e^- \rightarrow \gamma(4S)$ 331 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .332 ALBRECHT 89G reports < 0.0020 assuming the  $\gamma(4S)$  decays 45% to  $B^0\bar{B}^0$ . We rescale to 50%.

$\Gamma(K_2^*(1430)^+\gamma)/\Gamma_{\text{total}}$	$\Gamma_{205}/\Gamma$			
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.45 \pm 0.40 \pm 0.15$	333 AUBERT,B	04U BABR	$e^+e^- \rightarrow \gamma(4S)$	■

• • • We do not use the following data for averages, fits, limits, etc. • • •

<140 90 334 ALBRECHT 89G ARG  $e^+e^- \rightarrow \gamma(4S)$ 333 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .334 ALBRECHT 89G reports < 0.0013 assuming the  $\gamma(4S)$  decays 45% to  $B^0\bar{B}^0$ . We rescale to 50%.

**$\Gamma(K^*(1680)^+\gamma)/\Gamma_{\text{total}}$**   **$\Gamma_{206}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.0019</b>	90	335 ALBRECHT	89G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

335 ALBRECHT 89G reports  $< 0.0017$  assuming the  $\Upsilon(4S)$  decays 45% to  $B^0 \bar{B}^0$ . We rescale to 50%.

**$\Gamma(K_3^*(1780)^+\gamma)/\Gamma_{\text{total}}$**   **$\Gamma_{207}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.0055</b>	90	336 ALBRECHT	89G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

336 ALBRECHT 89G reports  $< 0.005$  assuming the  $\Upsilon(4S)$  decays 45% to  $B^0 \bar{B}^0$ . We rescale to 50%.

**$\Gamma(K_4^*(2045)^+\gamma)/\Gamma_{\text{total}}$**   **$\Gamma_{208}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.0099</b>	90	337 ALBRECHT	89G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

337 ALBRECHT 89G reports  $< 0.0090$  assuming the  $\Upsilon(4S)$  decays 45% to  $B^0 \bar{B}^0$ . We rescale to 50%.

**$\Gamma(\rho^+\gamma)/\Gamma_{\text{total}}$**   **$\Gamma_{209}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.8 × 10<sup>-6</sup></b>	90	338 AUBERT	05 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.1 \times 10^{-6}$	90	338 AUBERT	04C BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$<1.3 \times 10^{-5}$	90	338,339 COAN	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

338 Assumes equal production of  $B^+$  and  $B^0$  at  $\Upsilon(4S)$ .

339 No evidence for a nonresonant  $K\pi\gamma$  contamination was seen; the central value assumes no contamination.

**$\Gamma(\pi^+\pi^0)/\Gamma_{\text{total}}$**   **$\Gamma_{210}/\Gamma$**

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>0.51<sup>+0.08</sup><sub>-0.09</sub> OUR AVERAGE</b>				

0.50 $\pm 0.12 \pm 0.05$		340 CHAO	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
0.55 $\pm 0.10 \pm 0.06$		340 AUBERT	03L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.46 $\pm 0.18 \pm 0.06$		340 BORNHEIM	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.74 $\pm 0.23 \pm 0.09$		340 CASEY	02 BELL	Repl. by CHAO 04
< 1.34	90	340 ABE	01H BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
< 0.96	90	340 AUBERT	01E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
< 1.27	90	340 CRONIN-HEN.	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
< 2.0	90	GODANG	98 CLE2	Repl. by CRONIN-HENNESSY 00
< 1.7	90	ASNER	96 CLE2	Repl. by GODANG 98
< 24	90	341 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
<230	90	342 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

340 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

341 ALBRECHT 90B limit assumes equal production of  $B^0 \bar{B}^0$  and  $B^+ B^-$  at  $\Upsilon(4S)$ .

342 BEBEK 87 assume the  $\Upsilon(4S)$  decays 43% to  $B^0 \bar{B}^0$ .

$\Gamma(\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{211}/\Gamma$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>10.9 \pm 3.3 \pm 1.6</math></b>		343 AUBERT	03M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<130	90	344 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
<220	90	345 ABREU	95N DLPH	Sup. by ADAM 96D
<450	90	346 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
<190	90	347 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

343 Assumes equal production of  $B^0$  and  $B^+$  at the  $\Upsilon(4S)$ ; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

344 ADAM 96D assumes  $f_{B^0} = f_{B^-} = 0.39$  and  $f_{B_s} = 0.12$ .

345 Assumes a  $B^0$ ,  $B^-$  production fraction of 0.39 and a  $B_s$  production fraction of 0.12.

346 ALBRECHT 90B limit assumes equal production of  $B^0\bar{B}^0$  and  $B^+B^-$  at  $\Upsilon(4S)$ .

347 BORTOLETTO 89 reports  $< 1.7 \times 10^{-4}$  assuming the  $\Upsilon(4S)$  decays 43% to  $B^0\bar{B}^0$ . We rescale to 50%.

 $\Gamma(\rho^0\pi^+)/\Gamma_{\text{total}}$  $\Gamma_{212}/\Gamma$ 

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.92 \pm 0.11</math> OUR AVERAGE</b>					
0.95 $\pm 0.11 \pm 0.09$			348 AUBERT	04Z BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.80 $^{+0.23}_{-0.20} \pm 0.07$			348 GORDON	02 BELL	$e^+ e^- \rightarrow \Upsilon(rS)$
1.04 $^{+0.33}_{-0.34} \pm 0.21$			348 JESSOP	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 8.3	90	349 ABE	00C SLD	$e^+ e^- \rightarrow Z$
<16	90	350 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
< 4.3	90	ASNER	96 CLE2	Repl. by JESSOP 00
<26	90	351 ABREU	95N DLPH	Sup. by ADAM 96D
<15	90	352 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
<17	90	353 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
<23	90	353 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
<60	90	0 GILES	84 CLEO	Repl. by BEBEK 87

348 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

349 ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$ .

350 ADAM 96D assumes  $f_{B^0} = f_{B^-} = 0.39$  and  $f_{B_s} = 0.12$ .

351 Assumes a  $B^0$ ,  $B^-$  production fraction of 0.39 and a  $B_s$  production fraction of 0.12.

352 ALBRECHT 90B limit assumes equal production of  $B^0\bar{B}^0$  and  $B^+B^-$  at  $\Upsilon(4S)$ .

353 Papers assume the  $\Upsilon(4S)$  decays 43% to  $B^0\bar{B}^0$ . We rescale to 50%.

 $[\Gamma(K^*(892)^0\pi^+) + \Gamma(\rho^0\pi^+)]/\Gamma_{\text{total}}$  $(\Gamma_{162} + \Gamma_{212})/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>(17 \pm 12 \pm 2) \times 10^{-5}</math></b>	354 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$

354 ADAM 96D assumes  $f_{B^0} = f_{B^-} = 0.39$  and  $f_{B_s} = 0.12$ .

$\Gamma(\pi^+ f_0(980))/\Gamma_{\text{total}}$   $\Gamma_{213}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-4}$	90	355 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

355 BORTOLETTO 89 reports  $< 1.2 \times 10^{-4}$  assuming the  $\gamma(4S)$  decays 43% to  $B^0 \bar{B}^0$ .  
We rescale to 50%.

$\Gamma(\pi^+ f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_{214}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.4 \times 10^{-4}$	90	356 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

356 BORTOLETTO 89 reports  $< 2.1 \times 10^{-4}$  assuming the  $\gamma(4S)$  decays 43% to  $B^0 \bar{B}^0$ .  
We rescale to 50%.

$\Gamma(\pi^+ \pi^- \pi^+ \text{nonresonant})/\Gamma_{\text{total}}$   $\Gamma_{215}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.1 \times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\pi^+ \pi^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{216}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.9 \times 10^{-4}$	90	357 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$

357 ALBRECHT 90B limit assumes equal production of  $B^0 \bar{B}^0$  and  $B^+ B^-$  at  $\gamma(4S)$ .

$\Gamma(\rho^+ \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{217}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>10.9 \pm 1.9 \pm 1.9</math></b>	90	358 AUBERT	04Z BABR	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 43	90	359 JESSOP	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
< 77	90	ASNER	96 CLE2	Repl. by JESSOP 00
< 550	90	360 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$

358 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

359 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . Assumes no nonresonant contributions of  $B^+ \rightarrow \pi^+ \pi^0 \pi^0$ .

360 ALBRECHT 90B limit assumes equal production of  $B^0 \bar{B}^0$  and  $B^+ B^-$  at  $\gamma(4S)$ .

$\Gamma(\pi^+ \pi^- \pi^+ \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{218}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.0 \times 10^{-3}$	90	361 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$

361 ALBRECHT 90B limit assumes equal production of  $B^0 \bar{B}^0$  and  $B^+ B^-$  at  $\gamma(4S)$ .

$\Gamma(\rho^+ \rho^0)/\Gamma_{\text{total}}$   $\Gamma_{219}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>2.6 \pm 0.6</math> OUR AVERAGE</b>				

$2.25^{+0.57}_{-0.54} \pm 0.58$  362 AUBERT 03V BABR  $e^+ e^- \rightarrow \gamma(4S)$

$3.17 \pm 0.71^{+0.38}_{-0.67}$  363 ZHANG 03B BELL  $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<100 90 364 ALBRECHT 90B ARG  $e^+ e^- \rightarrow \gamma(4S)$

362 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

363 Assumes equal production of  $B^0$  and  $B^+$  at the  $\Upsilon(4S)$  and the systematic error includes the error associated with the helicity-mix uncertainty.

364 ALBRECHT 90B limit assumes equal production of  $B^0\bar{B}^0$  and  $B^+B^-$  at  $\Upsilon(4S)$ .

### $\Gamma(a_1(1260)^+\pi^0)/\Gamma_{\text{total}}$ $\Gamma_{220}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-3}$	90	365 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

365 ALBRECHT 90B limit assumes equal production of  $B^0\bar{B}^0$  and  $B^+B^-$  at  $\Upsilon(4S)$ .

### $\Gamma(a_1(1260)^0\pi^+)/\Gamma_{\text{total}}$ $\Gamma_{221}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.0 \times 10^{-4}$	90	366 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

366 ALBRECHT 90B limit assumes equal production of  $B^0\bar{B}^0$  and  $B^+B^-$  at  $\Upsilon(4S)$ .

### $\Gamma(\omega\pi^+)/\Gamma_{\text{total}}$ $\Gamma_{222}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>0.59±0.10 OUR AVERAGE</b>				Error includes scale factor of 1.2.
$0.55 \pm 0.09 \pm 0.05$		367 AUBERT	04H BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.57^{+0.14}_{-0.13} \pm 0.06$		367 WANG	04A BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$1.13^{+0.33}_{-0.29} \pm 0.14$		367 JESSOP	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.42^{+0.20}_{-0.18} \pm 0.05$       367 LU      02 BELL      Repl. by WANG 04A

$0.66^{+0.21}_{-0.18} \pm 0.07$       367 AUBERT      01G BABR      Repl. by AUBERT 04H

$< 2.3$       90      367 BERGFELD      98 CLE2      Repl. by JESSOP 00

$<40$       90      368 ALBRECHT      90B ARG       $e^+e^- \rightarrow \Upsilon(4S)$

367 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

368 ALBRECHT 90B limit assumes equal production of  $B^0\bar{B}^0$  and  $B^+B^-$  at  $\Upsilon(4S)$ .

### $\Gamma(\omega\rho^+)/\Gamma_{\text{total}}$ $\Gamma_{223}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN
$<6.1 \times 10^{-5}$	90	369 BERGFELD	98 CLE2

369 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

### $\Gamma(\eta\pi^+)/\Gamma_{\text{total}}$ $\Gamma_{224}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>4.8^{+1.3}_{-1.1} OUR AVERAGE</b>				Error includes scale factor of 1.4.

$5.3 \pm 1.0 \pm 0.3$       370 AUBERT      04H BABR       $e^+e^- \rightarrow \Upsilon(4S)$

$1.2^{+2.8}_{-1.2}$       370 RICHICHI      00 CLE2       $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 15$       90      BEHRENS      98 CLE2      Repl. by RICHICHI 00

$<700$       90      371 ALBRECHT      90B ARG       $e^+e^- \rightarrow \Upsilon(4S)$

370 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

371 ALBRECHT 90B limit assumes equal production of  $B^0\bar{B}^0$  and  $B^+B^-$  at  $\Upsilon(4S)$ .

$\Gamma(\eta'\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{225}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.5 \times 10^{-6}$	90	372 AUBERT	04H BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<7.0 \times 10^{-6}$	90	372 ABE	01M BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$<1.2 \times 10^{-5}$	90	372 AUBERT	01G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$<1.2 \times 10^{-5}$	90	372 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$<3.1 \times 10^{-5}$	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

372 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . $\Gamma(\eta'\rho^+)/\Gamma_{\text{total}}$   $\Gamma_{226}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	90	373 AUBERT,B	04D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<3.3 \times 10^{-5}$	90	373 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$<4.7 \times 10^{-5}$	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

373 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . $\Gamma(\eta\rho^+)/\Gamma_{\text{total}}$   $\Gamma_{227}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-5}$	90	374 AUBERT,B	04D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<1.5 \times 10^{-5}$	90	375 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$<3.2 \times 10^{-5}$	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

374 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .375 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . $\Gamma(\phi\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{228}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.1 \times 10^{-7}$	90	376 AUBERT	04A BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<1.4 \times 10^{-6}$	90	376 AUBERT	01D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$<1.53 \times 10^{-4}$	90	377 ABE	00C SLD	$e^+ e^- \rightarrow Z$
$<0.5 \times 10^{-5}$	90	376 BERGFELD	98 CLE2	

376 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .377 ABE 00C assumes  $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$  and the  $B$  fractions  $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$  and  $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$ . $\Gamma(\phi\rho^+)/\Gamma_{\text{total}}$   $\Gamma_{229}/\Gamma$ 

VALUE	DOCUMENT ID	TECN
$<1.6 \times 10^{-5}$	378 BERGFELD	98 CLE2

378 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . $\Gamma(a_0^0\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{230}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<5.8$	90	379 AUBERT,BE	04 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

379 Assumes equal production of charged and neutral  $B$  mesons from  $\Upsilon(4S)$  decays.

### $\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$ $\Gamma_{231}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.6 \times 10^{-4}$	90	380 ALBRECHT	90B ARG	$e^+e^- \rightarrow \gamma(4S)$
380 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and $B^+\bar{B}^-$ at $\gamma(4S)$ .				

### $\Gamma(\rho^0 a_1(1260)^+)/\Gamma_{\text{total}}$ $\Gamma_{232}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.2 \times 10^{-4}$	90	381 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				

$<6.0 \times 10^{-4}$	90	382 ALBRECHT	90B ARG	$e^+e^- \rightarrow \gamma(4S)$
$<3.2 \times 10^{-3}$	90	381 BEBEK	87 CLEO	$e^+e^- \rightarrow \gamma(4S)$

381 BORTOLETTO 89 reports  $< 5.4 \times 10^{-4}$  assuming the  $\gamma(4S)$  decays 43% to  $B^0\bar{B}^0$ .

We rescale to 50%.

382 ALBRECHT 90B limit assumes equal production of  $B^0\bar{B}^0$  and  $B^+\bar{B}^-$  at  $\gamma(4S)$ .

### $\Gamma(\rho^0 a_2(1320)^+)/\Gamma_{\text{total}}$ $\Gamma_{233}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.2 \times 10^{-4}$	90	383 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				

$<2.6 \times 10^{-3}$	90	384 BEBEK	87 CLEO	$e^+e^- \rightarrow \gamma(4S)$
383 BORTOLETTO 89 reports $< 6.3 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$ . We rescale to 50%.				

384 BEBEK 87 reports  $< 2.3 \times 10^{-3}$  assuming the  $\gamma(4S)$  decays 43% to  $B^0\bar{B}^0$ . We rescale to 50%.

### $\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ $\Gamma_{234}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.3 \times 10^{-3}$	90	385 ALBRECHT	90B ARG	$e^+e^- \rightarrow \gamma(4S)$
385 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and $B^+\bar{B}^-$ at $\gamma(4S)$ .				

### $\Gamma(a_1(1260)^+ a_1(1260)^0)/\Gamma_{\text{total}}$ $\Gamma_{235}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-2}$	90	386 ALBRECHT	90B ARG	$e^+e^- \rightarrow \gamma(4S)$
386 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and $B^+\bar{B}^-$ at $\gamma(4S)$ .				

### $\Gamma(h^+\pi^0)/\Gamma_{\text{total}}$ $\Gamma_{236}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
$(1.6^{+0.6}_{-0.5} \pm 0.36) \times 10^{-5}$	GODANG	98 CLE2	$e^+e^- \rightarrow \gamma(4S)$

### $\Gamma(\omega h^+)/\Gamma_{\text{total}}$ $\Gamma_{237}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$1.38^{+0.27}_{-0.24}$ OUR AVERAGE			

$1.34^{+0.33}_{-0.29} \pm 0.11$

387 LU 02 BELL  $e^+e^- \rightarrow \gamma(4S)$

$1.43^{+0.36}_{-0.32} \pm 0.20$

387 JESSOP 00 CLE2  $e^+e^- \rightarrow \gamma(4S)$

$\bullet \bullet \bullet$  We do not use the following data for averages, fits, limits, etc.  $\bullet \bullet \bullet$

$2.5^{+0.8}_{-0.7} \pm 0.3$       387 BERGFELD    98 CLE2    Repl. by JESSOP 00

387 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

### $\Gamma(h^+ X^0 (\text{Familon})) / \Gamma_{\text{total}}$      $\Gamma_{238}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.9 \times 10^{-5}$	90	388 AMMAR	01B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

388 AMMAR 01B searched for the two-body decay of the  $B$  meson to a massless neutral feebly-interacting particle  $X^0$  such as the familon, the Nambu-Goldstone boson associated with a spontaneously broken global family symmetry.

### $\Gamma(p\bar{p}\pi^+)/\Gamma_{\text{total}}$      $\Gamma_{239}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$3.06^{+0.73}_{-0.62} \pm 0.37$	389,390 WANG	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.7$	90	389,391 ABE	02K BELL	Repl. by WANG 04
$< 500$	90	392 ABREU	95N DLPH	Sup. by ADAM 96D
$< 160$	90	393 BEBEK	89 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
$570 \pm 150 \pm 210$	394 ALBRECHT	88F ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	

389 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

390 The branching fraction for  $M_{p\bar{p}} < 2.85$  is also reported.

391 Explicitly vetoes resonant production of  $p\bar{p}$  from Charmonium states.

392 Assumes a  $B^0, B^-$  production fraction of 0.39 and a  $B_s$  production fraction of 0.12.

393 BEBEK 89 reports  $< 1.4 \times 10^{-4}$  assuming the  $\Upsilon(4S)$  decays 43% to  $B^0\bar{B}^0$ . We rescale to 50%.

394 ALBRECHT 88F reports  $(5.2 \pm 1.4 \pm 1.9) \times 10^{-4}$  assuming the  $\Upsilon(4S)$  decays 45% to  $B^0\bar{B}^0$ . We rescale to 50%.

### $\Gamma(p\bar{p}\pi^+ \text{nonresonant}) / \Gamma_{\text{total}}$      $\Gamma_{240}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.3 \times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

### $\Gamma(p\bar{p}\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$      $\Gamma_{241}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-4}$	90	395 ALBRECHT	88F ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

395 ALBRECHT 88F reports  $< 4.7 \times 10^{-4}$  assuming the  $\Upsilon(4S)$  decays 45% to  $B^0\bar{B}^0$ . We rescale to 50%.

### $\Gamma(p\bar{p}K^+)/\Gamma_{\text{total}}$      $\Gamma_{242}/\Gamma$

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
$5.66^{+0.67}_{-0.57} \pm 0.62$	396,397 WANG	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.3^{+1.1}_{-0.9} \pm 0.5$       398,399 ABE      02K BELL      Repl. by WANG 04

396 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

397 The branching fraction for  $M_{p\bar{p}} < 2.85$  is also reported.

398 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

399 Explicitly vetoes resonant production of  $p\bar{p}$  from Charmonium states.

$\Gamma(p\bar{p}K^+\text{ nonresonant})/\Gamma_{\text{total}}$   $\Gamma_{243}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.9 \times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(p\bar{p}K^*(892)^+)/\Gamma_{\text{total}}$   $\Gamma_{244}/\Gamma$

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
$10.3^{+3.6+1.3}_{-2.8-1.7}$	400,401 WANG	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

400 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

401 The branching fraction for  $M_{p\bar{p}} < 2.85$  is also reported.

$\Gamma(p\bar{\Lambda})/\Gamma_{\text{total}}$   $\Gamma_{245}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-6}$	90	402 BORNHEIM	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.2 \times 10^{-6}$	90	402 ABE	020 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$<2.6 \times 10^{-6}$	90	403 COAN	99 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$<6 \times 10^{-5}$	90	404 Avery	89B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
$<9.3 \times 10^{-5}$	90	405 ALBRECHT	88F ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

402 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

403 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

404 Avery 89B reports  $< 5 \times 10^{-5}$  assuming the  $\Upsilon(4S)$  decays 43% to  $B^0\bar{B}^0$ . We rescale to 50%.

405 ALBRECHT 88F reports  $< 8.5 \times 10^{-5}$  assuming the  $\Upsilon(4S)$  decays 45% to  $B^0\bar{B}^0$ . We rescale to 50%.

$[\Gamma(p\bar{\Lambda}\gamma) + \Gamma(p\bar{\Sigma}\gamma)]/\Gamma_{\text{total}}$   $(\Gamma_{246}+0.3\Gamma_{247})/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.9 \times 10^{-6}$	90	406 EDWARDS	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

406 Corresponds to  $E_\gamma > 1.5$  GeV. The limit changes to  $3.3 \times 10^{-6}$  for  $E_\gamma > 2.0$  GeV.

$[\Gamma(p\bar{\Lambda}\gamma) + \Gamma(p\bar{\Sigma}\gamma)]/\Gamma_{\text{total}}$   $(0.4\Gamma_{246}+\Gamma_{247})/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.9 \times 10^{-6}$	90	407 EDWARDS	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

407 Corresponds to  $E_\gamma > 1.5$  GeV. The limit changes to  $6.4 \times 10^{-6}$  for  $E_\gamma > 2.0$  GeV.

$\Gamma(p\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{248}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-4}$	90	408 ALBRECHT	88F ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

408 ALBRECHT 88F reports  $< 1.8 \times 10^{-4}$  assuming the  $\Upsilon(4S)$  decays 45% to  $B^0\bar{B}^0$ . We rescale to 50%.

$\Gamma(\Lambda\bar{\Lambda}\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{249}/\Gamma$

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
$2.91^{+0.9}_{-0.70} \pm 0.38$	409 LEE	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

409 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ .

$\Gamma(\Lambda\bar{\Lambda}K^+)/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{250}/\Gamma$
<b>&lt;2.8</b>	90	410 LEE	04 BELL	$e^+e^- \rightarrow \Upsilon(4S)$	

410 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . | $\Gamma(\Delta^0 p)/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{251}/\Gamma$
<b>&lt;3.8 × 10<sup>-4</sup></b>	90	411 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

411 BORTOLETTO 89 reports  $< 3.3 \times 10^{-4}$  assuming the  $\Upsilon(4S)$  decays 43% to  $B^0\bar{B}^0$ . We rescale to 50%. | $\Gamma(\Delta^{++}\bar{p})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{252}/\Gamma$
<b>&lt;1.5 × 10<sup>-4</sup></b>	90	412 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

412 BORTOLETTO 89 reports  $< 1.3 \times 10^{-4}$  assuming the  $\Upsilon(4S)$  decays 43% to  $B^0\bar{B}^0$ . We rescale to 50%. | $\Gamma(D^+ p\bar{p})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{253}/\Gamma$
<b>&lt;1.5 × 10<sup>-5</sup></b>	90	413 ABE	02W BELL	$e^+e^- \rightarrow \Upsilon(4S)$	

413 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . | $\Gamma(D^*(2010)^+ p\bar{p})/\Gamma_{\text{total}}$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{254}/\Gamma$
<b>&lt;1.5 × 10<sup>-5</sup></b>	90	414 ABE	02W BELL	$e^+e^- \rightarrow \Upsilon(4S)$	

414 Assumes equal production of  $B^+$  and  $B^0$  at the  $\Upsilon(4S)$ . | $\Gamma(\bar{\Lambda}_c^- p\pi^+)/\Gamma_{\text{total}}$ 

<u>VALUE</u> (units $10^{-4}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{255}/\Gamma$
---------------------------------	--------------------	-------------	----------------	-----------------------

**2.1±0.7 OUR AVERAGE**

2.4±0.6±0.6	415 DYTMAN	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
1.9±0.5±0.5	416 GABYSHEV	02 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.2 <sup>+2.3</sup> <sub>-2.0</sub> ±1.6	417 FU	97 CLE2	Repl. by DYTMAN 02
--	--------	---------	--------------------

415 DYTMAN 02 reports  $(2.4^{+0.63}_{-0.62}) \times 10^{-4}$  for  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$ . We rescale to our best value  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.416 GABYSHEV 02 reports  $(1.87^{+0.51}_{-0.49}) \times 10^{-4}$  for  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$ . We rescale to our best value  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.417 FU 97 uses PDG 96 values of  $\Lambda_c$  branching fraction.

$\Gamma(\bar{\Lambda}_c^- p \pi^+ \pi^0)/\Gamma_{\text{total}}$		$\Gamma_{256}/\Gamma$		
VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>1.81 \pm 0.29^{+0.52}_{-0.50}</math></b>	418,419	DYTMAN	02	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.12 90 420 FU 97 CLE2  $e^+ e^- \rightarrow \gamma(4S)$

418 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

419 DYTGAN 02 measurement uses  $B(\bar{\Lambda}_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$ . The second error includes the systematic and the uncertainty of the branching ratio.

420 FU 97 uses PDG 96 values of  $\Lambda_c$  branching ratio.

$\Gamma(\bar{\Lambda}_c^- p \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$		$\Gamma_{257}/\Gamma$		
VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>2.25 \pm 0.25^{+0.63}_{-0.61}</math></b>	421,422	DYTMAN	02	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.46 90 423 FU 97 CLE2  $e^+ e^- \rightarrow \gamma(4S)$

421 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

422 DYTGAN 02 measurement uses  $B(\bar{\Lambda}_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$ . The second error includes the systematic and the uncertainty of the branching ratio.

423 FU 97 uses PDG 96 values of  $\Lambda_c$  branching ratio.

$\Gamma(\bar{\Lambda}_c^- p \pi^+ \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$		$\Gamma_{258}/\Gamma$		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;1.34 \times 10^{-2}</math></b>	90	424 FU	97	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

424 FU 97 uses PDG 96 values of  $\Lambda_c$  branching ratio.

$\Gamma(\bar{\Sigma}_c(2455)^0 p)/\Gamma_{\text{total}}$		$\Gamma_{259}/\Gamma$		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;0.8 \times 10^{-4}</math></b>	90	425,426 DYTGAN	02	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< $9.3 \times 10^{-5}$  90 425,427 GABYSHEV 02 BELL  $e^+ e^- \rightarrow \gamma(4S)$

425 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

426 DYTGAN 02 measurement uses  $B(\bar{\Lambda}_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$ . The second error includes the systematic and the uncertainty of the branching ratio.

427 Uses the value for  $\Lambda_c \rightarrow p K^- \pi^+$  branching ratio ( $5.0 \pm 1.3\%$ ).

$\Gamma(\bar{\Sigma}_c(2520)^0 p)/\Gamma_{\text{total}}$		$\Gamma_{260}/\Gamma$		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;4.6 \times 10^{-5}</math></b>	90	428,429 GABYSHEV	02	BELL $e^+ e^- \rightarrow \gamma(4S)$

428 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

429 Uses the value for  $\Lambda_c \rightarrow p K^- \pi^+$  branching ratio ( $5.0 \pm 1.3\%$ ).

$\Gamma(\bar{\Sigma}_c(2455)^0 p \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{261}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.4±1.4±1.1</b>	430,431 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

430 DYTMAN 02 reports  $(4.4 \pm 1.4) \times 10^{-4}$  for  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$ . We rescale to our best value  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.  
 431 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(\bar{\Sigma}_c(2455)^0 p \pi^- \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{262}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.4±1.3±1.1</b>	432,433 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

432 DYTMAN 02 reports  $(4.4 \pm 1.3) \times 10^{-4}$  for  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$ . We rescale to our best value  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.  
 433 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(\bar{\Sigma}_c(2455)^{--} p \pi^+ \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{263}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.8±1.0±0.7</b>	434,435 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

434 DYTMAN 02 reports  $(2.8 \pm 1.0) \times 10^{-4}$  for  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$ . We rescale to our best value  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.  
 435 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(\bar{\Lambda}_c(2593)^- / \bar{\Lambda}_c(2625)^- p \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{264}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.9 × 10<sup>-4</sup></b>	90	436,437 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

436 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

437 DYTMAN 02 measurement uses  $B(\Lambda_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$ . The second error includes the systematic and the uncertainty of the branching ratio.

$\Gamma(\pi^+ e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{265}/\Gamma$

Test for  $\Delta B = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.0039</b>	90	438 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$

438 WEIR 90B assumes  $B^+$  production cross section from LUND.

$\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{266}/\Gamma$

Test for  $\Delta B = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.0091</b>	90	439 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$

439 WEIR 90B assumes  $B^+$  production cross section from LUND.

$\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_{267}/\Gamma$ 

Test for  $\Delta B = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u> (units $10^{-7}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

**8.0 $^{+2.2}_{-1.9}$  OUR AVERAGE** Error includes scale factor of 1.4.

$10.5^{+2.5}_{-2.2} \pm 0.7$	440	AUBERT	03U BABR	$e^+ e^- \rightarrow \gamma(4S)$
$6.3^{+1.9}_{-1.7} \pm 0.3$	441	ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 14	90	440 ABE	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$
< 9	90	440 AUBERT	02L BABR	$e^+ e^- \rightarrow \gamma(4S)$
< 24	90	442 ANDERSON	01B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
< 990	90	443 ALBRECHT	91E ARG	$e^+ e^- \rightarrow \gamma(4S)$
< 68000	90	444 WEIR	90B MRK2	$e^+ e^- \rightarrow \gamma(4S)$ 29 GeV
< 600	90	445 Avery	89B CLEO	$e^+ e^- \rightarrow \gamma(4S)$
< 2500	90	446 Avery	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

440 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

441 Assumes equal production of  $B^0$  and  $B^+$  at  $\gamma(4S)$ . The second error is a total of systematic uncertainties including model dependence.

442 The result is for di-lepton masses above 0.5 GeV.

443 ALBRECHT 91E reports  $< 9.0 \times 10^{-5}$  assuming the  $\gamma(4S)$  decays 45% to  $B^0 \bar{B}^0$ . We rescale to 50%.

444 WEIR 90B assumes  $B^+$  production cross section from LUND.

445 Avery 89B reports  $< 5 \times 10^{-5}$  assuming the  $\gamma(4S)$  decays 43% to  $B^0 \bar{B}^0$ . We rescale to 50%.

446 Avery 87 reports  $< 2.1 \times 10^{-4}$  assuming the  $\gamma(4S)$  decays 40% to  $B^0 \bar{B}^0$ . We rescale to 50%.

 $\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$  $\Gamma_{268}/\Gamma$ 

Test for  $\Delta B = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

**0.34 $^{+0.19}_{-0.14}$  OUR AVERAGE** Error includes scale factor of 1.7.

$0.07^{+0.19}_{-0.11} \pm 0.02$	447	AUBERT	03U BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.45^{+0.14}_{-0.12} \pm 0.03$	448	ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.98^{+0.46}_{-0.36} \pm 0.16$	447 ABE	02 BELL	Repl. by ISHIKAWA 03
< 1.2	90	447 AUBERT	02L BABR
< 3.68	90	449 ANDERSON	01B CLE2
< 5.2	90	450 AFFOLDER	99B CDF
< 10	90	451 ABE	96L CDF
			Repl. by AF-FOLDER 99B
< 240	90	452 ALBRECHT	91E ARG
< 6400	90	453 WEIR	90B MRK2
< 170	90	454 Avery	89B CLEO
< 380	90	455 Avery	87 CLEO

- 447 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .  
 448 Assumes equal production of  $B^0$  and  $B^+$  at  $\gamma(4S)$ . The second error is a total of systematic uncertainties including model dependence.  
 449 The result is for di-lepton masses above 0.5 GeV.  
 450 AFFOLDER 99B measured relative to  $B^+ \rightarrow J/\psi(1S) K^+$ .  
 451 ABE 96L measured relative to  $B^+ \rightarrow J/\psi(1S) K^+$  using PDG 94 branching ratios.  
 452 ALBRECHT 91E reports  $< 2.2 \times 10^{-4}$  assuming the  $\gamma(4S)$  decays 45% to  $B^0 \bar{B}^0$ . We rescale to 50%.  
 453 WEIR 90B assumes  $B^+$  production cross section from LUND.  
 454 Avery 89B reports  $< 1.5 \times 10^{-4}$  assuming the  $\gamma(4S)$  decays 43% to  $B^0 \bar{B}^0$ . We rescale to 50%.  
 455 Avery 87 reports  $< 3.2 \times 10^{-4}$  assuming the  $\gamma(4S)$  decays 40% to  $B^0 \bar{B}^0$ . We rescale to 50%.

$\Gamma(K^+ \ell^+ \ell^-)/\Gamma_{\text{total}}$	$\Gamma_{269}/\Gamma$		
VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>5.3^{+1.1}_{-1.0} \pm 0.3</math></b>	456 ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$

456 Assumes equal production of  $B^0$  and  $B^+$  at  $\gamma(4S)$ .

$\Gamma(K^+ \bar{\nu}\nu)/\Gamma_{\text{total}}$	$\Gamma_{270}/\Gamma$			
TEST FOR $\Delta B = 1$ WEAK NEUTRAL CURRENT. ALLOWED BY HIGHER-ORDER ELECTROWEAK INTERACTIONS.				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.4 \times 10^{-4}$	90	457 BROWDER	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

457 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(K^*(892)^+ e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{271}/\Gamma$			
TEST FOR $\Delta B = 1$ WEAK NEUTRAL CURRENT. ALLOWED BY HIGHER-ORDER ELECTROWEAK INTERACTIONS.				
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.6$	90	458 ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
$0.20^{+1.34}_{-0.87} \pm 0.28$		459 AUBERT	03U BABR	$e^+ e^- \rightarrow \gamma(4S)$
$< 8.9$	90	459 ABE	02 BELL	Repl. by ISHIKAWA 03
$< 9.5$	90	459 AUBERT	02L BABR	$e^+ e^- \rightarrow \gamma(4S)$
$< 690$	90	460 ALBRECHT	91E ARG	$e^+ e^- \rightarrow \gamma(4S)$

458 Assumes equal production of  $B^0$  and  $B^+$  at  $\gamma(4S)$ . The second error is a total of systematic uncertainties including model dependence.

459 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

460 ALBRECHT 91E reports  $< 6.3 \times 10^{-4}$  assuming the  $\gamma(4S)$  decays 45% to  $B^0 \bar{B}^0$ . We rescale to 50%.

$\Gamma(K^*(892)^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$	$\Gamma_{272}/\Gamma$			
TEST FOR $\Delta B = 1$ WEAK NEUTRAL CURRENT. ALLOWED BY HIGHER-ORDER ELECTROWEAK INTERACTIONS.				
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.2$	90	461 ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.07^{+2.58}_{-1.78} \pm 0.42$	462	AUBERT	03U BABR	$e^+ e^- \rightarrow \gamma(4S)$
< 3.9	90	462	ABE	02 BELL Repl. by ISHIKAWA 03
< 17.0	90	462	AUBERT	02L BABR $e^+ e^- \rightarrow \gamma(4S)$
<1200	90	463	ALBRECHT	91E ARG $e^+ e^- \rightarrow \gamma(4S)$

461 Assumes equal production of  $B^0$  and  $B^+$  at  $\gamma(4S)$ . The second error is a total of systematic uncertainties including model dependence.

462 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

463 ALBRECHT 91E reports  $< 1.1 \times 10^{-3}$  assuming the  $\gamma(4S)$  decays 45% to  $B^0 \bar{B}^0$ . We rescale to 50%.

### $\Gamma(K^*(892)^+ \ell^+ \ell^-)/\Gamma_{\text{total}}$

### $\Gamma_{273}/\Gamma$

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<22	90	464	ISHIKAWA	03 BELL $e^+ e^- \rightarrow \gamma(4S)$

464 Assumes equal production of  $B^0$  and  $B^+$  at  $\gamma(4S)$ .

### $\Gamma(\pi^+ e^+ \mu^-)/\Gamma_{\text{total}}$

### $\Gamma_{274}/\Gamma$

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	465	WEIR	90B MRK2 $e^+ e^-$ 29 GeV

465 WEIR 90B assumes  $B^+$  production cross section from LUND.

### $\Gamma(\pi^+ e^- \mu^+)/\Gamma_{\text{total}}$

### $\Gamma_{275}/\Gamma$

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	466	WEIR	90B MRK2 $e^+ e^-$ 29 GeV

466 WEIR 90B assumes  $B^+$  production cross section from LUND.

### $\Gamma(K^+ e^+ \mu^-)/\Gamma_{\text{total}}$

### $\Gamma_{276}/\Gamma$

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.8 \times 10^{-6}$	90	467	AUBERT	02L BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0064 90 468 WEIR 90B MRK2  $e^+ e^-$  29 GeV

467 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

468 WEIR 90B assumes  $B^+$  production cross section from LUND.

### $\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$

### $\Gamma_{277}/\Gamma$

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	469	WEIR	90B MRK2 $e^+ e^-$ 29 GeV

469 WEIR 90B assumes  $B^+$  production cross section from LUND.

### $\Gamma(K^*(892)^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$

### $\Gamma_{278}/\Gamma$

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.9 \times 10^{-6}$	90	470	AUBERT	02L BABR $e^+ e^- \rightarrow \gamma(4S)$

470 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(\pi^- e^+ e^+)/\Gamma_{\text{total}}$  $\Gamma_{279}/\Gamma$ 

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.6 \times 10^{-6}$	90	471 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.0039$	90	472 WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-----------	----	----------	----------	------------------

471 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .472 WEIR 90B assumes  $B^+$  production cross section from LUND. $\Gamma(\pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$  $\Gamma_{280}/\Gamma$ 

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-6}$	90	473 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.0091$	90	474 WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-----------	----	----------	----------	------------------

473 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .474 WEIR 90B assumes  $B^+$  production cross section from LUND. $\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$  $\Gamma_{281}/\Gamma$ 

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-6}$	90	475 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.0064$	90	476 WEIR	90B MRK2	$e^+ e^-$ 29 GeV
-----------	----	----------	----------	------------------

475 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .476 WEIR 90B assumes  $B^+$  production cross section from LUND. $\Gamma(\rho^- e^+ e^+)/\Gamma_{\text{total}}$  $\Gamma_{282}/\Gamma$ 

Test of total lepton number conservation.

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<2.6$	90	477 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

477 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . $\Gamma(\rho^- \mu^+ \mu^+)/\Gamma_{\text{total}}$  $\Gamma_{283}/\Gamma$ 

Test of total lepton number conservation.

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<5.0$	90	478 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

478 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . $\Gamma(\rho^- e^+ \mu^+)/\Gamma_{\text{total}}$  $\Gamma_{284}/\Gamma$ 

Test of lepton family number conservation.

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<3.3$	90	479 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

479 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

$\Gamma(K^- e^+ e^+)/\Gamma_{\text{total}}$  $\Gamma_{285}/\Gamma$ 

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-6}$	90	480 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0039	90	481 WEIR	90B MRK2	$e^+ e^-$ 29 GeV
---------	----	----------	----------	------------------

480 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .481 WEIR 90B assumes  $B^+$  production cross section from LUND. $\Gamma(K^- \mu^+ \mu^+)/\Gamma_{\text{total}}$  $\Gamma_{286}/\Gamma$ 

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	90	482 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0091	90	483 WEIR	90B MRK2	$e^+ e^-$ 29 GeV
---------	----	----------	----------	------------------

482 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .483 WEIR 90B assumes  $B^+$  production cross section from LUND. $\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$  $\Gamma_{287}/\Gamma$ 

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-6}$	90	484 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0064	90	485 WEIR	90B MRK2	$e^+ e^-$ 29 GeV
---------	----	----------	----------	------------------

484 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .485 WEIR 90B assumes  $B^+$  production cross section from LUND. $\Gamma(K^*(892)^- e^+ e^+)/\Gamma_{\text{total}}$  $\Gamma_{288}/\Gamma$ 

Test of total lepton number conservation.

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.8	90	486 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

486 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . $\Gamma(K^*(892)^- \mu^+ \mu^+)/\Gamma_{\text{total}}$  $\Gamma_{289}/\Gamma$ 

Test of total lepton number conservation.

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<8.3	90	487 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

487 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ . $\Gamma(K^*(892)^- e^+ \mu^+)/\Gamma_{\text{total}}$  $\Gamma_{290}/\Gamma$ 

Test of lepton family number conservation.

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<4.4	90	488 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

488 Assumes equal production of  $B^+$  and  $B^0$  at the  $\gamma(4S)$ .

## POLARIZATION IN $B^+$ DECAY

### $\Gamma_L / \Gamma$ in $B^+ \rightarrow \bar{D}^{*0} \rho^+$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.892±0.018±0.016</b>	CSORNA	03	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

### $\Gamma_L / \Gamma$ in $B^+ \rightarrow \bar{D}^{*0} K^{*+}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.86±0.06±0.03</b>	AUBERT	04K	BABR $e^+ e^- \rightarrow \gamma(4S)$

### $\Gamma_L / \Gamma$ in $B^+ \rightarrow \phi K^*(892)^+$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.46±0.12±0.03</b>	AUBERT	03v	BABR $e^+ e^- \rightarrow \gamma(4S)$

### $\Gamma_L / \Gamma$ in $B^+ \rightarrow \rho^0 K^*(892)^+$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.96±0.04±0.04</b>	AUBERT	03v	BABR $e^+ e^- \rightarrow \gamma(4S)$

### $\Gamma_L / \Gamma$ in $B^+ \rightarrow \rho^+ \rho^0$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.96 ± 0.05</b> OUR AVERAGE			
0.97 ± 0.03 ± 0.04	AUBERT	03v	BABR $e^+ e^- \rightarrow \gamma(4S)$
0.948±0.106±0.021	ZHANG	03B	BELL $e^+ e^- \rightarrow \gamma(4S)$

## **CP VIOLATION**

$A_{CP}$  is defined as

$$\frac{B(B^- \rightarrow \bar{f}) - B(B^+ \rightarrow f)}{B(B^- \rightarrow \bar{f}) + B(B^+ \rightarrow f)},$$

the  $CP$ -violation charge asymmetry of exclusive  $B^-$  and  $B^+$  decay.

### $A_{CP}(B^+ \rightarrow J/\psi(1S) K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.016±0.016 OUR AVERAGE</b>			Error includes scale factor of 1.2.
0.03 ± 0.015 ± 0.006	AUBERT	04P	BABR $e^+ e^- \rightarrow \gamma(4S)$
-0.026±0.022±0.017	ABE	03B	BELL $e^+ e^- \rightarrow \gamma(4S)$
0.018±0.043±0.004	489 BONVICINI	00	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.003±0.030±0.004                  AUBERT    02F   BABR   Repl. by AUBERT 04P

489 A +0.3% correction is applied due to a slightly higher reconstruction efficiency for the positive kaons.

### $A_{CP}(B^+ \rightarrow J/\psi(1S)\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.09 ± 0.08 OUR AVERAGE</b>			
0.123 ± 0.085 ± 0.004	AUBERT	04P BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.023 ± 0.164 ± 0.015	ABE	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.01 ± 0.22 ± 0.01	AUBERT	02F BABR	Repl. by AUBERT 04P

### $A_{CP}(B^+ \rightarrow \psi(2S)K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.037 ± 0.025 OUR AVERAGE</b>			
-0.042 ± 0.020 ± 0.017	ABE	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.02 ± 0.091 ± 0.01	490 BONVICINI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

490 A + 0.3% correction is applied due to a slightly higher reconstruction efficiency for the positive kaons.

### $A_{CP}(B^+ \rightarrow \bar{D}^0 K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.04 ± 0.06 ± 0.03</b>	491 SWAIN	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.003 ± 0.080 ± 0.037	492 ABE	03D BELL	Repl. by SWAIN 03
491 Corresponds to 90% confidence range $-0.07 < A_{CP} < 0.15$ .			
492 Corresponds to 90% confidence range $-0.15 < A_{CP} < 0.16$ .			

### $A_{CP}(B^+ \rightarrow D_{CP(+1)} K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.07 ± 0.13 OUR AVERAGE</b>			
0.07 ± 0.17 ± 0.06	AUBERT	04N BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.06 ± 0.19 ± 0.04	493 SWAIN	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.29 ± 0.26 ± 0.05	494 ABE	03D BELL	Repl. by SWAIN 03
493 Corresponds to 90% confidence range $-0.26 < A_{CP} < 0.38$ .			
494 Corresponds to 90% confidence range $-0.14 < A_{CP} < 0.73$ .			

### $A_{CP}(B^+ \rightarrow D_{CP(-1)} K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.19 ± 0.17 ± 0.05</b>	495 SWAIN	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.22 ± 0.24 ± 0.04	496 ABE	03D BELL	Repl. by SWAIN 03
495 Corresponds to 90% confidence range $-0.47 < A_{CP} < 0.11$ .			
496 Corresponds to 90% confidence range $-0.62 < A_{CP} < 0.18$ .			

**$A_{CP}(B^+ \rightarrow \pi^+\pi^0)$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.02 ± 0.09 OUR AVERAGE</b>			
-0.02 ± 0.10 ± 0.01	497 CHAO	04B BELL	$e^+e^- \rightarrow \gamma(4S)$
-0.03 <sup>+0.18</sup> <sub>-0.17</sub> ± 0.02	498 AUBERT	03L BABR	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.30 ± 0.30 <sup>+0.06</sup> <sub>-0.04</sub>	499 CASEY	02 BELL	Repl. by CHAO 04B

497 This corresponds to 90% CL interval of  $-0.18 < A_{CP} < 0.14$ .498 Corresponds to 90% confidence range  $-0.32 < A_{CP} < 0.27$ .499 Corresponds to 90% confidence range  $-0.23 < A_{CP} < +0.86$ . **$A_{CP}(B^+ \rightarrow K^+\pi^0)$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.01 ± 0.06 OUR AVERAGE</b>			
			Error includes scale factor of 1.2.
0.04 ± 0.05 ± 0.02	500 CHAO	04B BELL	$e^+e^- \rightarrow \gamma(4S)$
-0.09 ± 0.09 ± 0.01	501 AUBERT	03L BABR	$e^+e^- \rightarrow \gamma(4S)$
-0.29 ± 0.23	502 CHEN	00 CLE2	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.02 ± 0.19 ± 0.02	503 CASEY	02 BELL	Repl. by CHAO 04B
-0.059 <sup>+0.222</sup> <sub>-0.196</sub> <sup>+0.055</sup> <sub>-0.017</sub>	504 ABE	01K BELL	Repl. by CASEY 02
0.00 ± 0.18 ± 0.04	505 AUBERT	01E BABR	Repl. by AUBERT 03L
500 Corresponds to 90% CL interval of $-0.05 < A_{CP} < 0.13$ .			
501 Corresponds to 90% confidence range $-0.24 < A_{CP} < 0.06$ .			
502 Corresponds to 90% confidence range $-0.67 < A_{CP} < 0.09$ .			
503 Corresponds to 90% confidence range $-0.35 < A_{CP} < +0.30$ .			
504 Corresponds to 90% confidence range $-0.40 < A_{CP} < 0.36$ .			
505 Corresponds to 90% confidence range $-0.30 < A_{CP} < +0.30$ .			

 **$A_{CP}(B^+ \rightarrow K_S^0\pi^+)$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.02 ± 0.06 OUR AVERAGE</b>			
-0.05 ± 0.08 ± 0.01	506 AUBERT	04M BABR	$e^+e^- \rightarrow \gamma(4S)$
0.07 <sup>+0.09</sup> <sub>-0.08</sub> <sup>+0.01</sup> <sub>-0.03</sub>	507 UNNO	03 BELL	$e^+e^- \rightarrow \gamma(4S)$
0.18 ± 0.24	508 CHEN	00 CLE2	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.46 ± 0.15 ± 0.02	509 CASEY	02 BELL	Repl. by UNNO 03
0.098 <sup>+0.430</sup> <sub>-0.343</sub> <sup>+0.020</sup> <sub>-0.063</sub>	510 ABE	01K BELL	Repl. by CASEY 02
-0.21 ± 0.18 ± 0.03	511 AUBERT	01E BABR	Repl. by AUBERT 04M
506 90% CL interval $-0.18 < A_{CP} < 0.08$			
507 Corresponds to 90% confidence range $-0.10 < A_{CP} < +0.22$ .			
508 Corresponds to 90% confidence range $-0.22 < A_{CP} < 0.56$ .			
509 Corresponds to 90% confidence range $+0.19 < A_{CP} < +0.72$ .			
510 Corresponds to 90% confidence range $-0.53 < A_{CP} < 0.82$ .			
511 Corresponds to 90% confidence range $-0.51 < A_{CP} < 0.09$ .			

$A_{CP}(B^+ \rightarrow \pi^+ \pi^- \pi^+)$

VALUE

**-0.39±0.33±0.12**

$A_{CP}(B^+ \rightarrow \rho^0 \pi^+)$

VALUE

**-0.19±0.11±0.02**

$A_{CP}(B^+ \rightarrow \rho^+ \pi^0)$

VALUE

**0.24±0.16±0.06**

$A_{CP}(B^+ \rightarrow \rho^+ \rho^0)$

VALUE

**-0.09±0.16 OUR AVERAGE**

$-0.19\pm0.23\pm0.03$

$0.00\pm0.22\pm0.03$

$A_{CP}(B^+ \rightarrow K^+ \pi^- \pi^+)$

VALUE

**0.01±0.07±0.03**

$A_{CP}(B^+ \rightarrow K^+ K^- K^+)$

VALUE

**0.02±0.07±0.03**

$A_{CP}(B^+ \rightarrow K^+ K_S^0 K_S^0)$

VALUE

**-0.04±0.11±0.02**

512 Corresponds to 90% confidence range  $-0.23 < A_{CP} < 0.15$ .

$A_{CP}(B^+ \rightarrow \eta \pi^+)$

VALUE

**-0.44±0.18±0.01**

$A_{CP}(B^+ \rightarrow \eta K^+)$

VALUE

**-0.52±0.24±0.01**

$A_{CP}(B^+ \rightarrow \eta K^*(892)^+)$

VALUE

**0.13±0.14±0.02**

DOCUMENT ID	TECN	COMMENT
AUBERT	03M BABR	$e^+ e^- \rightarrow \gamma(4S)$

DOCUMENT ID	TECN	COMMENT
AUBERT	04Z BABR	$e^+ e^- \rightarrow \gamma(4S)$

DOCUMENT ID	TECN	COMMENT
AUBERT	04Z BABR	$e^+ e^- \rightarrow \gamma(4S)$

DOCUMENT ID	TECN	COMMENT
AUBERT	03V BABR	$e^+ e^- \rightarrow \gamma(4S)$
ZHANG	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$

DOCUMENT ID	TECN	COMMENT
AUBERT	03M BABR	$e^+ e^- \rightarrow \gamma(4S)$

DOCUMENT ID	TECN	COMMENT
AUBERT	03M BABR	$e^+ e^- \rightarrow \gamma(4S)$

DOCUMENT ID	TECN	COMMENT
512 AUBERT,B	04V BABR	$e^+ e^- \rightarrow \gamma(4S)$

DOCUMENT ID	TECN	COMMENT
AUBERT	04H BABR	$e^+ e^- \rightarrow \gamma(4S)$

DOCUMENT ID	TECN	COMMENT
AUBERT	04H BABR	$e^+ e^- \rightarrow \gamma(4S)$


DOCUMENT ID	TECN	COMMENT
AUBERT,B	04D BABR	$e^+ e^- \rightarrow \gamma(4S)$

### $A_{CP}(B^+ \rightarrow K^+ \eta')$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.009±0.035 OUR AVERAGE</b>			
0.037±0.045±0.011	513 AUBERT	03W BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.11 ± 0.11 ± 0.02	514 AUBERT	02E BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.015±0.070±0.009	515 CHEN	02B BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.03 ± 0.12	516 CHEN	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.06 ± 0.15 ± 0.01	517 ABE	01M BELL	Repl. by CHEN 02B
513	Corresponds to 90% confidence range $-0.04 < A_{CP} < 0.11$ .		
514	Corresponds to 90% confidence range $-0.28 < A_{CP} < 0.07$ .		
515	Corresponds to 90% confidence range $-0.13 < A_{CP} < 0.10$ .		
516	Corresponds to 90% confidence range $-0.17 < A_{CP} < 0.23$ .		
517	Corresponds to 90% confidence range $-0.20 < A_{CP} < 0.32$ .		

### $A_{CP}(B^+ \rightarrow \omega\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.10±0.22 OUR AVERAGE</b>			
			Error includes scale factor of 1.9. See the ideogram below.
0.03±0.16±0.01	AUBERT	04H BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.50^{+0.23}_{-0.20} \pm 0.02$	518 WANG	04A BELL	$e^+ e^- \rightarrow \gamma(4S)$
-0.34±0.25	519 CHEN	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$-0.01^{+0.29}_{-0.31} \pm 0.03$	520 AUBERT	02E BABR	Repl. by AUBERT 04H
518	Corresponds to 90% CL interval $-0.25 < A_{CP} < 0.41$		
519	Corresponds to 90% confidence range $-0.75 < A_{CP} < 0.07$ .		
520	Corresponds to 90% confidence range $-0.50 < A_{CP} < 0.46$ .		



### $A_{CP}(B^+ \rightarrow \omega K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.02±0.13 OUR AVERAGE</b>			
-0.09±0.17±0.01	AUBERT	04H BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.06^{+0.21}_{-0.18} \pm 0.01$	521 WANG	04A BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.21±0.28±0.03                    522 LU                    02 BELL    Repl. by WANG 04A

521 Corresponds to 90% CL interval  $0.15 < A_{CP} < 0.90$

522 Corresponds to 90% confidence range  $-0.70 < A_{CP} < +0.38$ .

### $A_{CP}(B^+ \rightarrow \phi K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.03±0.07 OUR AVERAGE</b>			
0.04±0.09±0.01	523 AUBERT	04A BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.01^{+0.12}_{-0.05} \pm 0.05$	524 CHEN	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.05±0.20±0.03                    525 AUBERT                    02E BABR     $e^+ e^- \rightarrow \gamma(4S)$

523 Corresponds to 90% confidence range  $-0.10 < A_{CP} < 0.18$ .

524 Corresponds to 90% confidence range  $-0.20 < A_{CP} < 0.22$ .

525 Corresponds to 90% confidence range  $-0.37 < A_{CP} < 0.28$ .

### $A_{CP}(B^+ \rightarrow \phi K^*(892)^+)$

VALUE  
**0.09±0.15 OUR AVERAGE**

	DOCUMENT ID	TECN	COMMENT
0.16±0.17±0.03	AUBERT	03V BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.13±0.29±0.11	526 CHEN	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
-0.43±0.36±0.06	527 AUBERT	02E BABR	Repl. by AUBERT 03V

526 Corresponds to 90% confidence range  $-0.64 < A_{CP} < 0.36$ .

527 Corresponds to 90% confidence range  $-0.88 < A_{CP} < 0.18$ .

### $A_{CP}(B^+ \rightarrow \rho^0 K^*(892)^+)$

VALUE

**0.20±0.32±0.04**

DOCUMENT ID	TECN	COMMENT
AUBERT	03V BABR	$e^+ e^- \rightarrow \gamma(4S)$

### $A_{CP}(B^+ \rightarrow p\bar{p}\pi^+)$

VALUE

**-0.16±0.22±0.01**

DOCUMENT ID	TECN	COMMENT
WANG	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$

### $A_{CP}(B^+ \rightarrow p\bar{p}K^+)$

VALUE

**-0.05±0.11±0.01**

DOCUMENT ID	TECN	COMMENT
WANG	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$

### $\gamma(B^+ \rightarrow D^{(*)}K^+)$

For angle  $\gamma(\phi_3)$  of the CKM unitarity triangle, see the review on “CP Violation” in the Reviews section.

VALUE (°)

**77±17±17**

DOCUMENT ID	TECN	COMMENT
528 POLUEKTOV	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$

528 Uses a Dalitz plot analysis of the 3-body  $D \rightarrow K_S^0 \pi^+ \pi^-$  decays coming from  $B^\pm \rightarrow D K^\pm$  and  $B^\pm \rightarrow D^* K^\pm$  followed by  $D^* \rightarrow D \pi^0$ ; here we use  $D$  to denote that the neutral  $D$  meson produced in the decay is an admixture of  $D^0$  and  $\bar{D}^0$ . The corresponding 2 standard deviation interval for  $\gamma$  is  $26^\circ < \gamma < 126^\circ$ . POLUEKTOV 04 also reports the amplitude ratios and the strong phases.

## $B^\pm$ REFERENCES

ABE	05B	PR D71 072003	K. Abe <i>et al.</i>	(BELLE Collab.)
Also	05C	PR D71 079903 (Erratum)	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	05	PRL 94 011801	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABDALLAH	04E	EPJ C33 307	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	04A	PR D69 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04C	PRL 92 111801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	04H	PRL 92 061801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04K	PRL 92 141801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04M	PRL 92 201802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04N	PRL 92 202002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04O	PRL 92 221803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04P	PRL 92 241802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04Q	PR D69 051101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04T	PR D69 071103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04Y	PRL 93 041801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	04Z	PRL 93 051802	B. Aubert <i>et al.</i>	(BABAR Collab.)

AUBERT,B	04B	PR D70 011101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04D	PR D70 032006	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04L	PRL 93 131804	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04P	PR D70 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04U	PR D70 091105R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04V	PRL 93 181805	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	04	PR D70 111102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	04A	PR D70 112006	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	04B	PR D70 091106	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHAO	04	PR D69 111102R	Y. Chao <i>et al.</i>	(BELLE Collab.)
CHAO	04B	PRL 93 191802	Y. Chao <i>et al.</i>	(BELLE Collab.)
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	04	PRL 92 051801	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
GARMASH	04	PR D69 012001	A. Garmash <i>et al.</i>	(BELLE Collab.)
LEE	04	PRL 93 211801	Y.-J. Lee <i>et al.</i>	(BELLE Collab.)
MAJUMDER	04	PR D70 111103R	G. Majumder <i>et al.</i>	(BELLE Collab.)
NAKAO	04	PR D69 112001	M. Nakao <i>et al.</i>	(BELLE Collab.)
POLUEKTOV	04	PR D70 072003	A. Poluektov <i>et al.</i>	(BELLE Collab.)
SCHWANDA	04	PRL 93 131803	C. Schwanda <i>et al.</i>	(BELLE Collab.)
WANG	04	PRL 92 131801	M.Z. Wang <i>et al.</i>	(BELLE Collab.)
WANG	04A	PR D70 012001	C.H. Wang <i>et al.</i>	(BELLE Collab.)
ZANG	04	PR D69 017101	S.L. Zang <i>et al.</i>	(BELLE Collab.)
ABE	03B	PR D67 032003	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	03D	PRL 90 131803	K. Abe <i>et al.</i>	(BELLE Collab.)
ADAM	03	PR D67 032001	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ADAM	03B	PR D68 012004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ATHAR	03	PR D68 072003	S.B. Athar <i>et al.</i>	(CLEO Collab.)
AUBERT	03K	PRL 90 231801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03L	PRL 91 021801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03M	PRL 91 051801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03O	PRL 91 071801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03U	PRL 91 221802	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03V	PRL 91 171802	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03W	PRL 91 161801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03X	PR D68 092001	B. Aubert <i>et al.</i>	(BaBar Collab.)
BORNHEIM	03	PR D68 052002	A. Bornheim <i>et al.</i>	(CLEO Collab.)
CHEN	03B	PRL 91 201801	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
CHOI	03	PRL 91 262001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
CSORNA	03	PR D67 112002	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
EDWARDS	03	PR D68 011102R	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
FANG	03	PRL 90 071801	F. Fang <i>et al.</i>	(BELLE Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
ISHIKAWA	03	PRL 91 261601	A. Ishikawa <i>et al.</i>	(BELLE Collab.)
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)
SWAIN	03	PR D68 051101R	S.K. Swain <i>et al.</i>	(BELLE Collab.)
UNNO	03	PR D68 011103R	Y. Unno <i>et al.</i>	(BELLE Collab.)
ZHANG	03B	PRL 91 221801	J. Zhang <i>et al.</i>	(BELLE Collab.)
ABE	02	PRL 88 021801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02B	PRL 88 031802	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02H	PRL 88 171801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02K	PRL 88 181803	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02N	PL B538 11	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02O	PR D65 091103R	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02W	PRL 89 151802	K. Abe <i>et al.</i>	(BELLE Collab.)
ACOSTA	02C	PR D65 092009	D. Acosta <i>et al.</i>	(CDF Collab.)
ACOSTA	02F	PR D66 052005	D. Acosta <i>et al.</i>	(CDF Collab.)
AHMED	02B	PR D66 031101R	S. Ahmed <i>et al.</i>	(CLEO Collab.)
AUBERT	02	PR D65 032001	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02C	PRL 88 101805	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02E	PR D65 051101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02F	PR D65 091101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02L	PRL 88 241801	B. Aubert <i>et al.</i>	(BaBar Collab.)
BRIERE	02	PRL 89 081803	R. Briere <i>et al.</i>	(CLEO Collab.)
CASEY	02	PR D66 092002	B.C.K. Casey <i>et al.</i>	(BELLE Collab.)
CHEN	02B	PL B546 196	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	02	PL B542 171	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
DYTMAN	02	PR D66 091101R	S.A. Dytmam <i>et al.</i>	(CLEO Collab.)
ECKHART	02	PRL 89 251801	E. Eckhart <i>et al.</i>	(CLEO Collab.)
EDWARDS	02B	PR D65 111102R	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
GABYSHEV	02	PR D66 091102R	N. Gabyshev <i>et al.</i>	(BELLE Collab.)

GARMASH	02	PR D65 092005	A. Garmash <i>et al.</i>	(BELLE Collab.)
GODANG	02	PRL 88 021802	R. Godang <i>et al.</i>	(CLEO Collab.)
GORDON	02	PL B542 183	A. Gordon <i>et al.</i>	(BELLE Collab.)
LU	02	PRL 89 191801	R.-S. Lu <i>et al.</i>	(BELLE Collab.)
MAHAPATRA	02	PRL 88 101803	R. Mahapatra <i>et al.</i>	(CLEO Collab.)
NISHIDA	02	PRL 89 231801	S. Nishida <i>et al.</i>	(BELLE Collab.)
ABE	01H	PRL 87 101801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01I	PRL 87 111801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01K	PR D64 071101	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01L	PRL 87 161601	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01M	PL B517 309	K. Abe <i>et al.</i>	(BELLE Collab.)
ALEXANDER	01B	PR D64 092001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AMMAR	01B	PRL 87 271801	R. Ammar <i>et al.</i>	(CLEO Collab.)
ANDERSON	01B	PRL 87 181803	S. Anderson <i>et al.</i>	(CLEO Collab.)
AUBERT	01D	PRL 87 151801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01E	PRL 87 151802	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01F	PRL 87 201803	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01G	PRL 87 221802	B. Aubert <i>et al.</i>	(BaBar Collab.)
BARATE	01E	EPJ C19 213	R. Barate <i>et al.</i>	(ALEPH Collab.)
BRIERE	01	PRL 86 3718	R.A. Biere <i>et al.</i>	(CLEO Collab.)
BROWDER	01	PRL 86 2950	T.E. Browder <i>et al.</i>	(CLEO Collab.)
EDWARDS	01	PRL 86 30	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
GRITSAN	01	PR D64 077501	A. Gritsan <i>et al.</i>	(CLEO Collab.)
RICHICHI	01	PR D63 031103R	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ABBIENDI	00B	PL B476 233	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABE	00C	PR D62 071101R	K. Abe <i>et al.</i>	(SLD Collab.)
AHMED	00B	PR D62 112003	S. Ahmed <i>et al.</i>	(CLEO Collab.)
ANASTASSOV	00	PRL 84 1393	A. Anastassov <i>et al.</i>	(CLEO Collab.)
BARATE	00R	PL B492 275	R. Barate <i>et al.</i>	(ALEPH Collab.)
BEHRENS	00	PR D61 052001	B.H. Behrens <i>et al.</i>	(CLEO Collab.)
BONVICINI	00	PRL 84 5940	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
CHEN	00	PRL 85 525	S. Chen <i>et al.</i>	(CLEO Collab.)
COAN	00	PRL 84 5283	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN...	00	PRL 85 515	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
CSORNA	00	PR D61 111101	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
JESSOP	00	PRL 85 2881	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
RICHICHI	00	PRL 85 520	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ABBIENDI	99J	EPJ C12 609	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AFFOLDER	99B	PRL 83 3378	T. Affolder <i>et al.</i>	(CDF Collab.)
BARTELT	99	PRL 82 3746	J. Bartelt <i>et al.</i>	(CLEO Collab.)
COAN	99	PR D59 111101	T.E. Coan <i>et al.</i>	(CLEO Collab.)
ABE	98B	PR D57 5382	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98O	PR D58 072001	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98Q	PR D58 092002	F. Abe <i>et al.</i>	(CDF Collab.)
ACCIARRI	98S	PL B438 417	M. Acciari <i>et al.</i>	(L3 Collab.)
ANASTASSOV	98	PRL 80 4127	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ATHANAS	98	PRL 80 5493	M. Athanas <i>et al.</i>	(CLEO Collab.)
BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
BEHRENS	98	PRL 80 3710	B.H. Behrens <i>et al.</i>	(CLEO Collab.)
BERGFELD	98	PRL 81 272	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BRANDENB...	98	PRL 80 2762	G. Brandenbrug <i>et al.</i>	(CLEO Collab.)
GODANG	98	PRL 80 3456	R. Godang <i>et al.</i>	(CLEO Collab.)
ABE	97J	PRL 79 590	K. Abe <i>et al.</i>	(SLD Collab.)
ACCIARRI	97F	PL B396 327	M. Acciari <i>et al.</i>	(L3 Collab.)
ARTUSO	97	PL B399 321	M. Artuso <i>et al.</i>	(CLEO Collab.)
ATHANAS	97	PRL 79 2208	M. Athanas <i>et al.</i>	(CLEO Collab.)
BROWDER	97	PR D56 11	T. Browder <i>et al.</i>	(CLEO Collab.)
FU	97	PRL 79 3125	X. Fu <i>et al.</i>	(CLEO Collab.)
JESSOP	97	PRL 79 4533	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
ABE	96B	PR D53 3496	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96C	PRL 76 4462	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96H	PRL 76 2015	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96L	PRL 76 4675	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96Q	PR D54 6596	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96R	PRL 77 5176	F. Abe <i>et al.</i>	(CDF Collab.)
ADAM	96D	ZPHY C72 207	W. Adam <i>et al.</i>	(DELPHI Collab.)
ALEXANDER	96T	PRL 77 5000	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ASNER	96	PR D53 1039	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BERGFELD	96B	PRL 77 4503	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BISHAI	96	PL B369 186	M. Bishai <i>et al.</i>	(CLEO Collab.)

BUSKULIC	96J	ZPHY C71 31	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
GIBAUT	96	PR D53 4734	D. Gibaut <i>et al.</i>	(CLEO Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	
ABREU	95N	PL B357 255	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95Q	ZPHY C68 13	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ADAM	95	ZPHY C68 363	W. Adam <i>et al.</i>	(DELPHI Collab.)
AKERS	95T	ZPHY C67 379	R. Akers <i>et al.</i>	(OPAL Collab.)
ALBRECHT	95D	PL B353 554	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	95	PL B341 435	J. Alexander <i>et al.</i>	(CLEO Collab.)
Also	95C	PL B347 469 (erratum)	J. Alexander <i>et al.</i>	(CLEO Collab.)
ARTUSO	95	PRL 75 785	M. Artuso <i>et al.</i>	(CLEO Collab.)
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	95	PL B343 444	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABE	94D	PRL 72 3456	F. Abe <i>et al.</i>	(CDF Collab.)
ALAM	94	PR D50 43	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	94D	PL B335 526	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ATHANAS	94	PRL 73 3503	M. Athanas <i>et al.</i>	(CLEO Collab.)
Also	95	PRL 74 3090 (erratum)	M. Athanas <i>et al.</i>	(CLEO Collab.)
PDG	94	PR D50 1173	L. Montanet <i>et al.</i>	(CERN, LBL, BOST+)
STONE	94	HEPSY 93-11	S. Stone	
Published in <i>B Decays</i> , 2nd Edition, World Scientific, Singapore				
ABREU	93D	ZPHY C57 181	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	93G	PL B312 253	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	93C	PL B307 247	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ALBRECHT	93E	ZPHY C60 11	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	93B	PL B319 365	J. Alexander <i>et al.</i>	(CLEO Collab.)
AMMAR	93	PRL 71 674	R. Ammar <i>et al.</i>	(CLEO Collab.)
BEAN	93B	PRL 70 2681	A. Bean <i>et al.</i>	(CLEO Collab.)
BUSKULIC	93D	PL B307 194	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
Also	94H	PL B325 537 (errata)	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
SANGHERA	93	PR D47 791	S. Sanghera <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92C	PL B275 195	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92E	PL B277 209	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92G	ZPHY C54 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
BUSKULIC	92G	PL B295 396	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ALBRECHT	91B	PL B254 288	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91C	PL B255 297	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91E	PL B262 148	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BERKELMAN	91	ARNPS 41 1	K. Berkelman, S. Stone	(CORN, SYRA)
"Decays of <i>B</i> Mesons"				
FULTON	91	PR D43 651	R. Fulton <i>et al.</i>	(CLEO Collab.)
ALBRECHT	90B	PL B241 278	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	90J	ZPHY C48 543	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANTREASYAN	90B	ZPHY C48 553	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
BORTOLETTO	90	PRL 64 2117	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
Also	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
WEIR	90B	PR D41 1384	A.J. Weir <i>et al.</i>	(Mark II Collab.)
ALBRECHT	89G	PL B229 304	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AVERY	89B	PL B223 470	P. Avery <i>et al.</i>	(CLEO Collab.)
BEBEK	89	PRL 62 8	C. Bebek <i>et al.</i>	(CLEO Collab.)
BORTOLETTO	89	PRL 62 2436	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
ALBRECHT	88F	PL B209 119	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88K	PL B215 424	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87C	PL B185 218	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87D	PL B199 451	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AVERY	87	PL B183 429	P. Avery <i>et al.</i>	(CLEO Collab.)
BEBEK	87	PR D36 1289	C. Bebek <i>et al.</i>	(CLEO Collab.)
ALAM	86	PR D34 3279	M.S. Alam <i>et al.</i>	(CLEO Collab.)
PDG	86	PL 170B	M. Aguilar-Benitez <i>et al.</i>	(CERN, CIT+)
GILES	84	PR D30 2279	R. Giles <i>et al.</i>	(CLEO Collab.)