

B^\pm/B^0 ADMIXTURE

B DECAY MODES

The branching fraction measurements are for an admixture of B mesons at the $\Upsilon(4S)$. The values quoted assume that $B(\Upsilon(4S) \rightarrow B\bar{B}) = 100\%$.

For inclusive branching fractions, e.g., $B \rightarrow D^\pm$ anything, the values usually are multiplicities, not branching fractions. They can be greater than one.

\bar{B} modes are charge conjugates of the modes below. Reactions indicate the weak decay vertex and do not include mixing.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Semileptonic and leptonic modes		
Γ_1 $B \rightarrow e^+ \nu_e$ anything	[a] $(10.87 \pm 0.17) \%$	
Γ_2 $B \rightarrow \bar{p} e^+ \nu_e$ anything	$< 5.9 \times 10^{-4}$ CL=90%	
Γ_3 $B \rightarrow \mu^+ \nu_\mu$ anything	[a] $(10.87 \pm 0.17) \%$	
Γ_4 $B \rightarrow \ell^+ \nu_\ell$ anything	[a,b] $(10.87 \pm 0.17) \%$	
Γ_5 $B \rightarrow D^- \ell^+ \nu_\ell$ anything	[b] $(2.8 \pm 0.9) \%$	
Γ_6 $B \rightarrow \bar{D}^0 \ell^+ \nu_\ell$ anything	[b] $(7.3 \pm 1.5) \%$	
Γ_7 $B \rightarrow D^{*-} \ell^+ \nu_\ell$ anything	[c] $(6.7 \pm 1.3) \times 10^{-3}$	
Γ_8 $B \rightarrow D^{*0} \ell^+ \nu_\ell$ anything		
Γ_9 $B \rightarrow \bar{D}^{**} \ell^+ \nu_\ell$	[b,d] $(2.7 \pm 0.7) \%$	
Γ_{10} $B \rightarrow \bar{D}_1(2420) \ell^+ \nu_\ell$ anything	$(7.4 \pm 1.6) \times 10^{-3}$	
Γ_{11} $B \rightarrow D \pi \ell^+ \nu_\ell$ anything + $D^* \pi \ell^+ \nu_\ell$ anything	$(2.6 \pm 0.5) \%$	S=1.5
Γ_{12} $B \rightarrow D \pi \ell^+ \nu_\ell$ anything	$(1.5 \pm 0.6) \%$	
Γ_{13} $B \rightarrow D^* \pi \ell^+ \nu_\ell$ anything	$(1.9 \pm 0.4) \%$	
Γ_{14} $B \rightarrow \bar{D}_2^*(2460) \ell^+ \nu_\ell$ anything	$< 6.5 \times 10^{-3}$ CL=95%	
Γ_{15} $B \rightarrow D^{*-} \pi^+ \ell^+ \nu_\ell$ anything	$(1.00 \pm 0.34) \%$	
Γ_{16} $B \rightarrow D_s^- \ell^+ \nu_\ell$ anything	[b] $< 9 \times 10^{-3}$ CL=90%	
Γ_{17} $B \rightarrow D_s^- \ell^+ \nu_\ell K^+$ anything	[b] $< 6 \times 10^{-3}$ CL=90%	
Γ_{18} $B \rightarrow D_s^- \ell^+ \nu_\ell K^0$ anything	[b] $< 9 \times 10^{-3}$ CL=90%	
Γ_{19} $B \rightarrow \ell^+ \nu_\ell$ charm	$(10.61 \pm 0.17) \%$	
Γ_{20} $B \rightarrow \ell^+ \nu_\ell$ noncharmed	[b] $(2.2 \pm 0.5) \times 10^{-3}$	
Γ_{21} $B \rightarrow X_u \ell^+ \nu_\ell$	$(2.2 \pm 0.5) \%$	
Γ_{22} $B \rightarrow K^+ \ell^+ \nu_\ell$ anything	[b] $(6.3 \pm 0.6) \%$	
Γ_{23} $B \rightarrow K^- \ell^+ \nu_\ell$ anything	[b] $(10 \pm 4) \times 10^{-3}$	
Γ_{24} $B \rightarrow K^0/\bar{K}^0 \ell^+ \nu_\ell$ anything	[b] $(4.6 \pm 0.5) \%$	

D, D*, or D_s modes

Γ_{25}	$B \rightarrow D^\pm$ anything	(23.5 \pm 1.9) %	
Γ_{26}	$B \rightarrow D^0/\overline{D}{}^0$ anything	(63.7 \pm 3.0) %	S=1.1
Γ_{27}	$B \rightarrow D^*(2010)^\pm$ anything	(22.5 \pm 1.5) %	
Γ_{28}	$B \rightarrow D^*(2007)^0$ anything	(26.0 \pm 2.7) %	
Γ_{29}	$B \rightarrow D_s^\pm$ anything	[e] (10.5 \pm 2.6) %	
Γ_{30}	$B \rightarrow D_s^{*\pm}$ anything	(7.9 \pm 2.2) %	
Γ_{31}	$B \rightarrow D_s^{*\pm}\overline{D}{}^{(*)}$	(4.2 \pm 1.2) %	
Γ_{32}	$B \rightarrow \overline{D}{} D_{sJ}(2317)$		
Γ_{33}	$B \rightarrow \overline{D}{} D_{sJ}(2457)$		
Γ_{34}	$B \rightarrow D^{(*)}\overline{D}{}^{(*)}K^0 + D^{(*)}\overline{D}{}^{(*)}K^\pm$	[e,f] (7.1 \pm 2.7) %	
Γ_{35}	$b \rightarrow c\bar{c}s$	(22 \pm 4) %	
Γ_{36}	$B \rightarrow D_s^{(*)}\overline{D}{}^{(*)}$	[e,f] (4.9 \pm 1.2) %	
Γ_{37}	$B \rightarrow D^*D^*(2010)^\pm$	[e] < 5.9 $\times 10^{-3}$ CL=90%	
Γ_{38}	$B \rightarrow DD^*(2010)^\pm + D^*D^\pm$	[e] < 5.5 $\times 10^{-3}$ CL=90%	
Γ_{39}	$B \rightarrow DD^\pm$	[e] < 3.1 $\times 10^{-3}$ CL=90%	
Γ_{40}	$B \rightarrow D_s^{(*)\pm}\overline{D}{}^{(*)}X(n\pi^\pm)$	[e,f] (9 \pm 5) %	
Γ_{41}	$B \rightarrow D^*(2010)\gamma$	< 1.1 $\times 10^{-3}$ CL=90%	
Γ_{42}	$B \rightarrow D_s^+\pi^-$, $D_s^{*+}\pi^-$, $D_s^+\rho^-$, [e] < 5 $D_s^{*+}\rho^-$, $D_s^+\pi^0$, $D_s^{*+}\pi^0$, $D_s^+\eta$, $D_s^{*+}\eta$, $D_s^+\rho^0$, $D_s^{*+}\rho^0$, $D_s^+\omega$, $D_s^{*+}\omega$	$\times 10^{-4}$ CL=90%	
Γ_{43}	$B \rightarrow D_{s1}(2536)^+$ anything	< 9.5 $\times 10^{-3}$ CL=90%	

Charmonium modes

Γ_{44}	$B \rightarrow J/\psi(1S)$ anything	(1.094 \pm 0.032) %	S=1.1
Γ_{45}	$B \rightarrow J/\psi(1S)$ (direct) anything	(7.8 \pm 0.4) $\times 10^{-3}$	S=1.1
Γ_{46}	$B \rightarrow \psi(2S)$ anything	(3.07 \pm 0.21) $\times 10^{-3}$	
Γ_{47}	$B \rightarrow \chi_{c1}(1P)$ anything	(3.86 \pm 0.27) $\times 10^{-3}$	
Γ_{48}	$B \rightarrow \chi_{c1}(1P)$ (direct) anything	(3.34 \pm 0.27) $\times 10^{-3}$	
Γ_{49}	$B \rightarrow \chi_{c2}(1P)$ anything	(1.4 \pm 0.4) $\times 10^{-3}$	S=1.7
Γ_{50}	$B \rightarrow \chi_{c2}(1P)$ (direct) anything	(1.65 \pm 0.31) $\times 10^{-3}$	
Γ_{51}	$B \rightarrow \eta_c(1S)$ anything	< 9 $\times 10^{-3}$ CL=90%	

K or K* modes

Γ_{52}	$B \rightarrow K^\pm$ anything	[e] (78.9 \pm 2.5) %	
Γ_{53}	$B \rightarrow K^+$ anything	(66 \pm 5) %	
Γ_{54}	$B \rightarrow K^-$ anything	(13 \pm 4) %	
Γ_{55}	$B \rightarrow K^0/\overline{K}{}^0$ anything	[e] (64 \pm 4) %	

Γ_{56}	$B \rightarrow K^*(892)^{\pm} \text{anything}$	(18 \pm 6) %
Γ_{57}	$B \rightarrow K^*(892)^0 / \bar{K}^*(892)^0 \text{any-}$	[e] (14.6 \pm 2.6) %
	thing	
Γ_{58}	$B \rightarrow K^*(892)\gamma$	(4.2 \pm 0.6) $\times 10^{-5}$
Γ_{59}	$B \rightarrow K_1(1400)\gamma$	< 1.27 $\times 10^{-4}$ CL=90%
Γ_{60}	$B \rightarrow K_2^*(1430)\gamma$	(1.7 \pm 0.6) $\times 10^{-5}$
Γ_{61}	$B \rightarrow K_2(1770)\gamma$	< 1.2 $\times 10^{-3}$ CL=90%
Γ_{62}	$B \rightarrow K_3^*(1780)\gamma$	< 3.0 $\times 10^{-3}$ CL=90%
Γ_{63}	$B \rightarrow K_4^*(2045)\gamma$	< 1.0 $\times 10^{-3}$ CL=90%
Γ_{64}	$B \rightarrow K\eta'(958)$	(8.3 \pm 1.1) $\times 10^{-5}$
Γ_{65}	$B \rightarrow K^*(892)\eta'(958)$	< 2.2 $\times 10^{-5}$ CL=90%
Γ_{66}	$B \rightarrow K\eta$	< 5.2 $\times 10^{-6}$ CL=90%
Γ_{67}	$B \rightarrow K^*(892)\eta$	(1.8 \pm 0.5) $\times 10^{-5}$
Γ_{68}	$B \rightarrow K\phi\phi$	(2.3 \pm 0.9) $\times 10^{-6}$
Γ_{69}	$B \rightarrow \bar{b} \rightarrow \bar{s}\gamma$	(3.41 \pm 0.34) $\times 10^{-4}$
Γ_{70}	$B \rightarrow \bar{b} \rightarrow \bar{s}\text{gluon}$	< 6.8 % CL=90%
Γ_{71}	$B \rightarrow \eta \text{anything}$	< 4.4 $\times 10^{-4}$ CL=90%
Γ_{72}	$B \rightarrow \eta' \text{anything}$	(4.2 \pm 0.9) $\times 10^{-4}$

Light unflavored meson modes

Γ_{73}	$B \rightarrow \rho\gamma$	< 1.9 $\times 10^{-6}$ CL=90%
Γ_{74}	$B \rightarrow \rho/\omega\gamma$	< 1.2 $\times 10^{-6}$ CL=90%
Γ_{75}	$B \rightarrow \pi^{\pm} \text{anything}$	[e,g] (358 \pm 7) %
Γ_{76}	$B \rightarrow \pi^0 \text{anything}$	(235 \pm 11) %
Γ_{77}	$B \rightarrow \eta \text{anything}$	(17.6 \pm 1.6) %
Γ_{78}	$B \rightarrow \rho^0 \text{anything}$	(21 \pm 5) %
Γ_{79}	$B \rightarrow \omega \text{anything}$	< 81 % CL=90%
Γ_{80}	$B \rightarrow \phi \text{anything}$	(3.42 \pm 0.13) %
Γ_{81}	$B \rightarrow \phi K^*(892)$	< 2.2 $\times 10^{-5}$ CL=90%

Baryon modes

Γ_{82}	$B \rightarrow \Lambda_c^+ / \bar{\Lambda}_c^- \text{anything}$	(6.4 \pm 1.1) %
Γ_{83}	$B \rightarrow \Lambda_c^+ \text{anything}$	
Γ_{84}	$B \rightarrow \bar{\Lambda}_c^- \text{anything}$	
Γ_{85}	$B \rightarrow \bar{\Lambda}_c^- e^+ \text{anything}$	< 3.2 $\times 10^{-3}$ CL=90%
Γ_{86}	$B \rightarrow \bar{\Lambda}_c^- p \text{anything}$	(3.6 \pm 0.7) %
Γ_{87}	$B \rightarrow \bar{\Lambda}_c^- p e^+ \nu_e$	< 1.5 $\times 10^{-3}$ CL=90%
Γ_{88}	$B \rightarrow \bar{\Sigma}_c^{*-} \text{anything}$	(4.2 \pm 2.4) $\times 10^{-3}$
Γ_{89}	$B \rightarrow \bar{\Sigma}_c^- \text{anything}$	< 9.6 $\times 10^{-3}$ CL=90%
Γ_{90}	$B \rightarrow \bar{\Sigma}_c^0 \text{anything}$	(4.6 \pm 2.4) $\times 10^{-3}$
Γ_{91}	$B \rightarrow \bar{\Sigma}_c^0 N (N = p \text{ or } n)$	< 1.5 $\times 10^{-3}$ CL=90%
Γ_{92}	$B \rightarrow \Xi_c^0 \text{anything}$	(1.4 \pm 0.5) $\times 10^{-4}$
	$\times B(\Xi_c^0 \rightarrow \Xi^- \pi^+)$	

Γ_{93}	$B \rightarrow \Xi_c^+ \text{anything}$	$(4.5 \pm 1.3) \times 10^{-4}$
	$\times B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+)$	
Γ_{94}	$B \rightarrow p/\bar{p} \text{anything}$	[e] (8.0 ± 0.4)%
Γ_{95}	$B \rightarrow p/\bar{p}(\text{direct}) \text{anything}$	[e] (5.5 ± 0.5)%
Γ_{96}	$B \rightarrow \Lambda/\bar{\Lambda} \text{anything}$	[e] (4.0 ± 0.5)%
Γ_{97}	$B \rightarrow \Lambda \text{anything}$	
Γ_{98}	$B \rightarrow \bar{\Lambda} \text{anything}$	
Γ_{99}	$B \rightarrow \Xi^-/\Xi^+ \text{anything}$	[e] (2.7 ± 0.6) × 10 ⁻³
Γ_{100}	$B \rightarrow \text{baryons anything}$	(6.8 ± 0.6)%
Γ_{101}	$B \rightarrow p\bar{p} \text{anything}$	(2.47 ± 0.23)%
Γ_{102}	$B \rightarrow \Lambda\bar{p}/\bar{\Lambda}p \text{anything}$	[e] (2.5 ± 0.4)%
Γ_{103}	$B \rightarrow \Lambda\bar{\Lambda} \text{anything}$	< 5 × 10 ⁻³ CL=90%

**Lepton Family number (*LF*) violating modes or
 $\Delta B = 1$ weak neutral current (*B1*) modes**

Γ_{104}	$B \rightarrow s e^+ e^-$	<i>B1</i>	(5.6 ± 1.7) × 10 ⁻⁶
Γ_{105}	$B \rightarrow s \mu^+ \mu^-$	<i>B1</i>	(6.6 ± 2.0) × 10 ⁻⁶
Γ_{106}	$B \rightarrow s \ell^+ \ell^-$	<i>B1</i>	[b] (5.9 ± 1.4) × 10 ⁻⁶
Γ_{107}	$B \rightarrow K e^+ e^-$	<i>B1</i>	(6.0 ± 1.4) × 10 ⁻⁷ S=1.1
Γ_{108}	$B \rightarrow K^*(892) e^+ e^-$	<i>B1</i>	(1.24 ± 0.37) × 10 ⁻⁶
Γ_{109}	$B \rightarrow K \mu^+ \mu^-$	<i>B1</i>	(4.7 ± 1.1) × 10 ⁻⁷
Γ_{110}	$B \rightarrow K^*(892) \mu^+ \mu^-$	<i>B1</i>	(1.19 ± 0.34) × 10 ⁻⁶
Γ_{111}	$B \rightarrow K \ell^+ \ell^-$	<i>B1</i>	(5.4 ± 0.8) × 10 ⁻⁷
Γ_{112}	$B \rightarrow K^*(892) \ell^+ \ell^-$	<i>B1</i>	(1.05 ± 0.20) × 10 ⁻⁶
Γ_{113}	$B \rightarrow e^\pm \mu^\mp s$	<i>LF</i>	[e] < 2.2 × 10 ⁻⁵ CL=90%
Γ_{114}	$B \rightarrow \pi e^\pm \mu^\mp$	<i>LF</i>	< 1.6 × 10 ⁻⁶ CL=90%
Γ_{115}	$B \rightarrow \rho e^\pm \mu^\mp$	<i>LF</i>	< 3.2 × 10 ⁻⁶ CL=90%
Γ_{116}	$B \rightarrow K e^\pm \mu^\mp$	<i>LF</i>	< 1.6 × 10 ⁻⁶ CL=90%
Γ_{117}	$B \rightarrow K^*(892) e^\pm \mu^\mp$	<i>LF</i>	< 6.2 × 10 ⁻⁶ CL=90%

[a] These values are model dependent.

[b] An ℓ indicates an e or a μ mode, not a sum over these modes.

[c] Here “anything” means at least one particle observed.

[d] D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances.

[e] The value is for the sum of the charge states or particle/antiparticle states indicated.

[f] $D^{(*)}\bar{D}^{(*)}$ stands for the sum of $D^*\bar{D}^*$, $D^*\bar{D}$, $D\bar{D}^*$, and $D\bar{D}$.

[g] Inclusive branching fractions have a multiplicity definition and can be greater than 100%.

B^\pm/B^0 ADMIXTURE BRANCHING RATIOS

$\Gamma(\ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$

Γ_4/Γ

These branching fraction values are model dependent.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account corrections between the measurements.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.1087±0.0017 OUR EVALUATION

0.1081±0.0014 OUR AVERAGE Includes data from the 2 datablocks that follow this one.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.108 ±0.002 ±0.0056 ¹ HENDERSON 92 CLEO $e^+ e^- \rightarrow \gamma(4S)$

¹ HENDERSON 92 measurement employs e and μ . The systematic error contains 0.004 in quadrature from model dependence. The authors average a variation of the Isgur, Scora, Grinstein, and Wise model with that of the Altarelli-Cabibbo-Corbò-Maiani-Martinelli model for semileptonic decays to correct the acceptance.

$\Gamma(e^+ \nu_e \text{anything})/\Gamma_{\text{total}}$

Γ_1/Γ

These branching fraction values are model dependent.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account corrections between the measurements.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

0.1087±0.0017 OUR EVALUATION

0.1081±0.0014 OUR AVERAGE

0.1083±0.0016±0.0006 ² AUBERT 04X BABR $e^+ e^- \rightarrow \gamma(4S)$

0.1091±0.0009±0.0024 ³ MAHMOOD 04 CLEO $e^+ e^- \rightarrow \gamma(4S)$

0.109 ±0.0012±0.0049 ⁴ ABE 02Y BELL $e^+ e^- \rightarrow \gamma(4S)$

0.097 ±0.005 ±0.004 ⁵ ALBRECHT 93H ARG $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1036±0.0006±0.0023 ⁶ AUBERT,B 04A BABR $e^+ e^- \rightarrow \gamma(4S)$

0.1087±0.0018±0.0030 ⁷ AUBERT 03 BABR Repl. by AUBERT 04X

0.1049±0.0017±0.0043 ⁸ BARISH 96B CLE2 Repl. by MAHMOOD 04

0.100 ±0.004 ±0.003 ⁹ YANAGISAWA 91 CSB2 $e^+ e^- \rightarrow \gamma(4S)$

0.103 ±0.006 ±0.002 ¹⁰ ALBRECHT 90H ARG $e^+ e^- \rightarrow \gamma(4S)$

0.117 ±0.004 ±0.010 ¹¹ WACHS 89 CBAL Direct e at $\gamma(4S)$

0.120 ±0.007 ±0.005 CHEN 84 CLEO Direct e at $\gamma(4S)$

0.132 ±0.008 ±0.014 ¹² KLOPFEN... 83B CUSB Direct e at $\gamma(4S)$

- ² The semileptonic branching ratio, $|V_{cb}|$ and other heavy-quark parameters are determined from a simultaneous fit to moments of the hadronic-mass and lepton-energy distribution.
- ³ Uses charge and angular correlations in $\Upsilon(4S)$ events with a high-momentum lepton and an additional electron.
- ⁴ Uses the high-momentum lepton tag method. ABE 02Y also reports $|V_{cb}| = 0.0408 \pm 0.0010(\text{exp}) \pm 0.0025(\text{theo.})$. The second error is due to uncertainties of theoretical inputs.
- ⁵ ALBRECHT 93H analysis performed using tagged semileptonic decays of the B . This technique is almost model independent for the lepton branching ratio.
- ⁶ Uses the high-momentum lepton tag method and requires the electron energy above 0.6 GeV.
- ⁷ Uses the high-momentum lepton tag method. They also report $|V_{cb}| = 0.0423 \pm 0.0007(\text{exp}) \pm 0.0020(\text{theo.})$.
- ⁸ BARISH 96B analysis performed using tagged semileptonic decays of the B . This technique is almost model independent for the lepton branching ratio.
- ⁹ YANAGISAWA 91 also measures an average semileptonic branching ratio at the $\Upsilon(5S)$ of 9.6–10.5% depending on assumptions about the relative production of different B meson species.
- ¹⁰ ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta. 0.099 ± 0.006 is obtained using ISGUR 89B.
- ¹¹ Using data above $p(e) = 2.4$ GeV, WACHS 89 determine $\sigma(B \rightarrow e\nu\text{up})/\sigma(B \rightarrow e\nu\text{charm}) < 0.065$ at 90% CL.
- ¹² Ratio $\sigma(b \rightarrow e\nu\text{up})/\sigma(b \rightarrow e\nu\text{charm}) < 0.055$ at CL = 90%.

$\Gamma(\mu^+ \nu_\mu \text{anything})/\Gamma_{\text{total}}$

Γ_3/Γ

These branching fraction values are model dependent.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account corrections between the measurements.

VALUE	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.			

0.1087 ± 0.0017 OUR EVALUATION

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.100 $\pm 0.006 \pm 0.002$	13 ALBRECHT 90H ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.108 $\pm 0.006 \pm 0.01$	CHEN 84 CLEO	Direct μ at $\Upsilon(4S)$
0.112 $\pm 0.009 \pm 0.01$	LEVMAN 84 CUSB	Direct μ at $\Upsilon(4S)$

¹³ ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta. 0.097 ± 0.006 is obtained using ISGUR 89B.

$\Gamma(\bar{p}e^+ \nu_e \text{anything})/\Gamma_{\text{total}}$

Γ_2/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.9 \times 10^{-4}$	90	14 ADAM 03B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0016	90	ALBRECHT 90H ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	------------------	------------------------------------

¹⁴ Based on $V-A$ model.

$\Gamma(D^- \ell^+ \nu_\ell \text{anything})/\Gamma(\ell^+ \nu_\ell \text{anything})$
 $\ell = e \text{ or } \mu.$
 Γ_5/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
0.26±0.07±0.04	15 FULTON	91 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

15 FULTON 91 uses $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.1 \pm 1.3 \pm 0.4)\%$ as measured by MARK III.

 $\Gamma(\bar{D}^0 \ell^+ \nu_\ell \text{anything})/\Gamma(\ell^+ \nu_\ell \text{anything})$
 $\ell = e \text{ or } \mu.$
 Γ_6/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
0.67±0.09±0.10	16 FULTON	91 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

16 FULTON 91 uses $B(D^0 \rightarrow K^- \pi^+) = (4.2 \pm 0.4 \pm 0.4)\%$ as measured by MARK III.

 $\Gamma(D^{*-} \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$
 Γ_7/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
0.67±0.08±0.10	ABDALLAH 04D DLPH	$e^+ e^- \rightarrow Z^0$	■

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.3 ± 0.1	17 BARISH	95 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
-----------------	-----------	---------	----------------------------------

17 BARISH 95 use $B(D^0 \rightarrow K^- \pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*+} \rightarrow D^0 \pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$.

 $\Gamma(D^{*0} \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$
 Γ_8/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

0.6±0.6±0.1

0.6±0.6±0.1	18 BARISH	95 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
-------------	-----------	---------	----------------------------------

18 BARISH 95 use $B(D^0 \rightarrow K^- \pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$, $B(D^{*+} \rightarrow D^0 \pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$, $B(D^{*0} \rightarrow D^0 \pi^0) = (63.6 \pm 2.3 \pm 3.3)\%$.

 $\Gamma(\bar{D}^{**} \ell^+ \nu_\ell)/\Gamma_{\text{total}}$
 Γ_9/Γ

D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances. $\ell = e \text{ or } \mu$, not sum over e and μ modes.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.027±0.005±0.005	63	19 ALBRECHT	93 ARG	$e^+ e^- \rightarrow \gamma(4S)$	■

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.028	95	20 BARISH	95 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
--------	----	-----------	---------	----------------------------------

19 ALBRECHT 93 assumes the GISW model to correct for unseen modes. Using the BHKT model, the result becomes $0.023 \pm 0.006 \pm 0.004$. Assumes $B(D^{*+} \rightarrow D^0 \pi^+) = 68.1\%$, $B(D^0 \rightarrow K^- \pi^+) = 3.65\%$, $B(D^0 \rightarrow K^- \pi^+ \pi^- \pi^+) = 7.5\%$. We have taken their average e and μ value.

20 BARISH 95 use $B(D^0 \rightarrow K^- \pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$, assume all nonresonant channels are zero, and use GISW model for relative abundances of D^{**} states.

$\Gamma(\overline{D}_1(2420)\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0074±0.0016	21 BUSKULIC	97B ALEP	$e^+e^- \rightarrow Z$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
seen	22 BUSKULIC	95B ALEP	Repl. by BUSKULIC 97B
21 BUSKULIC 97B assumes $B(D_1(2420) \rightarrow D^*\pi) = 1$, $B(D_1(2420) \rightarrow D^*\pi^\pm) = 2/3$, and $B(b \rightarrow B) = 0.378 \pm 0.022$.			
22 BUSKULIC 95B reports $f_B \times B(B \rightarrow \overline{D}_1(2420)^0\ell^+\nu_\ell\text{anything}) \times B(\overline{D}_1(2420)^0 \rightarrow \overline{D}^*(2010)^-\pi^+) = (2.04 \pm 0.58 \pm 0.34)10^{-3}$, where f_B is the production fraction for a single B charge state.			

$[\Gamma(D\pi\ell^+\nu_\ell\text{anything}) + \Gamma(D^*\pi\ell^+\nu_\ell\text{anything})]/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.026 ± 0.005 OUR AVERAGE			Error includes scale factor of 1.5.
0.0340±0.0052±0.0032	23 ABREU	00R DLPH	$e^+e^- \rightarrow Z$
0.0226±0.0029±0.0033	24 BUSKULIC	97B ALEP	$e^+e^- \rightarrow Z$
23 Assumes no contribution from B_s and b baryons. Further assumes contributions from single pion ($D\pi$ and $D^*\pi$) states only, allowing isospin conservation to relate the relative π^0 and π^+ rates.			
24 BUSKULIC 97B assumes $B(b \rightarrow B) = 0.378 \pm 0.022$ and uses isospin invariance by assuming that all observed $D^0\pi^+$, $D^{*0}\pi^+$, $D^+\pi^-$, and $D^{*+}\pi^-$ are from D^{**} states. A correction has been applied to account for the production of B_s^0 and Λ_b^0 .			

$\Gamma(D\pi\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0154±0.0061	ABREU	00R DLPH	$e^+e^- \rightarrow Z$

$\Gamma(D^*\pi\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0186±0.0038	ABREU	00R DLPH	$e^+e^- \rightarrow Z$

$\Gamma(\overline{D}_2^*(2460)\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0065	95	25 BUSKULIC	97B ALEP	$e^+e^- \rightarrow Z$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
not seen		26 BUSKULIC	95B ALEP	$e^+e^- \rightarrow Z$
25 A revised number based on BUSKULIC 97B which assumes $B(D_2^*(2460) \rightarrow D^*\pi^\pm) = 0.20$ and $B(b \rightarrow B) = 0.378 \pm 0.022$.				
26 BUSKULIC 95B reports $f_B \times B(B \rightarrow \overline{D}_2^*(2460)^0\ell^+\nu_\ell\text{anything}) \times B(\overline{D}_2^*(2460)^0 \rightarrow \overline{D}^*(2010)^-\pi^+) \leq 0.81 \times 10^{-3}$ at CL=95%, where f_B is the production fraction for a single B charge state.				

$\Gamma(D^{*-}\pi^+\ell^+\nu_\ell\text{anything})/\Gamma_{\text{total}}$ Γ_{15}/Γ

Includes resonant and nonresonant contributions.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
10.0±2.7±2.1	27 BUSKULIC	95B ALEP	$e^+e^- \rightarrow Z$

27 BUSKULIC 95B reports $f_B \times B(B \rightarrow \overline{D}^*(2010)^-\pi^+\ell^+\nu_\ell\text{anything}) = (3.7 \pm 1.0 \pm 0.7)10^{-3}$. Above value assumes $f_B = 0.37 \pm 0.03$.

$\Gamma(D_s^- \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$					Γ_{16}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.009	90	28 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$	

28 ALBRECHT 93E reports < 0.012 for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(D_s^- \ell^+ \nu_\ell K^+ \text{anything})/\Gamma_{\text{total}}$					Γ_{17}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.006	90	29 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$	

29 ALBRECHT 93E reports < 0.008 for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(D_s^- \ell^+ \nu_\ell K^0 \text{anything})/\Gamma_{\text{total}}$					Γ_{18}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.009	90	30 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$	

30 ALBRECHT 93E reports < 0.012 for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$.

$\Gamma(\ell^+ \nu_\ell \text{charm})/\Gamma_{\text{total}}$					Γ_{19}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
0.1061 ± 0.0016 ± 0.0006	31 AUBERT	04x BABR	$e^+ e^- \rightarrow \gamma(4S)$		

31 The semileptonic branching ratio, $|V_{cb}|$ and other heavy-quark parameters are determined from a simultaneous fit to moments of the hadronic-mass and lepton-energy distribution.

$\Gamma(\ell^+ \nu_\ell \text{noncharmed})/\Gamma(\ell^+ \nu_\ell \text{anything})$					Γ_{20}/Γ_4
ℓ denotes e or μ , not the sum. These experiments measure this ratio in very limited momentum intervals.					
VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.06 ± 0.25 ± 0.42			32 AUBERT	04I BABR	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.0	90	33 ALBRECHT	94C ARG	$e^+ e^- \rightarrow \gamma(4S)$	
<4.0	90	34 BARTELTT	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
<5.5	90	77 ALBRECHT	91C ARG	$e^+ e^- \rightarrow \gamma(4S)$	
		41 ALBRECHT	90 ARG	$e^+ e^- \rightarrow \gamma(4S)$	
		76 FULTON	90 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
<4.0	90	38 BEHRENDS	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
<4.0	90	CHEN	84 CLEO	Direct e at $\gamma(4S)$	
		KLOPFEN...	83B CUSB	Direct e at $\gamma(4S)$	

32 The third error includes the systematics and theoretical errors summed in quadrature.

33 ALBRECHT 94C find $\Gamma(b \rightarrow c)/\Gamma(b \rightarrow \text{all}) = 0.99 \pm 0.02 \pm 0.04$.

34 BARTELTT 93B (CLEO II) measures an excess of $107 \pm 15 \pm 11$ leptons in the lepton momentum interval 2.3–2.6 GeV/c which is attributed to $b \rightarrow u\ell\nu_\ell$. This corresponds to a model-dependent partial branching ratio ΔB_{ub} between $(1.15 \pm 0.16 \pm 0.15) \times 10^{-4}$, as evaluated using the KS model (KOERNER 88), and $(1.54 \pm 0.22 \pm 0.20) \times 10^{-4}$ using the ACCMM model (ARTUSO 93). The corresponding values of $|V_{ub}|/|V_{cb}|$ are 0.056 ± 0.006 and 0.076 ± 0.008 , respectively.

- ³⁵ ALBRECHT 91C result supersedes ALBRECHT 90. Two events are fully reconstructed providing evidence for the $b \rightarrow u$ transition. Using the model of ALTARELLI 82, they obtain $|V_{ub}/V_{cb}| = 0.11 \pm 0.012$ from 77 leptons in the 2.3–2.6 GeV momentum range.
- ³⁶ ALBRECHT 90 observes 41 ± 10 excess e and μ (lepton) events in the momentum interval $p = 2.3\text{--}2.6$ GeV signaling the presence of the $b \rightarrow u$ transition. The events correspond to a model-dependent measurement of $|V_{ub}/V_{cb}| = 0.10 \pm 0.01$.
- ³⁷ FULTON 90 observe 76 ± 20 excess e and μ (lepton) events in the momentum interval $p = 2.4\text{--}2.6$ GeV signaling the presence of the $b \rightarrow u$ transition. The average branching ratio, $(1.8 \pm 0.4 \pm 0.3) \times 10^{-4}$, corresponds to a model-dependent measurement of approximately $|V_{ub}/V_{cb}| = 0.1$ using $B(b \rightarrow c\ell\nu) = 10.2 \pm 0.2 \pm 0.7\%$.
- ³⁸ The quoted possible limits range from 0.018 to 0.04 for the ratio, depending on which model or momentum range is chosen. We select the most conservative limit they have calculated. This corresponds to a limit on $|V_{ub}|/|V_{cb}| < 0.20$. While the endpoint technique employed is more robust than their previous results in CHEN 84, these results do not provide a numerical improvement in the limit.

 $\Gamma(X_u \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
2.24 \pm 0.27 \pm 0.47	39,40 AUBERT	04I BABR	$e^+ e^- \rightarrow \gamma(4S)$

³⁹ Used BaBar measurement of Semileptonic branching fraction $B(B \rightarrow X\ell\nu_\ell) = (10.87 \pm 0.18 \pm 0.30)\%$ to convert the ratio of rates to branching fraction.

⁴⁰ The third error includes the systematics and theoretical errors summed in quadrature.

 $\Gamma(K^+ \ell^+ \nu_\ell \text{anything})/\Gamma(\ell^+ \nu_\ell \text{anything})$ Γ_{22}/Γ_4

ℓ denotes e or μ , not the sum.

VALUE	DOCUMENT ID	TECN	COMMENT
0.58 \pm 0.05 OUR AVERAGE			
0.594 \pm 0.021 \pm 0.056	ALBRECHT	94C ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.54 \pm 0.07 \pm 0.06	41 ALAM	87B CLEO	$e^+ e^- \rightarrow \gamma(4S)$

⁴¹ ALAM 87B measurement relies on lepton-kaon correlations.

 $\Gamma(K^- \ell^+ \nu_\ell \text{anything})/\Gamma(\ell^+ \nu_\ell \text{anything})$ Γ_{23}/Γ_4

ℓ denotes e or μ , not the sum.

VALUE	DOCUMENT ID	TECN	COMMENT
0.092 \pm 0.035 OUR AVERAGE			
0.086 \pm 0.011 \pm 0.044	ALBRECHT	94C ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.10 \pm 0.05 \pm 0.02	42 ALAM	87B CLEO	$e^+ e^- \rightarrow \gamma(4S)$

⁴² ALAM 87B measurement relies on lepton-kaon correlations.

 $\Gamma(K^0/\bar{K}^0 \ell^+ \nu_\ell \text{anything})/\Gamma(\ell^+ \nu_\ell \text{anything})$ Γ_{24}/Γ_4

ℓ denotes e or μ , not the sum. Sum over K^0 and \bar{K}^0 states.

VALUE	DOCUMENT ID	TECN	COMMENT
0.42 \pm 0.05 OUR AVERAGE			
0.452 \pm 0.038 \pm 0.056	43 ALBRECHT	94C ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.39 \pm 0.06 \pm 0.04	44 ALAM	87B CLEO	$e^+ e^- \rightarrow \gamma(4S)$

⁴³ ALBRECHT 94C assume a K^0/\bar{K}^0 multiplicity twice that of K_S^0 .

⁴⁴ ALAM 87B measurement relies on lepton-kaon correlations.

$\langle n_c \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
1.10±0.05	45 GIBBONS	97B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.98±0.16±0.12	46 ALAM	87B CLEO	$e^+ e^- \rightarrow \gamma(4S)$
45 GIBBONS 97B from charm counting using $B(D_s^+ \rightarrow \phi\pi) = 0.036 \pm 0.009$ and $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = 0.044 \pm 0.006$.			
46 From the difference between K^- and K^+ widths. ALAM 87B measurement relies on lepton-kaon correlations. It does not consider the possibility of $B\bar{B}$ mixing. We have thus removed it from the average.			

 $\Gamma(D^\pm \text{anything})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{25}/Γ
0.235±0.019 OUR AVERAGE					
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.234±0.012±0.015		47 GIBBONS	97B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
0.25 ± 0.04 ± 0.02		48 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
0.23 ± 0.05 ± 0.01		49 ALBRECHT	91H ARG	$e^+ e^- \rightarrow \gamma(4S)$	
0.21 ± 0.05 ± 0.01	20k	50 BORTOLETTO87	CLEO	Sup. by BORTOLETTO 92	
47 GIBBONS 97B reports $[B(B \rightarrow D^\pm \text{anything}) \times B(D^+ \rightarrow K^-\pi^+\pi^+)] = 0.0216 \pm 0.0008 \pm 0.00082$. We divide by our best value $B(D^+ \rightarrow K^-\pi^+\pi^+) = (9.2 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
48 BORTOLETTO 92 reports $[B(B \rightarrow D^\pm \text{anything}) \times B(D^+ \rightarrow K^-\pi^+\pi^+)] = 0.0226 \pm 0.0030 \pm 0.0018$. We divide by our best value $B(D^+ \rightarrow K^-\pi^+\pi^+) = (9.2 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
49 ALBRECHT 91H reports $[B(B \rightarrow D^\pm \text{anything}) \times B(D^+ \rightarrow K^-\pi^+\pi^+)] = 0.0209 \pm 0.0027 \pm 0.0040$. We divide by our best value $B(D^+ \rightarrow K^-\pi^+\pi^+) = (9.2 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
50 BORTOLETTO 87 reports $[B(B \rightarrow D^\pm \text{anything}) \times B(D^+ \rightarrow K^-\pi^+\pi^+)] = 0.019 \pm 0.004 \pm 0.002$. We divide by our best value $B(D^+ \rightarrow K^-\pi^+\pi^+) = (9.2 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(D^0/\bar{D}^0 \text{anything})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{26}/Γ
0.637±0.030 OUR AVERAGE				Error includes scale factor of 1.1.	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.658±0.025±0.015		51 GIBBONS	97B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
0.61 ± 0.05 ± 0.01		52 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
0.51 ± 0.08 ± 0.01		53 ALBRECHT	91H ARG	$e^+ e^- \rightarrow \gamma(4S)$	
0.55 ± 0.07 ± 0.01	21k	54 BORTOLETTO87	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
0.63 ± 0.19 ± 0.01		55 GREEN	83 CLEO	Repl. by BORTOLETTO 87	

- ⁵¹ GIBBONS 97B reports $[B(B \rightarrow D^0/\bar{D}^0 \text{ anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0251 \pm 0.0006 \pm 0.00075$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.81 \pm 0.09) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁵² BORTOLETTO 92 reports $[B(B \rightarrow D^0/\bar{D}^0 \text{ anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0233 \pm 0.0012 \pm 0.0014$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.81 \pm 0.09) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁵³ ALBRECHT 91H reports $[B(B \rightarrow D^0/\bar{D}^0 \text{ anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0194 \pm 0.0015 \pm 0.0025$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.81 \pm 0.09) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁵⁴ BORTOLETTO 87 reports $[B(B \rightarrow D^0/\bar{D}^0 \text{ anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.0210 \pm 0.0015 \pm 0.0021$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.81 \pm 0.09) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁵⁵ GREEN 83 reports $[B(B \rightarrow D^0/\bar{D}^0 \text{ anything}) \times B(D^0 \rightarrow K^- \pi^+)] = 0.024 \pm 0.006 \pm 0.004$. We divide by our best value $B(D^0 \rightarrow K^- \pi^+) = (3.81 \pm 0.09) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^*(2010)^{\pm} \text{ anything})/\Gamma_{\text{total}}$

Γ_{27}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.225±0.015 OUR AVERAGE				
0.247±0.019±0.01		56 GIBBONS 97B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
0.205±0.019±0.007		57 ALBRECHT 96D ARG	$e^+ e^- \rightarrow \gamma(4S)$	
0.230±0.028±0.009		58 BORTOLETTO92 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.283±0.053±0.002		59 ALBRECHT 91H ARG	Sup. by ALBRECHT 96D	
0.22 ± 0.04 + 0.07 - 0.04	5200	60 BORTOLETTO87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
0.27 ± 0.06 + 0.08 - 0.06	510	61 CSORNA 85 CLEO	Repl. by BORTOLETTO 87	
56 GIBBONS 97B reports $B(B \rightarrow D^*(2010)^+ \text{ anything}) = 0.239 \pm 0.015 \pm 0.014 \pm 0.009$ using CLEO measured D and D^* branching fractions. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
57 ALBRECHT 96D reports $B(B \rightarrow D^*(2010)^+ \text{ anything}) = 0.196 \pm 0.019$ using CLEO measured $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.681 \pm 0.01 \pm 0.013$, $B(D^0 \rightarrow K^- \pi^+) = 0.0401 \pm 0.0014$, $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-) = 0.081 \pm 0.005$. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
58 BORTOLETTO 92 reports $B(B \rightarrow D^*(2010)^+ \text{ anything}) = 0.25 \pm 0.03 \pm 0.04$ using MARK II $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$ and $B(D^0 \rightarrow K^- \pi^+) = 0.042 \pm 0.008$. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
59 ALBRECHT 91H reports $0.348 \pm 0.060 \pm 0.035$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.55 \pm 0.04$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Uses the PDG 90 $B(D^0 \rightarrow K^- \pi^+) = 0.0371 \pm 0.0025$.				

⁶⁰ BORTOLETTO 87 uses old MARK III (BALTRUSAITIS 86E) branching ratios $B(D^0 \rightarrow K^- \pi^+) = 0.056 \pm 0.004 \pm 0.003$ and also assumes $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.60^{+0.08}_{-0.15}$. The product branching ratio for $B(B \rightarrow D^*(2010)^+) B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ is $0.13 \pm 0.02 \pm 0.012$. Superseded by BORTOLETTO 92.

⁶¹ $V-A$ momentum spectrum used to extrapolate below $p = 1$ GeV. We correct the value assuming $B(D^0 \rightarrow K^- \pi^+) = 0.042 \pm 0.006$ and $B(D^{*+} \rightarrow D^0 \pi^+) = 0.6^{+0.08}_{-0.15}$. The product branching fraction is $B(B \rightarrow D^{*+} X) \cdot B(D^{*+} \rightarrow \pi^+ D^0) \cdot B(D^0 \rightarrow K^- \pi^+) = (68 \pm 15 \pm 9) \times 10^{-4}$.

$\Gamma(D^*(2007)^0 \text{anything})/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.260±0.023±0.015	62 GIBBONS	97B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

⁶² GIBBONS 97B reports $B(B \rightarrow D^*(2007)^0 \text{anything}) = 0.247 \pm 0.012 \pm 0.018 \pm 0.018$ using CLEO measured D and D^* branching fractions. We rescale to our PDG 96 values of D and D^* branching ratios. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^\pm \text{anything})/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.105±0.026 OUR AVERAGE				

$0.109 \pm 0.006^{+0.026}_{-0.027}$ ⁶³ AUBERT 02G BABR $e^+ e^- \rightarrow \gamma(4S)$

$0.117 \pm 0.009^{+0.028}_{-0.029}$ ⁶⁴ GIBAUT 96 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

$0.081 \pm 0.014^{+0.019}_{-0.020}$ ⁶⁵ ALBRECHT 92G ARG $e^+ e^- \rightarrow \gamma(4S)$

$0.085 \pm 0.013^{+0.020}_{-0.021}$ ²⁵⁷ ⁶⁶ BORTOLETTO90 CLEO $e^+ e^- \rightarrow \gamma(4S)$

$0.105 \pm 0.028^{+0.025}_{-0.026}$ ⁶⁷ HAAS 86 CLEO $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.116 \pm 0.030 \pm 0.028$ ⁶⁸ ALBRECHT 87H ARG $e^+ e^- \rightarrow \gamma(4S)$

⁶³ AUBERT 02G reports $[B(B \rightarrow D_s^\pm \text{anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.00393 \pm 0.00007 \pm 0.00021$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶⁴ GIBAUT 96 reports $0.1211 \pm 0.0039 \pm 0.0088$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶⁵ ALBRECHT 92G reports $[B(B \rightarrow D_s^\pm \text{anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.00292 \pm 0.00039 \pm 0.00031$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶⁶ BORTOLETTO 90 reports $[B(B \rightarrow D_s^\pm \text{anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.00306 \pm 0.00047$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶⁷ HAAS 86 reports $[B(B \rightarrow D_s^\pm \text{anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.0038 \pm 0.0010$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their

experiment's error and our second error is the systematic error from using our best value.
 $64 \pm 22\%$ decays are 2-body.

68 ALBRECHT 87H reports $[B(B \rightarrow D_s^{\pm} \text{anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.0042 \pm 0.0009 \pm 0.0006$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. 46 \pm 16% of $B \rightarrow D_s X$ decays are 2-body. Superseded by ALBRECHT 92G.

$\Gamma(D_s^{*\pm} \text{anything})/\Gamma_{\text{total}}$	Γ_{30}/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.079 \pm 0.011 \pm 0.019	69 AUBERT	02G BABR	$e^+ e^- \rightarrow \gamma(4S)$

69 AUBERT 02G reports $[B(B \rightarrow D_s^{*\pm} \text{anything}) \times B(D_s^+ \rightarrow \phi\pi^+)] = 0.00284 \pm 0.00029 \pm 0.00025$. We divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.6 \pm 0.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^{*\pm} \bar{D}^{(*)})/\Gamma(D_s^{*\pm} \text{anything})$	Γ_{31}/Γ_{30}		
Sum over modes	DOCUMENT ID	TECN	COMMENT
0.533 \pm 0.037 \pm 0.037	AUBERT	02G BABR	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\bar{D} D_{sJ}(2317))/\Gamma_{\text{total}}$	Γ_{32}/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	70 KROKOVNY	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$

70 The product branching ratio for $B(B \rightarrow \bar{D} D_{sJ}(2317)^+) \times B(D_{sJ}(2317)^+ \rightarrow D_s \pi^0)$ is measured to be $(8.5^{+2.1}_{-1.9} \pm 2.6) \times 10^{-4}$.

$\Gamma(\bar{D} D_{sJ}(2457))/\Gamma_{\text{total}}$	Γ_{33}/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	71 KROKOVNY	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$

71 The product branching ratio for $B(B \rightarrow \bar{D} D_{sJ}(2457)^+) \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0, D_s^+ \gamma)$ are measured to be $(17.8^{+4.5}_{-3.9} \pm 5.3) \times 10^{-4}$ and $(6.7^{+1.3}_{-1.2} \pm 2.0) \times 10^{-4}$, respectively.

$[\Gamma(D^{(*)} \bar{D}^{(*)} K^0) + \Gamma(D^{(*)} \bar{D}^{(*)} K^\pm)]/\Gamma_{\text{total}}$	Γ_{34}/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.071 \pm 0.025 \pm 0.010 -0.015 -0.009	72 BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

72 The systematic error includes the uncertainties due to the charm branching ratios.

$\Gamma(c\bar{s})/\Gamma_{\text{total}}$	Γ_{35}/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.219 \pm 0.037	73 COAN	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

73 COAN 98 uses D - ℓ correlation.

$\Gamma(D_s^*(*)\bar{D}^*)/\Gamma(D_s^\pm \text{anything})$
 Sum over modes.
 Γ_{36}/Γ_{29}

VALUE	DOCUMENT ID	TECN	COMMENT
0.469±0.017 OUR AVERAGE			

0.464±0.013±0.015

AUBERT 02G BABR $e^+e^- \rightarrow \gamma(4S)$ 0.56 +0.21 +0.09
-0.15 -0.0874 BARATE 98Q ALEP $e^+e^- \rightarrow Z$

0.457±0.019±0.037

GIBAUT 96 CLE2 $e^+e^- \rightarrow \gamma(4S)$

0.58 ±0.07 ±0.09

ALBRECHT 92G ARG $e^+e^- \rightarrow \gamma(4S)$

0.56 ±0.10

BORTOLETTI90 CLEO $e^+e^- \rightarrow \gamma(4S)$

74 BARATE 98Q measures $B(B \rightarrow D_s^*(*)\bar{D}^*) = 0.056^{+0.021+0.009+0.019}_{-0.015-0.008-0.011}$, where the third error results from the uncertainty on the different D branching ratios and is dominated by the uncertainty on $B(D_s^+ \rightarrow \phi\pi^+)$. We divide $B(B \rightarrow D_s^*(*)\bar{D}^*)$ by our best value of $B(B \rightarrow D_s \text{anything}) = 0.1 \pm 0.025$.

 $\Gamma(D^*D^*(2010)^\pm)/\Gamma_{\text{total}}$
 Γ_{37}/Γ

VALUE	CL%
$<5.9 \times 10^{-3}$	90

DOCUMENT ID	TECN	COMMENT
BARATE	98Q ALEP	$e^+e^- \rightarrow Z$

 $[\Gamma(DD^*(2010)^\pm) + \Gamma(D^*D^\pm)]/\Gamma_{\text{total}}$
 Γ_{38}/Γ

VALUE	CL%
$<5.5 \times 10^{-3}$	90

DOCUMENT ID	TECN	COMMENT
BARATE	98Q ALEP	$e^+e^- \rightarrow Z$

 $\Gamma(DD^\pm)/\Gamma_{\text{total}}$
 Γ_{39}/Γ

VALUE	CL%
$<3.1 \times 10^{-3}$	90

DOCUMENT ID	TECN	COMMENT
BARATE	98Q ALEP	$e^+e^- \rightarrow Z$

 $\Gamma(D_s^*(*)\bar{D}^* X(n\pi^\pm))/\Gamma_{\text{total}}$
 Γ_{40}/Γ

VALUE	CL%
$0.094^{+0.040+0.034}_{-0.031-0.024}$	90

DOCUMENT ID	TECN	COMMENT
75 BARATE	98Q ALEP	$e^+e^- \rightarrow Z$

75 The systematic error includes the uncertainties due to the charm branching ratios.

 $\Gamma(D^*(2010)\gamma)/\Gamma_{\text{total}}$
 Γ_{41}/Γ

VALUE	CL%
$<1.1 \times 10^{-3}$	90

DOCUMENT ID	TECN	COMMENT
76 LESIAK	92 CBAL	$e^+e^- \rightarrow \gamma(4S)$

76 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

 $\Gamma(D_s^+\pi^-, D_s^{*+}\pi^-, D_s^+\rho^-, D_s^{*+}\rho^-, D_s^+\pi^0, D_s^{*+}\pi^0, D_s^+\eta, D_s^{*+}\eta, D_s^+\rho^0, D_s^{*+}\rho^0, D_s^+\omega, D_s^{*+}\omega)/\Gamma_{\text{total}}$
 Sum over modes.
 Γ_{42}/Γ

VALUE	CL%
<0.0005	90

DOCUMENT ID	TECN	COMMENT
77 ALEXANDER	93B CLE2	$e^+e^- \rightarrow \gamma(4S)$

77 ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$. This branching ratio limit provides a model-dependent upper limit $|V_{ub}|/|V_{cb}| < 0.16$ at CL=90%.

$\Gamma(D_{s1}(2536)^+ \text{anything})/\Gamma_{\text{total}}$ Γ_{43}/Γ

$D_{s1}(2536)^+$ is the narrow P -wave D_s^+ meson with $J^P = 1^+$.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0095	90	78 BISHAI	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

⁷⁸ Assuming factorization, the decay constant $f_{D_{s1}^+}$ is at least a factor of 2.5 times smaller than $f_{D_s^+}$.

 $\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.094 ± 0.032 OUR AVERAGE Error includes scale factor of 1.1.

1.057 ± 0.012 ± 0.040 ⁷⁹ AUBERT 03F BABR $e^+ e^- \rightarrow \gamma(4S)$

1.121 ± 0.013 ± 0.042 ANDERSON 02 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

1.30 ± 0.45 ± 0.02 ⁸⁰ MASCHMANN 90 CBAL $e^+ e^- \rightarrow \gamma(4S)$

1.24 ± 0.27 ± 0.02 ⁸¹ ALBRECHT 87D ARG $e^+ e^- \rightarrow \gamma(4S)$

1.37 ± 0.25 ± 0.02 ⁸² ALAM 86 CLEO $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.13 ± 0.06 ± 0.02 ⁸³ BAILEST 95B CLE2 $e^+ e^- \rightarrow \gamma(4S)$

1.4 ^{+0.6} _{-0.5} ⁷ ⁸⁴ ALBRECHT 85H ARG $e^+ e^- \rightarrow \gamma(4S)$

1.1 ± 0.21 ± 0.23 ⁴⁶ ⁸⁵ HAAS 85 CLEO Repl. by ALAM 86

⁷⁹ AUBERT 03F also reports the momentum distribution and helicity of $J/\psi \rightarrow \ell^+ \ell^-$ in the $\gamma(4S)$ center-of-mass frame.

⁸⁰ MASCHMANN 90 reports $(1.12 \pm 0.33 \pm 0.25) \times 10^{-2}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁸¹ ALBRECHT 87D reports $(1.07 \pm 0.16 \pm 0.22) \times 10^{-2}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. ALBRECHT 87D find the branching ratio for J/ψ not from $\psi(2S)$ to be 0.0081 ± 0.0023 .

⁸² ALAM 86 reports $(1.09 \pm 0.16 \pm 0.21) \times 10^{-2}$ for $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.074 \pm 0.012$. We rescale to our best value $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = (5.88 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁸³ BAILEST 95B reports $(1.12 \pm 0.04 \pm 0.06) \times 10^{-2}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0599 \pm 0.0025$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.93 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.. They measure $J/\psi(1S) \rightarrow e^+ e^-$ and $\mu^+ \mu^-$ and use PDG 1994 values for the branching fractions. The rescaling is the same for either mode so we use $e^+ e^-$.

⁸⁴ Statistical and systematic errors were added in quadrature. ALBRECHT 85H also report a CL = 90% limit of 0.007 for $B \rightarrow J/\psi(1S) + X$ where $m_X < 1$ GeV.

⁸⁵ Dimuon and dielectron events used.

$\Gamma(J/\psi(1S)(\text{direct}) \text{ anything})/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0078 ± 0.0004 OUR AVERAGE			Error includes scale factor of 1.1.
0.00740 ± 0.00023 ± 0.00043	86 AUBERT	03F BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.00813 ± 0.00017 ± 0.00037	87 ANDERSON	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0080 ± 0.0008	88 BALEST	95B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
86 AUBERT 03F also reports the helicity of $J/\psi \rightarrow \ell^+ \ell^-$ produced directly in B decay.			
87 Also reports the measurement of $J/\psi \rightarrow \ell^+ \ell^-$ polarization produced directly from B decay.			
88 BALEST 95B assume PDG 1994 values for sub mode branching ratios. $J/\psi(1S)$ mesons are reconstructed in $J/\psi(1S) \rightarrow e^+ e^-$ and $J/\psi(1S) \rightarrow \mu^+ \mu^-$. The $B \rightarrow J/\psi(1S)X$ branching ratio contains $J/\psi(1S)$ mesons directly from B decays and also from feeddown through $\psi(2S) \rightarrow J/\psi(1S)$, $\chi_{c1}(1P) \rightarrow J/\psi(1S)$, or $\chi_{c2}(1P) \rightarrow J/\psi(1S)$. Using the measured inclusive rates, BALEST 95B corrects for the feeddown and finds the $B \rightarrow J/\psi(1S)(\text{direct}) X$ branching ratio.			

$\Gamma(\psi(2S) \text{ anything})/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.00307 ± 0.00021 OUR AVERAGE				
0.00297 ± 0.00020 ± 0.00020		AUBERT	03F BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.00316 ± 0.00014 ± 0.00028		89 ANDERSON	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.0046 ± 0.0017 ± 0.0011	8	ALBRECHT	87D ARG	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0034 ± 0.0004 ± 0.0003	240	90 BALEST	95B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
89 Also reports the measurement of $\psi(2S) \rightarrow \ell^+ \ell^-$ polarization produced directly from B decay.				
90 BALEST 95B assume PDG 1994 values for sub mode branching ratios. They find $B(B \rightarrow \psi(2S)X, \psi(2S) \rightarrow \ell^+ \ell^-) = 0.30 \pm 0.05 \pm 0.04$ and $B(B \rightarrow \psi(2S)X, \psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = 0.37 \pm 0.05 \pm 0.05$. Weighted average is quoted for $B(B \rightarrow \psi(2S)X)$.				

$\Gamma(\chi_{c1}(1P) \text{ anything})/\Gamma_{\text{total}}$ Γ_{47}/Γ


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.00386 ± 0.00027 OUR AVERAGE				
0.00367 ± 0.00035 ± 0.00044		AUBERT	03F BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.00363 ± 0.00022 ± 0.00034		91 ABE	02L BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.00435 ± 0.00029 ± 0.00040		ANDERSON	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0036 ± 0.0004 ± 0.0003		92 CHEN	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.0040 ± 0.0006 ± 0.0004	112	93 BALEST	95B CLE2	Repl. by CHEN 01
0.0105 ± 0.0035 ± 0.0025		94 ALBRECHT	92E ARG	$e^+ e^- \rightarrow \gamma(4S)$
91 ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+ \ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.				
92 CHEN 01 reports $0.00414 \pm 0.00031 \pm 0.00040$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (31.6 \pm 2.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				
93 BALEST 95B assume $B(\chi_{c1}(1P) \rightarrow J/\psi(1S)\gamma) = (27.3 \pm 1.6) \times 10^{-2}$, the PDG 1994 value. Fit to ψ -photon invariant mass distribution allows for a $\chi_{c1}(1P)$ and a $\chi_{c2}(1P)$ component.				
94 ALBRECHT 92E assumes no $\chi_{c2}(1P)$ production.				

$\Gamma(\chi_{c1}(1P)(\text{direct}) \text{ anything})/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.00334±0.00027 OUR AVERAGE			
0.00341±0.00035±0.00042	AUBERT	03F BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.00332±0.00022±0.00034	95 ABE	02L BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.0033 ±0.0004 ±0.0002	96 CHEN	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0037 ±0.0007	97 BALEST	95B CLE2	Repl. by CHEN 01
95 ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+ \ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.			
96 CHEN 01 reports $0.00383 \pm 0.00031 \pm 0.00040$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (31.6 \pm 2.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.			
97 BALEST 95B assume PDG 1994 values. $J/\psi(1S)$ mesons are reconstructed in the $e^+ e^-$ and $\mu^+ \mu^-$ modes. The $B \rightarrow \chi_{c1}(1P)X$ branching ratio contains $\chi_{c1}(1P)$ mesons directly from B decays and also from feeddown through $\psi(2S) \rightarrow \chi_{c1}(1P)\gamma$. Using the measured inclusive rates, BALEST 95B corrects for the feeddown and finds the $B \rightarrow \chi_{c1}(1P)(\text{direct}) X$ branching ratio.			

$\Gamma(\chi_{c2}(1P)\text{anything})/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
14 ±4 OUR AVERAGE Error includes scale factor of 1.7. See the ideogram below.					
21.0±4.5±3.1			AUBERT	03F BABR	$e^+ e^- \rightarrow \gamma(4S)$
18.0 ^{+2.3} _{-2.8} ±2.6			98 ABE	02L BELL	$e^+ e^- \rightarrow \gamma(4S)$
7.4±3.8±0.3			99 CHEN	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<38	90	35	100 BALEST	95B CLE2	Repl. by CHEN 01
98 ABE 02L uses PDG 01 values for $B(J/\psi(1S) \rightarrow \ell^+ \ell^-)$ and $B(\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma)$.					
99 CHEN 01 reports $(9.8 \pm 4.8 \pm 1.5) \times 10^{-4}$ for $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$. We rescale to our best value $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (17.8 \pm 0.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.					
100 BALEST 95B assume $B(\chi_{c2}(1P) \rightarrow J/\psi(1S)\gamma) = (13.5 \pm 1.1) \times 10^{-2}$, the PDG 1994 value. $J/\psi(1S)$ mesons are reconstructed in the $e^+ e^-$ and $\mu^+ \mu^-$ modes, and PDG 1994 branching fractions are used. If interpreted as signal, the 35 ± 13 events correspond to $B(B \rightarrow \chi_{c2}(1P)X) = (0.25 \pm 0.10 \pm 0.03) \times 10^{-2}$.					



$$\Gamma(\chi_{c2}(1P) \text{anything})/\Gamma_{\text{total}}$$

$$\Gamma_{49}/\Gamma$$

$$\Gamma(\chi_{c2}(1P) \text{(direct anything)})/\Gamma_{\text{total}}$$

$$\Gamma_{50}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.00165±0.00031 OUR AVERAGE

0.00190±0.00045±0.00029 AUBERT 03F BABR e⁺ e⁻ → $\gamma(4S)$

0.00153^{+0.00023}_{-0.00028}±0.00027 101 ABE 02L BELL e⁺ e⁻ → $\gamma(4S)$

101 ABE 02L uses PDG 01 values for B($J/\psi(1S) \rightarrow \ell^+ \ell^-$) and B($\chi_{c1,c2} \rightarrow J/\psi(1S)\gamma$).

$$\Gamma(\eta_c(1S) \text{anything})/\Gamma_{\text{total}}$$

$$\Gamma_{51}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.009	90	102 BAEST	95B CLE2	e ⁺ e ⁻ → $\gamma(4S)$

102 BAEST 95B assume PDG 1994 values for sub mode branching ratios. $J/\psi(1S)$ mesons are reconstructed in $J/\psi(1S) \rightarrow e^+ e^-$ and $J/\psi(1S) \rightarrow \mu^+ \mu^-$. Search region $2960 < m_{\eta_c(1S)} < 3010$ MeV/ c^2 .

$$\Gamma(K^\pm \text{anything})/\Gamma_{\text{total}}$$

$$\Gamma_{52}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.789±0.025 OUR AVERAGE

0.82 ± 0.01 ± 0.05 ALBRECHT 94C ARG e⁺ e⁻ → $\gamma(4S)$

0.775±0.015±0.025 103 ALBRECHT 93I ARG e⁺ e⁻ → $\gamma(4S)$

0.85 ± 0.07 ± 0.09 ALAM 87B CLEO e⁺ e⁻ → $\gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 104 BRODY 82 CLEO e⁺ e⁻ → $\gamma(4S)$

seen 105 GIANNINI 82 CUSB e⁺ e⁻ → $\gamma(4S)$

- 103 ALBRECHT 93I value is not independent of the sum of $B \rightarrow K^+$ anything and $B \rightarrow K^-$ anything ALBRECHT 94C values.
 104 Assuming $\Upsilon(4S) \rightarrow B\bar{B}$, a total of $3.38 \pm 0.34 \pm 0.68$ kaons per $\Upsilon(4S)$ decay is found (the second error is systematic). In the context of the standard B -decay model, this leads to a value for $(b\text{-quark} \rightarrow c\text{-quark})/(b\text{-quark} \rightarrow \text{all})$ of $1.09 \pm 0.33 \pm 0.13$.
 105 GIANNINI 82 at CESR-CUSB observed 1.58 ± 0.35 K^0 per hadronic event much higher than 0.82 ± 0.10 below threshold. Consistent with predominant $b \rightarrow cX$ decay.

$\Gamma(K^+\text{ anything})/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.66 ±0.05	106 ALBRECHT 94C ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.620 \pm 0.013 \pm 0.038$	107 ALBRECHT 94C ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
$0.66 \pm 0.05 \pm 0.07$	107 ALAM 87B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

- 106 Measurement relies on lepton-kaon correlations. It is for the weak decay vertex and does not include mixing of the neutral B meson. Mixing effects were corrected for by assuming a mixing parameter r of $(18.1 \pm 4.3)\%$.
 107 Measurement relies on lepton-kaon correlations. It includes production through mixing of the neutral B meson.

$\Gamma(K^-\text{ anything})/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.13 ±0.04	108 ALBRECHT 94C ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.165 \pm 0.011 \pm 0.036$	109 ALBRECHT 94C ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
$0.19 \pm 0.05 \pm 0.02$	109 ALAM 87B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

108 Measurement relies on lepton-kaon correlations. It is for the weak decay vertex and does not include mixing of the neutral B meson. Mixing effects were corrected for by assuming a mixing parameter r of $(18.1 \pm 4.3)\%$.
 109 Measurement relies on lepton-kaon correlations. It includes production through mixing of the neutral B meson.

$\Gamma(K^0/\bar{K}^0\text{ anything})/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.64 ±0.04 OUR AVERAGE			
$\bullet \bullet \bullet$			
$0.642 \pm 0.010 \pm 0.042$	110 ALBRECHT 94C ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
$0.63 \pm 0.06 \pm 0.06$	ALAM 87B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

110 ALBRECHT 94C assume a K^0/\bar{K}^0 multiplicity twice that of K_S^0 .

$\Gamma(K^*(892)^\pm\text{ anything})/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.182±0.054±0.024	ALBRECHT 94J ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

$\Gamma(K^*(892)^0/\bar{K}^*(892)^0\text{ anything})/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.146±0.016±0.020	ALBRECHT 94J ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

$\Gamma(K^*(892)\gamma)/\Gamma_{\text{total}}$ Γ_{58}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.24 ± 0.54 ± 0.32	111	COAN	00	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 150	90	112	LESIAK	92 CBAL $e^+ e^- \rightarrow \gamma(4S)$
< 24	90		ALBRECHT	88H ARG $e^+ e^- \rightarrow \gamma(4S)$

111 An average of $B(B^+ \rightarrow K^*(892)^+\gamma)$ and $B(B^0 \rightarrow K^*(892)^0\gamma)$ measurements reported in COAN 00 by assuming full correlated systematic errors.

112 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

 $\Delta_{0+}(B \rightarrow K^*(892)\gamma)$

Δ_{0+} describes the isospin asymmetry between $\Gamma(B^0 \rightarrow K^*(892)^0\gamma)$ and $\Gamma(B^+ \rightarrow K^*(892)^+\gamma)$.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.050 ± 0.045 ± 0.037	113 AUBERT,BE	04A BABR	$e^+ e^- \rightarrow \gamma(4S)$

113 Uses the production ratio of charged and neutral B from $\gamma(4S)$ decays $R^{+/0} = 1.006 \pm 0.048$ and the lifetime ratio of $\tau_{B^+} / \tau_{B^0} = 1.083 \pm 0.017$. The 90% CL interval is $-0.046 < \Delta_{0+} < 0.146$.

 $\Gamma(K_1(1400)\gamma)/\Gamma_{\text{total}}$ Γ_{59}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 12.7 × 10⁻⁵	90	114 COAN	00	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 1.6×10^{-3}	90	115 LESIAK	92 CBAL	$e^+ e^- \rightarrow \gamma(4S)$
< 4.1×10^{-4}	90		ALBRECHT	88H ARG $e^+ e^- \rightarrow \gamma(4S)$

114 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

115 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

 $\Gamma(K_2^*(1430)\gamma)/\Gamma_{\text{total}}$ Γ_{60}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.66^{+0.59}_{-0.53} ± 0.13	116 COAN	00	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

< 83 90 ALBRECHT 88H ARG $e^+ e^- \rightarrow \gamma(4S)$

116 COAN 00 obtains a fitted signal yield of $15.9^{+5.7}_{-5.2}$ events. A search for contamination by $K^*(1410)$ yielded a rate consistent with 0; the central value assumes no contamination.

 $\Gamma(K_2(1770)\gamma)/\Gamma_{\text{total}}$ Γ_{61}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.2 × 10⁻³	90	117 LESIAK	92 CBAL	$e^+ e^- \rightarrow \gamma(4S)$

117 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

$\Gamma(K_3^*(1780)\gamma)/\Gamma_{\text{total}}$		Γ_{62}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-3}$	90	ALBRECHT	88H ARG	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(K_4^*(2045)\gamma)/\Gamma_{\text{total}}$		Γ_{63}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-3}$	90	118 LESIAK	92 CBAL	$e^+ e^- \rightarrow \gamma(4S)$

118 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s -quark hadronization.

$\Gamma(K\eta'(958))/\Gamma_{\text{total}}$		Γ_{64}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$(8.3^{+0.9}_{-0.8} \pm 0.7) \times 10^{-5}$	119 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	

119 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)\eta'(958))/\Gamma_{\text{total}}$		Γ_{65}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	90	120 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

120 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K\eta)/\Gamma_{\text{total}}$		Γ_{66}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-6}$	90	121 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

121 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)\eta)/\Gamma_{\text{total}}$		Γ_{67}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$(1.80^{+0.49}_{-0.43} \pm 0.18) \times 10^{-5}$	122 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	

122 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K\phi\phi)/\Gamma_{\text{total}}$		Γ_{68}/Γ		
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$2.3^{+0.9}_{-0.8} \pm 0.3$	123 HUANG	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$	

123 Assumes equal production of charged and neutral B meson pairs and isospin symmetry.

$\Gamma(\bar{b} \rightarrow \bar{s}\gamma)/\Gamma_{\text{total}}$		Γ_{69}/Γ		
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
3.41 ± 0.34 OUR AVERAGE				
$3.55 \pm 0.32 \pm 0.32$	124 KOPPENBURG04	BELL	$e^+ e^- \rightarrow \gamma(4S)$	
$3.21 \pm 0.43^{+0.32}_{-0.29}$	125 CHEN	01C CLE2	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.36 \pm 0.53^{+0.65}_{-0.68}$	126 ABE	01F BELL	Repl. by KOPPEN-BURG 04
$2.32 \pm 0.57 \pm 0.35$	ALAM	95 CLE2	Repl. by CHEN 01C

124 KOPPENBURG 04 also report measurements of first and second moments of the photon energy spectrum above 1.8 GeV, $\langle E_\gamma \rangle = 2.292 \pm 0.026 \pm 0.034$ GeV, and $\langle E_\gamma^2 \rangle - \langle E_\gamma \rangle^2 = 0.0305 \pm 0.0074 \pm 0.0063$ GeV².

125 We have combined the experimental systematic theoretical uncertainties in quadrature. Also determined the first and second moments of the photon energy spectrum above 2.0 GeV: $\langle E_\gamma \rangle = 2.346 \pm 0.032 \pm 0.011$ GeV and $\langle E_\gamma^2 \rangle - \langle E_\gamma \rangle^2 = 0.0226 \pm 0.0066 \pm 0.0020$ GeV².

126 ABE 01F reports their systematic errors $\pm 0.42^{+0.50}_{-0.54}$, where the second error is due to the theoretical uncertainty. We combine them in quadrature.

$\Gamma(\bar{b} \rightarrow \bar{s}\text{gluon})/\Gamma_{\text{total}}$

Γ_{70}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.068	90	127	COAN	98	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.08	2	128	ALBRECHT	95D ARG	$e^+ e^- \rightarrow \gamma(4S)$

127 COAN 98 uses D - ℓ correlation.

128 ALBRECHT 95D use full reconstruction of one B decay as tag. Two candidate events for charmless B decay can be interpreted as either $b \rightarrow s\text{gluon}$ or $b \rightarrow u$ transition. If interpreted as $b \rightarrow s\text{gluon}$ they find a branching ratio of ~ 0.026 or the upper limit quoted above. Result is highly model dependent.

$\Gamma(\eta \text{anything})/\Gamma_{\text{total}}$

Γ_{71}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.4 × 10 ⁻⁴	90	129	BROWDER	98 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

129 BROWDER 98 search for high momentum $B \rightarrow \eta X_s$ between 2.1 and 2.7 GeV/c.

$\Gamma(\eta' \text{anything})/\Gamma_{\text{total}}$

Γ_{72}/Γ

VALUE (units 10 ⁻⁴)	DOCUMENT ID	TECN	COMMENT
4.2±0.9 OUR AVERAGE			
3.9 ± 0.8 ± 0.9	130 AUBERT,B	04F BABR	$e^+ e^- \rightarrow \gamma(4S)$
4.6 ± 1.1 ± 0.6	131 BONVICINI	03 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
6.2 ± 1.6 ^{+1.3} _{-2.0}	132 BROWDER	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

130 The reported branching ratio is for high momentum η between 2.0 and 2.7 GeV in the $\gamma(4S)$ center-of-mass frame. X_s represents a recoil system consisting of a kaon and zero to four pions.

131 BONVICINI 03 observed a signal of 61.2 ± 13.9 events in $B \rightarrow \eta' X_{nc}$ production for high momentum η' between 2.0 and 2.7 GeV/c in the $\gamma(4S)$ center-of-mass frame. The X_{nc} denotes “charmless” hadronic states recoiling against η' . The second error combines systematic and background subtraction uncertainties in quadrature.

132 BROWDER 98 observed a signal of 39.0 ± 11.6 events in high momentum $B \rightarrow \eta' X_s$ production between 2.0 and 2.7 GeV/c. The branching fraction is based on the interpretation of $b \rightarrow sg$, where the last error includes additional uncertainties due to the color-suppressed $b \rightarrow$ backgrounds.

$\Gamma(\rho\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{73}/Γ
$<1.9 \times 10^{-6}$	90	133 AUBERT	04C BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<1.4 \times 10^{-5}$	90	134 COAN	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
133 Assumes $\Gamma(B \rightarrow \rho\gamma) = \Gamma(B^+ \rightarrow \rho^+\gamma) = 2\Gamma(B^0 \rightarrow \rho^0\gamma)$ and uses lifetime ratio of $\tau_{B^+}/\tau_{B^0} = 1.083 \pm 0.017$.					
134 COAN 00 reports $B(B \rightarrow \rho\gamma)/B(B \rightarrow K^*(892)\gamma) < 0.32$ at 90%CL and scaled by the central value of $B(B \rightarrow K^*(892)\gamma) = (4.24 \pm 0.54 \pm 0.32) \times 10^{-5}$.					

 $\Gamma(\rho/\omega\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{74}/Γ
$<1.2 \times 10^{-6}$	90	AUBERT	05	BABR	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(\pi^\pm \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{75}/Γ
$3.585 \pm 0.025 \pm 0.070$	135 ALBRECHT	93I ARG	$e^+ e^- \rightarrow \gamma(4S)$	
135 ALBRECHT 93 excludes π^\pm from K_S^0 and Λ decays. If included, they find $4.105 \pm 0.025 \pm 0.080$.				

 $\Gamma(\pi^0 \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{76}/Γ
$2.35 \pm 0.02 \pm 0.11$	136 ABE	01J BELL	$e^+ e^- \rightarrow \gamma(4S)$	
136 From fully inclusive π^0 yield with no corrections from decays of K_S^0 or other particles.				

 $\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{77}/Γ
$0.176 \pm 0.011 \pm 0.012$	KUBOTA	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	

 $\Gamma(\rho^0 \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{78}/Γ
$0.208 \pm 0.042 \pm 0.032$	ALBRECHT	94J ARG	$e^+ e^- \rightarrow \gamma(4S)$	

 $\Gamma(\omega \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{79}/Γ
<0.81	ALBRECHT	94J ARG	$e^+ e^- \rightarrow \gamma(4S)$	

 $\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{80}/Γ
0.0342 ± 0.0013 OUR AVERAGE	AUBERT	04S BABR	$e^+ e^- \rightarrow \gamma(4S)$	
0.0341 $\pm 0.0006 \pm 0.0012$	ALBRECHT	94J ARG	$e^+ e^- \rightarrow \gamma(4S)$	
0.0390 $\pm 0.0030 \pm 0.0035$	BORTOLETTO86	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	

 $\Gamma(\phi K^*(892))/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	Γ_{81}/Γ
$<2.2 \times 10^{-5}$	90	137 BERGFELD	98 CLE2	

137 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.064±0.008±0.008		138 CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.14 ± 0.09		139 ALBRECHT	88E ARG	$e^+ e^- \rightarrow \gamma(4S)$
<0.112	90	140 ALAM	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
138 CRAWFORD 92 result derived from lepton baryon correlations. Assumes all charmed baryons in B^0 and B^\pm decay are Λ_c .				
139 ALBRECHT 88E measured $B(B \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (0.30 \pm 0.12 \pm 0.06)\%$ and used $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (2.2 \pm 1.0)\%$ from ABRAMS 80 to obtain above number.				
140 Assuming all baryons result from charmed baryons, ALAM 86 conclude the branching fraction is $7.4 \pm 2.9\%$. The limit given above is model independent.				

 $\Gamma(\Lambda_c^+ \text{ anything})/\Gamma(\bar{\Lambda}_c^- \text{ anything})$ Γ_{83}/Γ_{84}

VALUE	DOCUMENT ID	TECN	COMMENT
0.19±0.13±0.04	141 AMMAR	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
141 AMMAR 97 uses a high-momentum lepton tag ($P_\ell > 1.4 \text{ GeV}/c^2$).			

 $\Gamma(\bar{\Lambda}_c^- e^+ \text{ anything})/\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{ anything})$ Γ_{85}/Γ_{82}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.05	90	142 BONVICINI	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
142 BONVICINI 98 uses the electron with momentum above $0.6 \text{ GeV}/c$.				

 $\Gamma(\bar{\Lambda}_c^- p \text{ anything})/\Gamma(\Lambda_c^+ / \bar{\Lambda}_c^- \text{ anything})$ Γ_{86}/Γ_{82}

VALUE	DOCUMENT ID	TECN	COMMENT
0.57±0.05±0.05	BONVICINI 98	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(\bar{\Lambda}_c^- p e^+ \nu_e)/\Gamma(\bar{\Lambda}_c^- \text{ anything})$ Γ_{87}/Γ_{86}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	143 BONVICINI	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
143 BONVICINI 98 uses the electron with momentum above $0.6 \text{ GeV}/c$.				

 $\Gamma(\bar{\Sigma}_c^{--} \text{ anything})/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0042±0.0021±0.0011	77	144 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
144 PROCARIO 94 reports $[B(B \rightarrow \bar{\Sigma}_c^{--} \text{ anything}) \times B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = 0.00021 \pm 0.00008 \pm 0.00007$. We divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(\bar{\Sigma}_c^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.010	90	145 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
145 PROCARIO 94 reports $[B(B \rightarrow \bar{\Sigma}_c^- \text{ anything}) \times B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = < 0.00048$. We divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.050$.				

$\Gamma(\Sigma_c^0 \text{anything})/\Gamma_{\text{total}}$	Γ_{90}/Γ				
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.0046 ± 0.0021 ± 0.0012	76	146 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
146 PROCARIO 94 reports $[B(B \rightarrow \Sigma_c^0 \text{anything}) \times B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = 0.00023 \pm 0.00008 \pm 0.00007$. We divide by our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(\Sigma_c^0 N(N=p \text{ or } n))/\Gamma_{\text{total}}$	Γ_{91}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0015	90	147 PROCARIO	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
147 PROCARIO 94 reports < 0.0017 for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.043$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.050$.				

$\Gamma(\Xi_c^0 \text{anything} \times B(\Xi_c^0 \rightarrow \Xi^- \pi^+)) / \Gamma_{\text{total}}$	Γ_{92}/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.144 \pm 0.048 \pm 0.021	148 BARISH	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
148 BARISH 97 find $79 \pm 27 \Xi_c^0$ events.			

$\Gamma(\Xi_c^+ \text{anything} \times B(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+))/\Gamma_{\text{total}}$	Γ_{93}/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$0.453 \pm 0.096^{+0.085}_{-0.065}$	149 BARISH	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(p/\bar{p}\text{anything})/\Gamma_{\text{total}}$	Γ_{94}/Γ			
Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.				
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.080±0.004 OUR AVERAGE				
0.080±0.005±0.005		ALBRECHT	93I	ARG $e^+ e^- \rightarrow \gamma(4S)$
0.080±0.005±0.003		CRAWFORD	92	CLEO $e^+ e^- \rightarrow \gamma(4S)$
+0.013	150			

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.021 $^{151}\text{ALAM}$ 83B CLEO $e^+ e^- \rightarrow \gamma(4S)$

¹⁵⁰ALBRECHT 89K include direct and nondirect protons.

¹⁵¹ ALAM 83B reported their result as $> 0.036 \pm 0.006 \pm 0.009$. Data are consistent with equal yields of p and \bar{p} . Using assumed yields below cut, $B(B \rightarrow p + X) = 0.03$ not including protons from Λ decays.

$\Gamma(p/\bar{p}(\text{direct}) \text{ anything})/\Gamma_{\text{total}}$	Γ_{95}/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.055±0.005 OUR AVERAGE				
0.055±0.005±0.0035		ALBRECHT	93I ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.056±0.006±0.005		CRAWFORD	92 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.055±0.016	1220	152 ALBRECHT	89K ARG	$e^+ e^- \rightarrow \gamma(4S)$

¹⁵²ALBRECHT 89K subtract contribution of Λ decay from the inclusive proton yield.

$\Gamma(\Lambda/\bar{\Lambda}\text{anything})/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.040±0.005 OUR AVERAGE				
0.038±0.004±0.006	2998	CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.042±0.005±0.006	943	ALBRECHT 89K	ARG	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.022±0.003±0.0022	153	ACKERSTAFF 97N	OPAL	$e^+ e^- \rightarrow Z$
>0.011	154	ALAM 83B	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
153 ACKERSTAFF 97N assumes $B(b \rightarrow B) = 0.868 \pm 0.041$, i.e., an admixture of B^0 , B^\pm , and B_s .				
154 ALAM 83B reported their result as $> 0.022 \pm 0.007 \pm 0.004$. Values are for $(B(\Lambda X) + B(\bar{\Lambda} X))/2$. Data are consistent with equal yields of p and \bar{p} . Using assumed yields below cut, $B(B \rightarrow \Lambda X) = 0.03$.				

$\Gamma(\Lambda\text{anything})/\Gamma(\bar{\Lambda}\text{anything})$ Γ_{97}/Γ_{98}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.43±0.09±0.07				
155	AMMAR 97	97	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
155 AMMAR 97 uses a high-momentum lepton tag ($P_\ell > 1.4 \text{ GeV}/c^2$).				

$\Gamma(\Xi^-\bar{\Xi}^+\text{anything})/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0027±0.0006 OUR AVERAGE				
0.0027±0.0005±0.0004	147	CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.0028±0.0014	54	ALBRECHT 89K	ARG	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\text{baryons anything})/\Gamma_{\text{total}}$ Γ_{100}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.068±0.005±0.003				
156	ALBRECHT 920	920	ARG	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.076±0.014	157	ALBRECHT 89K	ARG	$e^+ e^- \rightarrow \gamma(4S)$
156 ALBRECHT 920 result is from simultaneous analysis of p and Λ yields, $p\bar{p}$ and $\Lambda\bar{p}$ correlations, and various lepton-baryon and lepton-baryon-antibaryon correlations. Supersedes ALBRECHT 89K.				
157 ALBRECHT 89K obtain this result by adding their measurements ($5.5 \pm 1.6\%$) for direct protons and ($4.2 \pm 0.5 \pm 0.6\%$) for inclusive Λ production. They then assume ($5.5 \pm 1.6\%$) for neutron production and add it in also. Since each B decay has two baryons, they divide by 2 to obtain ($7.6 \pm 1.4\%$).				

$\Gamma(p\bar{p}\text{anything})/\Gamma_{\text{total}}$ Γ_{101}/Γ

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0247±0.0023 OUR AVERAGE				
0.024 ± 0.001 ± 0.004		CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.025 ± 0.002 ± 0.002	918	ALBRECHT 89K	ARG	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(p\bar{p}\text{anything})/\Gamma(p/\bar{p}\text{anything})$ Γ_{101}/Γ_{94}

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.30±0.02±0.05	158	CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
158 CRAWFORD 92 value is not independent of their $\Gamma(p\bar{p}\text{anything})/\Gamma_{\text{total}}$ value.				

$\Gamma(\Lambda\bar{p}/\bar{\Lambda}p\text{anything})/\Gamma_{\text{total}}$

Γ_{102}/Γ

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.025±0.004 OUR AVERAGE				
0.029±0.005±0.005		CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
0.023±0.004±0.003	165	ALBRECHT 89K	ARG	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\Lambda\bar{p}/\bar{\Lambda}p\text{anything})/\Gamma(\Lambda/\bar{\Lambda}\text{anything})$

Γ_{102}/Γ_{96}

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.76±0.11±0.08	159 CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
159 CRAWFORD 92 value	is not independent of their $[\Gamma(\Lambda\bar{p}\text{anything})+\Gamma(\bar{\Lambda}p\text{anything})]/\Gamma_{\text{total}}$ value.		

$\Gamma(\Lambda\bar{\Lambda}\text{anything})/\Gamma_{\text{total}}$

Γ_{103}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.005	90		CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0088	90	12	ALBRECHT 89K	ARG	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\Lambda\bar{\Lambda}\text{anything})/\Gamma(\Lambda/\bar{\Lambda}\text{anything})$

Γ_{103}/Γ_{96}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.13	90	160 CRAWFORD 92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
160 CRAWFORD 92 value	is not independent of their $\Gamma(\Lambda\bar{\Lambda}\text{anything})/\Gamma_{\text{total}}$ value.			

$\Gamma(s e^+ e^-)/\Gamma_{\text{total}}$

Γ_{104}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
5.6±1.7 OUR AVERAGE				
6.0±1.7±1.3		161 AUBERT,B 04I	BABR	$e^+ e^- \rightarrow \gamma(4S)$
5.0±2.3 ^{+1.3} _{-1.1}		161 KANEKO 03	BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 57	90	GLENN 98	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
<50000	90	BEBEK 81	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

161 Requires $M_{e^+ e^-} > 0.2 \text{ GeV}/c^2$.

$\Gamma(s \mu^+ \mu^-)/\Gamma_{\text{total}}$

Γ_{105}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
6.6±2.0 OUR AVERAGE				
5.0±2.8±1.2		AUBERT,B 04I	BABR	$e^+ e^- \rightarrow \gamma(4S)$
7.9±2.1 ^{+2.1} _{-1.5}		KANEKO 03	BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 58	90	GLENN 98	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
<17000	90	CHADWICK 81	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(s e^+ e^-) + \Gamma(s \mu^+ \mu^-) / \Gamma_{\text{total}}$ $\Gamma_{104} + \Gamma_{105} / \Gamma$

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.2 \times 10^{-5}$	90	GLENN	98	CLEO $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0024	90	162 BEAN	87	CLEO Repl. by GLENN 98
<0.0062	90	163 AVERY	84	CLEO Repl. by BEAN 87

162 BEAN 87 reports $[(\mu^+ \mu^-) + (e^+ e^-)]/2$ and we converted it.

163 Determine ratio of B^+ to B^0 semileptonic decays to be in the range 0.25–2.9.

$\Gamma(s \ell^+ \ell^-) / \Gamma_{\text{total}}$ Γ_{106} / Γ

Test for $\Delta B = 1$ weak neutral current.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
5.9 ± 1.4 OUR AVERAGE			
5.6 ± 1.5 ± 1.3	164 AUBERT,B	04I BABR	$e^+ e^- \rightarrow \gamma(4S)$
6.1 ± 1.4 ± 1.4	164 KANEKO	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$

164 Requires $M_{e^+ e^-} > 0.2 \text{ GeV}/c^2$.

$\Gamma(K e^+ e^-) / \Gamma_{\text{total}}$ Γ_{107} / Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
6.0 ± 1.4 OUR AVERAGE Error includes scale factor of 1.1.				
7.4 ± 1.8 ± 0.5	165 AUBERT	03U BABR	$e^+ e^- \rightarrow \gamma(4S)$	
4.8 ± 1.5 ± 0.3	165,166 ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13 90 ABE 02 BELL Repl. by ISHIKAWA 03

165 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

166 The second error is a total of systematic uncertainties including model dependence.

$\Gamma(K^*(892) e^+ e^-) / \Gamma_{\text{total}}$ Γ_{108} / Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
1.24 ± 0.37 OUR AVERAGE				
0.98 ± 0.50 ± 0.11	167 AUBERT	03U BABR	$e^+ e^- \rightarrow \gamma(4S)$	
1.49 ± 0.52 ± 0.12	168 ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5.6 90 ABE 02 BELL Repl. by ISHIKAWA 03

167 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

168 Assumes equal production of B^0 and B^+ at $\gamma(4S)$. The second error is a total of systematic uncertainties including model dependence.

$\Gamma(K\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{109}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	DOCUMENT ID	TECN	COMMENT
(4.7 ± 1.1) $\times 10^{-7}$ OUR AVERAGE			
(4.5 ± 2.3 ± 0.4) $\times 10^{-7}$	169 AUBERT	03U BABR	$e^+ e^- \rightarrow \gamma(4S)$
(4.8 ± 1.2 ± 0.4) $\times 10^{-7}$	170 ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
(0.99 ± 0.40 ± 0.13) $\times 10^{-6}$	ABE	02 BELL	Repl. by ISHIKAWA 03

169 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

170 Assumes equal production of B^0 and B^+ at $\gamma(4S)$. The second error is a total of systematic uncertainties including model dependence.

 $\Gamma(K^*(892)\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{110}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
1.19 ± 0.34 OUR AVERAGE				
1.27 ± 0.76 ± 0.16		171 AUBERT	03U BABR	$e^+ e^- \rightarrow \gamma(4S)$
1.17 ± 0.36 ± 0.10		172 ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<3.1	90	ABE	02 BELL	Repl. by ISHIKAWA 03

171 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

172 Assumes equal production of B^0 and B^+ at $\gamma(4S)$. The second error is a total of systematic uncertainties including model dependence.

 $\Gamma(K\ell^+\ell^-)/\Gamma_{\text{total}}$ Γ_{111}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
0.54 ± 0.08 OUR AVERAGE				
0.65 ± 0.14 ± 0.04		173 AUBERT	03U BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.48 ± 0.10 ± 0.03		174 ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.75 ± 0.25 ± 0.06		175 ABE	02 BELL	Repl. by ISHIKAWA 03
<0.51	90	176 AUBERT	02L BABR	$e^+ e^- \rightarrow \gamma(4S)$
<1.7	90	177 ANDERSON	01B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

173 Assumes all four $B \rightarrow K\ell^+\ell^-$ modes having equal partial widths in the fit.

174 Assumes equal production rate for charge and neutral B meson pairs, isospin invariance, lepton universality for $B \rightarrow K\ell^+\ell^-$, and $B(B \rightarrow K^*(892)\mu^+\mu^-) = 1.33$. The second error is total systematic uncertainties including model dependence.

175 Assumes lepton universality.

176 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

177 The result is for di-lepton masses above 0.5 GeV.

$\Gamma(K^*(892)\ell^+\ell^-)/\Gamma_{\text{total}}$ Γ_{112}/Γ Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
1.05 ± 0.20 OUR AVERAGE				

0.88 $^{+0.33}_{-0.29} \pm 0.10$ 178 AUBERT 03U BABR $e^+ e^- \rightarrow \gamma(4S)$ 1.15 $^{+0.26}_{-0.24} \pm 0.08$ 179 ISHIKAWA 03 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.1 90 180,181 AUBERT 02L BABR Repl. by AUBERT 03U

<3.3 90 182 ANDERSON 01B CLE2 $e^+ e^- \rightarrow \gamma(4S)$ 178 Assumes the partial width ratio of electron and muon modes to be $\Gamma(B \rightarrow K^*(892)e^+e^-)/\Gamma(B \rightarrow K^*(892)\mu^+\mu^-) = 1.33$.179 Assumes equal production rate for charge and neutral B meson pairs, isospin invariance, lepton universality for $B \rightarrow K\ell^+\ell^-$, and $B(B \rightarrow K^*(892)\mu^+\mu^-) = 1.33$. The second error is total systematic uncertainties including model dependence.180 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.181 For averaging $K^*(892)\mu^+\mu^-$ and $K^*(892)e^+e^-$ modes, AUBERT 02L assumed $B(B \rightarrow K^*(892)e^+e^-)/B(B \rightarrow K^*(892)\mu^+\mu^-) = 1.2$.

182 The result is for di-lepton masses above 0.5 GeV.

 $\Gamma(e^\pm\mu^\mp s)/\Gamma_{\text{total}}$ Γ_{113}/Γ

Test for lepton family number conservation. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.2 × 10⁻⁵				

 $\Gamma(\pi e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{114}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.6 × 10⁻⁶				

183 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\rho e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{115}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.2 × 10⁻⁶				

184 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{116}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.6 × 10⁻⁶				

185 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K^*(892)e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{117}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.2 × 10⁻⁶				

186 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

***CP* VIOLATION**

A_{CP} is defined as

$$\frac{B(B \rightarrow \bar{f}) - B(\bar{B} \rightarrow f)}{B(B \rightarrow \bar{f}) + B(\bar{B} \rightarrow f)},$$

the CP -violation charge asymmetry of inclusive B^\pm and B^0 decay.

$A_{CP}(B \rightarrow K^*(892)\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.010 ± 0.028 OUR AVERAGE			
$-0.013 \pm 0.036 \pm 0.010$	187 AUBERT,BE	04A BABR	$e^+ e^- \rightarrow \gamma(4S)$
$-0.015 \pm 0.044 \pm 0.012$	188 NAKAO	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$
$+0.08 \pm 0.13 \pm 0.03$	188 COAN	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$-0.044 \pm 0.076 \pm 0.012$	189 AUBERT	02C BABR	Repl. by AUBERT,BE 04A

187 Corresponds to a 90% CL allowed region, $-0.074 < A_{CP} < 0.049$.

188 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

189 A 90% CL range is $-0.170 < A_{CP} < 0.082$.

$A_{CP}(B \rightarrow s\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.00 ± 0.04 OUR AVERAGE			
$0.025 \pm 0.050 \pm 0.015$	190 AUBERT,B	04E BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.002 \pm 0.050 \pm 0.030$	191 NISHIDA	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$
$-0.079 \pm 0.108 \pm 0.022$	192 COAN	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
190 Corresponds to $-0.06 < A_{CP} < +0.11$ at 90% CL.			
191 This measurement is performed inclusively for recoil mass X_s less than 2.1 GeV, which corresponds to $-0.093 < A_{CP} < 0.096$ at 90% CL.			
192 Corresponds to $-0.27 < A_{CP} < 0.10$ at 90% CL.			

$A_{CP}(b \rightarrow X_s \ell^+ \ell^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.22 \pm 0.26 \pm 0.02$			
193 AUBERT,B	04I BABR	$e^+ e^- \rightarrow \gamma(4S)$	
193 The final state flavor is determined by the kaon and pion charges where modes with $X_s = K_S^0, K_S^0 \pi^0$ or $K_S^0 \pi^+ \pi^-$ are not used.			

$B \rightarrow X_c \ell \nu$ HADRONIC MASS MOMENTS

$\langle M_X^2 - \bar{M}_D^2 \rangle$ (First Moments)

VALUE (GeV ²)	DOCUMENT ID	TECN	COMMENT
$0.293 \pm 0.012 \pm 0.058$			
194 CSORNA	04 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.251 $\pm 0.023 \pm 0.062$	195 CRONIN-HEN..01B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
194 Uses minimum lepton energy of 1.5 GeV and also reports moments with $E_\ell > 1.0$ GeV.			
195 The leptons are required to have $P_1 > 1.5$ GeV/c.			

$\langle (M_X^2 - \bar{M}_X^2)^2 \rangle$ (Second Moments)

VALUE (GeV ⁴)	DOCUMENT ID	TECN	COMMENT
0.629±0.031±0.143	196 CSORNA 04	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.576±0.048±0.168	197 CRONIN-HEN..01B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
196 Uses minimum lepton energy of 1.5 GeV and also reports moments with $E_\ell > 1.0$ GeV.			
197 The leptons are required to have $P_1 > 1.5$ GeV/c.			

$\langle (M_X^2 - \bar{M}_D^2)^2 \rangle$ (Second Moments)

VALUE (GeV ⁴)	DOCUMENT ID	TECN	COMMENT
0.639±0.056±0.178	198 CRONIN-HEN..01B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
198 The leptons are required to have $P_1 > 1.5$ GeV/c.			


$B \rightarrow X_c \ell \nu$ LEPTON MOMENTUM MOMENTS

$R_0 (\Gamma_{E_l > 1.7\text{GeV}} / \Gamma_{E_l > 1.5\text{GeV}})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.6187±0.0014±0.0016	199 MAHMOOD 03	CLE2	$e^+ e^- \rightarrow \gamma(4S)$
199 The leptons are required to have $E_l > 1.5$ GeV in the B rest frame.			

$R_1 (\langle E_l \rangle_{E_l > 1.5\text{GeV}})$

VALUE	DOCUMENT ID	TECN	COMMENT
1.7797±0.0018 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.		
1.7743±0.0019±0.0014			
1.7792±0.0021±0.0027	200 AUBERT,B 04A BABR	$e^+ e^- \rightarrow \gamma(4S)$	
1.7810±0.0007±0.0009	201 MAHMOOD 04 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
202 MAHMOOD 03 CLE2 $e^+ e^- \rightarrow \gamma(4S)$			
200 The leptons are required to have $E_l > 1.5$ GeV in the B rest frame. The result with $E_l > 0.6$ GeV is also given.			
201 Uses $E_e > 1.5$ GeV and also reports moments with other minimum minimum E_e conditions, as low as $E_e > 0.6$ GeV.			
202 The leptons are required to have $E_l > 1.5$ GeV in the B rest frame.			



$R_2 (\langle E_l^2 - \bar{E}_l^2 \rangle_{E_l > 1.5 GeV})$

VALUE (10^{-3} GeV^2)	DOCUMENT ID	TECN	COMMENT
30.8±0.8 OUR AVERAGE			
30.3±0.9±0.5	203 AUBERT,B	04A BABR	$e^+ e^- \rightarrow \gamma(4S)$
31.6±0.8±1.0	204 MAHMOOD	04 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
203 The leptons are required to have $E_l > 1.5 \text{ GeV}$ in the B rest frame. The result with $E_l > 0.6 \text{ GeV}$ is also given.			
204 Uses $E_e > 1.5 \text{ GeV}$ and also reports moments with other minimum minimum E_e conditions, as low as $E_e > 0.6 \text{ GeV}$.			

$R_3 (\langle E_l^3 - \bar{E}_l^3 \rangle_{E_l > 1.5 GeV})$

VALUE (10^{-3} GeV^3)	DOCUMENT ID	TECN	COMMENT
2.12±0.47±0.20			
2.05	205 AUBERT,B	04A BABR	$e^+ e^- \rightarrow \gamma(4S)$
205 The leptons are required to have $E_l > 1.5 \text{ GeV}$ in the B rest frame. The result with $E_l > 0.6 \text{ GeV}$ is also given.			

B^\pm/B^0 ADMIXTURE REFERENCES

AUBERT	05	PRL 94 011801	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABDALLAH	04D	EPJ C33 213	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
AUBERT	04C	PRL 92 111801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	04I	PRL 92 071802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04S	PR D69 052005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04X	PRL 93 011803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04A	PR D69 111104R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04E	PRL 93 021804	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04F	PRL 93 061801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04I	PRL 93 081802	B. Aubert <i>et al.</i>	(BABAR Collab.)

AUBERT,BE	04A	PR D70 112006	B. Aubert <i>et al.</i>	(BABAR Collab.)
CSORNA	04	PR D70 032002	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
KOPPENBURG	04	PRL 93 061803	P. Koppenburg <i>et al.</i>	(BELLE Collab.)
MAHMOOD	04	PR D70 032003	A.H. Mahmood <i>et al.</i>	(CLEO Collab.)
NAKAO	04	PR D69 112001	M. Nakao <i>et al.</i>	(BELLE Collab.)
NISHIDA	04	PRL 93 031803	S. Nishida <i>et al.</i>	(BELLE Collab.)
ADAM	03B	PR D68 012004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
AUBERT	03	PR D67 031101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03F	PR D67 032002	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03U	PRL 91 221802	B. Aubert <i>et al.</i>	(BaBar Collab.)
BONVICINI	03	PR D68 011101R	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
ISHIKAWA	03	PRL 91 261601	A. Ishikawa <i>et al.</i>	(BELLE Collab.)
KANEKO	03	PRL 90 021801	J. Kaneko <i>et al.</i>	(BELLE Collab.)
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)
MAHMOOD	03	PR D67 072001	A.H. Mahmood <i>et al.</i>	(CLEO Collab.)
ABE	02	PRL 88 021801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02L	PRL 89 011803	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02Y	PL B547 181	K. Abe <i>et al.</i>	(BELLE Collab.)
ANDERSON	02	PRL 89 282001	S. Anderson <i>et al.</i>	(CLEO Collab.)
AUBERT	02C	PRL 88 101805	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02G	PR D65 091104R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02L	PRL 88 241801	B. Aubert <i>et al.</i>	(BaBar Collab.)
EDWARDS	02B	PR D65 111102R	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ABE	01F	PL B511 151	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01J	PR D64 072001	K. Abe <i>et al.</i>	(BELLE Collab.)
ANDERSON	01B	PRL 87 181803	S. Anderson <i>et al.</i>	(CLEO Collab.)
CHEN	01	PR D63 031102	S. Chen <i>et al.</i>	(CLEO Collab.)
CHEN	01C	PRL 87 251807	S. Chen <i>et al.</i>	(CLEO Collab.)
COAN	01	PRL 86 5661	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HENNESSY	01B	PRL 87 251808	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PDG	01	Unofficial 2001 WWW edition		
ABREU	00R	PL B475 407	P. Abreu <i>et al.</i>	(DELPHI Collab.)
COAN	00	PRL 84 5283	T.E. Coan <i>et al.</i>	(CLEO Collab.)
RICHICHI	00	PRL 85 520	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERGFELD	98	PRL 81 272	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BISHAI	98	PR D57 3847	M. Bishai <i>et al.</i>	(CLEO Collab.)
BONVICINI	98	PR D57 6604	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BROWDER	98	PRL 81 1786	T.E. Browder <i>et al.</i>	(CLEO Collab.)
COAN	98	PRL 80 1150	T.E. Coan <i>et al.</i>	(CLEO Collab.)
GLENN	98	PRL 80 2289	S. Glenn <i>et al.</i>	(CLEO Collab.)
ACKERSTAFF	97N	ZPHY C74 423	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
AMMAR	97	PR D55 13	R. Ammar <i>et al.</i>	(CLEO Collab.)
BARISH	97	PRL 79 3599	B. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	97B	ZPHY C73 601	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
GIBBONS	97B	PR D56 3783	L. Gibbons <i>et al.</i>	(CLEO Collab.)
ALBRECHT	96D	PL B374 256	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)
GIBAUT	96	PR D53 4734	D. Gibaut <i>et al.</i>	(CLEO Collab.)
KUBOTA	96	PR D53 6033	Y. Kubota <i>et al.</i>	(CLEO Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	
ALAM	95	PRL 74 2885	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	95D	PL B353 554	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BALEST	95B	PR D52 2661	R. Balest <i>et al.</i>	(CLEO Collab.)
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	95B	PL B345 103	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ALBRECHT	94C	ZPHY C62 371	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	94J	ZPHY C61 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
PROCARIO	94	PRL 73 1472	M. Procario <i>et al.</i>	(CLEO Collab.)
ALBRECHT	93	ZPHY C57 533	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93E	ZPHY C60 11	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93H	PL B318 397	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	93I	ZPHY C58 191	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	93B	PL B319 365	J. Alexander <i>et al.</i>	(CLEO Collab.)
ARTUSO	93	PL B311 307	M. Artuso	(SYRA)
BARTELT	93B	PRL 71 4111	J.E. Bartelt <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92E	PL B277 209	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92G	ZPHY C54 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92O	ZPHY C56 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)

CRAWFORD	92	PR D45 752	G. Crawford <i>et al.</i>	(CLEO Collab.)
HENDERSON	92	PR D45 2212	S. Henderson <i>et al.</i>	(CLEO Collab.)
LESIAK	92	ZPHY C55 33	T. Lesiak <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	91C	PL B255 297	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91H	ZPHY C52 353	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
FULTON	91	PR D43 651	R. Fulton <i>et al.</i>	(CLEO Collab.)
YANAGISAWA	91	PRL 66 2436	C. Yanagisawa <i>et al.</i>	(CUSB II Collab.)
ALBRECHT	90	PL B234 409	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	90H	PL B249 359	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	90	PRL 64 2117	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
Also	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
FULTON	90	PRL 64 16	R. Fulton <i>et al.</i>	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
PDG	90	PL B239	J.J. Hernandez <i>et al.</i>	(IFIC, BOST, CIT+)
ALBRECHT	89K	ZPHY C42 519	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ISGUR	89B	PR D39 799	N. Isgur <i>et al.</i>	(TNTO, CIT)
WACHS	89	ZPHY C42 33	K. Wachs <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	88E	PL B210 263	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88H	PL B210 258	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KOERNER	88	ZPHY C38 511	J.G. Korner, G.A. Schuler	(MANZ, DESY)
ALAM	87	PRL 59 22	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALAM	87B	PRL 58 1814	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	87D	PL B199 451	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87H	PL B187 425	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BEAN	87	PR D35 3533	A. Bean <i>et al.</i>	(CLEO Collab.)
BEHRENDS	87	PRL 59 407	S. Behrends <i>et al.</i>	(CLEO Collab.)
BORTOLETTO	87	PR D35 19	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
ALAM	86	PR D34 3279	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BALTRUSAIT...	86E	PRL 56 2140	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BORTOLETTO	86	PRL 56 800	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
HAAS	86	PRL 56 2781	J. Haas <i>et al.</i>	(CLEO Collab.)
ALBRECHT	85H	PL 162B 395	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
CSORNA	85	PRL 54 1894	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
HAAS	85	PRL 55 1248	J. Haas <i>et al.</i>	(CLEO Collab.)
AVERY	84	PRL 53 1309	P. Avery <i>et al.</i>	(CLEO Collab.)
CHEN	84	PRL 52 1084	A. Chen <i>et al.</i>	(CLEO Collab.)
LEVMAN	84	PL 141B 271	G.M. Levman <i>et al.</i>	(CUSB Collab.)
ALAM	83B	PRL 51 1143	M.S. Alam <i>et al.</i>	(CLEO Collab.)
GREEN	83	PRL 51 347	J. Green <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83B	PL 130B 444	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALTARELLI	82	NP B208 365	G. Altarelli <i>et al.</i>	(ROMA, INFN, FRAS)
BRODY	82	PRL 48 1070	A.D. Brody <i>et al.</i>	(CLEO Collab.)
GIANNINI	82	NP B206 1	G. Giannini <i>et al.</i>	(CUSB Collab.)
BEBEK	81	PRL 46 84	C. Bebek <i>et al.</i>	(CLEO Collab.)
CHADWICK	81	PRL 46 88	K. Chadwick <i>et al.</i>	(CLEO Collab.)
ABRAMS	80	PRL 44 10	G.S. Abrams <i>et al.</i>	(SLAC, LBL)