

$f_0(600)$
or σ

$I^G(J^{PC}) = 0^+(0^{++})$

NOTE ON SCALAR MESONS

Updated January 2006 by S. Spanier (University of Tennessee)
and N.A. Törnqvist (Helsinki).

I. Introduction: In contrast to the vector and tensor mesons, the identification of the scalar mesons is a long-standing puzzle. Scalar resonances are difficult to resolve because of their large decay widths which cause a strong overlap between resonances and background, and also because several decay channels open up within a short mass interval. In addition, the $\bar{K}K$ and $\eta\eta$ thresholds produce sharp cusps in the energy dependence of the resonant amplitude. Furthermore, one expects non- $\bar{q}q$ scalar objects, like glueballs and multiquark states in the mass range below 1800 MeV. The number of experimental and theoretical publications since our last issue indicates great activity in this field. For some recent reviews see AMSLER 04, BUGG 04C, CLOSE 02B.

Scalars are produced, for example, in πN scattering on polarized/unpolarized targets, $\bar{p}p$ annihilation, central hadronic production, J/Ψ , B -, D - and K -meson decays, $\gamma\gamma$ formation, and ϕ radiative decays. Experiments are accompanied by the development of theoretical models for the reaction amplitudes, which are based on common fundamental principles of two-body unitarity, analyticity, Lorentz invariance, and chiral- and flavour-symmetry using different techniques (K -matrix formalism, N/D -method, Dalitz Tuan ansatz, unitarized quark models with coupled channels, effective chiral field theories like the linear sigma model, *etc.*). Dynamics near the lowest two-body thresholds in some analyses is described by crossed

channel (t, u) meson exchange or with an effective range parameterization instead of or in addition to resonant features in the s -channel, only. Furthermore, elastic S -wave scattering amplitudes involving soft pions have zeros close to threshold (ADLER 65, 65A), which may be shifted or removed in associated production processes.

The mass and width of a resonance are found from the position of the nearest pole in the process amplitude (T -matrix or S -matrix) at an unphysical sheet of the complex energy plane: $(E - i\Gamma/2)$. It is important to notice that only in the case of narrow well-separated resonances, far away from the opening of decay channels, does the naive Breit-Wigner parameterization (or K -matrix pole parametrization) agree with this pole position.

In this note, we discuss all light scalars organized in the listings under the entries ($I = 1/2$) $K_0^*(800)$ (or κ), $K^*(1430)$, ($I = 1$) $a_0(980)$, $a_0(1450)$, and ($I = 0$) $f_0(600)$ (or σ), $f_0(980)$, $f_0(1370)$, and $f_0(1500)$. This list is minimal and does not necessarily exhaust the list of actual resonances. The ($I = 2$) $\pi\pi$ and ($I = 3/2$) $K\pi$ phase shifts do not exhibit any resonant behavior. See also our notes in previous issues for further comments on *e.g.*, scattering lengths and older papers.

II. The $I = 1/2$ States: The $K^*(1430)$ (ASTON 88) is perhaps the least controversial of the light scalar mesons. The $K\pi$ S-wave scattering has two possible isospin channels, $I = 1/2$ and $I = 3/2$. The $I = 3/2$ wave is elastic and repulsive up to 1.7 GeV (ESTABROOKS 78) and contains no known resonances. The $I = 1/2$ $K\pi$ phase shift, measured from about 100 MeV above threshold in Kp production, rises smoothly, passes 90° at 1350 MeV, and continues to rise to about 170° at 1600 MeV. The first important inelastic threshold is $K\eta'(958)$. In the inelastic region the continuation of

the amplitude is uncertain since the partial-wave decomposition has several solutions. The data are extrapolated towards the $K\pi$ threshold using effective range type formulas (ASTON 88, ABELE 98) or chiral perturbation predictions (BERNARD 91, JAMIN 00, CHERRY 01). In analyses using unitarized amplitudes there is agreement on the presence of a resonance pole around 1410 MeV having a width of about 300 MeV. In recent years there has been controversy about the existence of a light and very broad “ κ ” meson in the 700-900 MeV region (*e.g.* D -meson decay analyses LINK 02, AITALA 02, 06). Some authors find this pole in their phenomenological analysis (see *e.g.* PALAEZ 04A, ZHENG 04, ISHIDA 03, BLACK 01,03, BUGG 03, DELBOURGO 98, OLLER 99, 99C, SCADRON 03, ANISOVICH 97C, JAMIN 00, SHAKIN 01), while others do not (*e.g.* CHERRY 01, KOPP 01, LINK 05I). Since it appears to be a very wide object ($\Gamma \approx 500$ MeV) near the $K\pi$ threshold, its presence and properties are difficult to establish on data.

In an important observation BES finds a κ like structure in J/ψ decays to $\bar{K}^{*0}(892)K^+\pi^-$ where κ recoils against the $K^*(892)$ (ABLIKIM 06C).

III. The $I = 1$ States: Two isovector states are known, the established $a_0(980)$ and the $a_0(1450)$. Independent of any model, the $\bar{K}K$ component in the $a_0(980)$ wave function must be large: it lies just below the opening of the $\bar{K}K$ channel to which it strongly couples. This generates an important cusp-like behavior in the resonant amplitude. Hence, its mass and width parameters are strongly distorted. To reveal its true coupling constants, a coupled channel model with energy-dependent widths and mass shift contributions is necessary. In all measurements in our listings, the mass position agrees on a value near 984 MeV, but the width takes values between 50 and

300 MeV, mostly due to the different models. For example, the analysis of the $\bar{p}p$ -annihilation data using an unitary K -matrix description finds a width as determined from the T -matrix pole of 92 ± 8 MeV, while the observed width of the peak in the $\pi\eta$ mass spectrum is about 45 MeV.

The relative coupling $\bar{K}K/\pi\eta$ is determined indirectly from $f_1(1285)$ (BARBERIS 98C, CORDEN 78, DEFOIX 72) or $\eta(1410)$ decays (BAI 90C, BOLTON 92B, AMSLER 95C), from the line shape observed in the $\pi\eta$ decay mode (FLATTE 76, AMSLER 94D, BUGG 94, JANSSEN 95), or from the coupled-channel analysis of $\pi\pi\eta$ and $\bar{K}K\pi$ final states of $\bar{p}p$ annihilation at rest (ABELE 98).

The $a_0(1450)$ is seen in $\bar{p}p$ annihilation experiments with stopped and higher momenta \bar{p} , with a mass of about 1450 MeV or close to the $a_2(1320)$ meson which is typically a dominant feature. The broad structure at about 1300 MeV observed in $\pi N \rightarrow \bar{K}KN$ reactions (MARTIN 79) needs further confirmation in its existence and isospin assignment.

IV. The $I = 0$ States: The $I = 0 J^{PC} = 0^{++}$ sector is the most complex one, both experimentally and theoretically. The data have been obtained from $\pi\pi$, $\bar{K}K$, $\eta\eta$, 4π , and $\eta\eta'(958)$ systems produced in S -wave. Analyses based on several different production processes conclude that probably four poles are needed in the mass range from $\pi\pi$ threshold to about 1600 MeV. The claimed isoscalar resonances are found under separate entries σ or $f_0(600)$, $f_0(980)$, $f_0(1370)$, and $f_0(1500)$.

Below 1100 MeV, the important data come from the $\pi\pi$ and $\bar{K}K$ final states. Information on the $\pi\pi$ S -wave phase shift $\delta_J^I = \delta_0^0$ was already extracted 30 years ago from the πN scattering with unpolarized (GRAYER 74) and polarized targets (BECKER 79), and near threshold from the K_{e4} -decay (ROSSELET 77). The $\pi\pi$ S -wave inelasticity is not

accurately known, and the reported $\pi\pi \rightarrow \bar{K}K$ cross sections (WETZEL 76, POLYCHRONAKOS 79, COHEN 80, and ETKIN 82B) may have large uncertainties. The πN data (GRAYER 74, BECKER 79) have been analyzed in combination with high-statistics data from $\bar{p}p$ annihilation at rest (see entries labeled as RVUE for re-analyses of the data). The re-analysis (KAMINSKI 97, 02, 03) finds two out of four relevant solutions, with the S -wave phase shift rising slower than the P -wave [$\rho(770)$], which is used as a reference. One of these corresponds to the well-known “down” solution of GRAYER 74. The other “up” solution shows a decrease of the modulus in the mass interval between 800-980 MeV. Both solutions exhibit a sudden drop in the modulus and inelasticity at 1 GeV, due to the appearance of $f_0(980)$ which is very close to the opening of the $\bar{K}K$ -threshold. The phase shift δ_0^0 rises smoothly up to this point, where it jumps by 120° (in the “up”) or 140° (in the “down”) solution to reach 230° , and then both continue to rise slowly.

The suggestion (SVEC 97) of the existence of a narrow f_0 state near 750 MeV, with a small width of 100 to 200 MeV, is excluded by unitarity as shown by (KAMINSKI 97, 00) using both the π - and $a_1(1260)$ -exchange in the reaction amplitudes. The $2\pi^0$ invariant mass spectra of the $\bar{p}p$ annihilation at rest (AMSLER 95D, ABELE 96) and the central collision (ALDE 97) do not show a distinct resonance structure below 900 MeV, and these data are consistently described with the standard “down” solution (GRAYER 74, KAMINSKI 97), which allows for the existence of the broad ($\Gamma \approx 500$ MeV) resonance called σ . An enhancement is observed in the $\pi^+\pi^-$ invariant mass near threshold in the decays $D^+ \rightarrow \pi^+\pi^-\pi^+$ (AITALA 01B, LINK 04) and $J/\psi \rightarrow \omega\pi^+\pi^-$ (AUGUSTIN 89,

ABLIKIM 04A). The σ pole is difficult to establish because of its large width, and can certainly not be modelled by a naive Breit-Wigner resonance. It can be distorted by background as required by chiral symmetry, and from crossed channel exchanges, the $f_0(1370)$, and other dynamical features; it may be generated by t -channel meson exchanges (LOHSE 90, ZOU 94). However, most analyses listed in our issue under $f_0(600)$ agree on a pole position near $500 - i 250$ MeV.

The $f_0(980)$ overlaps strongly with the σ and the above mentioned broad background. This can lead to a dip in the $\pi\pi$ spectrum at the $\bar{K}K$ threshold. It changes from a dip into a peak structure in the $\pi^0\pi^0$ invariant mass spectrum of the reaction $\pi^- p \rightarrow \pi^0\pi^0 n$ (ACHASOV 98E), with increasing four-momentum transfer to the $\pi^0\pi^0$ system, which means increasing the a_1 -exchange contribution in the amplitude, while the π -exchange decreases.

One also observes the σ , and the $a_0(980)$, in radiative decays ($\phi \rightarrow f_0\gamma$, $\phi \rightarrow a_0\gamma$) in SND data (ACHASOV 00F, ACHASOV 00H), CMD2 (AKHMETSHIN 99B), and in KLOE data (ALOISIO 02C, ALOISIO 02D). In addition to these observations of the σ , its existence is also supported by the reaction $e^+e^- \rightarrow \pi^0\pi^0\gamma$ in the vicinity of the ρ and ω peaks. Both SND (ACHASOV 02F) and CMD-2 (AKHMETSHIN 04B) conclude that their value for the branching ratio $\rho \rightarrow \pi^0\pi^0\gamma$ exceeds the expectations from vector dominance, and that their results are much better described if a direct coupling of $\rho \rightarrow \sigma\gamma$ is added.

A meson resonance that is very well studied experimentally, is the $f_0(1500)$ seen by the Crystal Barrel experiment in five decay modes: $\pi\pi$, $\bar{K}K$, $\eta\eta$, $\eta\eta'(958)$, and 4π (AMSLER 95D, ABELE 96, and ABELE 98). Due to its interference with the $f_0(1370)$ (and $f_0(1700)$), the peak attributed to $f_0(1500)$ can

appear shifted in invariant mass spectra. Therefore, the application of simple Breit-Wigner forms arrive at slightly different resonance masses for $f_0(1500)$. Analyses of central-production data of the likewise five decay modes (BABERIS 99D, BABERIS 00E) agree on the description of the S wave with the one above. The $\bar{p}p$, $\bar{n}p/\bar{p}n$ (GASPERO 93, ADAMO 93, AMSLER 94, ABELE 96) show a single enhancement at 1400 MeV in the invariant 4π mass spectra, which is resolved into $f_0(1370)$ and $f_0(1500)$ (ABELE 01, ABELE01B). The data on 4π from central production (BABERIS 00C) require both resonances, too, but disagree on the relative content of $\rho\rho$ and $\sigma\sigma$ in 4π . All investigations agree, that the 4π decay mode represents about half of the $f_0(1500)$ decay width and is dominant for $f_0(1370)$.

The determination of the $\pi\pi$ coupling of $f_0(1370)$ is aggravated by the strong overlap with the broad $f_0(600)$ and $f_0(1500)$. Since it does not show up prominently in the 2π spectra, its mass and width are difficult to determine. Multi-channel analyses of hadronically produced two- and three-body final states agree on a mass between 1300 MeV and 1400 MeV and a narrow $f_0(1500)$, but arrive at a somewhat smaller width for $f_0(1370)$.

Both Belle and BaBar have observed strong indications of scalars in B meson decays. For example, GARMASH 02 saw a broad structure between 1.0 and 1.5 GeV in $\pi^+\pi^-$, K^+K^- and $K\pi$ final states. It could be a result of interference of several resonances in this mass range, but lack of statistics prevent from an unambiguous identification of this effect.

V. Interpretation: What is the nature of the light scalars? In the literature, many suggestions are discussed such as conventional $q\bar{q}$ mesons, $q\bar{q}q\bar{q}$ or meson-meson bound states mixed with a scalar glueball. In reality, they can be superpositions

of these components, and one depends on models to determine the dominant one. Although we have seen progress in recent years, this question remains open. Here, we mention some of the present conclusions.

Almost every model on scalar states agrees that the $K^*(1430)$ is predominantly the quark model $s\bar{u}$ or $s\bar{d}$ state.

If one uses the naive quark model (which may be too naive because of lack of chiral symmetry constraints), it is natural to assume the $f_0(1370)$, $a_0(1450)$, and the $K^*(1430)$ are in the same SU(3) flavour nonet being the $(\bar{u}u + \bar{d}d)$, $u\bar{d}$ and $u\bar{s}$ state, respectively. In this picture, the choice of the ninth member of the nonet is ambiguous. The controversially discussed candidates are $f_0(1500)$ and $f_0(1700)$. Compared to the above states, the $f_0(1500)$ is very narrow. Thus, it is unlikely to be their isoscalar partner. It is also too light to be the first radial excitation. Assuming the three f_0 's in the 1300-1700 MeV region to be mixtures between an $\bar{u}u$, $\bar{s}s$, and a gluonium state, one can arrive at an arrangement of these states, although different analyses (CLOSE 01B, LI 01) do not agree in detail. See our note on non- $\bar{q}q$ states.

The $f_0(980)$ and $a_0(980)$ are often interpreted as multi-quark states (JAFFE 77, ALFORD 00, MAIANI 04A) or $\bar{K}K$ bound states (WEINSTEIN 90). The insight into their internal structure using two-photon widths (BARNES 85, LI 91, DEL-BOURGO 99, LUCIO 99, ACHASOV 00H) is not conclusive. The $f_0(980)$ appears as a peak structure in $J/\psi \rightarrow \phi\pi^+\pi^-$ and in D_s decays without $f_0(600)$ background. Based on that observation it is suggested that $f_0(980)$ has a large $\bar{s}s$ component, which according to (DEANDREA 01) is surrounded by a virtual $\bar{K}K$ cloud. Data on radiative decays ($\phi \rightarrow f_0\gamma$ and $\phi \rightarrow a_0\gamma$) from SND, CMD2, and KLOE (see above) favour a 4-quark picture of the $f_0(980)$ and $a_0(980)$. The underlying

model for this conclusion (BOGLIONE 03, OLLER 03B) however may be oversimplified. But it remains quite possible that the states $f_0(980)$ and $a_0(980)$, together with the $f_0(600)$ and the $K_0^*(800)$, form a new low-mass state nonet of predominantly four-quark states, where at larger distances the quarks recombine into a pair of pseudoscalar mesons forming by a meson cloud.

Attempts have been made to start directly from chiral Lagrangians (SCADRON 99, OLLER 99, ISHIDA 99, TORN-QVIST 99, OLLER 03B, NAPSUCIALE 04, 04A) which predict the existence of the σ meson near 500 MeV. Hence, *e.g.*, in the chiral linear sigma model with 3 flavours, the σ , $a_0(980)$, $f_0(980)$, and κ (or $K_0^*(1430)$) would form a nonet (not necessarily $\bar{q}q$), while the lightest pseudoscalars would be their chiral partners. In the approach of (OLLER 99) the above resonances are generated starting from chiral perturbation theory predictions near the first open channel, and then by extending the predictions to the resonance regions using unitarity.

In the unitarized quark model with coupled $q\bar{q}$ and meson-meson channels, the light scalars can be understood as additional manifestations of bare $\bar{q}q$ confinement states, strongly mass shifted from the 1.3 - 1.5 GeV region and very distorted due to the strong 3P_0 coupling to S -wave two-meson decay channels (TORNQVIST 95, 96, BEVEREN 86, 99, 01B). Thus, the light scalar nonet comprising the $f_0(600)$, $f_0(980)$, $K_0^*(800)$, and $a_0(980)$, as well as the regular nonet consisting of the $f_0(1370)$, $f_0(1500)$ (or $f_0(1700)$), $K^*(1430)$, and $a_0(1450)$, respectively, are two manifestations of the same bare input states (see also BOGLIONE 02).

Other models with different groupings of the observed resonances exist and may *e.g.* be found in earlier versions of this review and papers listed as other related papers below.

References

References may be found at the end of the $f_0(600)$ listing.

$f_0(600)$ T-MATRIX POLE \sqrt{s}

Note that $\Gamma \approx 2 \operatorname{Im}(\sqrt{s}_{\text{pole}})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(400–1200)–i(250–500) OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$(441^{+16}_{-8}) - i(272^{+9}_{-12.5})$	1 CAPRINI 06 RVUE	$\pi\pi \rightarrow \pi\pi$	
$(470 \pm 50) - i(285 \pm 25)$	2 ZHOU 05 RVUE		
$(541 \pm 39) - i(252 \pm 42)$	3 ABLIKIM 04A BES2	$J/\psi \rightarrow \omega\pi^+\pi^-$	
$(528 \pm 32) - i(207 \pm 23)$	4 GALLEGO 04 RVUE	Compilation	
$(440 \pm 8) - i(212 \pm 15)$	5 PELAEZ 04A RVUE	$\pi\pi \rightarrow \pi\pi$	
$(533 \pm 25) - i(247 \pm 25)$	6 BUGG 03 RVUE		
$532 - i272$	BLACK 01 RVUE	$\pi^0\pi^0 \rightarrow \pi^0\pi^0$	
$(470 \pm 30) - i(295 \pm 20)$	1 COLANGELO 01 RVUE	$\pi\pi \rightarrow \pi\pi$	
$(535^{+48}_{-36}) - i(155^{+76}_{-53})$	7 ISHIDA 01	$\Upsilon(3S) \rightarrow \Upsilon\pi\pi$	
$610 \pm 14 - i620 \pm 26$	8 SUROVTSEV 01 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
$(558^{+34}_{-27}) - i(196^{+32}_{-41})$	ISHIDA 00B	$p\bar{p} \rightarrow \pi^0\pi^0\pi^0$	
$445 - i235$	HANNAH 99 RVUE	π scalar form factor	
$(523 \pm 12) - i(259 \pm 7)$	KAMINSKI 99 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$	
$442 - i227$	OLLER 99 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
$469 - i203$	OLLER 99B RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
$445 - i221$	OLLER 99C RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta$	
$(1530^{+90}_{-250}) - i(560 \pm 40)$	ANISOVICH 98B RVUE	Compilation	
$420 - i212$	LOCHER 98 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
$(602 \pm 26) - i(196 \pm 27)$	9 ISHIDA 97	$\pi\pi \rightarrow \pi\pi$	
$(537 \pm 20) - i(250 \pm 17)$	10 KAMINSKI 97B RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, 4\pi$	
$470 - i250$	11,12 TORNQVIST 96 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$	
$\sim (1100 - i300)$	AMSLER 95B CBAR	$\bar{p}p \rightarrow 3\pi^0$	
$400 - i500$	12,13 AMSLER 95D CBAR	$\bar{p}p \rightarrow 3\pi^0$	
$1100 - i137$	12,14 AMSLER 95D CBAR	$\bar{p}p \rightarrow 3\pi^0$	
$387 - i305$	12,15 JANSEN 95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
$525 - i269$	16 ACHASOV 94 RVUE	$\pi\pi \rightarrow \pi\pi$	
$(506 \pm 10) - i(247 \pm 3)$	KAMINSKI 94 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
$370 - i356$	17 ZOU 94B RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
$408 - i342$	12,17 ZOU 93 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
$870 - i370$	12,18 AU 87 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
$470 - i208$	19 BEVEREN 86 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta, \dots$	
$(750 \pm 50) - i(450 \pm 50)$	20 ESTABROOKS 79 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
$(660 \pm 100) - i(320 \pm 70)$	PROTOPOP... 73 HBC	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
$650 - i370$	21 BASDEVANT 72 RVUE	$\pi\pi \rightarrow \pi\pi$	

- 1 From the solution of the Roy equation (ROY 71) for the isoscalar S-wave and using a phase-shift analysis of HYAMS 73 and PROTOPOPESCU 73 data.
- 2 Reanalysis of the data from PROTOPOPESCU 73, ESTABROOKS 74, GRAYER 74, ROSSELET 77, PISLAK 03, and AKHMETSHIN 04.
- 3 From a mean of six different analyses and $f_0(600)$ parameterizations.
- 4 Using data on $\psi(2S) \rightarrow J/\psi\pi\pi$ from BAI 00E and on $\Upsilon(nS) \rightarrow \Upsilon(mS)\pi\pi$ from BUTLER 94B and ALEXANDER 98.
- 5 Reanalysis of data from PROTOPOPESCU 73, ESTABROOKS 74, GRAYER 74, and COHEN 80 in the unitarized ChPT model.
- 6 From a combined analysis of HYAMS 73, AUGUSTIN 89, AITALA 01B, and PISLAK 01.
- 7 A similar analysis (KOMADA 01) finds $(580^{+79}_{-30}) - i(190^{+107}_{-49})$ MeV.
- 8 Coupled channel reanalysis of BATON 70, BENSINGER 71, BAILLON 72, HYAMS 73, HYAMS 75, ROSSELET 77, COHEN 80, and ETKIN 82B using the uniformizing variable.
- 9 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
- 10 Average and spread of 4 variants ("up" and "down") of KAMINSKI 97B 3-channel model.
- 11 Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.
- 12 Demonstrates explicitly that $f_0(600)$ and $f_0(1370)$ are two different poles.
- 13 Coupled channel analysis of $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$ and $\pi^0\pi^0\eta$ on sheet II.
- 14 Coupled channel analysis of $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$ and $\pi^0\pi^0\eta$ on sheet III.
- 15 Analysis of data from FALVARD 88.
- 16 Analysis of data from OCHS 73, ESTABROOKS 75, ROSSELET 77, and MUKHIN 80.
- 17 Analysis of data from OCHS 73, GRAYER 74, and ROSSELET 77.
- 18 Analysis of data from OCHS 73, GRAYER 74, BECKER 79, and CASON 83.
- 19 Coupled-channel analysis using data from PROTOPOPESCU 73, HYAMS 73, HYAMS 75, GRAYER 74, ESTABROOKS 74, ESTABROOKS 75, FROGGATT 77, CORDEN 79, BISWAS 81.
- 20 Analysis of data from APEL 73, GRAYER 74, CASON 76, PAWLICKI 77. Includes spread and errors of 4 solutions.
- 21 Analysis of data from BATON 70, BENSINGER 71, COLTON 71, BAILLON 72, PROTOPOPESCU 73, and WALKER 67.

$f_0(600)$ BREIT-WIGNER MASS OR K-MATRIX POLE PARAMETERS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(400–1200) OUR ESTIMATE			
513 ± 32	²² MURAMATSU 02	CLEO	$e^+e^- \approx 10$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$478^{+24}_{-23} \pm 17$	AITALA	01B E791	$D^+ \rightarrow \pi^-\pi^+\pi^+$
563 ± 58	²³ ISHIDA	01	$\Upsilon(3S) \rightarrow \Upsilon\pi\pi$
555	²⁴ ASNER	00 CLE2	$\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
540 \pm 36	ISHIDA	00B	$p\bar{p} \rightarrow \pi^0\pi^0\pi^0$
750 \pm 4	ALEKSEEV	99 SPEC	$1.78\pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
744 \pm 5	ALEKSEEV	98 SPEC	$1.78\pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
759 \pm 5	²⁵ TROYAN	98	$5.2 np \rightarrow np\pi^+\pi^-$

780 ± 30	ALDE	97	GAM2	$450 \text{ } pp \rightarrow pp\pi^0\pi^0$
585 ± 20	²⁶ ISHIDA	97		$\pi\pi \rightarrow \pi\pi$
761 ± 12	²⁷ SVEC	96	RVUE	$6-17 \pi N_{\text{polar}} \rightarrow \pi^+\pi^-N$
~ 860	^{28,29} TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1165 ± 50	^{30,31} ANISOVICH	95	RVUE	$\pi^- p \rightarrow \pi^0\pi^0 n,$ $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\pi^0\eta, \pi^0\eta\eta$
~ 1000	³² ACHASOV	94	RVUE	$\pi\pi \rightarrow \pi\pi$
414 ± 20	²⁷ AUGUSTIN	89	DM2	
²² Statistical uncertainty only.				
²³ A similar analysis (KOMADA 01) finds 526^{+48}_{-37} MeV.				
²⁴ From the best fit of the Dalitz plot.				
²⁵ 6σ effect, no PWA.				
²⁶ Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.				
²⁷ Breit-Wigner fit to S-wave intensity measured in $\pi N \rightarrow \pi^-\pi^+N$ on polarized targets. The fit does not include $f_0(980)$.				
²⁸ Uses data from ASTON 88, OCHS 73, HYAMS 73, ARMSTRONG 91B, GRAYER 74, CASON 83, ROSSELET 77, and BEIER 72B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.				
²⁹ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decays.				
³⁰ Uses $\pi^0\pi^0$ data from ANISOVICH 94, AMSLER 94D, and ALDE 95B, $\pi^+\pi^-$ data from OCHS 73, GRAYER 74 and ROSSELET 77, and $\eta\eta$ data from ANISOVICH 94.				
³¹ The pole is on Sheet III. Demonstrates explicitly that $f_0(600)$ and $f_0(1370)$ are two different poles.				
³² Analysis of data from OCHS 73, ESTABROOKS 75, ROSSELET 77, and MUKHIN 80.				

$f_0(600)$ BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(600–1000) OUR ESTIMATE			
335 ± 67	³³ MURAMATSU 02	CLEO	$e^+e^- \approx 10 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$324^{+42}_{-40} \pm 21$	AITALA	01B E791	$D^+ \rightarrow \pi^-\pi^+\pi^+$
372 ± 229	³⁴ ISHIDA	01	$\gamma(3S) \rightarrow \gamma\pi\pi$
540	³⁵ ASNER	00 CLE2	$\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
372 ± 80	ISHIDA	00B	$p\bar{p} \rightarrow \pi^0\pi^0\pi^0$
119 ± 13	ALEKSEEV	99 SPEC	$1.78 \pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
77 ± 22	ALEKSEEV	98 SPEC	$1.78 \pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
35 ± 12	³⁶ TROYAN	98	$5.2 np \rightarrow np\pi^+\pi^-$
780 ± 60	ALDE	97 GAM2	$450 pp \rightarrow pp\pi^0\pi^0$
385 ± 70	³⁷ ISHIDA	97	$\pi\pi \rightarrow \pi\pi$
290 ± 54	³⁸ SVEC	96 RVUE	$6-17 \pi N_{\text{polar}} \rightarrow \pi^+\pi^-N$
~ 880	^{39,40} TORNQVIST	96 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
460 ± 40	^{41,42} ANISOVICH	95 RVUE	$\pi^- p \rightarrow \pi^0\pi^0 n,$ $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\pi^0\eta, \pi^0\eta\eta$
~ 3200	⁴³ ACHASOV	94 RVUE	$\pi\pi \rightarrow \pi\pi$
494 ± 58	³⁸ AUGUSTIN	89 DM2	

- 33 Statistical uncertainty only.
 34 A similar analysis (KOMADA 01) finds 301^{+145}_{-100} MeV.
 35 From the best fit of the Dalitz plot.
 36 6σ effect, no PWA.
 37 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
 38 Breit-Wigner fit to S-wave intensity measured in $\pi^- N \rightarrow \pi^- \pi^+ N$ on polarized targets. The fit does not include $f_0(980)$.
 39 Uses data from ASTON 88, OCHS 73, HYAMS 73, ARMSTRONG 91B, GRAYER 74, CASON 83, ROSSELET 77, and BEIER 72B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.
 40 Also observed by ASNER 00 in $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$ decays.
 41 Uses $\pi^0 \pi^0$ data from ANISOVICH 94, AMSLER 94D, and ALDE 95B, $\pi^+ \pi^-$ data from OCHS 73, GRAYER 74 and ROSSELET 77, and $\eta\eta$ data from ANISOVICH 94.
 42 The pole is on Sheet III. Demonstrates explicitly that $f_0(600)$ and $f_0(1370)$ are two different poles.
 43 Analysis of data from OCHS 73, ESTABROOKS 75, ROSSELET 77, and MUKHIN 80.

$f_0(600)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	dominant
$\Gamma_2 \gamma\gamma$	seen

$f_0(600)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$			Γ_2
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3.8 ± 1.5	44,45 BOGLIONE	99 RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$
5.4 ± 2.3	44 MORGAN	90 RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$
10 ± 6	COURAU	86 DM1	$e^+ e^- \rightarrow \pi^+ \pi^- e^+ e^-$

44 This width could equally well be assigned to the $f_0(1370)$. The authors analyse data from BOYER 90 and MARSISKE 90 and report strong correlation with $\gamma\gamma$ width of $f_2(1270)$.

45 Supersedes MORGAN 90.

$f_0(600)$ REFERENCES

CAPRINI	06	PRL 96 132001	I. Caprini, G. Colangelo, H. Leutwyler	(BCIP+)
ZHOU	05	JHEP 0502 043	Z.Y. Zhou <i>et al.</i>	
ABLIKIM	04A	PL B598 149	M. Ablikim <i>et al.</i>	(BES Collab.)
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
GALLEGOS	04	PR D69 074033	A. Gallegos <i>et al.</i>	
PELAEZ	04A	MPL A19 2879	J.R. Pelaez	
BUGG	03	PL B572 1	D.V. Bugg	
PISLAK	03	PR D67 072004	S. Pislak <i>et al.</i>	(BNL E865 Collab.)
MURAMATSU	02	PRL 89 251802	H. Muramatsu <i>et al.</i>	(CLEO Collab.)
Also		PRL 90 059901 (erratum)	H. Muramatsu <i>et al.</i>	(CLEO Collab.)
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BLACK	01	PR D64 014031	D. Black <i>et al.</i>	
COLANGELO	01	NP B603 125	G. Colangelo, J. Gasser, H. Leutwyler	
ISHIDA	01	PL B518 47	M. Ishida <i>et al.</i>	
KOMADA	01	PL B508 31	T. Komada <i>et al.</i>	

PISLAK	01	PRL 87 221801	S. Pislak <i>et al.</i>	(BNL E865 Collab.)
Also		PR D67 072004	S. Pislak <i>et al.</i>	(BNL E865 Collab.)
SUROVTSEV	01	PR D63 054024	Y.S. Surovtsev, D. Krupa, M. Nagy	
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BAI	00E	PR D62 032002	J. Bai <i>et al.</i>	(BES Collab.)
ISHIDA	00B	PTP 104 203	M. Ishida <i>et al.</i>	
ALEKSEEV	99	NP B541 3	I.G. Alekseev <i>et al.</i>	
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington	
HANNAH	99	PR D60 017502	T. Hannah	
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)
OLLER	99	PR D60 099906 (erratum)	J.A. Oller <i>et al.</i>	
OLLER	99B	NP A652 407 (erratum)	J.A. Oller, E. Oset	
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset	
ALEKSEEV	98	PAN 61 174	I.G. Alekseev <i>et al.</i>	
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ANISOVICH	98B	UFN 41 419	V.V. Anisovich <i>et al.</i>	
LOCHER	98	EPJ C4 317	M.P. Locher <i>et al.</i>	(PSI)
TROYAN	98	JINRRC 5-91 33	Yu. Troyan <i>et al.</i>	
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)
ISHIDA	97	PTP 98 1005	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)
KAMINSKI	97B	PL B413 130	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, IPN)
Also		PTP 95 745	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)
SVEC	96	PR D53 2343	M. Svec	(MCGI)
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)
ALDE	95B	ZPHY C66 375	D.M. Alde <i>et al.</i>	(GAMS Collab.)
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	95	PL B355 363	V.V. Anisovich <i>et al.</i>	(PNPI, SERP)
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)
ACHASOV	94	PR D49 5779	N.N. Achasov, G.N. Shestakov	(NOVM)
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)
BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)
ZOU	94B	PR D50 591	B.S. Zou, D.V. Bugg	(LOQM)
ZOU	93	PR D48 R3948	B.S. Zou, D.V. Bugg	(LOQM)
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LAZO+)
AU	87	PR D35 1633	K.L. Au, D. Morgan, M.R. Pennington	(DURH, RAL)
BEVEREN	86	ZPHY C30 615	E. van Beveren <i>et al.</i>	(NIJM, BIEL)
COURAU	86	NP B271 1	A. Courau <i>et al.</i>	(CLER, LAZO)
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
BISWAS	81	PRL 47 1378	N.N. Biswas <i>et al.</i>	(NDAM, ANL)
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL) IJP
MUKHIN	80	JETPL 32 601	K.N. Mukhin <i>et al.</i>	(KIAE)
		Translated from ZETFP 32 616.		
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP
ESTABROOKS	79	PR D19 2678	P. Estabrooks	(CARL)
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL) IJ
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)
CASON	76	PRL 36 1485	N.M. Cason <i>et al.</i>	(NDAM, ANL) IJ
ESTABROOKS	75	NP B95 322	P.G. Estabrooks, A.D. Martin	(DURH)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
SRINIVASAN	75	PR D12 681	V. Srinivasan <i>et al.</i>	(NDAM, ANL)
ESTABROOKS	74	NP B79 301	P.G. Estabrooks, A.D. Martin	(DURH)
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)
APEL	73	PL 41B 542	W.D. Apel <i>et al.</i>	(KARL, PISA)
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)
PROTOPOP...	73	PR D7 1279	S.D. Protopopescu <i>et al.</i>	(LBL)
BAILLON	72	PL 38B 555	P.H. Baillon <i>et al.</i>	(SLAC)
BASDEVANT	72	PL 41B 178	J.L. Basdevant, C.D. Froggatt, J.L. Petersen	(CERN)
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)

BENSINGER	71	PL 36B 134	J.R. Bensinger <i>et al.</i>	(WISC)
COLTON	71	PR D3 2028	E.P. Colton <i>et al.</i>	(LBL, FNAL, UCLA+)
ROY	71	PL 36B 353	S.M. Roy	
BATON	70	PL 33B 528	J.P. Baton, G. Laurens, J. Reignier	(SACL)
WALKER	67	RMP 39 695	W.D. Walker	(WISC)

OTHER RELATED PAPERS

ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
AITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BEDIAGA	06	PL B633 167	I. Bediaga, J.M. de Miranda	
CHENG	06	PR D73 014017	H.-Y. Cheng, C.-K. Chua, K.-C. Yang	
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT,B	05G	PR D72 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRITO	05	PL B608 69	T.V. Brito <i>et al.</i>	
CRONIN-HEN...	05	PR D72 031102R	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
GIACOSA	05	PR C71 025202	F. Giacosa <i>et al.</i>	
JAFFE	05	PRPL 409 1	R.L. Jaffe	
LI	05B	EPJ A25 263	D.-M. Li, K.-W. Wei, H. Yu	
LINK	05I	PL B621 72	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
RODRIGUEZ	05	PR D71 074008	S. Rodriguez, M. Napsuciale	
VIJANDE	05	PR D72 034025	J. Vijande, A. Valarce, F. Fernandez	
AKHMETSHIN	04B	PL B580 119	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AMSLER	04	PRPL 389 61	C. Amsler, N.A. Tornqvist	
AUBERT,B	04O	PR D70 091103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04P	PR D70 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
BUGG	04C	PRPL 397 257	D.V. Bugg	
BUGG	04D	EPJ C37 433	D.V. Bugg	
FARIBORZ	04	IJMP A19 2095	A.H. Fariborz	
KALOSHIN	04	EPJ A20 475	A.E. Kaloshin <i>et al.</i>	
LINK	04	PL B585 200	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
MAIANI	04A	PRL 93 212002	L. Maiani <i>et al.</i>	
NAPSUCIALE	04	PL B603 195	M. Napsuciale, S. Rodriguez	(GUAN)
NAPSUCIALE	04A	PR D70 094043	N.N. Napsuciale, S. Rodriguez	
PELAEZ	04	PRL 92 102001	J.R. Pelaez	
PRAKHOB	04	PR C69 042202R	S. Prakhov <i>et al.</i>	(BNL Crystal Ball Collab.)
PRAKHOB	04A	PR C69 045202	S. Prakhov <i>et al.</i>	
PRAKHOB	04B	PR C70 034605	S. Prakhov <i>et al.</i>	
TESHIMA	04	JPG 30 663	T. Teshima <i>et al.</i>	
VANBEVEREN	04	MPL A19 1949	E. Vanbeveren, G. Rupp	
ZHENG	04	NP A733 235	H.Q. Zheng <i>et al.</i>	
ABDEL-REHIM	03	PR D67 054001	A. Abdel-Rehim <i>et al.</i>	
ABDEL-REHIM	03B	PR D68 013008	A. Abdel-Rehim <i>et al.</i>	
BOGLIONE	03	EPJ C30 503	M. Boglione, M.R. Pennington	
ISHIDA	03	PTPS 149 190	M. Ishida	
KAMINSKI	03	PL B551 241	R. Kaminski, L. Lesniak, B. Loiseau	
OLLER	03B	NP A714 161	J.A. Oller <i>et al.</i>	
SCADRON	03	NP A724 391	M.D. Scadron <i>et al.</i>	
SEMEONOV	03	PAN 66 526	S.V. Semenov	
		Translated from YAF 66	553.	
ACHASOV	02F	PL B537 201	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AITALA	02	PRL 89 121801	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
ALOISIO	02C	PL B536 209	A. Aloisio <i>et al.</i>	(KLOE Collab.)
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)
BEVEREN	02	MPL A17 1673	E. van Beveren <i>et al.</i>	
BLACK	02	PRL 88 181603	D. Black, M. Harada, J. Schechter	
BOGLIONE	02	PR D65 114010	M. Boglione, M.R. Pennington	
BRAMON	02	EPJ C26 253	A. Bramon <i>et al.</i>	
CLOSE	02B	JPG 28 R249	F.E. Close, N. Tornqvist	
GARMASH	02	PR D65 092005	A. Garmash <i>et al.</i>	(BELLE Collab.)
HE	02	PL B536 59	J. He, Z.G. Xiao, H.Q. Zheng	
HERNANDEZ	02	PR C66 065201	E. Hernandez, E. Oset, M.J. Vicente Vacas	
ISHIDA	02	PL B539 249	S. Ishida, M. Ishida	
KAMINSKI	02	EPJ Direct C4 1	R. Kaminski, L. Lesniak, K. Rybicki	
LINK	02	PL B525 205	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02E	PL B535 43	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
TESHIMA	02	JPG 28 1391	T. Teshima, I. Kitamura, N. Morisita	
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BEVEREN	01B	EPJ C22 493	E. van Beveren	

CHERRY	01	NP A688 823	S.N. Cherry, M.R. Pennington
CLOSE	01B	EPJ C21 531	F.E. Close, A. Kirk
DEANDREA	01	PL B502 79	A. Deandrea <i>et al.</i>
FAZIO	01	PL B521 15	F. De Fazio, M.R. Pennington
GOKALP	01	PR D64 053017	A. Gokalp, O. Yilmaz
KOPP	01	PR D63 092001	S. Kopp <i>et al.</i>
LI	01	JPG 27 807	D.-M. Li, H. Yu, Q.-X. Shen
NARISON	01C	NPBPS 96 244	S. Narison
SHAKIN	01	PR D63 014019	C.M. Shakin, H. Wang
XIAO	01	NP A695 273	Z. Xiao, H. Zheng
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i>
ACHASOV	00H	PL B485 349	(Novosibirsk SND Collab.)
ALFORD	00	NP B578 367	M.N. Achasov <i>et al.</i>
BARBERIS	00C	PL B471 440	(Novosibirsk SND Collab.)
BARBERIS	00E	PL B479 59	M. Alford, R.L. Jaffe
BLACK	00B	PR D61 074030	D. Barberis <i>et al.</i>
FANG	00	NP A671 416	D. Barberis <i>et al.</i>
JAMIN	00	NP B587 331	D. Black, A. Fariborz, J. Schechter
KAMINSKI	00	APP B31 895	Fang Shi <i>et al.</i>
MONTANET	00	NPBPS 86 381	M. Jamin <i>et al.</i>
ABREU	99J	PL B449 364	R. Kaminski, L. Lesniak, K. Rybicki
AKHMETSHIN	99B	PL B462 371	L. Montanet
BARBERIS	99D	PL B462 462	P. Abreu <i>et al.</i>
BEVEREN	99	EPJ C10 469	(DELPHI Collab.)
BLACK	99	PR D59 074026	R.R. Akhmetshin <i>et al.</i>
DELBOURGO	99	PL B446 332	(Novosibirsk CMD-2 Collab.)
IGI	99	PR D59 034005	D. Barberis <i>et al.</i>
ISHIDA	99	PTP 101 661	(Omega Expt.)
LUCIO	99	PL B454 365	E. Van Beveren, G. Rupp
MINKOWSKI	99	EPJ C9 283	D. Black <i>et al.</i>
SCADRON	99	EPJ C6 141	R. Delbourgo, D. Liu, M. Scadron
TAKAMATSU	99	PAN 62 435	K. Igi, K. Hikasa
TORNQVIST	99	EPJ C11 359	M. Ishida
ABELE	98	PR D57 3860	J.L. Lucio, M. Napsuciale
ACHASOV	98E	PR D58 054011	P. Minkowski, W. Ochs
ACKERSTAFF	98A	EPJ C5 411	M. Scadron
ANISOVICH	98	PL B437 209	K. Takamatsu
BARBERIS	98C	PL B440 225	N. Tornqvist
DELBOURGO	98	IJMP A13 657	A. Abele <i>et al.</i>
OLLER	98	PRL 80 3452	(Crystal Barrel Collab.)
ANISOVICH	97	PL B395 123	N.N. Achasov, G.N. Shestakov
ANISOVICH	97C	PL B413 137	K. Ackerstaff <i>et al.</i>
ANISOVICH	97D	ZPHY A359 173	(OPAL Collab.)
HARADA	97	PRL 78 1603	V.V. Anisovich <i>et al.</i>
ISHIDA	97B	PTP 98 621	D. Barberis <i>et al.</i>
KAMINSKI	97	ZPHY C74 79	(WA 102 Collab.)
MALTMAN	97	PL B393 19	R. Delbourgo <i>et al.</i>
OLLER	97	NP A620 438	J.A. Oller <i>et al.</i>
SVEC	97	PR D55 4355	A.V. Anisovich, A.V. Sarantsev
SVEC	97B	PR D55 5727	(PNPI)
ABELE	96	PL B380 453	A.V. Anisovich, A.V. Sarantsev
AMSLER	96	PR D53 295	A.V. Anisovich, V.V. Anisovich, A.V. Sarantsev
BIJNENS	96	PL B374 210	M. Harada, F. Sannino, J. Schechter
BONUTTI	96	PRL 77 603	S. Ishida <i>et al.</i>
BUGG	96	NP B471 59	R. Kaminski, L. Lesniak, K. Rybicki
HARADA	96	PR D54 1991	(CRAC)
ISHIDA	96	PTP 95 745	K. Maltman, C.E. Wolfe
AMSLER	95C	PL B353 571	(YORKC)
ANTINORI	95	PL B353 589	J.A. Oller <i>et al.</i>
GASPERO	95	NP A588 861	(VALE)
TORNQVIST	95	ZPHY C68 647	M. Svec
AMSLER	94	PL B322 431	M. Svec
BUGG	94	PR D50 4412	A. Abele <i>et al.</i>
ZOU	94	PL B329 519	(Crystal Barrel Collab.)
ADAMO	93	NP A558 13C	C. Amsler, F.E. Close
GASPERO	93	NP A562 407	(ZURI, RAL)
MORGAN	93	PR D48 1185	J. Bijnens <i>et al.</i>
Also		NC A Conf. Suppl.	(NORD, BERN, WIEN+)
BOLTON	92B	PRL 69 1328	F. Bonutti <i>et al.</i>
SVEC	92	PR D45 55	(TRSTI, TRSTT, TRIU)
SVEC	92B	PR D45 1518	D.V. Bugg, A.V. Sarantsev, B.S. Zou
			(LOQM, PNPI)
			M. Harasa <i>et al.</i>
			(SYRA)
			S. Ishida <i>et al.</i>
			(TOKY, MIYA, KEK)
			C. Amsler <i>et al.</i>
			(Crystal Barrel Collab.)
			F. Antinori <i>et al.</i>
			(ATHU, BARI, BIRM+)
			M. Gaspero
			(ROMA)
			N.A. Tornqvist
			(HELS)
			C. Amsler <i>et al.</i>
			(Crystal Barrel Collab.)
			D.V. Bugg <i>et al.</i>
			(LOQM)
			Y. Zou <i>et al.</i>
			(RUTG, MINN, MICH)
			A. Adamo <i>et al.</i>
			(OBELIX Collab.)
			M. Gaspero
			(ROMAI)
			D. Morgan, M.R. Pennington
			(RAL, DURH)
			D. Morgan
			(RAL)
			T. Bolton <i>et al.</i>
			(Mark III Collab.)
			M. Svec, A. de Lesquen, L. van Rossum
			(MCGI+)
			M. Svec, A. de Lesquen, L. van Rossum
			(MCGI+)

SVEC	92C	PR D46 949	M. Svec, A. de Lesquen, L. van Rossum	(MCGI+)
BERNARD	91	PR D43 2757	V. Bernard, N. Kaiser, U.G. Meissner	
LI	91	PR D43 2161	Z.P. Li <i>et al.</i>	(TENN)
RIGGENBACH	91	PR D43 127	C. Rigggenbach <i>et al.</i>	(BERN, CERN, MASA)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
LOHSE	90	PL B234 235	D. Lohse <i>et al.</i>	
WEINSTEIN	90	PR D41 2236	J. Weinstein, N. Isgur	(TNTO)
ASTON	88D	NP B301 525	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BARNES	85	PL B165 434	T. Barnes	
ACHASOV	84	ZPHY C22 53	N.N. Achasov, S.A. Devyanin, G.N. Shestakov	(NOVM)
GASSER	84	ANP 158 142	J. Gasser, H. Leutwyler	
TORNQVIST	82	PRL 49 624	N.A. Tornqvist	(HELS)
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)
BECKER	79B	NP B150 301	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH) IJP
NAGELS	79	PR D20 1633	M.M. Nagels, T.A. Rijken, J.J. de Swart	(NIJM)
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL) IJP
CORDEN	78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)
JAFFE	77	PR D15 267,281	R. Jaffe	(MIT)
FLATTE	76	PL 63B 224	S.M. Flatte	(CERN)
WETZEL	76	NP B115 208	W. Wetzel <i>et al.</i>	(ETH, CERN, LOIC)
DEFOIX	72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)
ADLER	65	PR 137 B1022	S.L. Adler	
ADLER	65A	PR 139 B1638	S.L. Adler	