

$\omega(1650)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\omega(1650)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1670± 30 OUR ESTIMATE					
1660± 10± 2		AUBERT,B	04N BABR		$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
1700± 20		EUGENIO	01 SPEC		$18 \pi^- p \rightarrow \omega \eta n$
1705± 26	612	¹ AKHMETSHIN	00D CMD2		$e^+ e^- \rightarrow \omega \pi^+ \pi^-$
1662± 13	750	² ANTONELLI	92 DM2		$1.34\text{--}2.4 e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1770± 50±60	1.2M	³ ACHASOV	03D RVUE		$0.44\text{--}2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1619± 5		⁴ HENNER	02 RVUE		$1.2\text{--}2.0 e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$
1820^{+190}_{-150}		⁵ ACHASOV	98H RVUE		$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1840^{+100}_{-70}		⁶ ACHASOV	98H RVUE		$e^+ e^- \rightarrow \omega \pi^+ \pi^-$
1780^{+170}_{-300}		⁷ ACHASOV	98H RVUE		$e^+ e^- \rightarrow K^+ K^-$
~2100		⁸ ACHASOV	98H RVUE		$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1606± 9		⁹ CLEGG	94 RVUE		
1670± 20		ATKINSON	83B OMEG		$20\text{--}70 \gamma p \rightarrow 3\pi X$
1657± 13		CORDIER	81 DM1		$e^+ e^- \rightarrow \omega 2\pi$
1679± 34	21	ESPOSITO	80 FRAM		$e^+ e^- \rightarrow 3\pi$
1652± 17		COSME	79 OSPK 0		$e^+ e^- \rightarrow 3\pi$

¹ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho \pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

² From the combined fit of the $\rho \pi$ and $\omega \pi \pi$ final states.

³ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁴ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

⁵ Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.

⁶ Using the data from ANTONELLI 92.

⁷ Using the data from IVANOV 81 and BISELLO 88B.

⁸ Using the data from BISELLO 91C.

⁹ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

$\omega(1650)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
315 ± 35 OUR ESTIMATE					
230 ± 30 ± 20		AUBERT,B	04N BABR	10.6	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
250 ± 50		EUGENIO	01 SPEC	18	$\pi^- p \rightarrow \omega \eta n$
370 ± 25	612	AKHMETSHIN	00D CMD2		$e^+ e^- \rightarrow \omega \pi^+ \pi^-$
280 ± 24	750	ANTONELLI	92 DM2	1.34–2.4	$e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
490 ⁺²⁰⁰ ₋₁₅₀ ± 130	1.2M	ACHASOV	03D RVUE	0.44–2.00	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
250 ± 14		HENNER	02 RVUE	1.2–2.0	$e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$
113 ± 20		CLEGG	94 RVUE		
160 ± 20		ATKINSON	83B OMEG	20–70	$\gamma p \rightarrow 3\pi X$
136 ± 46		CORDIER	81 DM1		$e^+ e^- \rightarrow \omega 2\pi$
99 ± 49	21	ESPOSITO	80 FRAM		$e^+ e^- \rightarrow 3\pi$
42 ± 17		COSME	79 OSPK 0		$e^+ e^- \rightarrow 3\pi$
10 Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho \pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.					
11 From the combined fit of the $\rho \pi$ and $\omega \pi \pi$ final states.					
12 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.					
13 Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.					
14 From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.					

$\omega(1650)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho \pi$	seen
$\Gamma_2 \omega \pi \pi$	seen
$\Gamma_3 \omega \eta$	seen
$\Gamma_4 e^+ e^-$	seen

$$\omega(1650) \Gamma(i) \Gamma(e^+ e^-) / \Gamma^2(\text{total})$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_1 \Gamma_4 / \Gamma^2$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.3 ± 0.1 ± 0.1		AUBERT,B	04N BABR	10.6	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
1.2 ^{+0.4} _{-0.1} ± 0.8	1.2M	ACHASOV ^{15,16}	03D RVUE	0.44–2.00	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.921 ± 0.230		CLEGG ^{17,18}	94 RVUE		
0.479 ± 0.050	750	ANTONELLI ^{19,20}	92 DM2	1.34–2.4	$e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$

$\Gamma(\omega\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}^2$ $\Gamma_2\Gamma_4/\Gamma^2$

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.41 ± 0.09 ± 0.13	1.2M	^{15,16} ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.540 ± 0.095		²¹ AKHMETSHIN	00D CMD2	$1.2-1.38 e^+e^- \rightarrow \omega\pi^+\pi^-$
0.318 ± 0.080		^{17,18} CLEGG	94 RVUE	
0.607 ± 0.061	750	^{19,20} ANTONELLI	92 DM2	$1.34-2.4 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

 $\Gamma(\omega\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}^2$ $\Gamma_3\Gamma_4/\Gamma^2$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6 90 ²² AKHMETSHIN 03B CMD2 $e^+e^- \rightarrow \eta\pi^0\gamma$

¹⁵ Calculated by us from the cross section at the peak.

¹⁶ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

¹⁷ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

¹⁸ From the partial and leptonic width given by the authors.

¹⁹ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

²⁰ From the product of the leptonic width and partial branching ratio given by the authors.

²¹ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

²² $\omega(1650)$ mass and width fixed at 1700 MeV and 250 MeV, respectively.

$\omega(1650)$ BRANCHING RATIOS

 $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 0.35	1.2M	²⁴ ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.620 ± 0.014		²⁵ HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 0.65	1.2M	²⁴ ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.380 ± 0.014		²⁵ HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE (units 10^{-7})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 18	1.2M	23,25 ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
32 ± 1	25 HENNER	02 RVUE		$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
23 Calculated by us from the cross section at the peak.				
24 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.				
25 Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.				

 $\omega(1650)$ REFERENCES

AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>	
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
EUGENIO	01	PL B497 190	P. Eugenio <i>et al.</i>	
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
BISELLA	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLA	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from ZETFP 46 132.		
ATKINSON	83B	PL 127B 132	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
ESPOSITO	80	LNC 28 195	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
COSME	79	NP B152 215	G. Cosme <i>et al.</i>	(IPN)

OTHER RELATED PAPERS

AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AKHMETSHIN	03	PL B551 27	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
Also		PAN 65 1222	E.V. Anashkin, V.M. Aulchenko, R.R. Akhmetshin	
		Translated from YAF 65 1255.		
ACHASOV	02B	PAN 65 153	N.N. Achasov, A.A. Kozhevnikov	
		Translated from YAF 65 158.		
CLOSE	02	PR D65 092003	F.E. Close, A. Donnachie, Yu.S. Kalashnikova	
ACHASOV	00J	PR D62 117503	N.N. Achasov, A.A. Kozhevnikov	
ANISOVICH	00H	PL B485 341	A.V. Anisovich <i>et al.</i>	
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BELOZEROVA	98	PPN 29 63	T.S. Belozerova, V.K. Henner	
		Translated from FECAY 29 148.		
ACHASOV	97F	PAN 60 2029	N.N. Achasov, A.A. Kozhevnikov	(NOVM)
		Translated from YAF 60 2212.		
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
ATKINSON	87	ZPHY C34 157	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	84	NP B231 15	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)