

$f_2(1950)$ $I^G(J^{PC}) = 0^+(2^{++})$ **$f_2(1950)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1944±12 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.			
1930±25	¹ BINON	05 GAMS	33	$\pi^- p \rightarrow \eta\eta n$
2010±25	ANISOVICH	00J SPEC		
1940±50	BAI	00A BES		$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
1980±22	² BARBERIS	00C		$450 \bar{p}p \rightarrow \bar{p}p4\pi$
1940±22	³ BARBERIS	00C		$450 \bar{p}p \rightarrow \bar{p}p2\pi2\pi^0$
1980±50	ANISOVICH	99B SPEC		$1.35\text{--}1.94 \bar{p}\bar{p} \rightarrow \eta\eta\pi^0$
1960±30	BARBERIS	97B OMEG		$450 \bar{p}p \rightarrow \bar{p}p2(\pi^+ \pi^-)$
1918±12	ANTINORI	95 OMEG		$300,450 \bar{p}p \rightarrow \bar{p}p2(\pi^+ \pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1980± 2±14	ABE	04 BELL	10.6 $e^+ e^- \rightarrow$ $e^+ e^- K^+ K^-$
1867±46	⁴ AMSLER	02 CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \eta\eta, \pi^0 \pi^0 \pi^0$
~1996	HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$
~1990	⁵ OAKDEN	94 RVUE	0.36\text{--}1.55 $\bar{p}p \rightarrow \pi\pi$
1950±15	⁶ ASTON	91 LASS 0	11 $K^- p \rightarrow \Lambda K\bar{K}\pi\pi$

¹ First solution, PWA is ambiguous.


² Decaying into $\pi^+ \pi^- 2\pi^0$.

³ Decaying into $2(\pi^+ \pi^-)$.

⁴ T-matrix pole.

⁵ From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

⁶ Cannot determine spin to be 2.



f₂(1950) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
472± 18 OUR AVERAGE				
450± 50	⁷ BINON	05	GAMS	$33 \pi^- p \rightarrow \eta\eta\pi$
495± 35	ANISOVICH	00J	SPEC	
380^{+120}_{-90}	BAI	00A	BES	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
520± 50	⁸ BARBERIS	00C		$450 pp \rightarrow pp4\pi$
485± 55	⁹ BARBERIS	00C		$450 pp \rightarrow pp4\pi$
500±100	ANISOVICH	99B	SPEC	$1.35\text{--}1.94 p\bar{p} \rightarrow \eta\eta\pi^0$
460± 40	BARBERIS	97B	OMEG	$450 pp \rightarrow pp2(\pi^+\pi^-)$
390± 60	ANTINORI	95	OMEG	$300,450 pp \rightarrow pp2(\pi^+\pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

297± 12±6	ABE	04	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
385± 58	¹⁰ AMSLER	02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
~134	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~100	¹¹ OAKDEN	94	RVUE	$0.36\text{--}1.55 \bar{p}p \rightarrow \pi\pi$
250± 50	¹² ASTON	91	LASS 0	$11 K^- p \rightarrow \Lambda K\bar{K}\pi\pi$

⁷ First solution, PWA is ambiguous.

⁸ Decaying into $\pi^+\pi^-2\pi^0$.

⁹ Decaying into $2(\pi^+\pi^-)$.

¹⁰ T-matrix pole.

¹¹ From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J=3$ to be important but not significantly resonant.

12 Cannot determine spin to be 2.

$f_2(1950)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K^*(892)\bar{K}^*(892)$	seen
$\Gamma_2 \pi^+ \pi^-$	seen
$\Gamma_3 4\pi$	seen
$\Gamma_4 \pi^+ \pi^- \pi^+ \pi^-$	
$\Gamma_5 a_2(1320)\pi$	
$\Gamma_6 f_2(1270)\pi\pi$	
$\Gamma_7 \eta\eta$	seen
$\Gamma_8 K\bar{K}$	seen
$\Gamma_9 \gamma\gamma$	seen

$f_2(1950) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_8\Gamma_9/\Gamma$
<u>VALUE</u> (eV)	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
$122 \pm 4 \pm 26$	¹³ ABE 04 BELL $10.6 \frac{e^+ e^-}{e^+ e^- K^+ K^-} \rightarrow$
13 Assuming spin 2.	

$f_2(1950)$ BRANCHING RATIOS

$\Gamma(K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
seen	ASTON 91 LASS 0 $11 \frac{K^-\bar{p}}{\Lambda K\bar{K}\pi\pi} \rightarrow$

$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$	Γ_5/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
not seen	BARBERIS 00B $450 \frac{pp}{p_f \eta \pi^+ \pi^- p_s} \rightarrow$
not seen	BARBERIS 00C $450 pp \rightarrow p_f 4\pi p_s$
possibly seen	BARBERIS 97B OMEG $450 pp \rightarrow pp 2(\pi^+ \pi^-)$

$\Gamma(\eta\eta)/\Gamma(4\pi)$	Γ_7/Γ_3
<u>VALUE</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
$<5.0 \times 10^{-3}$	90 BARBERIS 00E $450 pp \rightarrow p_f \eta\eta p_s$

$\Gamma(\eta\eta)/\Gamma(\pi^+\pi^-)$	Γ_7/Γ_2		
VALUE	DOCUMENT ID	TECN	COMMENT
0.14±0.05	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0\eta\eta,$ $\pi^0\pi^0\pi^0$

f₂(1950) REFERENCES

BINON	05	PAN 68 960 Translated from YAF 68	F. Binon <i>et al.</i> 998.
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington
ASTON	91	NPBPS B21 5	D. Aston <i>et al.</i>

OTHER RELATED PAPERS

ANISOVICH	05	JETPL 80 715 Translated from ZETFP 80	V.V. Anisovich 845.
ANISOVICH	05A	JETPL 81 417 Translated from ZETFP 81	V.V. Anisovich, A.V. Sarantsev 531.
ANISOVICH	05C	IJMP A20 6327	V.V. Anisovich, M.A. Matveev, A.V. Sarantsev
LONGACRE	04	PR D70 094041	R.S. Longacre, S.J. Lindenbaum
ALBRECHT	88N	PL B212 528	H. Albrecht <i>et al.</i>
ALBRECHT	87Q	PL B198 255	H. Albrecht <i>et al.</i>
ARMSTRONG	87C	ZPHY C34 33	T.A. Armstrong <i>et al.</i>