

$$I(J^P) = 0(0^-)$$


The angular distributions of the decays of the ϕ and $\bar{K}^*(892)^0$ in the $\phi\pi^+$ and $K^+\bar{K}^*(892)^0$ modes strongly indicate that the spin is zero. The parity given is that expected of a $c\bar{s}$ ground state.

D_s^\pm MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , and $D_s^{*\pm}$ mass and mass difference measurements. Measurements of the D_s^\pm mass with an error greater than 10 MeV are omitted from the fit and average. A number of early measurements have been omitted altogether.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1968.2 ± 0.5 OUR FIT	Error includes scale factor of 1.1.			
1969.0 ± 1.4 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.			
1967.0 ± 1.0 ± 1.0	54	BARLAG	90C ACCM	π^- Cu 230 GeV
1969.3 ± 1.4 ± 1.4		ALBRECHT	88 ARG	e^+e^- 9.4–10.6 GeV
1972.7 ± 1.5 ± 1.0	21	BECKER	87B SILI	200 GeV π, K, p
1972.4 ± 3.7 ± 3.7	27	BLAYLOCK	87 MRK3	e^+e^- 4.14 GeV
1963 ± 3 ± 3	30	DERRICK	85B HRS	e^+e^- 29 GeV
1970 ± 5 ± 5	104	CHEN	83C CLEO	e^+e^- 10.5 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1968.3 ± 0.7 ± 0.7	290	¹ ANJOS	88 E691	Photoproduction
1980 ± 15	6	USHIDA	86 EMUL	ν wideband
1973.6 ± 2.6 ± 3.0	163	ALBRECHT	85D ARG	e^+e^- 10 GeV
1948 ± 28 ± 10	65	AIHARA	84D TPC	e^+e^- 29 GeV
1975 ± 9 ± 10	49	ALTHOFF	84 TASS	e^+e^- 14–25 GeV
1975 ± 4	3	BAILEY	84 ACCM	hadron ⁺ Be → $\phi\pi^+X$

¹ ANJOS 88 enters the fit via $m_{D_s^\pm} - m_{D^\pm}$ (see below).



$m_{D_s^\pm} - m_{D^\pm}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
98.85 ± 0.30 OUR FIT	Error includes scale factor of 1.4.			
98.85 ± 0.25 OUR AVERAGE	Error includes scale factor of 1.1.			
$99.41 \pm 0.38 \pm 0.21$		ACOSTA	03D CDF2	$\bar{p}p$, $\sqrt{s} = 1.96$ TeV
$98.4 \pm 0.1 \pm 0.3$	48k	AUBERT	02G BABR	$e^+e^- \approx \Upsilon(4S)$
$99.5 \pm 0.6 \pm 0.3$		BROWN	94 CLE2	$e^+e^- \approx \Upsilon(4S)$
98.5 ± 1.5	555	CHEN	89 CLEO	e^+e^- 10.5 GeV
99.0 ± 0.8	290	ANJOS	88 E691	Photoproduction


D_s^\pm MEAN LIFE

Measurements with an error greater than 100×10^{-15} s or with fewer than 100 events have been omitted from the Listings.

VALUE (10^{-15} s)	EVTS	DOCUMENT ID	TECN	COMMENT
500 ± 7 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
$507.4 \pm 5.5 \pm 5.1$	13.6k	LINK	05J FOCS	$\phi\pi^+$ and $\bar{K}^{*0}K^+$
$472.5 \pm 17.2 \pm 6.6$	760	IORI	01 SELX	600 GeV Σ^- , π^- , p
$518 \pm 14 \pm 7$	1662	AITALA	99 E791	π^- nucleus, 500 GeV

$486.3 \pm 15.0^{+4.9}_{-5.1}$	2167	² BONVICINI	99	CLE2	$e^+e^- \approx \Upsilon(4S)$
$475 \pm 20 \pm 7$	900	FRABETTI	93F	E687	$\gamma\text{Be}, \phi\pi^+$
$500 \pm 60 \pm 30$	104	FRABETTI	90	E687	$\gamma\text{Be}, \phi\pi^+$
$470 \pm 40 \pm 20$	228	RAAB	88	E691	Photoproduction

² BONVICINI 99 obtains 1.19 ± 0.04 for the ratio of D_s^+ to D^0 lifetimes.



D_s^+ DECAY MODES

Unless otherwise noted, the branching fractions for modes with a resonance in the final state include all the decay modes of the resonance. D_s^- modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
------	--------------------------------	-----------------------------------

Inclusive modes

Γ_1	K^- anything	(13 $\begin{smallmatrix} +14 \\ -12 \end{smallmatrix}$) %
Γ_2	\bar{K}^0 anything + K^0 anything	(39 ± 28) %
Γ_3	K^+ anything	(20 $\begin{smallmatrix} +18 \\ -14 \end{smallmatrix}$) %
Γ_4	(non- K \bar{K}) anything	(64 ± 17) %
Γ_5	e^+ anything	(8 $\begin{smallmatrix} + 6 \\ - 5 \end{smallmatrix}$) %
Γ_6	ϕ anything	(18 $\begin{smallmatrix} +15 \\ -10 \end{smallmatrix}$) %

Leptonic and semileptonic modes

Γ_7	$\mu^+ \nu_\mu$		$(6.1 \pm 1.9) \times 10^{-3}$	S=1.4
Γ_8	$\tau^+ \nu_\tau$		$(6.4 \pm 1.5) \%$	
Γ_9	$\phi \ell^+ \nu_\ell$	[a]	$(2.4 \pm 0.4) \%$	S=1.1
Γ_{10}	$\eta \ell^+ \nu_\ell + \eta'(958) \ell^+ \nu_\ell$	[a]	$(4.2 \pm 0.8) \%$	
Γ_{11}	$\eta \ell^+ \nu_\ell$	[a]	$(3.1 \pm 0.6) \%$	
Γ_{12}	$\eta'(958) \ell^+ \nu_\ell$	[a]	$(1.08 \pm 0.35) \%$	

Hadronic modes with a $K\bar{K}$ pair

Γ_{13}	$K^+ \bar{K}^0$		$(4.4 \pm 0.9) \%$	
Γ_{14}	$K^+ K^- \pi^+$	[b]	$(5.2 \pm 0.9) \%$	S=1.1
Γ_{15}	$\phi \pi^+$	[c]	$(4.4 \pm 0.6) \%$	S=1.1
Γ_{16}	$\phi \pi^+, \phi \rightarrow K^+ K^-$		$(2.16 \pm 0.28) \%$	S=1.1
Γ_{17}	$K^+ \bar{K}^*(892)^0$			
Γ_{18}	$K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+$		$(2.5 \pm 0.5) \%$	
Γ_{19}	$f_0(980) \pi^+, f_0 \rightarrow K^+ K^-$		$(5.7 \pm 2.5) \times 10^{-3}$	
Γ_{20}	$K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^{*0} \rightarrow K^- \pi^+$		$(4.8 \pm 2.5) \times 10^{-3}$	
Γ_{21}	$f_0(1710) \pi^+, f_0 \rightarrow K^+ K^-$			
Γ_{22}	$K^+ K^- \pi^+$ nonresonant			
Γ_{23}	$K^0 \bar{K}^0 \pi^+$		—	
Γ_{24}	$K^*(892)^+ \bar{K}^0$	[c]	$(5.3 \pm 1.3) \%$	
Γ_{25}	$K^+ K^- \pi^+ \pi^0$		—	
Γ_{26}	$\phi \pi^+ \pi^0$	[c]	$(11 \pm 5) \%$	
Γ_{27}	$\phi \rho^+$	[c]	$(8.2 \pm 2.0) \%$ $(8.2 \pm 2.4) \%$	
Γ_{28}	$\phi \pi^+ \pi^0$ 3-body	[c]	$< 3.1 \%$	CL=90%
Γ_{29}	$K^+ K^- \pi^+ \pi^0$ non- ϕ		$< 11 \%$	CL=90%
Γ_{30}	$K^+ \bar{K}^0 \pi^+ \pi^-$		$(3.1 \pm 0.9) \%$	
Γ_{31}	$K^0 K^- \pi^+ \pi^+$		$(5.3 \pm 1.4) \%$	
Γ_{32}	$K^*(892)^+ \bar{K}^*(892)^0$	[c]	$(7.0 \pm 2.7) \%$	
Γ_{33}	$K^0 K^- \pi^+ \pi^+ (\text{non-}K^* \bar{K}^{*0})$		$< 3.5 \%$	CL=90%
Γ_{34}	$K^+ K^- \pi^+ \pi^+ \pi^-$		$(8.3 \pm 2.0) \times 10^{-3}$	
Γ_{35}	$\phi \pi^+ \pi^+ \pi^-$	[c]	$(1.18 \pm 0.20) \%$	
Γ_{36}	$K^+ K^- \rho^0 \pi^+$ non- ϕ		$< 2.5 \times 10^{-4}$	CL=90%
Γ_{37}	$\phi \rho^0 \pi^+$	[c]	$(1.24 \pm 0.33) \%$	
Γ_{38}	$\phi a_1(1260)^+$	[c]	$(2.9 \pm 0.7) \%$	
Γ_{39}	$K^+ K^- \pi^+ \pi^+ \pi^-$ nonresonant		$(8 \pm 7) \times 10^{-4}$	
Γ_{40}	$K_S^0 K_S^0 \pi^+ \pi^+ \pi^-$		$(2.7 \pm 1.3) \times 10^{-3}$	

Hadronic modes without K 's

Γ_{41}	$\pi^+ \pi^+ \pi^-$	(1.22 ± 0.23) %	S=1.2
Γ_{42}	$\rho^0 \pi^+$		
Γ_{43}	$\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$	[d] (1.06 ± 0.22) %	
Γ_{44}	$f_0(980) \pi^+, f_0 \rightarrow \pi^+ \pi^-$		
Γ_{45}	$f_0(1370) \pi^+, f_0 \rightarrow \pi^+ \pi^-$		
Γ_{46}	$f_0(1500) \pi^+, f_0 \rightarrow \pi^+ \pi^-$		
Γ_{47}	$f_2(1270) \pi^+, f_2 \rightarrow \pi^+ \pi^-$	(1.2 ± 0.7) × 10 ⁻³	
Γ_{48}	$\rho(1450)^0 \pi^+, \rho^0 \rightarrow \pi^+ \pi^-$	(8 ± 7) × 10 ⁻⁴	
Γ_{49}	$\pi^+ \pi^+ \pi^-$ nonresonant		
Γ_{50}	$\pi^+ \pi^+ \pi^- \pi^0$	< 15 %	CL=90%
Γ_{51}	$\eta \pi^+$	[c] (2.11 ± 0.35) %	
Γ_{52}	$\omega \pi^+$	[c] (3.4 ± 1.2) × 10 ⁻³	
Γ_{53}	$3\pi^+ 2\pi^-$	(7.6 ± 1.6) × 10 ⁻³	
Γ_{54}	$\pi^+ \pi^+ \pi^- \pi^0 \pi^0$	—	
Γ_{55}	$\eta \rho^+$	[c] (13.1 ± 2.6) %	
Γ_{56}	$\eta \pi^+ \pi^0$ 3-body	[c] < 5 %	CL=90%
Γ_{57}	$3\pi^+ 2\pi^- \pi^0$	(4.9 ± 3.2) %	
Γ_{58}	$\eta'(958) \pi^+$	[c] (4.7 ± 0.7) %	
Γ_{59}	$3\pi^+ 2\pi^- 2\pi^0$	—	
Γ_{60}	$\eta'(958) \rho^+$	[c] (12.2 ± 2.4) %	
Γ_{61}	$\eta'(958) \pi^+ \pi^0$ 3-body	[c] < 1.8 %	CL=90%

Modes with one or three K 's

Γ_{62}	$K^0 \pi^+$	< 9 × 10 ⁻³	CL=90%
Γ_{63}	$K^+ \pi^+ \pi^-$	(6.6 ± 1.4) × 10 ⁻³	
Γ_{64}	$K^+ \rho^0$	(2.6 ± 0.7) × 10 ⁻³	
Γ_{65}	$K^+ \rho(1450)^0, \rho^0 \rightarrow \pi^+ \pi^-$	(7.0 ± 2.9) × 10 ⁻⁴	
Γ_{66}	$K^*(892)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$	(1.4 ± 0.4) × 10 ⁻³	
Γ_{67}	$K^*(1410)^0 \pi^+, K^{*0} \rightarrow$ $K^+ \pi^-$	(1.2 ± 0.4) × 10 ⁻³	
Γ_{68}	$K^*(1430)^0 \pi^+, K^{*0} \rightarrow$ $K^+ \pi^-$	(5 ± 4) × 10 ⁻⁴	
Γ_{69}	$K^+ \pi^+ \pi^-$ nonresonant	(1.0 ± 0.4) × 10 ⁻³	
Γ_{70}	$K^+ K^+ K^-$	(4.6 ± 1.8) × 10 ⁻⁴	
Γ_{71}	ϕK^+	[c] < 6 × 10 ⁻⁴	CL=90%

Doubly Cabibbo-suppressed modes

Γ_{72}	$K^+ K^+ \pi^-$	(2.7 ± 1.2) × 10 ⁻⁴	
---------------	-----------------	----------------------------------	--

$\Delta C = 1$ weak neutral current ($C1$) modes, Lepton family number (LF), or Lepton number (L) violating modes

Γ_{73}	$\pi^+ e^+ e^-$	[e] < 2.7 × 10 ⁻⁴	CL=90%
Γ_{74}	$\pi^+ \mu^+ \mu^-$	[e] < 2.6 × 10 ⁻⁵	CL=90%
Γ_{75}	$K^+ e^+ e^-$	$C1$ < 1.6 × 10 ⁻³	CL=90%

Γ_{76}	$K^+ \mu^+ \mu^-$	CI	< 3.6	$\times 10^{-5}$	CL=90%
Γ_{77}	$K^*(892)^+ \mu^+ \mu^-$	CI	< 1.4	$\times 10^{-3}$	CL=90%
Γ_{78}	$\pi^+ e^\pm \mu^\mp$	LF	$[f] < 6.1$	$\times 10^{-4}$	CL=90%
Γ_{79}	$K^+ e^\pm \mu^\mp$	LF	$[f] < 6.3$	$\times 10^{-4}$	CL=90%
Γ_{80}	$\pi^- e^+ e^+$	L	< 6.9	$\times 10^{-4}$	CL=90%
Γ_{81}	$\pi^- \mu^+ \mu^+$	L	< 2.9	$\times 10^{-5}$	CL=90%
Γ_{82}	$\pi^- e^+ \mu^+$	L	< 7.3	$\times 10^{-4}$	CL=90%
Γ_{83}	$K^- e^+ e^+$	L	< 6.3	$\times 10^{-4}$	CL=90%
Γ_{84}	$K^- \mu^+ \mu^+$	L	< 1.3	$\times 10^{-5}$	CL=90%
Γ_{85}	$K^- e^+ \mu^+$	L	< 6.8	$\times 10^{-4}$	CL=90%
Γ_{86}	$K^*(892)^- \mu^+ \mu^+$	L	< 1.4	$\times 10^{-3}$	CL=90%
Γ_{87}	A dummy mode used by the fit.		$(79.8 \pm 2.8) \%$		S=1.1

- [a] For now, we average together measurements of the $X e^+ \nu_e$ and $X \mu^+ \nu_\mu$ branching fractions. This is the *average*, not the *sum*.
- [b] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.
- [c] This branching fraction includes all the decay modes of the final-state resonance.
- [d] This comes from a K -matrix parametrization of the $\pi^+ \pi^-$ S -wave and is a sum over the $f_0(980)$, $f_0(1300)$, $f_0(1200-1600)$, $f_0(1500)$, and $f_0(1750)$. Not all of these correspond to particles in our Tables.
- [e] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [f] The value is for the sum of the charge states or particle/antiparticle states indicated.

CONSTRAINED FIT INFORMATION

An overall fit to 11 branching ratios uses 19 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 11.4$ for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_9	46							
x_{11}	32	69						
x_{12}	21	44	31					
x_{14}	36	64	44	28				
x_{15}	49	88	61	39	73			
x_{16}	49	87	61	39	73	100		
x_{41}	35	62	43	27	67	71	71	
x_{87}	-52	-90	-74	-50	-85	-93	-93	-75
	x_7	x_9	x_{11}	x_{12}	x_{14}	x_{15}	x_{16}	x_{41}

D_s^+ BRANCHING RATIOS

A few older, now obsolete results have been omitted. They may be found in earlier editions.

Inclusive modes

$\Gamma(K^- \text{ anything}) / \Gamma_{\text{total}}$ Γ_1 / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.13^{+0.14}_{-0.12} \pm 0.02$	COFFMAN	91	MRK3 $e^+ e^-$ 4.14 GeV

$[\Gamma(\bar{K}^0 \text{ anything}) + \Gamma(K^0 \text{ anything})] / \Gamma_{\text{total}}$ Γ_2 / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.39^{+0.28}_{-0.27} \pm 0.04$	COFFMAN	91	MRK3 $e^+ e^-$ 4.14 GeV

$\Gamma(K^+ \text{ anything}) / \Gamma_{\text{total}}$ Γ_3 / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.20^{+0.18}_{-0.13} \pm 0.04$	COFFMAN	91	MRK3 $e^+ e^-$ 4.14 GeV

$\Gamma(\text{(non-}K\bar{K}\text{) anything}) / \Gamma_{\text{total}}$ Γ_4 / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.64 \pm 0.17 \pm 0.03$	³ COFFMAN	91	MRK3 $e^+ e^-$ 4.14 GeV

³COFFMAN 91 uses the direct measurements of the kaon content to determine this non- $K\bar{K}$ fraction. This number implies that a large fraction of D_s^+ decays involve η , η' , and/or non-spectator decays.

$\Gamma(e^+ \text{ anything})/\Gamma_{\text{total}}$					Γ_5/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$0.077^{+0.057+0.024}_{-0.043-0.021}$		BAI	97 BES	$e^+ e^- \rightarrow D_s^+ D_s^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.20 90 ⁴BAI 90 MRK3 $e^+ e^-$ 4.14 GeV

⁴Expressed as a value, the BAI 90 result is $\Gamma(e^+ \text{ anything})/\Gamma_{\text{total}} = 0.05 \pm 0.05 \pm 0.02$.

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$					Γ_6/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$0.178^{+0.151+0.006}_{-0.072-0.063}$	3	BAI	98 BES	$e^+ e^- \rightarrow D_s^+ D_s^-$	

———— Leptonic and semileptonic modes ————

D_s^+ DECAY CONSTANT

Revised March 2006 by A. Edwards and P. Burchat (Stanford University)

In the Standard Model, the D_s^+ leptonic branching fractions are related to the D_s^+ decay constant f_{D_s} by the equation [1]

$$B(D_s^+ \rightarrow \ell^+ \nu_\ell) = \frac{G_F^2}{8\pi} |V_{cs}|^2 f_{D_s}^2 \frac{\tau_{D_s}}{\hbar} m_{D_s} m_\ell^2 \left(1 - \frac{m_\ell^2}{m_{D_s}^2}\right)^2. \quad (1)$$

Hence, measurements of $B(D_s^+ \rightarrow \ell^+ \nu_\ell)$ can be used to extract f_{D_s} . The most precise measurements of $D_s^+ \rightarrow \ell^+ \nu_\ell$ branching fractions come from L3 (ACCIARRI 97F), CLEO (CHADHA 98), BEATRICE (ALEXANDROV 00), OPAL (ABBIENDI 01L), and ALEPH (HEISTER 02I); see the end of the D_s^+ Data Listings for the references. All of these measurements either explicitly or implicitly measure the leptonic branching fraction relative to the branching fraction for $D_s^+ \rightarrow \phi\pi^+$. This fraction has, since our 2004 edition, changed from $3.6 \pm 0.9\%$ to $4.4 \pm 0.6\%$. The $D_s^+ \rightarrow \ell^+ \nu_\ell$ measurements of CLEO and BEATRICE are explicitly normalized to $D_s^+ \rightarrow \phi\pi^+$, and so can be easily updated. The LEP experiments (L3, OPAL, ALEPH) share a 23% correlated uncertainty in the normalization of the

leptonic branching fraction. They use the partial decay rate for $Z \rightarrow c\bar{c}$ and the D_s^+ production rate in $Z \rightarrow c\bar{c}$ events, which in turn depends on the assumed value of $B(D_s^+ \rightarrow \phi\pi^+)$.

We determine an average value of f_{D_s} from the above-mentioned five most precise measurements of the $D_s^+ \rightarrow \ell^+\nu_\ell$ branching fractions, assuming lepton universality, taking into account correlated uncertainties, and using a consistent and up-to-date set of input parameters [2] for the μ , τ , and D_s^+ masses, the D_s^+ lifetime, V_{cs} , $B(D_s^+ \rightarrow \phi\pi^+)$, and other relevant branching fractions. Although the uncertainty on $B(D_s^+ \rightarrow \phi\pi^+)$ is by far the largest uncertainty, we also take into account correlated uncertainties in other input parameters. Using both $D_s^+ \rightarrow \mu^+\nu_\mu$ and $D_s^+ \rightarrow \tau^+\nu_\tau$ branching fractions, and assuming lepton universality, we obtain

$$B(D_s^+ \rightarrow \mu^+\nu_\mu) = 0.0074 \pm 0.0013 . \quad (2)$$

Using this value (which is not the same as the $D_s^+ \rightarrow \mu^+\nu_\mu$ branching fraction in our Summary Tables, because we do not there use lepton universality), and including the relatively minor uncertainties on the other parameters in Eq. (1), we extract the world average D_s^+ decay constant:

$$f_{D_s} = (294 \pm 27) \text{ MeV} . \quad (3)$$

References

1. See the note on ‘‘Pseudoscalar-Meson Decay Constants’’ at the beginning of the Meson Particle Listings.
2. This *Review*.

$\Gamma(\mu^+\nu_\mu)/\Gamma_{\text{total}}$	Γ_7/Γ
<u>VALUE</u>	<u>TECN</u>
<u>EVTS</u>	<u>DOCUMENT ID</u>
<u>COMMENT</u>	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.0068 \pm 0.0011 \pm 0.0018$	553	⁵ HEISTER	02l	ALEP	Z decays
$0.015^{+0.013}_{-0.006} +^{+0.003}_{-0.002}$	3	⁶ BAI	95	BES	$e^+ e^- \rightarrow D_s^+ D_s^-$
$0.004^{+0.0018}_{-0.0014} +^{+0.0020}_{-0.0019}$	8	⁷ AOKI	93	WA75	π^- emulsion 350 GeV
<0.03	0	⁸ AUBERT	83	SPEC	μ^+ Fe, 250 GeV

⁵ This HEISTER 02l result is not actually an independent measurement of the absolute $\mu^+ \nu_\mu$ branching fraction, but is in fact based on our $\phi \pi^+$ branching fraction of $3.6 \pm 0.9\%$, so it cannot be included in our overall fit. HEISTER 02l combines its $D_s^+ \rightarrow \tau^+ \nu_\tau$ and $\mu^+ \nu_\mu$ branching fractions to get $f_{D_s} = (285 \pm 19 \pm 40)$ MeV.

⁶ BAI 95 uses one actual $D_s^+ \rightarrow \mu^+ \nu_\mu$ event together with two $D_s^+ \rightarrow \tau^+ \nu_\tau$ events and assumes μ - τ universality. This value of $\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$ gives a pseudoscalar decay constant of $(430^{+150}_{-130} \pm 40)$ MeV.

⁷ AOKI 93 assumes the ratio of production cross sections of the D_s^+ and D^0 is 0.27. The value of $\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$ gives a pseudoscalar decay constant $f_{D_s} = (232 \pm 45 \pm 52)$ MeV.

⁸ AUBERT 83 assume that the D_s^\pm production rate is 20% of total charm production rate.

$\Gamma(\mu^+ \nu_\mu)/\Gamma(\phi \pi^+)$

Γ_7/Γ_{15}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.14 ± 0.04 OUR FIT				Error includes scale factor of 1.4.
0.19 ± 0.04 OUR AVERAGE				

0.23 ± 0.06 ± 0.04 18 ⁹ ALEXANDROV00 BEAT π^- nucleus, 350 GeV

0.173 ± 0.023 ± 0.035 182 ¹⁰ CHADHA 98 CLE2 $e^+ e^- \approx \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.245 ± 0.052 ± 0.074 39 ¹¹ ACOSTA 94 CLE2 See CHADHA 98

⁹ ALEXANDROV 00 uses $f_D^2/f_{D_s}^2 = 0.82 \pm 0.09$ from a lattice-gauge-theory calculation to get the relative numbers of $D^+ \rightarrow \mu^+ \nu_\mu$ and $D_s^+ \rightarrow \mu^+ \nu_\mu$ events. The present result leads to $f_{D_s} = (323 \pm 44 \pm 36)$ MeV.

¹⁰ CHADHA 98 obtains $f_{D_s} = (280 \pm 19 \pm 28 \pm 34)$ MeV from this measurement, using $\Gamma(D_s^+ \rightarrow \phi \pi^+)/\Gamma(\text{total}) = 0.036 \pm 0.009$.

¹¹ ACOSTA 94 obtains $f_{D_s} = (344 \pm 37 \pm 52 \pm 42)$ MeV from this measurement, using $\Gamma(D_s^+ \rightarrow \phi \pi^+)/\Gamma(\text{total}) = 0.037 \pm 0.009$.

$\Gamma(\mu^+ \nu_\mu)/\Gamma(\phi \ell^+ \nu_\ell)$

Γ_7/Γ_9

$\Gamma(\phi \ell^+ \nu_\ell)$ is an average of $\Gamma(\phi e^+ \nu_e)$ and $\Gamma(\phi \mu^+ \nu_\mu)$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.25 ± 0.07 OUR FIT				Error includes scale factor of 1.5.
0.16 ± 0.06 ± 0.03	23	¹² KODAMA	96	E653 π^- emulsion, 600 GeV

¹² KODAMA 96 obtains $f_{D_s} = (194 \pm 35 \pm 20 \pm 14)$ MeV from this measurement, using $\Gamma(D_s^+ \rightarrow \phi \ell^+ \nu)/\Gamma_{\text{total}} = 0.0188 \pm 0.0029$. The third error is from the uncertainty on $\phi \ell^+ \nu_\ell$ branching fraction.

$\Gamma(\tau^+ \nu_\tau)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.064 ± 0.015 OUR AVERAGE				
0.0579 ± 0.0077 ± 0.0184	881	¹³ HEISTER	02I ALEP	Z decays
0.070 ± 0.021 ± 0.020	22	¹⁴ ABBIENDI	01L OPAL	$D_S^{*+} \rightarrow \gamma D_S^+$ from Z's
0.074 ± 0.028 ± 0.024	16	¹⁵ ACCIARRI	97F L3	$D_S^{*+} \rightarrow \gamma D_S^+$ from Z's

¹³ HEISTER 02I combines its $D_S^+ \rightarrow \tau^+ \nu_\tau$ and $\mu^+ \nu_\mu$ branching fractions to get $f_{D_S} = (285 \pm 19 \pm 40)$ MeV.

¹⁴ This ABBIENDI 01L value gives a decay constant f_{D_S} of $(286 \pm 44 \pm 41)$ MeV.

¹⁵ The second ACCIARRI 97F error here combines in quadrature systematic (0.016) and normalization (0.018) errors. The branching fraction gives $f_{D_S} = (309 \pm 58 \pm 33 \pm 38)$ MeV.

$\Gamma(\phi \ell^+ \nu_\ell)/\Gamma(\phi \pi^+)$ Γ_9/Γ_{15}

For now, we average together measurements of the $\Gamma(\phi e^+ \nu_e)/\Gamma(\phi \pi^+)$ and $\Gamma(\phi \mu^+ \nu_\mu)/\Gamma(\phi \pi^+)$ ratios. See the end of the D_S^+ Listings for measurements of $D_S^+ \rightarrow \phi \ell^+ \nu_\ell$ form-factor ratios.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.55 ± 0.04 OUR FIT				
0.54 ± 0.04 OUR AVERAGE				
0.540 ± 0.033 ± 0.048	793	LINK	02J FOCS	Uses $\phi \mu^+ \nu_\mu$
0.54 ± 0.05 ± 0.04	367	BUTLER	94 CLE2	Uses $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$
0.58 ± 0.17 ± 0.07	97	FRABETTI	93G E687	Uses $\phi \mu^+ \nu_\mu$
0.57 ± 0.15 ± 0.15	104	ALBRECHT	91 ARG	Uses $\phi e^+ \nu_e$
0.49 ± 0.10 ^{+0.10} / _{-0.14}	54	ALEXANDER	90B CLEO	Uses $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$

$\Gamma(\eta \ell^+ \nu_\ell)/\Gamma(\phi \ell^+ \nu_\ell)$ Γ_{11}/Γ_9

Unseen decay modes of the η and the ϕ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.27 ± 0.19 OUR FIT				
1.24 ± 0.12 ± 0.15	440	¹⁶ BRANDENB...	95 CLE2	$e^+ e^- \approx \Upsilon(4S)$

¹⁶ BRANDENBURG 95 uses both e^+ and μ^+ events and makes a phase-space adjustment to use the μ^+ events as e^+ events.

$\Gamma(\eta'(958) \ell^+ \nu_\ell)/\Gamma(\phi \ell^+ \nu_\ell)$ Γ_{12}/Γ_9

Unseen decay modes of the resonances are included.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.44 ± 0.13 OUR FIT					
0.43 ± 0.11 ± 0.07		29	¹⁷ BRANDENB...	95 CLE2	$e^+ e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.6		90	¹⁸ KODAMA	93B E653	π^- emulsion 600 GeV

¹⁷ BRANDENBURG 95 uses both e^+ and μ^+ events and makes a phase-space adjustment to use the μ^+ events as e^+ events.

¹⁸ KODAMA 93B uses μ^+ events.

$$[\Gamma(\eta\ell^+\nu_\ell) + \Gamma(\eta'(958)\ell^+\nu_\ell)]/\Gamma(\phi\ell^+\nu_\ell) \quad \Gamma_{10}/\Gamma_9 = (\Gamma_{11} + \Gamma_{12})/\Gamma_9$$

Unseen decay modes of the resonances are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

1.72 ± 0.23 OUR FIT

3.9 ± 1.6 13 ¹⁹KODAMA 93 E653 π⁻ emulsion 600 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.67 ± 0.17 ± 0.17 ²⁰BRANDENB... 95 CLE2 e⁺e⁻ ≈ γ(4S)

¹⁹KODAMA 93 uses μ⁺ events.

²⁰This BRANDENBURG 95 data is redundant with data in previous blocks.

————— **Hadronic modes with a K \bar{K} pair.** —————

$$\Gamma(K^+\bar{K}^0)/\Gamma(\phi\pi^+) \quad \Gamma_{13}/\Gamma_{15}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

1.01 ± 0.16 OUR AVERAGE

1.15 ± 0.31 ± 0.19 68 ANJOS 90C E691 γ Be

0.92 ± 0.32 ± 0.20 ADLER 89B MRK3 e⁺e⁻ 4.14 GeV

0.99 ± 0.17 ± 0.10 CHEN 89 CLEO e⁺e⁻ 10 GeV

$$\Gamma(\phi\pi^+)/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

We now have model-independent measurements of this branching fraction, and so we no longer use the earlier, model-dependent results.

VALUE (units 10 ⁻²)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	------	-------------	------	---------

4.4 ± 0.6 OUR FIT Error includes scale factor of 1.1.

4.4 ± 0.6 OUR AVERAGE Error includes scale factor of 1.1.

4.81 ± 0.52 ± 0.38 212 ± 19 ²¹AUBERT 05v BABR e⁺e⁻ ≈ γ(4S)

3.59 ± 0.77 ± 0.48 ²²ARTUSO 96 CLE2 e⁺e⁻ at γ(4S)

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.9 ^{+5.1} _{-1.9} ^{+1.8} _{-1.1} ²³BAI 95C BES e⁺e⁻ 4.03 GeV

5.1 ± 0.4 ± 0.8 ²⁴BUTLER 94 CLE2 e⁺e⁻ ≈ γ(4S)

<4.8 90 MUHEIM 94

4.6 ± 1.5 ²⁵MUHEIM 94

3.1 ± 0.9 ²⁵MUHEIM 94

3.1 ± 0.9 ± 0.6 ²⁴FRABETTI 93G E687 γ Be $\bar{E}_\gamma = 220$ GeV

2.4 ± 1.0 ²⁴ALBRECHT 91 ARG e⁺e⁻ ≈ 10.4 GeV

<4.1 90 0 ²³ADLER 90B MRK3 e⁺e⁻ 4.14 GeV

3.1 ± 0.6 ^{+1.1} _{-0.9} ²⁴ALEXANDER 90B CLEO e⁺e⁻ 10.5–11 GeV

4.8 ± 1.7 ± 1.9 ²⁶ALVAREZ 90C NA14 Photoproduction

>3.4 90 ²⁴ANJOS 90B E691 γ Be, $\bar{E}_\gamma \approx 145$ GeV

2 ± 1 405 ²⁷CHEN 89 CLEO e⁺e⁻ 10 GeV

3.3 ± 1.6 ± 1.0 9 ²⁷BRAUNSCH... 87 TASS e⁺e⁻ 35–44 GeV

3.3 ± 0.1 30 ²⁷DERRICK 85B HRS e⁺e⁻ 29 GeV

- ²¹ AUBERT 05V uses the ratio of $B^0 \rightarrow D^{*-} D_s^{*+}$ events seen in two different ways, in both of which the $D^{*-} \rightarrow \bar{D}^0 \pi^-$ decay is fully reconstructed: (1) The $D_s^{*+} \rightarrow D_s^+ \gamma$, $D_s^+ \rightarrow \phi \pi^+$ decay is fully reconstructed. (2) The number of events in the D_s^+ peak in the missing mass spectrum against the $D^{*-} \gamma$ is measured.
- ²² ARTUSO 96 uses partially reconstructed $\bar{B}^0 \rightarrow D^{*+} D_s^{*-}$ decays to get a model-independent value for $\Gamma(D_s^- \rightarrow \phi \pi^-)/\Gamma(D^0 \rightarrow K^- \pi^+)$ of $0.92 \pm 0.20 \pm 0.11$.
- ²³ BAI 95C uses $e^+ e^- \rightarrow D_s^+ D_s^-$ events in which one or both of the D_s^\pm are observed to obtain the first model-independent measurement of the $D_s^+ \rightarrow \phi \pi^+$ branching fraction, without assumptions about $\sigma(D_s^\pm)$. However, with only two “doubly-tagged” events, the statistical error is very large. ADLER 90B used the same method to set a limit.
- ²⁴ BUTLER 94, FRABETTI 93G, ALBRECHT 91, ALEXANDER 90B, and ANJOS 90B measure the ratio $\Gamma(D_s^+ \rightarrow \phi \ell^+ \nu_\ell)/\Gamma(D_s^+ \rightarrow \phi \pi^+)$, where $\ell = e$ and/or μ , and then use a theoretical calculation of the ratio of widths $\Gamma(D_s^+ \rightarrow \phi \ell^+ \nu_\ell)/\Gamma(D^+ \rightarrow \bar{K}^{*0} \ell^+ \nu)$. Not everyone uses the same value for this ratio.
- ²⁵ The two MUHEIM 94 values here are model-dependent calculations based on distinct data sets. The first uses measurements of the $D_2^*(2460)^0$ and $D_{s1}(2536)^+$, the second uses B -decay factorization and $\Gamma(D_s^+ \rightarrow \mu^+ \nu_\mu)/\Gamma(D_s^+ \rightarrow \phi \ell^+ \nu_\ell)$. A third calculation using the semileptonic width of $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$ is not independent of other results listed here. Note also the upper limit, based on the sum of established D_s^+ branching ratios.
- ²⁶ ALVAREZ 90C relies on the Lund model to estimate the ratio of D_s^+ to D^+ cross sections.
- ²⁷ Values based on crude estimates of the D_s^\pm production level. DERRICK 85B errors are statistical only.

$\Gamma(\phi \pi^+, \phi \rightarrow K^+ K^-)/\Gamma(\phi \pi^+)$

Γ_{16}/Γ_{15}

<u>VALUE</u>	<u>DOCUMENT ID</u>
0.491 ± 0.006 OUR FIT	
0.491 ± 0.006	28 PDG 06

²⁸ This is, of course, just the $\phi \rightarrow K^+ K^-$ branching fraction, but we need it to connect other modes in the fit.

$\Gamma(\phi \pi^+, \phi \rightarrow K^+ K^-)/\Gamma(K^+ K^- \pi^+)$

Γ_{16}/Γ_{14}

This is the “fit fraction” from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.42 ± 0.05 OUR FIT			
0.396 ± 0.033 ± 0.047	FRABETTI	95B E687	Dalitz fit, 701 evts

$\Gamma(K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+)/\Gamma(K^+ K^- \pi^+)$

Γ_{18}/Γ_{14}

This is the “fit fraction” from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.478 ± 0.046 ± 0.040	FRABETTI	95B E687	Dalitz fit, 701 evts

$\Gamma(K^+\bar{K}^*(892)^0)/\Gamma(\phi\pi^+)$ Γ_{17}/Γ_{15}

Unseen decay modes of the resonances are included. However, we now get branching fractions for resonant submodes of 3-body decays from Dalitz-plot analyses.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.85 \pm 0.34 \pm 0.20$	9	ALVAREZ	90C NA14	Photoproduction
$0.84 \pm 0.30 \pm 0.22$		ADLER	89B MRK3	e^+e^- 4.14 GeV
$1.05 \pm 0.17 \pm 0.12$		CHEN	89 CLEO	e^+e^- 10 GeV
$0.87 \pm 0.13 \pm 0.05$	117	ANJOS	88 E691	Photoproduction
1.44 ± 0.37	87	ALBRECHT	87F ARG	e^+e^- 10 GeV

 $\Gamma(f_0(980)\pi^+, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$ Γ_{19}/Γ_{14}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.11 \pm 0.035 \pm 0.026$	FRABETTI	95B E687	Dalitz fit, 701 evts

 $\Gamma(f_0(1710)\pi^+, f_0 \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$ Γ_{21}/Γ_{14}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.034 \pm 0.023 \pm 0.035$	²⁹ FRABETTI	95B E687	Dalitz fit, 701 evts

²⁹ In other words, FRABETTI 95B doesn't see this resonance.

 $\Gamma(K^+\bar{K}_0^*(1430)^0, \bar{K}_0^* \rightarrow K^-\pi^+)/\Gamma(K^+K^-\pi^+)$ Γ_{20}/Γ_{14}

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.093 \pm 0.032 \pm 0.032$	FRABETTI	95B E687	Dalitz fit, 701 evts

 $\Gamma(K^+K^-\pi^+ \text{ nonresonant})/\Gamma(\phi\pi^+)$ Γ_{22}/Γ_{15}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.25 \pm 0.07 \pm 0.05$	48	ANJOS	88 E691	Photoproduction

 $\Gamma(K^*(892)^+\bar{K}^0)/\Gamma(\phi\pi^+)$ Γ_{24}/Γ_{15}

Unseen decay modes of the resonances are included.

VALUE	DOCUMENT ID	TECN	COMMENT
$1.20 \pm 0.21 \pm 0.13$	CHEN	89 CLEO	e^+e^- 10 GeV

 $\Gamma(K^*(892)^+\bar{K}^0)/\Gamma(K^+\bar{K}^0)$ Γ_{24}/Γ_{13}

Unseen decay modes of the $K^*(892)^+$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.9	90	FRABETTI	95 E687	$\gamma\text{Be } \bar{E}_\gamma \approx 200 \text{ GeV}$

 $\Gamma(\phi\pi^+\pi^0)/\Gamma(\phi\pi^+)$ Γ_{26}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.4 \pm 1.0 \pm 0.5$		11	ANJOS	89E E691	Photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.6		90	ALVAREZ	90C NA14	Photoproduction

$\Gamma(\phi\rho^+)/\Gamma(\phi\pi^+)$					Γ_{27}/Γ_{15}
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.86 \pm 0.26^{+0.29}_{-0.40}$	253	AVERY	92 CLE2	$e^+e^- \simeq 10.5$ GeV	
$\Gamma(\phi\pi^+\pi^0\text{3-body})/\Gamma(\phi\pi^+)$					Γ_{28}/Γ_{15}
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.71	90	DAOUDI	92 CLE2	$e^+e^- \approx 10.5$ GeV	
$\Gamma(K^+K^-\pi^+\pi^0\text{non-}\phi)/\Gamma(\phi\pi^+)$					Γ_{29}/Γ_{15}
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<2.4	90	ANJOS	89E E691	Photoproduction	
$\Gamma(K^+\bar{K}^0\pi^+\pi^-)/\Gamma(K^0K^-\pi^+\pi^+)$					Γ_{30}/Γ_{31}
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.586 \pm 0.052 \pm 0.043$	476	LINK	01C FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV	
$\Gamma(K^0K^-\pi^+\pi^+)/\Gamma(\phi\pi^+)$					Γ_{31}/Γ_{15}
VALUE	DOCUMENT ID	TECN	COMMENT		
$1.2 \pm 0.2 \pm 0.2$	ALBRECHT	92B ARG	$e^+e^- \simeq 10.4$ GeV		
$\Gamma(K^*(892)^+\bar{K}^*(892)^0)/\Gamma(\phi\pi^+)$					Γ_{32}/Γ_{15}
Unseen decay modes of the resonances are included.					
VALUE	DOCUMENT ID	TECN	COMMENT		
$1.6 \pm 0.4 \pm 0.4$	ALBRECHT	92B ARG	$e^+e^- \simeq 10.4$ GeV		
$\Gamma(K^0K^-\pi^+\pi^+(\text{non-}K^{*+}\bar{K}^{*0}))/\Gamma(\phi\pi^+)$					Γ_{33}/Γ_{15}
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.80	90	ALBRECHT	92B ARG	$e^+e^- \simeq 10.4$ GeV	
$\Gamma(K^+K^-\pi^+\pi^+\pi^-)/\Gamma(K^+K^-\pi^+)$					Γ_{34}/Γ_{14}
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.160 ± 0.027 OUR AVERAGE					
0.150 ± 0.019 ± 0.025	240	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV	
0.188 ± 0.036 ± 0.040	75	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV	
$\Gamma(\phi\pi^+\pi^+\pi^-)/\Gamma(\phi\pi^+)$					Γ_{35}/Γ_{15}
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.269 ± 0.027 OUR AVERAGE					
0.249 ± 0.024 ± 0.021		136	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
0.28 ± 0.06 ± 0.01		40	FRABETTI	97C E687	γ Be, $\bar{E}_\gamma \approx 200$ GeV
0.58 ± 0.21 ± 0.10		21	FRABETTI	92 E687	γ Be
0.42 ± 0.13 ± 0.07		19	ANJOS	88 E691	Photoproduction
1.11 ± 0.37 ± 0.28		62	ALBRECHT	85D ARG	$e^+e^- 10$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.24	90	ALVAREZ	90C NA14	Photoproduction	

$\Gamma(\phi\pi^+\pi^+\pi^-)/\Gamma(K^+K^-\pi^+\pi^-\pi^-)$ Γ_{35}/Γ_{34}

Unseen decay modes of the ϕ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.42±0.10±0.12	136	³⁰ LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
----------------	-----	--------------------	----------	--

³⁰ This LINK 03D result is redundant with its $\Gamma(\phi\pi^+\pi^+\pi^-)/\Gamma(\phi\pi^+)$ result above.

$\Gamma(K^+K^-\rho^0\pi^+\text{non-}\phi)/\Gamma(K^+K^-\pi^+\pi^+\pi^-)$ Γ_{36}/Γ_{34}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.03	90	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
-------	----	------	----------	--

$\Gamma(\phi\rho^0\pi^+)/\Gamma(K^+K^-\pi^+\pi^+\pi^-)$ Γ_{37}/Γ_{34}

Unseen decay modes of the ϕ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

1.50±0.12±0.08	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
----------------	------	----------	--

$\Gamma(\phi a_1(1260)^+)/\Gamma(K^+K^-\pi^+)$ Γ_{38}/Γ_{14}

Unseen decay modes of the ϕ and $a_1(1260)^+$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.559±0.078±0.044	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
-------------------	------	----------	--

$\Gamma(K^+K^-\pi^+\pi^+\pi^-\text{nonresonant})/\Gamma(K^+K^-\pi^+\pi^+\pi^-)$ Γ_{39}/Γ_{34}

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.10±0.06±0.05	LINK	03D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
----------------	------	----------	--

$\Gamma(K_S^0 K_S^0 \pi^+\pi^+\pi^-)/\Gamma(K^0 K^-\pi^+\pi^+)$ Γ_{40}/Γ_{31}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.051±0.015±0.015	37 ± 10	LINK	04D FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV
-------------------	---------	------	----------	--

————— Pionic modes —————

$\Gamma(\pi^+\pi^+\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{41}/Γ_{14}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.235±0.035 OUR FIT	Error includes scale factor of 1.1.			
---------------------	-------------------------------------	--	--	--

0.265±0.041±0.031	98	FRABETTI	97D E687	γ Be ≈ 200 GeV
-------------------	----	----------	----------	-------------------------------

$\Gamma(\pi^+\pi^+\pi^-)/\Gamma(\phi\pi^+)$ Γ_{41}/Γ_{15}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.277±0.035 OUR FIT	Error includes scale factor of 1.3.			
---------------------	-------------------------------------	--	--	--


0.27 ±0.04 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
------------------------	---	--	--	--

0.245±0.028 ^{+0.019} _{-0.012}	848	AITALA	01A E791	π^- nucleus, 500 GeV
---	-----	--------	----------	--------------------------

0.33 ±0.10 ±0.04	29	ADAMOVICH	93 WA82	π^- 340 GeV
------------------	----	-----------	---------	-----------------

0.44 ±0.10 ±0.04	68	ANJOS	89 E691	Photoproduction
------------------	----	-------	---------	-----------------

WEIGHTED AVERAGE
 0.27 ± 0.04 (Error scaled by 1.3)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(\rho^0 \pi^+) / \Gamma(\pi^+ \pi^+ \pi^-)$

$\Gamma_{42} / \Gamma_{41}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.058 \pm 0.023 \pm 0.037$		AITALA	01A E791	Dalitz fit, 848 evts
<0.073	90	FRABETTI	97D E687	γ Be \approx 200 GeV

$\Gamma(\rho^0 \pi^+) / \Gamma(\phi \pi^+)$

$\Gamma_{42} / \Gamma_{15}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.08	90	ANJOS	89 E691	Photoproduction
<0.22	90	ALBRECHT	87G ARG	$e^+ e^-$ 10 GeV

$\Gamma(\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}) / \Gamma(\pi^+ \pi^+ \pi^-)$

$\Gamma_{43} / \Gamma_{41}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.8704 \pm 0.0560 \pm 0.0438$	³¹ LINK	04 FOCS	Dalitz fit, 1475 ± 50 evts

³¹LINK 04 borrows a K-matrix parametrization from ANISOVICH 03 of the full $\pi-\pi$ S-wave isoscalar scattering amplitude to describe the $\pi^+ \pi^-$ S-wave component of the $\pi^+ \pi^+ \pi^-$ state. The fit fraction given above is a sum over five f_0 mesons, the $f_0(980)$, $f_0(1300)$, $f_0(1200-1600)$, $f_0(1500)$, and $f_0(1750)$. See LINK 04 for details and discussion.

$\Gamma(f_0(980)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{44}/Γ_{41}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.565 \pm 0.043 \pm 0.047$	AITALA	01A E791	Dalitz fit, 848 evts
$1.074 \pm 0.140 \pm 0.043$	FRABETTI	97D E687	γ Be \approx 200 GeV

$\Gamma(f_2(1270)\pi^+, f_2 \rightarrow \pi^+\pi^-)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{47}/Γ_{41}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.0974 \pm 0.0449 \pm 0.0294$	LINK	04 FOCS	Dalitz fit, 1475 ± 50 evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.197 \pm 0.033 \pm 0.006$	AITALA	01A E791	Dalitz fit, 848 evts
$0.123 \pm 0.056 \pm 0.018$	FRABETTI	97D E687	γ Be \approx 200 GeV

$\Gamma(f_0(1370)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{45}/Γ_{41}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.324 \pm 0.077 \pm 0.017$	AITALA	01A E791	Dalitz fit, 848 evts

$\Gamma(\rho(1450)^0\pi^+, \rho^0 \rightarrow \pi^+\pi^-)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{48}/Γ_{41}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.0656 \pm 0.0343 \pm 0.0440$	LINK	04 FOCS	Dalitz fit, 1475 ± 50 evts
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.044 \pm 0.021 \pm 0.002$	AITALA	01A E791	Dalitz fit, 848 evts

$\Gamma(f_0(1500)\pi^+, f_0 \rightarrow \pi^+\pi^-)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{46}/Γ_{41}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.274 \pm 0.114 \pm 0.019$	³² FRABETTI	97D E687	γ Be \approx 200 GeV
³² FRABETTI 97D calls this mode $S(1475)\pi^+$, but finds the mass and width of this $S(1475)$ to be in excellent agreement with those of the $f_0(1500)$.			

$\Gamma(\pi^+\pi^+\pi^- \text{ nonresonant})/\Gamma(\pi^+\pi^+\pi^-)$ Γ_{49}/Γ_{41}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.005 \pm 0.014 \pm 0.017$		AITALA	01A E791	π^- nucleus, 500 GeV
<0.269	90	FRABETTI	97D E687	γ Be \approx 200 GeV

$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma(\phi\pi^+)$ Γ_{50}/Γ_{15}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.3	90	ANJOS	89E E691	Photoproduction

$\Gamma(\eta\pi^+)/\Gamma(\phi\pi^+)$ Γ_{51}/Γ_{15}

Unseen decay modes of the resonances are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.48±0.03±0.04	920	JESSOP	98 CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.54±0.09±0.06	165	ALEXANDER	92 CLE2	See JESSOP 98

$\Gamma(\omega\pi^+)/\Gamma(\eta\pi^+)$ Γ_{52}/Γ_{51}

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.16±0.04±0.03	BALEST	97 CLE2	$e^+e^- \approx \Upsilon(4S)$

$\Gamma(3\pi^+2\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{53}/Γ_{14}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.146±0.014 OUR AVERAGE				
0.145±0.011±0.010	671	LINK	03D FOCS	$\gamma A, \bar{E}_\gamma \approx 180$ GeV
0.158±0.042±0.031	37	FRABETTI	97C E687	$\gamma Be, \bar{E}_\gamma \approx 200$ GeV

$\Gamma(\eta\rho^+)/\Gamma(\phi\pi^+)$ Γ_{55}/Γ_{15}

Unseen decay modes of the resonances are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.98±0.20±0.39	447	JESSOP	98 CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.86±0.38 ^{+0.36} _{-0.38}	217	AVERY	92 CLE2	See JESSOP 98

$\Gamma(\eta\pi^+\pi^0\text{-body})/\Gamma(\phi\pi^+)$ Γ_{56}/Γ_{15}

Unseen decay modes of the resonances are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.1	90	JESSOP	98 CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.82	90	³³ DAUDI	92 CLE2	See JESSOP 98

³³We use the JESSOP 98 limit, even though the DAUDI 92 limit, from the same experiment but with a much smaller data sample, is more restrictive.

$\Gamma(3\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{57}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.049^{+0.033}_{-0.030}	BARLAG	92C ACCM	π^- 230 GeV

$\Gamma(\eta'(958)\pi^+)/\Gamma(\phi\pi^+)$ Γ_{58}/Γ_{15}

Unseen decay modes of the resonances are included.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.08±0.09 OUR AVERAGE					
1.03±0.06±0.07		537	JESSOP	98 CLE2	$e^+e^- \approx \Upsilon(4S)$
2.5 ±1.0 ^{+1.5} _{-0.4}		22	ALVAREZ	91 NA14	Photoproduction
2.5 ±0.5 ±0.3		215	ALBRECHT	90D ARG	$e^+e^- \approx 10.4$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.20±0.15±0.11		281	ALEXANDER	92 CLE2	See JESSOP 98
<1.3	90		ANJOS	91B E691	$\gamma Be, \bar{E}_\gamma \approx 145$ GeV

$\Gamma(\eta'(958)\rho^+)/\Gamma(\phi\pi^+)$ Γ_{60}/Γ_{15}

Unseen decay modes of the resonances are included.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.78±0.28±0.30	137	JESSOP	98 CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$3.44 \pm 0.62^{+0.44}_{-0.46}$	68	AVERY	92 CLE2	See JESSOP 98

 $\Gamma(\eta'(958)\pi^+\pi^0\text{3-body})/\Gamma(\phi\pi^+)$ Γ_{61}/Γ_{15}

Unseen decay modes of the resonances are included.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.4	90	JESSOP	98 CLE2	$e^+e^- \approx \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.85	90	DAOUDI	92 CLE2	See JESSOP 98

———— Modes with one or three K's ————

 $\Gamma(K^0\pi^+)/\Gamma(\phi\pi^+)$ Γ_{62}/Γ_{15}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.21	90	ADLER	89B MRK3	e^+e^- 4.14 GeV

 $\Gamma(K^0\pi^+)/\Gamma(K^+\bar{K}^0)$ Γ_{62}/Γ_{13}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.53	90	FRABETTI	95 E687	$\gamma\text{Be}, \bar{E}_\gamma \approx 200$ GeV

 $\Gamma(K^+\pi^+\pi^-)/\Gamma(K^+K^-\pi^+)$ Γ_{63}/Γ_{14}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.127±0.007±0.014	567 ± 31	LINK	04F FOCS	$\gamma\text{A}, \bar{E}_\gamma \approx 180$ GeV

 $\Gamma(K^+\pi^+\pi^-)/\Gamma(\phi\pi^+)$ Γ_{63}/Γ_{15}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.28 \pm 0.06 \pm 0.05$	85	FRABETTI	95E E687	$\gamma\text{Be}, \bar{E}_\gamma = 220$ GeV

 $\Gamma(K^+\rho^0)/\Gamma(K^+\pi^+\pi^-)$ Γ_{64}/Γ_{63}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.3883±0.0531±0.0261	LINK	04F FOCS	Dalitz fit, 567 evts

 $\Gamma(K^+\rho(1450)^0, \rho^0 \rightarrow \pi^+\pi^-)/\Gamma(K^+\pi^+\pi^-)$ Γ_{65}/Γ_{63}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.1062±0.0351±0.0104	LINK	04F FOCS	Dalitz fit, 567 evts

 $\Gamma(K^*(892)^0\pi^+, K^{*0} \rightarrow K^+\pi^-)/\Gamma(K^+\pi^+\pi^-)$ Γ_{66}/Γ_{63}

This is the "fit fraction" from the Dalitz-plot analysis.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.2164±0.0321±0.0114	LINK	04F FOCS	Dalitz fit, 567 evts

$\Gamma(K^*(1410)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{67} / \Gamma_{63}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1882 ± 0.0403 ± 0.0122	LINK	04F FOCS	Dalitz fit, 567 evts

$\Gamma(K^*(1430)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{68} / \Gamma_{63}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0765 ± 0.0500 ± 0.0170	LINK	04F FOCS	Dalitz fit, 567 evts

$\Gamma(K^+ \pi^+ \pi^- \text{ nonresonant}) / \Gamma(K^+ \pi^+ \pi^-)$ $\Gamma_{69} / \Gamma_{63}$

This is the "fit fraction" from the Dalitz-plot analysis.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1588 ± 0.0492 ± 0.0153	LINK	04F FOCS	Dalitz fit, 567 evts

$\Gamma(K^+ K^+ K^-) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{70} / \Gamma_{14}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.95 ± 2.12^{+2.24}_{-2.31}	31	LINK	02I FOCS	γ nucleus, ≈ 180 GeV

$\Gamma(\phi K^+) / \Gamma(\phi \pi^+)$ $\Gamma_{71} / \Gamma_{15}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.013	90	FRABETTI	95F E687	γ Be, $\bar{E}_\gamma \approx 220$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.071	90	ANJOS	92D E691	γ Be, $\bar{E}_\gamma = 145$ GeV

————— Doubly-Cabibbo-suppressed modes —————

$\Gamma(K^+ K^+ \pi^-) / \Gamma(K^+ K^- \pi^+)$ $\Gamma_{72} / \Gamma_{14}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0052 ± 0.0017 ± 0.0011	27 ± 9	LINK	05K FOCS	<0.78%, CL = 90%

————— Rare or forbidden modes —————

$\Gamma(\pi^+ e^+ e^-) / \Gamma_{\text{total}}$ Γ_{73} / Γ

This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.7 × 10⁻⁴	90	AITALA	99G E791	π^- N 500 GeV

$\Gamma(\pi^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$ Γ_{74} / Γ

This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<2.6 × 10⁻⁵	90		LINK	03F FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.4 × 10 ⁻⁴	90		AITALA	99G E791	π^- N 500 GeV
<4.3 × 10 ⁻⁴	90	0	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{75}/Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.6 \times 10^{-3}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

$\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{76}/Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.6 \times 10^{-5}$	90		LINK	03F FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.4 \times 10^{-4}$	90		AITALA	99G E791	$\pi^- N$ 500 GeV
$<5.9 \times 10^{-4}$	90	0	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(K^*(892)^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{77}/Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.4 \times 10^{-3}$	90	0	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(\pi^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{78}/Γ**

A test of lepton-family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.1 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

$\Gamma(K^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ **Γ_{79}/Γ**

A test of lepton-family-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.3 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

$\Gamma(\pi^- e^+ e^+)/\Gamma_{\text{total}}$ **Γ_{80}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.9 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

$\Gamma(\pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{81}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.9 \times 10^{-5}$	90		LINK	03F FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<8.2 \times 10^{-5}$	90		AITALA	99G E791	$\pi^- N$ 500 GeV
$<4.3 \times 10^{-4}$	90	0	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$ **Γ_{82}/Γ**

A test of lepton-number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.3 \times 10^{-4}$	90	AITALA	99G E791	$\pi^- N$ 500 GeV

$\Gamma(K^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{83}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.3 \times 10^{-4}$	90	AITALA	99G E791	π^- N 500 GeV

$\Gamma(K^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{84}/Γ

A test of lepton-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-5}$	90		LINK	03F FOCS	γ nucleus, $\bar{E}_\gamma \approx 180$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.8 \times 10^{-4}$	90		AITALA	99G E791	π^- N 500 GeV
$<5.9 \times 10^{-4}$	90	0	KODAMA	95 E653	π^- emulsion 600 GeV

$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{85}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.8 \times 10^{-4}$	90	AITALA	99G E791	π^- N 500 GeV

$\Gamma(K^*(892)^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{86}/Γ

A test of lepton-number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-3}$	90	0	KODAMA	95 E653	π^- emulsion 600 GeV

$D_s^+ - D_s^-$ T-VIOLATING DECAY-RATE ASYMMETRIES

$A_{Tviol}(K_S^0 K^\pm \pi^+ \pi^-)$ in $D_s^\pm \rightarrow K_S^0 K^\pm \pi^+ \pi^-$

$C_T \equiv \vec{p}_{K^+} \cdot (\vec{p}_{\pi^+} \times \vec{p}_{\pi^-})$ is a T -odd correlation of the K^+ , π^+ , and π^- momenta for the D_s^+ . $\bar{C}_T \equiv \vec{p}_{K^-} \cdot (\vec{p}_{\pi^-} \times \vec{p}_{\pi^+})$ is the corresponding quantity for the D_s^- . $A_T \equiv [\Gamma(C_T > 0) - \Gamma(C_T < 0)] / [\Gamma(C_T > 0) + \Gamma(C_T < 0)]$ would, in the absence of strong phases, test for T violation in D_s^+ decays (the Γ 's are partial widths). With $\bar{A}_T \equiv [\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)] / [\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)]$, the asymmetry $A_{Tviol} \equiv \frac{1}{2}(A_T - \bar{A}_T)$ tests for T violation even with nonzero strong phases.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$-0.036 \pm 0.067 \pm 0.023$	508 ± 34	LINK	05E FOCS	γ A, $\bar{E}_\gamma \approx 180$ GeV

$D_s^+ \rightarrow \phi \ell^+ \nu_\ell$ FORM FACTORS

$r_2 \equiv A_2(0)/A_1(0)$ in $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.32 ± 0.24 OUR AVERAGE				Error includes scale factor of 1.2.
$0.713 \pm 0.202 \pm 0.284$	793	LINK	04C FOCS	$\phi \mu^+ \nu_\mu$
$1.57 \pm 0.25 \pm 0.19$	271	AITALA	99D E791	$\phi e^+ \nu_e, \phi \mu^+ \nu_\mu$
$1.4 \pm 0.5 \pm 0.3$	308	AVERY	94B CLE2	$\phi e^+ \nu_e$
$1.1 \pm 0.8 \pm 0.1$	90	FRABETTI	94F E687	$\phi \mu^+ \nu_\mu$
$2.1 \begin{smallmatrix} +0.6 \\ -0.5 \end{smallmatrix} \pm 0.2$	19	KODAMA	93 E653	$\phi \mu^+ \nu_\mu$

$r_\nu \equiv V(0)/A_1(0)$ in $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
1.72 ± 0.21 OUR AVERAGE				
1.549 ± 0.250 ± 0.148	793	LINK	04C FOCS	$\phi \mu^+ \nu_\mu$
2.27 ± 0.35 ± 0.22	271	AITALA	99D E791	$\phi e^+ \nu_e, \phi \mu^+ \nu_\mu$
0.9 ± 0.6 ± 0.3	308	AVERY	94B CLE2	$\phi e^+ \nu_e$
1.8 ± 0.9 ± 0.2	90	FRABETTI	94F E687	$\phi \mu^+ \nu_\mu$
2.3 $^{+1.1}_{-0.9}$ ± 0.4	19	KODAMA	93 E653	$\phi \mu^+ \nu_\mu$

Γ_L/Γ_T in $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.72 ± 0.18 OUR AVERAGE				
1.0 ± 0.3 ± 0.2	308	AVERY	94B CLE2	$\phi e^+ \nu_e$
1.0 ± 0.5 ± 0.1	90	³⁴ FRABETTI	94F E687	$\phi \mu^+ \nu_\mu$
0.54 ± 0.21 ± 0.10	19	³⁴ KODAMA	93 E653	$\phi \mu^+ \nu_\mu$

³⁴ FRABETTI 94F and KODAMA 93 evaluate Γ_L/Γ_T for a lepton mass of zero.

D_s^\pm REFERENCES

PDG	06	JPG 33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
AUBERT	05V	PR D71 091104R	B. Aubert <i>et al.</i>	(BABAR Collab.)
LINK	05E	PL B622 239	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05J	PRL 95 052003	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	05K	PL B624 166	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04	PL B585 200	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04C	PL B586 183	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04D	PL B586 191	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	04F	PL B601 10	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ACOSTA	03D	PR D68 072004	D. Acosta <i>et al.</i>	(FNAL CDF-II Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
LINK	03D	PL B561 225	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	03F	PL B572 21	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
AUBERT	02G	PR D65 091104R	B. Aubert <i>et al.</i>	(BaBar Collab.)
HEISTER	02I	PL B528 1	A. Heister <i>et al.</i>	(ALEPH Collab.)
LINK	02I	PL B541 227	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	02J	PL B541 243	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABBIENDI	01L	PL B516 236	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
IORI	01	PL B523 22	M. Iori <i>et al.</i>	(FNAL SELEX Collab.)
LINK	01C	PRL 87 162001	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ALEXANDROV	00	PL B478 31	Y. Alexandrov <i>et al.</i>	(CERN BEATRICE Collab.)
AITALA	99	PL B445 449	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	99D	PL B450 294	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AITALA	99G	PL B462 401	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BONVICINI	99	PRL 82 4586	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BAI	98	PR D57 28	J.Z. Bai <i>et al.</i>	(BEP C BES Collab.)
CHADHA	98	PR D58 032002	M. Chada <i>et al.</i>	(CLEO Collab.)
JESSOP	98	PR D58 052002	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
ACCIARRI	97F	PL B396 327	M. Acciarri <i>et al.</i>	(L3 Collab.)
BAI	97	PR D56 3779	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALEST	97	PRL 79 1436	R. Balest <i>et al.</i>	(CLEO Collab.)
FRABETTI	97C	PL B401 131	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	97D	PL B407 79	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ARTUSO	96	PL B378 364	M. Artuso <i>et al.</i>	(CLEO Collab.)
KODAMA	96	PL B382 299	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
BAI	95	PRL 74 4599	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	95C	PR D52 3781	J.Z. Bai <i>et al.</i>	(BES Collab.)
BRANDENB...	95	PRL 75 3804	G.W. Brandenburg <i>et al.</i>	(CLEO Collab.)
FRABETTI	95	PL B346 199	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)

FRABETTI	95B	PL B351 591	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	95E	PL B359 403	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	95F	PL B363 259	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	95	PL B345 85	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ACOSTA	94	PR D49 5690	D. Acosta <i>et al.</i>	(CLEO Collab.)
AVERY	94B	PL B337 405	P. Avery <i>et al.</i>	(CLEO Collab.)
BROWN	94	PR D50 1884	D. Brown <i>et al.</i>	(CLEO Collab.)
BUTLER	94	PL B324 255	F. Butler <i>et al.</i>	(CLEO Collab.)
FRABETTI	94F	PL B328 187	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
MUHEIM	94	PR D49 3767	F. Muheim, S. Stone	(SYRA)
ADAMOVIICH	93	PL B305 177	M.I. Adamovich <i>et al.</i>	(CERN WA82 Collab.)
AOKI	93	PTP 89 131	S. Aoki <i>et al.</i>	(CERN WA75 Collab.)
FRABETTI	93F	PRL 71 827	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
FRABETTI	93G	PL B313 253	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
KODAMA	93	PL B309 483	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
KODAMA	93B	PL B313 260	K. Kodama <i>et al.</i>	(FNAL E653 Collab.)
ALBRECHT	92B	ZPHY C53 361	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	92	PRL 68 1275	J. Alexander <i>et al.</i>	(CLEO Collab.)
ANJOS	92D	PRL 69 2892	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
AVERY	92	PRL 68 1279	P. Avery <i>et al.</i>	(CLEO Collab.)
BARLAG	92C	ZPHY C55 383	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
Also		ZPHY C48 29	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
DAOUDI	92	PR D45 3965	M. Daoudi <i>et al.</i>	(CLEO Collab.)
FRABETTI	92	PL B281 167	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALBRECHT	91	PL B255 634	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALVAREZ	91	PL B255 639	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ANJOS	91B	PR D43 R2063	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
COFFMAN	91	PL B263 135	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
ADLER	90B	PRL 64 169	J.C. Adler <i>et al.</i>	(Mark III Collab.)
ALBRECHT	90D	PL B245 315	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	90B	PRL 65 1531	J. Alexander <i>et al.</i>	(CLEO Collab.)
ALVAREZ	90C	PL B246 261	M.P. Alvarez <i>et al.</i>	(CERN NA14/2 Collab.)
ANJOS	90B	PRL 64 2885	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ANJOS	90C	PR D41 2705	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
BAI	90	PRL 65 686	Z. Bai <i>et al.</i>	(Mark III Collab.)
BARLAG	90C	ZPHY C46 563	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
FRABETTI	90	PL B251 639	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ADLER	89B	PRL 63 1211	J. Adler <i>et al.</i>	(Mark III Collab.)
Also		PRL 63 2858 (erratum)	J. Adler <i>et al.</i>	(Mark III Collab.)
ANJOS	89	PRL 62 125	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
ANJOS	89E	PL B223 267	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
CHEN	89	PL B226 192	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
ALBRECHT	88	PL B207 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	88	PRL 60 897	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)
RAAB	88	PR D37 2391	J.R. Raab <i>et al.</i>	(FNAL E691 Collab.)
ALBRECHT	87F	PL B179 398	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87G	PL B195 102	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BECKER	87B	PL B184 277	H. Becker <i>et al.</i>	(NA11 and NA32 Collab.)
BLAYLOCK	87	PRL 58 2171	G.T. Blaylock <i>et al.</i>	(Mark III Collab.)
BRAUNSCH...	87	ZPHY C35 317	W. Braunschweig <i>et al.</i>	(TASSO Collab.)
USHIDA	86	PRL 56 1767	N. Ushida <i>et al.</i>	(FNAL E531 Collab.)
ALBRECHT	85D	PL 153B 343	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
DERRICK	85B	PRL 54 2568	M. Derrick <i>et al.</i>	(HRS Collab.)
AIHARA	84D	PRL 53 2465	H. Aihara <i>et al.</i>	(TPC Collab.)
ALTHOFF	84	PL 136B 130	M. Althoff <i>et al.</i>	(TASSO Collab.)
BAILEY	84	PL 139B 320	R. Bailey <i>et al.</i>	(ACCMOR Collab.)
AUBERT	83	NP B213 31	J.J. Aubert <i>et al.</i>	(EMC Collab.)
CHEN	83C	PRL 51 634	A. Chen <i>et al.</i>	(CLEO Collab.)

OTHER RELATED PAPERS

RICHMAN	95	RMP 67 893	J.D. Richman, P.R. Burchat	(UCSB, STAN)
---------	----	------------	----------------------------	--------------