

B^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

Quantum numbers not measured. Values shown are quark-model predictions.

See also the B^\pm/B^0 ADMIXTURE and $B^\pm/B_s^0/B_s^0/b$ -baryon ADMIXTURE sections.

B^\pm MASS

The fit uses m_{B^+} , $(m_{B^0} - m_{B^+})$, and m_{B^0} to determine m_{B^+} , m_{B^0} , and the mass difference.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5279.0±0.5 OUR FIT				
5279.1±0.5 OUR AVERAGE				
5279.1±0.4 ±0.4	526	1 CSORNA	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
5279.1±1.7 ±1.4	147	ABE	96B CDF	$p\bar{p}$ at 1.8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5278.8±0.54±2.0	362	ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
5278.3±0.4 ±2.0		BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
5280.5±1.0 ±2.0		2 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$
5275.8±1.3 ±3.0	32	ALBRECHT	87C ARG	$e^+ e^- \rightarrow \gamma(4S)$
5278.2±1.8 ±3.0	12	3 ALBRECHT	87D ARG	$e^+ e^- \rightarrow \gamma(4S)$
5278.6±0.8 ±2.0		BEBEK	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$
1 CSORNA 00 uses fully reconstructed $526 B^+ \rightarrow J/\psi(') K^+$ events and invariant masses without beam constraint.				
2 ALBRECHT 90J assumes 10580 for $\gamma(4S)$ mass. Supersedes ALBRECHT 87C and ALBRECHT 87D.				
3 Found using fully reconstructed decays with $J/\psi(1S)$. ALBRECHT 87D assume $m\gamma(4S) = 10577$ MeV.				

B^\pm MEAN LIFE

See $B^\pm/B_s^0/b$ -baryon ADMIXTURE section for data on B -hadron mean life averaged over species of bottom particles.

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account corrections between the measurements and asymmetric lifetime errors.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.638±0.011 OUR EVALUATION				
1.635±0.011±0.011	4 ABE	05B BELL	$e^+ e^- \rightarrow \gamma(4S)$	
1.624±0.014±0.018	5 ABDALLAH	04E DLPH	$e^+ e^- \rightarrow Z$	
1.636±0.058±0.025	6 ACOSTA	02C CDF	$p\bar{p}$ at 1.8 TeV	
1.673±0.032±0.023	7 AUBERT	01F BABR	$e^+ e^- \rightarrow \gamma(4S)$	
1.648±0.049±0.035	8 BARATE	00R ALEP	$e^+ e^- \rightarrow Z$	
1.643±0.037±0.025	9 ABBIENDI	99J OPAL	$e^+ e^- \rightarrow Z$	

$1.637 \pm 0.058^{+0.045}_{-0.043}$	⁸ ABE	98Q CDF	$p\bar{p}$ at 1.8 TeV
$1.66 \pm 0.06 \pm 0.03$	⁹ ACCIARRI	98S L3	$e^+ e^- \rightarrow Z$
$1.66 \pm 0.06 \pm 0.05$	⁹ ABE	97J SLD	$e^+ e^- \rightarrow Z$
$1.58^{+0.21}_{-0.18} \pm 0.04$	94	⁶ BUSKULIC	$e^+ e^- \rightarrow Z$
$1.61 \pm 0.16 \pm 0.12$		^{8,10} ABREU	$e^+ e^- \rightarrow Z$
$1.72 \pm 0.08 \pm 0.06$		¹¹ ADAM	$e^+ e^- \rightarrow Z$
$1.52 \pm 0.14 \pm 0.09$		⁸ AKERS	$e^+ e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.695 \pm 0.026 \pm 0.015$	⁷ ABE	02H BELL	Repl. by ABE 05B
$1.68 \pm 0.07 \pm 0.02$	⁶ ABE	98B CDF	Repl. by ACOSTA 02C
$1.56 \pm 0.13 \pm 0.06$	⁸ ABE	96C CDF	Repl. by ABE 98Q
$1.58 \pm 0.09 \pm 0.03$		¹² BUSKULIC	$e^+ e^- \rightarrow Z$
$1.58 \pm 0.09 \pm 0.04$		⁸ BUSKULIC	Repl. by BARATE 00R
1.70 ± 0.09		¹³ ADAM	$e^+ e^- \rightarrow Z$
$1.61 \pm 0.16 \pm 0.05$	148	⁶ ABE	94D CDF Repl. by ABE 98B
$1.30^{+0.33}_{-0.29} \pm 0.16$	92	⁸ ABREU	93D DLPN Sup. by ABREU 95Q
$1.56 \pm 0.19 \pm 0.13$	134	¹¹ ABREU	93G DLPN Sup. by ADAM 95
$1.51^{+0.30}_{-0.28} \pm 0.12_{-0.14}$	59	⁸ ACTON	93C OPAL Sup. by AKERS 95T
$1.47^{+0.22}_{-0.19} \pm 0.15_{-0.14}$	77	⁸ BUSKULIC	93D ALEP Sup. by BUSKULIC 96J

⁴ Measurement performed using a combined fit of CP -violation, mixing and lifetimes.

⁵ Measurement performed using an inclusive reconstruction and B flavor identification technique.

⁶ Measured mean life using fully reconstructed decays.

⁷ Events are selected in which one B meson is fully reconstructed while the second B meson is reconstructed inclusively.

⁸ Data analyzed using $D/D^*\ell X$ event vertices.

⁹ Data analyzed using charge of secondary vertex.

¹⁰ ABREU 95Q assumes $B(B^0 \rightarrow D^{**-} \ell^+ \nu_\ell) = 3.2 \pm 1.7\%$.

¹¹ Data analyzed using vertex-charge technique to tag B charge.

¹² Combined result of $D/D^*\ell X$ analysis and fully reconstructed B analysis.

¹³ Combined ABREU 95Q and ADAM 95 result.

B^+ DECAY MODES

B^- modes are charge conjugates of the modes below. Modes which do not identify the charge state of the B are listed in the B^\pm/B^0 ADMIXTURE section.

The branching fractions listed below assume 50% $B^0\bar{B}^0$ and 50% B^+B^- production at the $\Upsilon(4S)$. We have attempted to bring older measurements up to date by rescaling their assumed $\Upsilon(4S)$ production ratio to 50:50 and their assumed D , D_s , D^* , and ψ branching ratios to current values whenever this would affect our averages and best limits significantly.

Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.

For inclusive branching fractions, e.g., $B \rightarrow D^\pm$ anything, the values usually are multiplicities, not branching fractions. They can be greater than one.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Semileptonic and leptonic modes		
$\Gamma_1 \ell^+ \nu_\ell$ anything	[a] (10.9 \pm 0.4) %	
$\Gamma_2 \bar{D}^0 \ell^+ \nu_\ell$	[a] (2.15 \pm 0.22) %	
$\Gamma_3 \bar{D}^*(2007)^0 \ell^+ \nu_\ell$	[a] (6.5 \pm 0.5) %	
$\Gamma_4 \bar{D}_1(2420)^0 \ell^+ \nu_\ell$	(5.6 \pm 1.6) $\times 10^{-3}$	
$\Gamma_5 \bar{D}_2^*(2460)^0 \ell^+ \nu_\ell$	< 8 $\times 10^{-3}$ CL=90%	
$\Gamma_6 D^- \pi^+ \ell^+ \nu_\ell$	(5.3 \pm 1.0) $\times 10^{-3}$	
$\Gamma_7 D^{*-} \pi^+ \ell^+ \nu_\ell$	(6.4 \pm 1.5) $\times 10^{-3}$	
$\Gamma_8 \pi^0 \ell^+ \nu_\ell$	(7.4 \pm 1.1) $\times 10^{-5}$	
$\Gamma_9 \pi^0 e^+ \nu_e$		
$\Gamma_{10} \eta \ell^+ \nu_\ell$	(8 \pm 4) $\times 10^{-5}$	
$\Gamma_{11} \omega \ell^+ \nu_\ell$	[a] (1.3 \pm 0.6) $\times 10^{-4}$	
$\Gamma_{12} \omega \mu^+ \nu_\mu$		
$\Gamma_{13} \rho^0 \ell^+ \nu_\ell$	[a] (1.24 \pm 0.23) $\times 10^{-4}$	
$\Gamma_{14} p\bar{p} e^+ \nu_e$	< 5.2 $\times 10^{-3}$ CL=90%	
$\Gamma_{15} e^+ \nu_e$	< 1.5 $\times 10^{-5}$ CL=90%	
$\Gamma_{16} \mu^+ \nu_\mu$	< 6.6 $\times 10^{-6}$ CL=90%	
$\Gamma_{17} \tau^+ \nu_\tau$	< 2.6 $\times 10^{-4}$ CL=90%	
$\Gamma_{18} e^+ \nu_e \gamma$	< 2.0 $\times 10^{-4}$ CL=90%	
$\Gamma_{19} \mu^+ \nu_\mu \gamma$	< 5.2 $\times 10^{-5}$ CL=90%	
Inclusive modes		
$\Gamma_{20} D^0 X$	(9.8 \pm 1.1) %	
$\Gamma_{21} \bar{D}^0 X$	(79 \pm 5) %	
$\Gamma_{22} D^+ X$	(3.8 \pm 1.0) %	
$\Gamma_{23} D^- X$	(9.8 \pm 1.8) %	
$\Gamma_{24} D_s^+ X$	(14 \pm 5) %	
$\Gamma_{25} D_s^- X$	< 2.2 %	CL=90%
$\Gamma_{26} \Lambda_c^+ X$	(2.9 \pm 1.4) %	
$\Gamma_{27} \bar{\Lambda}_c^- X$	(3.5 \pm 1.5) %	
$\Gamma_{28} \bar{c} X$	(98 \pm 6) %	
$\Gamma_{29} c X$	(33 \pm 6) %	
$\Gamma_{30} \bar{c} c X$	(131 \pm 10) %	

D, D*, or D_s modes

Γ_{31}	$\overline{D}^0 \pi^+$	(4.92 \pm 0.20) $\times 10^{-3}$
Γ_{32}	$D_{CP(+1)} \pi^+$	[b] (4.0 \pm 0.8) $\times 10^{-3}$
Γ_{33}	$D_{CP(-1)} \pi^+$	[b] (3.6 \pm 0.8) $\times 10^{-3}$
Γ_{34}	$\overline{D}^0 \rho^+$	(1.34 \pm 0.18) %
Γ_{35}	$\overline{D}^0 K^+$	(4.08 \pm 0.24) $\times 10^{-4}$
Γ_{36}	$D_{CP(+1)} K^+$	[b] (3.7 \pm 0.6) $\times 10^{-4}$
Γ_{37}	$D_{CP(-1)} K^+$	[b] (3.5 \pm 0.5) $\times 10^{-4}$
Γ_{38}	$[K^- \pi^+]_D K^+$	[c]
Γ_{39}	$[K^+ \pi^-]_D K^+$	[c]
Γ_{40}	$[K^- \pi^+]_D K^*(892)^+$	[c]
Γ_{41}	$[K^+ \pi^-]_D K^*(892)^+$	[c]
Γ_{42}	$[K^- \pi^+]_D \pi^+$	[c] (1.7 \pm 0.5) $\times 10^{-5}$
Γ_{43}	$[\pi^+ \pi^- \pi^0]_D K^-$	(5.5 \pm 1.2) $\times 10^{-6}$
Γ_{44}	$\overline{D}^0 K^*(892)^+$	(6.3 \pm 0.8) $\times 10^{-4}$
Γ_{45}	$D_{CP(-1)} K^*(892)^+$	[b] (2.0 \pm 0.9) $\times 10^{-4}$
Γ_{46}	$D_{CP(+1)} K^*(892)^+$	[b] (6.2 \pm 1.5) $\times 10^{-4}$
Γ_{47}	$\overline{D}^0 K^+ \overline{K}^0$	(5.5 \pm 1.6) $\times 10^{-4}$
Γ_{48}	$\overline{D}^0 K^+ \overline{K}^*(892)^0$	(7.5 \pm 1.7) $\times 10^{-4}$
Γ_{49}	$\overline{D}^0 \pi^+ \pi^+ \pi^-$	(1.1 \pm 0.4) %
Γ_{50}	$\overline{D}^0 \pi^+ \pi^+ \pi^-$ nonresonant	(5 \pm 4) $\times 10^{-3}$
Γ_{51}	$\overline{D}^0 \pi^+ \rho^0$	(4.2 \pm 3.0) $\times 10^{-3}$
Γ_{52}	$\overline{D}^0 a_1(1260)^+$	(4 \pm 4) $\times 10^{-3}$
Γ_{53}	$\overline{D}^0 \omega \pi^+$	(4.1 \pm 0.9) $\times 10^{-3}$
Γ_{54}	$D^*(2010)^- \pi^+ \pi^+$	(1.35 \pm 0.22) $\times 10^{-3}$
Γ_{55}	$D^- \pi^+ \pi^+$	(1.02 \pm 0.16) $\times 10^{-3}$
Γ_{56}	$D^+ K^0$	< 5.0 $\times 10^{-6}$ CL=90%
Γ_{57}	$\overline{D}^*(2007)^0 \pi^+$	(4.6 \pm 0.4) $\times 10^{-3}$
Γ_{58}	$\overline{D}_{CP(+1)}^{*0} \pi^+$	[d]
Γ_{59}	$\overline{D}_{CP(-1)}^{*0} \pi^+$	[d]
Γ_{60}	$\overline{D}^*(2007)^0 \omega \pi^+$	(4.5 \pm 1.2) $\times 10^{-3}$
Γ_{61}	$\overline{D}^*(2007)^0 \rho^+$	(9.8 \pm 1.7) $\times 10^{-3}$
Γ_{62}	$\overline{D}^*(2007)^0 K^+$	(3.7 \pm 0.4) $\times 10^{-4}$
Γ_{63}	$\overline{D}_{CP(+1)}^{*0} K^+$	[d]
Γ_{64}	$\overline{D}_{CP(-1)}^{*0} K^+$	[d]
Γ_{65}	$\overline{D}^*(2007)^0 K^*(892)^+$	(8.1 \pm 1.4) $\times 10^{-4}$
Γ_{66}	$\overline{D}^*(2007)^0 K^+ \overline{K}^0$	< 1.06 $\times 10^{-3}$ CL=90%
Γ_{67}	$\overline{D}^*(2007)^0 K^+ K^*(892)^0$	(1.5 \pm 0.4) $\times 10^{-3}$
Γ_{68}	$\overline{D}^*(2007)^0 \pi^+ \pi^+ \pi^-$	(1.03 \pm 0.12) %
Γ_{69}	$\overline{D}^*(2007)^0 a_1(1260)^+$	(1.9 \pm 0.5) %
Γ_{70}	$\overline{D}^*(2007)^0 \pi^- \pi^+ \pi^+ \pi^0$	(1.8 \pm 0.4) %
Γ_{71}	$\overline{D}^{*0} 3\pi^+ 2\pi^-$	(5.7 \pm 1.2) $\times 10^{-3}$

Γ_{72}	$D^*(2010)^+ \pi^0$	< 1.7	$\times 10^{-4}$	CL=90%
Γ_{73}	$D^*(2010)^+ K^0$	< 9.0	$\times 10^{-6}$	CL=90%
Γ_{74}	$D^*(2010)^- \pi^+ \pi^+ \pi^0$	(1.5 \pm 0.7) %		
Γ_{75}	$D^*(2010)^- \pi^+ \pi^+ \pi^+ \pi^-$	(2.6 \pm 0.4) $\times 10^{-3}$		
Γ_{76}	$\overline{D}_1^*(2420)^0 \pi^+$	(1.5 \pm 0.6) $\times 10^{-3}$	S=1.3	
Γ_{77}	$\overline{D}_1(2420)^0 \pi^+ \times B(\overline{D}_1^0 \rightarrow \overline{D}^0 \pi^+ \pi^-)$	(1.9 \pm 0.5) $\times 10^{-4}$		
Γ_{78}	$\overline{D}_2^*(2462)^0 \pi^+ \times B(\overline{D}_2^*(2462)^0 \rightarrow D^- \pi^+)$	(3.4 \pm 0.8) $\times 10^{-4}$		
Γ_{79}	$\overline{D}_0^*(2308)^0 \pi^+ \times B(\overline{D}_0^*(2308)^0 \rightarrow D^- \pi^+)$	(6.1 \pm 1.9) $\times 10^{-4}$		
Γ_{80}	$\overline{D}_1(2421)^0 \pi^+ \times B(\overline{D}_1(2421)^0 \rightarrow D^{*-} \pi^+)$	(6.8 \pm 1.5) $\times 10^{-4}$		
Γ_{81}	$\overline{D}_2^*(2462)^0 \pi^+ \times B(\overline{D}_2^*(2462)^0 \rightarrow D^{*-} \pi^+)$	(1.8 \pm 0.5) $\times 10^{-4}$		
Γ_{82}	$\overline{D}'_1(2427)^0 \pi^+ \times B(\overline{D}'_1(2427)^0 \rightarrow D^{*-} \pi^+)$	(5.0 \pm 1.2) $\times 10^{-4}$		
Γ_{83}	$\overline{D}_1(2420)^0 \pi^+ \times B(\overline{D}_1^0 \rightarrow \overline{D}^{*0} \pi^+ \pi^-)$	< 6	$\times 10^{-6}$	CL=90%
Γ_{84}	$\overline{D}_1^*(2420)^0 \rho^+$	< 1.4	$\times 10^{-3}$	CL=90%
Γ_{85}	$\overline{D}_2^*(2460)^0 \pi^+$	< 1.3	$\times 10^{-3}$	CL=90%
Γ_{86}	$\overline{D}_2^*(2460)^0 \pi^+ \times B(\overline{D}_2^{*0} \rightarrow \overline{D}^{*0} \pi^+ \pi^-)$	< 2.2	$\times 10^{-5}$	CL=90%
Γ_{87}	$\overline{D}_2^*(2460)^0 \rho^+$	< 4.7	$\times 10^{-3}$	CL=90%
Γ_{88}	$\overline{D}^0 D_s^+$	(1.09 \pm 0.27) %		
Γ_{89}	$D_{s0}(2317)^+ \overline{D}^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0)$	(7.4 \pm 2.3) $\times 10^{-4}$		
Γ_{90}	$D_{s0}(2317)^+ \overline{D}^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^{*+} \gamma)$	< 7.6	$\times 10^{-4}$	CL=90%
Γ_{91}	$D_{s0}(2317)^+ \overline{D}^*(2010)^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^+ \pi^0)$	(9 \pm 7) $\times 10^{-4}$		
Γ_{92}	$D_{sJ}(2457)^+ \overline{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0)$	(1.4 \pm 0.6) $\times 10^{-3}$	S=1.3	
Γ_{93}	$D_{sJ}(2457)^+ \overline{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)$	(4.7 \pm 1.4) $\times 10^{-4}$		
Γ_{94}	$D_{sJ}(2457)^+ \overline{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^+ \pi^-)$	< 2.2	$\times 10^{-4}$	CL=90%
Γ_{95}	$D_{sJ}(2457)^+ \overline{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^0)$	< 2.7	$\times 10^{-4}$	CL=90%

Γ_{96}	$D_{sJ}(2457)^+ \bar{D}^0 \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \gamma)$	< 9.8	$\times 10^{-4}$ CL=90%
Γ_{97}	$D_{sJ}(2457)^+ \bar{D}^*(2010)^0 \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0)$	(7.6 \pm 3.6)	$\times 10^{-3}$
Γ_{98}	$D_{sJ}(2457)^+ \bar{D}^*(2010)^0 \times$ $B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)$	(1.4 \pm 0.7)	$\times 10^{-3}$
Γ_{99}	$\bar{D}^0 D_{sJ}(2536)^+ \times$ $B(D_{sJ}(2536)^+ \rightarrow D^*(2007)^0 K^+)$	< 2	$\times 10^{-4}$ CL=90%
Γ_{100}	$\bar{D}^*(2007)^0 D_{sJ}(2536)^+ \times$ $B(D_{sJ}(2536)^+ \rightarrow D^*(2007)^0 K^+)$	< 7	$\times 10^{-4}$ CL=90%
Γ_{101}	$\bar{D}^0 D_{sJ}(2573)^+ \times$ $B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)$	< 2	$\times 10^{-4}$ CL=90%
Γ_{102}	$\bar{D}^*(2007)^0 D_{sJ}(2573)^+ \times$ $B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)$	< 5	$\times 10^{-4}$ CL=90%
Γ_{103}	$\bar{D}^0 D_s^{*+}$	(7.2 \pm 2.6)	$\times 10^{-3}$
Γ_{104}	$\bar{D}^*(2007)^0 D_s^+$	(10 \pm 4)	$\times 10^{-3}$
Γ_{105}	$\bar{D}^*(2007)^0 D_s^{*+}$	(2.2 \pm 0.7)	%
Γ_{106}	$D_s^{(*)+} \bar{D}^{**0}$	(2.7 \pm 1.2)	%
Γ_{107}	$\bar{D}^*(2007)^0 D^*(2010)^+$	< 1.1	% CL=90%
Γ_{108}	$\bar{D}^0 D^*(2010)^+ +$ $\bar{D}^*(2007)^0 D^+$	< 1.3	% CL=90%
Γ_{109}	$\bar{D}^0 D^*(2010)^+$	(4.6 \pm 0.9)	$\times 10^{-4}$
Γ_{110}	$\bar{D}^0 D^+$	(4.8 \pm 1.0)	$\times 10^{-4}$
Γ_{111}	$\bar{D}^0 D^+ K^0$	< 2.8	$\times 10^{-3}$ CL=90%
Γ_{112}	$\bar{D}^*(2007)^0 D^+ K^0$	< 6.1	$\times 10^{-3}$ CL=90%
Γ_{113}	$\bar{D}^0 \bar{D}^*(2010)^+ K^0$	(5.2 \pm 1.2)	$\times 10^{-3}$
Γ_{114}	$\bar{D}^*(2007)^0 D^*(2010)^+ K^0$	(7.8 \pm 2.6)	$\times 10^{-3}$
Γ_{115}	$\bar{D}^0 D^0 K^+$	(1.37 \pm 0.32)	$\times 10^{-3}$ S=1.5
Γ_{116}	$\bar{D}^*(2010)^0 D^0 K^+$	< 3.8	$\times 10^{-3}$ CL=90%
Γ_{117}	$\bar{D}^0 D^*(2007)^0 K^+$	(4.7 \pm 1.0)	$\times 10^{-3}$
Γ_{118}	$\bar{D}^*(2007)^0 D^*(2007)^0 K^+$	(5.3 \pm 1.6)	$\times 10^{-3}$
Γ_{119}	$D^- D^+ K^+$	< 4	$\times 10^{-4}$ CL=90%
Γ_{120}	$D^- D^*(2010)^+ K^+$	< 7	$\times 10^{-4}$ CL=90%
Γ_{121}	$D^*(2010)^- D^+ K^+$	(1.5 \pm 0.4)	$\times 10^{-3}$
Γ_{122}	$D^*(2010)^- D^*(2010)^+ K^+$	< 1.8	$\times 10^{-3}$ CL=90%
Γ_{123}	$(\bar{D} + \bar{D}^*)(D + D^*)K$	(3.5 \pm 0.6)	%
Γ_{124}	$D_s^+ \pi^0$	< 1.7	$\times 10^{-4}$ CL=90%
Γ_{125}	$D_s^{*+} \pi^0$	< 2.7	$\times 10^{-4}$ CL=90%
Γ_{126}	$D_s^+ \eta$	< 4	$\times 10^{-4}$ CL=90%
Γ_{127}	$D_s^{*+} \eta$	< 6	$\times 10^{-4}$ CL=90%

Γ_{128}	$D_s^+ \rho^0$	<	3.1	$\times 10^{-4}$ CL=90%
Γ_{129}	$D_s^{*+} \rho^0$	<	4	$\times 10^{-4}$ CL=90%
Γ_{130}	$D_s^+ \omega$	<	4	$\times 10^{-4}$ CL=90%
Γ_{131}	$D_s^{*+} \omega$	<	6	$\times 10^{-4}$ CL=90%
Γ_{132}	$D_s^+ a_1(1260)^0$	<	1.8	$\times 10^{-3}$ CL=90%
Γ_{133}	$D_s^{*+} a_1(1260)^0$	<	1.3	$\times 10^{-3}$ CL=90%
Γ_{134}	$D_s^+ \phi$	<	1.9	$\times 10^{-6}$ CL=90%
Γ_{135}	$D_s^{*+} \phi$	<	1.2	$\times 10^{-5}$ CL=90%
Γ_{136}	$D_s^+ \bar{K}^0$	<	9	$\times 10^{-4}$ CL=90%
Γ_{137}	$D_s^{*+} \bar{K}^0$	<	9	$\times 10^{-4}$ CL=90%
Γ_{138}	$D_s^+ \bar{K}^*(892)^0$	<	4	$\times 10^{-4}$ CL=90%
Γ_{139}	$D_s^{*+} \bar{K}^*(892)^0$	<	4	$\times 10^{-4}$ CL=90%
Γ_{140}	$D_s^- \pi^+ K^+$	<	7	$\times 10^{-4}$ CL=90%
Γ_{141}	$D_s^{*-} \pi^+ K^+$	<	9.8	$\times 10^{-4}$ CL=90%
Γ_{142}	$D_s^- \pi^+ K^*(892)^+$	<	5	$\times 10^{-3}$ CL=90%
Γ_{143}	$D_s^{*-} \pi^+ K^*(892)^+$	<	7	$\times 10^{-3}$ CL=90%

Charmonium modes

Γ_{144}	$\eta_c K^+$	(9.1 \pm 1.3)	$\times 10^{-4}$
Γ_{145}	$\eta'_c K^+$	(3.4 \pm 1.8)	$\times 10^{-4}$
Γ_{146}	$J/\psi(1S) K^+$	(1.008 \pm 0.035)	$\times 10^{-3}$
Γ_{147}	$J/\psi(1S) K^+ \pi^+ \pi^-$	(1.07 \pm 0.19)	$\times 10^{-3}$ S=1.9
Γ_{148}	$h_c(1P) K^+ \times B(h_c(1P) \rightarrow J/\psi \pi^+ \pi^-)$	<	3.4	$\times 10^{-6}$ CL=90%
Γ_{149}	$X(3872) K^+$	<	3.2	$\times 10^{-4}$ CL=90%
Γ_{150}	$X(3872) K^+ \times B(X \rightarrow J/\psi \pi^+ \pi^-)$	(1.14 \pm 0.20)	$\times 10^{-5}$
Γ_{151}	$X(3872) K^+ \times B(X(3872) \rightarrow D^0 \bar{D}^0)$	<	6.0	$\times 10^{-5}$ CL=90%
Γ_{152}	$X(3872) K^+ \times B(X(3872) \rightarrow D^+ D^-)$	<	4.0	$\times 10^{-5}$ CL=90%
Γ_{153}	$X(3872) K^+ \times B(X(3872) \rightarrow D^0 \bar{D}^0 \pi^0)$	<	6.0	$\times 10^{-5}$ CL=90%
Γ_{154}	$X(3872) K^+ \times B(X(3872) \rightarrow J/\psi(1S) \eta)$	<	7.7	$\times 10^{-6}$ CL=90%
Γ_{155}	$X(3872)^+ K^0 \times B(X(3872)^+ \rightarrow J/\psi(1S) \pi^+ \pi^0)$	[e] <	2.2	$\times 10^{-5}$ CL=90%
Γ_{156}	$Y(4260)^0 K^+ \times B(Y^0 \rightarrow J/\psi \pi^+ \pi^-)$	<	2.9	$\times 10^{-5}$ CL=95%
Γ_{157}	$J/\psi(1S) K^*(892)^+$	(1.41 \pm 0.08)	$\times 10^{-3}$
Γ_{158}	$J/\psi(1S) K(1270)^+$	(1.8 \pm 0.5)	$\times 10^{-3}$
Γ_{159}	$J/\psi(1S) K(1400)^+$	<	5	$\times 10^{-4}$ CL=90%
Γ_{160}	$J/\psi(1S) \eta K^+$	(1.08 \pm 0.33)	$\times 10^{-4}$

Γ_{161}	$J/\psi(1S)\phi K^+$	$(5.2 \pm 1.7) \times 10^{-5}$	S=1.2
Γ_{162}	$J/\psi(1S)\pi^+$	$(4.9 \pm 0.6) \times 10^{-5}$	S=1.5
Γ_{163}	$J/\psi(1S)\rho^+$	$< 7.7 \times 10^{-4}$	CL=90%
Γ_{164}	$J/\psi(1S)a_1(1260)^+$	$< 1.2 \times 10^{-3}$	CL=90%
Γ_{165}	$J/\psi(1S)p\bar{\Lambda}$	$(1.18 \pm 0.31) \times 10^{-5}$	
Γ_{166}	$J/\psi(1S)\bar{\Sigma}^0 p$	$< 1.1 \times 10^{-5}$	CL=90%
Γ_{167}	$J/\psi(1S)D^+$	$< 1.2 \times 10^{-4}$	CL=90%
Γ_{168}	$J/\psi(1S)\bar{D}^0\pi^+$	$< 2.5 \times 10^{-5}$	CL=90%
Γ_{169}	$\psi(2S)K^+$	$(6.48 \pm 0.35) \times 10^{-4}$	
Γ_{170}	$\psi(2S)K^*(892)^+$	$(6.7 \pm 1.4) \times 10^{-4}$	S=1.3
Γ_{171}	$\psi(2S)K^+\pi^+\pi^-$	$(1.9 \pm 1.2) \times 10^{-3}$	
Γ_{172}	$\psi(3770)K^+$	$(4.9 \pm 1.3) \times 10^{-4}$	
Γ_{173}	$\psi(3770)K^+ \times B(\psi(3770) \rightarrow D^0\bar{D}^0)$	$(3.4 \pm 0.9) \times 10^{-4}$	
Γ_{174}	$\psi(3770)K^+ \times B(\psi(3770) \rightarrow D^+D^-K^+)$	$(1.4 \pm 0.8) \times 10^{-4}$	
Γ_{175}	$\chi_{c0}\pi^+ \times B(\chi_{c0} \rightarrow \pi^+\pi^-)$	$< 3 \times 10^{-7}$	CL=90%
Γ_{176}	$\chi_{c0}(1P)K^+$	$(1.6 \pm 0.5) \times 10^{-4}$	
Γ_{177}	$\chi_{c0}K^*(892)^+$	$< 2.86 \times 10^{-3}$	CL=90%
Γ_{178}	$\chi_{c2}K^+$	$< 2.9 \times 10^{-5}$	CL=90%
Γ_{179}	$\chi_{c2}K^*(892)^+$	$< 1.2 \times 10^{-5}$	CL=90%
Γ_{180}	$\chi_{c1}(1P)K^+$	$(5.3 \pm 0.7) \times 10^{-4}$	S=1.7
Γ_{181}	$\chi_{c1}(1P)K^*(892)^+$	$(3.6 \pm 0.9) \times 10^{-4}$	

K or K^* modes

Γ_{182}	$K^0\pi^+$	$(2.41 \pm 0.17) \times 10^{-5}$	S=1.4
Γ_{183}	$K^+\pi^0$	$(1.21 \pm 0.08) \times 10^{-5}$	
Γ_{184}	$\eta'K^+$	$(7.05 \pm 0.35) \times 10^{-5}$	
Γ_{185}	$\eta'K^*(892)^+$	$< 1.4 \times 10^{-5}$	CL=90%
Γ_{186}	ηK^+	$(2.6 \pm 0.6) \times 10^{-6}$	S=1.3
Γ_{187}	$\eta K^*(892)^+$	$(2.6 \pm 0.4) \times 10^{-5}$	
Γ_{188}	ωK^+	$(5.1 \pm 0.7) \times 10^{-6}$	
Γ_{189}	$\omega K^*(892)^+$	$< 7.4 \times 10^{-6}$	CL=90%
Γ_{190}	$a_0^+K^0$	$< 3.9 \times 10^{-6}$	CL=90%
Γ_{191}	$a_0^0K^+$	$< 2.5 \times 10^{-6}$	CL=90%
Γ_{192}	$K^*(892)^0\pi^+$	$(1.16 \pm 0.19) \times 10^{-5}$	S=1.8
Γ_{193}	$K^*(892)^+\pi^0$	$(6.9 \pm 2.4) \times 10^{-6}$	
Γ_{194}	$K^+\pi^-\pi^+$	$(5.6 \pm 0.9) \times 10^{-5}$	S=2.6
Γ_{195}	$K^+\pi^-\pi^+$ nonresonant	$(3.1 \pm 1.0) \times 10^{-6}$	
Γ_{196}	$K^+f_0(980) \times B(f_0 \rightarrow \pi^+\pi^-)$	$(8.9 \pm 1.0) \times 10^{-6}$	
Γ_{197}	$f_2(1270)^0K^+$	$< 2.3 \times 10^{-6}$	CL=90%
Γ_{198}	$f_0^*(1370)^0K^+ \times B(f_0^*(1370)^0 \rightarrow \pi^+\pi^-)$	$< 1.07 \times 10^{-5}$	CL=90%

Γ_{199}	$\rho^0(1450) K^+ \times B(\rho^0(1450) \rightarrow \pi^+ \pi^-)$	< 1.17	$\times 10^{-5}$ CL=90%
Γ_{200}	$f_0(1500) K^+ \times B(f_0(1500) \rightarrow \pi^+ \pi^-)$	< 4.4	$\times 10^{-6}$ CL=90%
Γ_{201}	$f'_2(1525) K^+ \times B(f'_2(1525) \rightarrow \pi^+ \pi^-)$	< 3.4	$\times 10^{-6}$ CL=90%
Γ_{202}	$K^+ \rho^0$	(5.0 \pm 0.7)	$\times 10^{-6}$
Γ_{203}	$K_0^*(1430)^0 \pi^+$	(3.8 \pm 0.5)	$\times 10^{-5}$
Γ_{204}	$K_2^*(1430)^0 \pi^+$	< 6.9	$\times 10^{-6}$ CL=90%
Γ_{205}	$K^*(1410)^0 \pi^+$	< 4.5	$\times 10^{-5}$ CL=90%
Γ_{206}	$K^*(1680)^0 \pi^+$	< 1.2	$\times 10^{-5}$ CL=90%
Γ_{207}	$K^- \pi^+ \pi^+$	< 1.8	$\times 10^{-6}$ CL=90%
Γ_{208}	$K^- \pi^+ \pi^+$ nonresonant	< 5.6	$\times 10^{-5}$ CL=90%
Γ_{209}	$K_1(1400)^0 \pi^+$	< 2.6	$\times 10^{-3}$ CL=90%
Γ_{210}	$K^0 \pi^+ \pi^0$	< 6.6	$\times 10^{-5}$ CL=90%
Γ_{211}	$K^0 \rho^+$	< 4.8	$\times 10^{-5}$ CL=90%
Γ_{212}	$K^*(892)^+ \pi^+ \pi^-$	< 1.1	$\times 10^{-3}$ CL=90%
Γ_{213}	$K^*(892)^+ \rho^0$	(1.1 \pm 0.4)	$\times 10^{-5}$
Γ_{214}	$K^*(892)^0 \rho^+$	(8.9 \pm 2.1)	$\times 10^{-6}$
Γ_{215}	$K^*(892)^+ K^*(892)^0$	< 7.1	$\times 10^{-5}$ CL=90%
Γ_{216}	$K_1(1400)^+ \rho^0$	< 7.8	$\times 10^{-4}$ CL=90%
Γ_{217}	$K_2^*(1430)^+ \rho^0$	< 1.5	$\times 10^{-3}$ CL=90%
Γ_{218}	$K^+ \overline{K}^0$	(1.20 \pm 0.32)	$\times 10^{-6}$
Γ_{219}	$\overline{K}^0 K^+ \pi^0$	< 2.4	$\times 10^{-5}$ CL=90%
Γ_{220}	$K^+ K_S^0 K_S^0$	(1.15 \pm 0.13)	$\times 10^{-5}$
Γ_{221}	$K_S^0 K_S^0 \pi^+$	< 3.2	$\times 10^{-6}$ CL=90%
Γ_{222}	$K^+ K^- \pi^+$	< 6.3	$\times 10^{-6}$ CL=90%
Γ_{223}	$K^+ K^- \pi^+$ nonresonant	< 7.5	$\times 10^{-5}$ CL=90%
Γ_{224}	$K^+ K^+ \pi^-$	< 1.3	$\times 10^{-6}$ CL=90%
Γ_{225}	$K^+ K^+ \pi^-$ nonresonant	< 8.79	$\times 10^{-5}$ CL=90%
Γ_{226}	$K^+ K^*(892)^0$	< 5.3	$\times 10^{-6}$ CL=90%
Γ_{227}	$K^+ f_J(2220)$		
Γ_{228}	$K^+ K^- K^+$	(3.01 \pm 0.19)	$\times 10^{-5}$
Γ_{229}	$K^+ \phi$	(9.0 \pm 0.8)	$\times 10^{-6}$ S=1.3
Γ_{230}	$f_0(980) K^+ \times B(f_0(980) \rightarrow K^+ K^-)$	< 2.9	$\times 10^{-6}$ CL=90%
Γ_{231}	$a_2(1320) K^+ \times B(a_2(1320) \rightarrow K^+ K^-)$	< 1.1	$\times 10^{-6}$ CL=90%
Γ_{232}	$f'_2(1525) K^+ \times B(f'_2(1525) \rightarrow K^+ K^-)$	< 4.9	$\times 10^{-6}$ CL=90%
Γ_{233}	$\phi(1680) K^+ \times B(\phi(1680) \rightarrow K^+ K^-)$	< 8	$\times 10^{-7}$ CL=90%

Γ_{234}	$K^+ K^- K^+$ nonresonant	$(-2.40 \pm 0.30) \times 10^{-5}$
Γ_{235}	$K^*(892)^+ K^+ K^-$	$< 1.6 \times 10^{-3} \text{ CL}=90\%$
Γ_{236}	$K^*(892)^+ \phi$	$(9.6 \pm 3.0) \times 10^{-6} \text{ S}=1.9$
Γ_{237}	$K_1(1400)^+ \phi$	$< 1.1 \times 10^{-3} \text{ CL}=90\%$
Γ_{238}	$K_2^*(1430)^+ \phi$	$< 3.4 \times 10^{-3} \text{ CL}=90\%$
Γ_{239}	$K^+ \phi \phi$	$(2.6 \pm 1.1) \times 10^{-6}$
Γ_{240}	$K^*(892)^+ \gamma$	$(4.03 \pm 0.26) \times 10^{-5}$
Γ_{241}	$K_1(1270)^+ \gamma$	$(4.3 \pm 1.3) \times 10^{-5}$
Γ_{242}	$\eta K^+ \gamma$	$(8.4 \pm 1.8) \times 10^{-6}$
Γ_{243}	$\phi K^+ \gamma$	$(3.4 \pm 1.0) \times 10^{-6}$
Γ_{244}	$K^+ \pi^- \pi^+ \gamma$	$(2.50 \pm 0.28) \times 10^{-5}$
Γ_{245}	$K^*(892)^0 \pi^+ \gamma$	$(2.0 \pm 0.7) \times 10^{-5}$
Γ_{246}	$K^+ \rho^0 \gamma$	$< 2.0 \times 10^{-5} \text{ CL}=90\%$
Γ_{247}	$K^+ \pi^- \pi^+ \gamma$ nonresonant	$< 9.2 \times 10^{-6} \text{ CL}=90\%$
Γ_{248}	$K_1(1400)^+ \gamma$	$< 1.5 \times 10^{-5}$
Γ_{249}	$K_2^*(1430)^+ \gamma$	$(1.4 \pm 0.4) \times 10^{-5}$
Γ_{250}	$K^*(1680)^+ \gamma$	$< 1.9 \times 10^{-3} \text{ CL}=90\%$
Γ_{251}	$K_3^*(1780)^+ \gamma$	$< 3.9 \times 10^{-5} \text{ CL}=90\%$
Γ_{252}	$K_4^*(2045)^+ \gamma$	$< 9.9 \times 10^{-3} \text{ CL}=90\%$

Light unflavored meson modes

Γ_{253}	$\rho^+ \gamma$	$< 1.8 \times 10^{-6} \text{ CL}=90\%$
Γ_{254}	$\pi^+ \pi^0$	$(5.5 \pm 0.6) \times 10^{-6}$
Γ_{255}	$\pi^+ \pi^+ \pi^-$	$(1.62 \pm 0.15) \times 10^{-5}$
Γ_{256}	$\rho^0 \pi^+$	$(8.7 \pm 1.1) \times 10^{-6}$
Γ_{257}	$\pi^+ f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-)$	$< 3.0 \times 10^{-6} \text{ CL}=90\%$
Γ_{258}	$\pi^+ f_2(1270)$	$(8.2 \pm 2.5) \times 10^{-6}$
Γ_{259}	$\rho(1450)^0 \pi^+$	$< 2.3 \times 10^{-6} \text{ CL}=90\%$
Γ_{260}	$f_0(1370) \pi^+ \times B(f_0(1370) \rightarrow \pi^+ \pi^-)$	$< 3.0 \times 10^{-6} \text{ CL}=90\%$
Γ_{261}	$f_0(600) \pi^+ \times B(f_0(600) \rightarrow \pi^+ \pi^-)$	$< 4.1 \times 10^{-6} \text{ CL}=90\%$
Γ_{262}	$\pi^+ \pi^- \pi^+ \text{ nonresonant}$	$< 4.6 \times 10^{-6} \text{ CL}=90\%$
Γ_{263}	$\pi^+ \pi^0 \pi^0$	$< 8.9 \times 10^{-4} \text{ CL}=90\%$
Γ_{264}	$\rho^+ \pi^0$	$(1.20 \pm 0.19) \times 10^{-5}$
Γ_{265}	$\pi^+ \pi^- \pi^+ \pi^0$	$< 4.0 \times 10^{-3} \text{ CL}=90\%$
Γ_{266}	$\rho^+ \rho^0$	$(2.6 \pm 0.6) \times 10^{-5}$
Γ_{267}	$a_1(1260)^+ \pi^0$	$< 1.7 \times 10^{-3} \text{ CL}=90\%$
Γ_{268}	$a_1(1260)^0 \pi^+$	$< 9.0 \times 10^{-4} \text{ CL}=90\%$
Γ_{269}	$\omega \pi^+$	$(5.9 \pm 1.0) \times 10^{-6} \text{ S}=1.2$
Γ_{270}	$\omega \rho^+$	$(1.3 \pm 0.4) \times 10^{-5}$

Γ_{271}	$\eta\pi^+$	$(4.9 \pm 0.5) \times 10^{-6}$
Γ_{272}	$\eta'\pi^+$	$(4.0 \pm 0.9) \times 10^{-6}$
Γ_{273}	$\eta'\rho^+$	$< 2.2 \times 10^{-5} \text{ CL}=90\%$
Γ_{274}	$\eta\rho^+$	$(8.4 \pm 2.2) \times 10^{-6}$
Γ_{275}	$\phi\pi^+$	$< 4.1 \times 10^{-7} \text{ CL}=90\%$
Γ_{276}	$\phi\rho^+$	$< 1.6 \times 10^{-5}$
Γ_{277}	$a_0^0\pi^+$	$< 5.8 \times 10^{-6} \text{ CL}=90\%$
Γ_{278}	$\pi^+\pi^+\pi^-\pi^-$	$< 8.6 \times 10^{-4} \text{ CL}=90\%$
Γ_{279}	$\rho^0 a_1(1260)^+$	$< 6.2 \times 10^{-4} \text{ CL}=90\%$
Γ_{280}	$\rho^0 a_2(1320)^+$	$< 7.2 \times 10^{-4} \text{ CL}=90\%$
Γ_{281}	$\pi^+\pi^+\pi^-\pi^-\pi^0$	$< 6.3 \times 10^{-3} \text{ CL}=90\%$
Γ_{282}	$a_1(1260)^+ a_1(1260)^0$	$< 1.3 \% \text{ CL}=90\%$

Charged particle (h^\pm) modes

	$h^\pm = K^\pm \text{ or } \pi^\pm$	
Γ_{283}	$h^+\pi^0$	$(1.6 \pm 0.7) \times 10^{-5}$
Γ_{284}	ωh^+	$(1.38 \pm 0.27) \times 10^{-5}$
Γ_{285}	$h^+ X^0 \text{ (Familon)}$	$< 4.9 \times 10^{-5} \text{ CL}=90\%$

Baryon modes

Γ_{286}	$p\bar{p}\pi^+$	$(3.1 \pm 0.8) \times 10^{-6}$
Γ_{287}	$p\bar{p}\pi^+ \text{ nonresonant}$	$< 5.3 \times 10^{-5} \text{ CL}=90\%$
Γ_{288}	$p\bar{p}\pi^+\pi^+\pi^-$	$< 5.2 \times 10^{-4} \text{ CL}=90\%$
Γ_{289}	$p\bar{p}K^+$	$(5.6 \pm 1.0) \times 10^{-6} \text{ S}=2.4$
Γ_{290}	$\Theta(1710)^{++}\bar{p} \times B(\Theta(1710)^{++} \rightarrow pK^+)$	$[f] < 9.1 \times 10^{-8} \text{ CL}=90\%$
Γ_{291}	$f_J(2220)K^+ \times B(f_J(2220) \rightarrow p\bar{p})$	$[f] < 4.1 \times 10^{-7} \text{ CL}=90\%$
Γ_{292}	$p\bar{\Lambda}(1520)$	$< 1.5 \times 10^{-6} \text{ CL}=90\%$
Γ_{293}	$p\bar{p}K^+ \text{ nonresonant}$	$< 8.9 \times 10^{-5} \text{ CL}=90\%$
Γ_{294}	$p\bar{p}K^*(892)^+$	$(1.03 \pm 0.38) \times 10^{-5}$
Γ_{295}	$p\bar{\Lambda}$	$< 4.9 \times 10^{-7} \text{ CL}=90\%$
Γ_{296}	$p\bar{\Lambda}\gamma$	$(2.2 \pm 0.6) \times 10^{-6}$
Γ_{297}	$p\bar{\Sigma}\gamma$	$< 4.6 \times 10^{-6} \text{ CL}=90\%$
Γ_{298}	$p\bar{\Lambda}\pi^+\pi^-$	$< 2.0 \times 10^{-4} \text{ CL}=90\%$
Γ_{299}	$\Lambda\bar{\Lambda}\pi^+$	$< 2.8 \times 10^{-6} \text{ CL}=90\%$
Γ_{300}	$\Lambda\bar{\Lambda}K^+$	$(2.9 \pm 1.0) \times 10^{-6}$
Γ_{301}	$\bar{\Delta}^0 p$	$< 3.8 \times 10^{-4} \text{ CL}=90\%$
Γ_{302}	$\Delta^{++}\bar{p}$	$< 1.5 \times 10^{-4} \text{ CL}=90\%$
Γ_{303}	$D^+ p\bar{p}$	$< 1.5 \times 10^{-5} \text{ CL}=90\%$
Γ_{304}	$D^*(2010)^+ p\bar{p}$	$< 1.5 \times 10^{-5} \text{ CL}=90\%$

Γ_{305}	$\bar{\Lambda}_c^- p \pi^+$	(2.1 \pm 0.7) $\times 10^{-4}$
Γ_{306}	$\bar{\Lambda}_c^- p \pi^+ \pi^0$	(1.8 \pm 0.6) $\times 10^{-3}$
Γ_{307}	$\bar{\Lambda}_c^- p \pi^+ \pi^+ \pi^-$	(2.3 \pm 0.7) $\times 10^{-3}$
Γ_{308}	$\bar{\Lambda}_c^- p \pi^+ \pi^+ \pi^- \pi^0$	< 1.34 % CL=90%
Γ_{309}	$\bar{\Sigma}_c(2455)^0 p$	< 8 $\times 10^{-5}$ CL=90%
Γ_{310}	$\bar{\Sigma}_c(2520)^0 p$	< 4.6 $\times 10^{-5}$ CL=90%
Γ_{311}	$\bar{\Sigma}_c(2455)^0 p \pi^0$	(4.4 \pm 1.8) $\times 10^{-4}$
Γ_{312}	$\bar{\Sigma}_c(2455)^0 p \pi^- \pi^+$	(4.4 \pm 1.7) $\times 10^{-4}$
Γ_{313}	$\bar{\Sigma}_c(2455)^{--} p \pi^+ \pi^+$	(2.8 \pm 1.2) $\times 10^{-4}$
Γ_{314}	$\bar{\Lambda}_c(2593)^- / \bar{\Lambda}_c(2625)^- p \pi^+$	< 1.9 $\times 10^{-4}$ CL=90%

**Lepton Family number (*LF*) or Lepton number (*L*) violating modes, or
 $\Delta B = 1$ weak neutral current (*B1*) modes**

Γ_{315}	$\pi^+ e^+ e^-$	<i>B1</i>	< 3.9	$\times 10^{-3}$ CL=90%
Γ_{316}	$\pi^+ \mu^+ \mu^-$	<i>B1</i>	< 9.1	$\times 10^{-3}$ CL=90%
Γ_{317}	$\pi^+ \nu \bar{\nu}$	<i>B1</i>	< 1.0	$\times 10^{-4}$ CL=90%
Γ_{318}	$K^+ e^+ e^-$	<i>B1</i>	(8.0 \pm 2.2) $\times 10^{-7}$	S=1.4
Γ_{319}	$K^+ \mu^+ \mu^-$	<i>B1</i>	(3.4 \pm 1.9) $\times 10^{-7}$	S=1.7
Γ_{320}	$K^+ \ell^+ \ell^-$	<i>B1</i>	[a] (5.3 \pm 1.1) $\times 10^{-7}$	
Γ_{321}	$K^+ \bar{\nu} \nu$	<i>B1</i>	< 5.2	$\times 10^{-5}$ CL=90%
Γ_{322}	$K^*(892)^+ e^+ e^-$	<i>B1</i>	< 4.6	$\times 10^{-6}$ CL=90%
Γ_{323}	$K^*(892)^+ \mu^+ \mu^-$	<i>B1</i>	< 2.2	$\times 10^{-6}$ CL=90%
Γ_{324}	$K^*(892)^+ \ell^+ \ell^-$	<i>B1</i>	[a] < 2.2	$\times 10^{-6}$ CL=90%
Γ_{325}	$\pi^+ e^+ \mu^-$	<i>LF</i>	< 6.4	$\times 10^{-3}$ CL=90%
Γ_{326}	$\pi^+ e^- \mu^+$	<i>LF</i>	< 6.4	$\times 10^{-3}$ CL=90%
Γ_{327}	$K^+ e^+ \mu^-$	<i>LF</i>	< 8	$\times 10^{-7}$ CL=90%
Γ_{328}	$K^+ e^- \mu^+$	<i>LF</i>	< 6.4	$\times 10^{-3}$ CL=90%
Γ_{329}	$K^*(892)^+ e^\pm \mu^\mp$	<i>LF</i>	< 7.9	$\times 10^{-6}$ CL=90%
Γ_{330}	$\pi^- e^+ e^+$	<i>L</i>	< 1.6	$\times 10^{-6}$ CL=90%
Γ_{331}	$\pi^- \mu^+ \mu^+$	<i>L</i>	< 1.4	$\times 10^{-6}$ CL=90%
Γ_{332}	$\pi^- e^+ \mu^+$	<i>L</i>	< 1.3	$\times 10^{-6}$ CL=90%
Γ_{333}	$\rho^- e^+ e^+$	<i>L</i>	< 2.6	$\times 10^{-6}$ CL=90%
Γ_{334}	$\rho^- \mu^+ \mu^+$	<i>L</i>	< 5.0	$\times 10^{-6}$ CL=90%
Γ_{335}	$\rho^- e^+ \mu^+$	<i>L</i>	< 3.3	$\times 10^{-6}$ CL=90%
Γ_{336}	$K^- e^+ e^+$	<i>L</i>	< 1.0	$\times 10^{-6}$ CL=90%
Γ_{337}	$K^- \mu^+ \mu^+$	<i>L</i>	< 1.8	$\times 10^{-6}$ CL=90%
Γ_{338}	$K^- e^+ \mu^+$	<i>L</i>	< 2.0	$\times 10^{-6}$ CL=90%
Γ_{339}	$K^*(892)^- e^+ e^+$	<i>L</i>	< 2.8	$\times 10^{-6}$ CL=90%
Γ_{340}	$K^*(892)^- \mu^+ \mu^-$	<i>L</i>	< 8.3	$\times 10^{-6}$ CL=90%
Γ_{341}	$K^*(892)^- e^+ \mu^+$	<i>L</i>	< 4.4	$\times 10^{-6}$ CL=90%

- [a] An ℓ indicates an e or a μ mode, not a sum over these modes.
- [b] An $CP(\pm 1)$ indicates the $CP=+1$ and $CP=-1$ eigenstates of the D^0 - \overline{D}^0 system.
- [c] D denotes D^0 or \overline{D}^0 .
- [d] D_{CP+}^{*0} decays into $D^0\pi^0$ with the D^0 reconstructed in CP -even eigenstates K^+K^- and $\pi^+\pi^-$.
- [e] $X(3872)^+$ is a hypothetical charged partner of the $X(3872)$.
- [f] $\Theta(1710)^{++}$ is a possible narrow pentaquark state and $G(2220)$ is a possible glueball resonance.

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 11 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 7.2$ for 9 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|c} x_{162} & 24 \\ \hline & x_{146} \end{array}$$

B^+ BRANCHING RATIOS

$\Gamma(\ell^+ \nu_\ell \text{anything}) / \Gamma_{\text{total}}$	Γ_1 / Γ		
<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>

10.9 ± 0.4 OUR AVERAGE

$11.15 \pm 0.26 \pm 0.41$	14 OKABE	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$
$10.25 \pm 0.57 \pm 0.65$	15 ARTUSO	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.1 \pm 1.8 \pm 1.5$	ATHANAS	94 CLE2	Sup. by ARTUSO 97
------------------------	---------	---------	-------------------

14 The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame, and their ratio of $B(B^+ \rightarrow e^+ \nu_e X) / B(B^0 \rightarrow e^+ \nu_e X) = 1.08 \pm 0.05 \pm 0.02$.

15 ARTUSO 97 uses partial reconstruction of $B \rightarrow D^* \ell \nu_\ell$ and inclusive semileptonic branching ratio from BARISH 96B ($0.1049 \pm 0.0017 \pm 0.0043$).

$\Gamma(D^0 \ell^+ \nu_\ell) / \Gamma_{\text{total}}$	Γ_2 / Γ
---	---------------------

$\ell = e$ or μ , not sum over e and μ modes.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.0215±0.0022 OUR AVERAGE

$0.0221 \pm 0.0013 \pm 0.0019$	16 BARTELT	99 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$0.016 \pm 0.006 \pm 0.003$	17 FULTON	91 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.0194 \pm 0.0015 \pm 0.0034$	18 ATHANAS	97 CLE2	Repl. by BARTELT 99
--------------------------------	------------	---------	---------------------

¹⁶ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹⁷ FULTON 91 assumes equal production of $B^0\bar{B}^0$ and B^+B^- at the $\Upsilon(4S)$.

¹⁸ ATHANAS 97 uses missing energy and missing momentum to reconstruct neutrino.

$\Gamma(\overline{D}^*(2007)^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_3/Γ

$\ell = e$ or μ , not sum over e and μ modes.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.065 ± 0.005 OUR AVERAGE				
0.0650 ± 0.0020 ± 0.0043		19 ADAM 03	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.066 ± 0.016 ± 0.015		20 ALBRECHT 92C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0650 ± 0.0020 ± 0.0043		21 BRIERE 02	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.0513 ± 0.0054 ± 0.0064	302	22 BARISH 95	CLE2	Repl. by ADAM 03
seen	398	23 SANGHERA 93	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.041 ± 0.008 +0.008 -0.009		24 FULTON 91	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.070 ± 0.018 ± 0.014		25 ANTREASYAN 90B	CBAL	$e^+e^- \rightarrow \Upsilon(4S)$

¹⁹ Simultaneous measurements of both $B^0 \rightarrow D^*(2010)^- \ell \nu$ and $B^+ \rightarrow \overline{D}(2007)^0 \ell \nu$.

²⁰ ALBRECHT 92C reports $0.058 \pm 0.014 \pm 0.013$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^-\pi^+)$. Assumes equal production of $B^0\bar{B}^0$ and B^+B^- at the $\Upsilon(4S)$.

²¹ The results are based on the same analysis and data sample reported in ADAM 03.

²² BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*0} \rightarrow D^0\pi^0) = (63.6 \pm 2.3 \pm 3.3)\%$.

²³ Combining $\overline{D}^{*0}\ell^+\nu_\ell$ and $\overline{D}^{*-}\ell^+\nu_\ell$ SANGHERA 93 test $V-A$ structure and fit the decay angular distributions to obtain $A_{FB} = 3/4 * (\Gamma^- - \Gamma^+)/\Gamma = 0.14 \pm 0.06 \pm 0.03$. Assuming a value of V_{cb} , they measure V , A_1 , and A_2 , the three form factors for the $D^*\ell\nu_\ell$ decay, where results are slightly dependent on model assumptions.

²⁴ Assumes equal production of $B^0\bar{B}^0$ and B^+B^- at the $\Upsilon(4S)$. Uncorrected for D and D^* branching ratio assumptions.

²⁵ ANTREASYAN 90B is average over B and $\overline{D}^*(2010)$ charge states.

$\Gamma(\overline{D}_1(2420)^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0056 ± 0.0013 ± 0.0009	26 ANASTASSOV 98	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
²⁶ ANASTASSOV 98 result is derived from the measurement of $B(B^+ \rightarrow \overline{D}_1^0 \ell^+\nu_\ell) \times B(\overline{D}_1^0 \rightarrow D^{*+}\pi^-) = (0.373 \pm 0.085 \pm 0.052 \pm 0.024)\%$ by assuming $B(\overline{D}_1^0 \rightarrow D^{*+}\pi^-) = 67\%$, where the third error includes theoretical uncertainties.			

$\Gamma(\overline{D}_2^*(2460)^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8 × 10⁻³	90	27 ANASTASSOV 98	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
²⁷ ANASTASSOV 98 result is derived from the measurement of $B(B^+ \rightarrow \overline{D}_2^{*0}\ell^+\nu_\ell) \times B(\overline{D}_2^{*0} \rightarrow D^{*+}\pi^-) < 0.16\%$ at 90% CL by assuming $B(\overline{D}_2^{*0} \rightarrow D^{*+}\pi^-) = 20\%$.				

$\Gamma(D^- \pi^+ \ell^+ \nu_\ell)/\Gamma_{\text{total}}$	Γ_6/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
5.3±0.9±0.5	28 LIVENTSEV 05 BELL	e ⁺ e ⁻ → $\gamma(4S)$	

²⁸ LIVENTSEV 05 reports $[B(B^+ \rightarrow D^- \pi^+ \ell^+ \nu_\ell) / B(B^0 \rightarrow D^- \ell^+ \nu_\ell)] = 0.25 \pm 0.03 \pm 0.03$. We multiply by our best value $B(B^0 \rightarrow D^- \ell^+ \nu_\ell) = (2.12 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^{*-} \pi^+ \ell^+ \nu_\ell)/\Gamma_{\text{total}}$	Γ_7/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
6.4±1.5±0.2	29,30 LIVENTSEV 05 BELL	e ⁺ e ⁻ → $\gamma(4S)$	

²⁹ Excludes D^{*+} contribution to $D\pi$ modes.
³⁰ LIVENTSEV 05 reports $[B(B^+ \rightarrow D^{*-} \pi^+ \ell^+ \nu_\ell) / B(B^0 \rightarrow D^{*(2010)}^- \ell^+ \nu_\ell)] = 0.12 \pm 0.02 \pm 0.02$. We multiply by our best value $B(B^0 \rightarrow D^{*(2010)}^- \ell^+ \nu_\ell) = (5.35 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$	Γ_8/Γ		
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.74±0.05±0.10	31 AUBERT,B 050 BABR	e ⁺ e ⁻ → $\gamma(4S)$	

³¹ B^+ and B^0 decays combined assuming isospin symmetry. Systematic errors include both experimental and form-factor uncertainties.

$\Gamma(\pi^0 e^+ \nu_e)/\Gamma_{\text{total}}$	Γ_9/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.9±0.2±0.2		32 ALEXANDER 96T CLE2	e ⁺ e ⁻ → $\gamma(4S)$	
<22	90	ANTREASYAN 90B CBAL	e ⁺ e ⁻ → $\gamma(4S)$	
32	Derived based in the reported B^0 result by assuming isospin symmetry: $\Gamma(B^0 \rightarrow \pi^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \pi^0 \ell^+ \nu)$.			

$\Gamma(\eta \ell^+ \nu_\ell)/\Gamma_{\text{total}}$	Γ_{10}/Γ		
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.84±0.31±0.18	33 ATHAR 03 CLE2	e ⁺ e ⁻ → $\gamma(4S)$	

³³ ATHAR 03 reports systematic errors 0.16 ± 0.09 , which are experimental systematic and systematic due to model dependence. We combine these in quadrature.

$\Gamma(\omega \ell^+ \nu_\ell)/\Gamma_{\text{total}}$	Γ_{11}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
1.3±0.4±0.4		34 SCHWANDA 04 BELL	e ⁺ e ⁻ → $\gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1	90	35 BEAN	93B CLE2	e ⁺ e ⁻ → $\gamma(4S)$
------	----	---------	----------	--

³⁴ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.
³⁵ BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\rho^0 \ell^+ \nu_\ell)$ and $\Gamma(\rho^- \ell^+ \nu_\ell)$ with this result, they obtain a limit $<(1.6-2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow \omega \ell^+ \nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.8-0.13$ at 90% CL is derived as well.

$\Gamma(\omega\mu^+\nu_\mu)/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
--------------	--------------------	-------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 36 ALBRECHT 91C ARG

36 In ALBRECHT 91C, one event is fully reconstructed providing evidence for the $b \rightarrow u$ transition.

 $\Gamma(\rho^0\ell^+\nu_\ell)/\Gamma_{\text{total}}$ Γ_{13}/Γ

$\ell = e$ or μ , not sum over e and μ modes.

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

1.24±0.23 OUR AVERAGE

37 AUBERT,B 050 BABR $e^+e^- \rightarrow \gamma(4S)$

38 BEHRENS 00 CLE2 $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

38 BEHRENS 00 CLE2 $e^+e^- \rightarrow \gamma(4S)$

38 ALEXANDER 96T CLE2 $e^+e^- \rightarrow \gamma(4S)$

<2.1 90 39 BEAN 93B CLE2 $e^+e^- \rightarrow \gamma(4S)$

37 B^+ and B^0 decays combined assuming isospin symmetry. Systematic errors include both experimental and form-factor uncertainties.

38 Derived based in the reported B^0 result by assuming isospin symmetry: $\Gamma(B^0 \rightarrow \rho^-\ell^+\nu) = 2\Gamma(B^+ \rightarrow \rho^0\ell^+\nu) \approx 2\Gamma(B^+ \rightarrow \omega\ell^+\nu)$.

39 BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\omega\ell^+\nu_\ell)$ and $\Gamma(\rho^-\ell^+\nu_\ell)$ with this result, they obtain a limit $<(1.6\text{--}2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow \rho^0\ell^+\nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.8\text{--}0.13$ at 90% CL is derived as well.

 $\Gamma(p\bar{p}e^+\nu_e)/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.2 × 10⁻³	90	40 ADAM 03B CLE2	$e^+e^- \rightarrow \gamma(4S)$	

40 Based on phase-space model; if V-A model is used, the 90% CL upper limit becomes $< 1.2 \times 10^{-3}$.

 $\Gamma(e^+\nu_e)/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.5 × 10⁻⁵	90	ARTUSO 95	CLE2	$e^+e^- \rightarrow \gamma(4S)$

 $\Gamma(\mu^+\nu_\mu)/\Gamma_{\text{total}}$ Γ_{16}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.6 × 10⁻⁶	90	AUBERT 040 BABR	$e^+e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1 × 10⁻⁵ 90 ARTUSO 95 CLE2 $e^+e^- \rightarrow \gamma(4S)$

$\Gamma(\tau^+ \nu_\tau)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.6 \times 10^{-4}$	90	41 AUBERT	06K BABR	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<4.2 \times 10^{-4}$	90	41 AUBERT,B	05B BABR	Repl. by AUBERT 06K
$<8.3 \times 10^{-4}$	90	42 BARATE	01E ALEP	$e^+ e^- \rightarrow Z$
$<8.4 \times 10^{-4}$	90	41 BROWDER	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$<5.7 \times 10^{-4}$	90	43 ACCIARRI	97F L3	$e^+ e^- \rightarrow Z$
$<1.04 \times 10^{-2}$	90	44 ALBRECHT	95D ARG	$e^+ e^- \rightarrow \gamma(4S)$
$<2.2 \times 10^{-3}$	90	ARTUSO	95 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$<1.8 \times 10^{-3}$	90	45 BUSKULIC	95 ALEP	$e^+ e^- \rightarrow Z$

41 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.42 The energy-flow and b -tagging algorithms were used.43 ACCIARRI 97F uses missing-energy technique and $f(b \rightarrow B^-) = (38.2 \pm 2.5)\%$.44 ALBRECHT 95D use full reconstruction of one B decay as tag.45 BUSKULIC 95 uses same missing-energy technique as in $\bar{b} \rightarrow \tau^+ \nu_\tau X$, but analysis is restricted to endpoint region of missing-energy distribution. $\Gamma(e^+ \nu_e \gamma)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-4}$	90	46 BROWDER	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

46 BROWDER 97 uses the hermiticity of the CLEO II detector to reconstruct the neutrino energy and momentum.

 $\Gamma(\mu^+ \nu_\mu \gamma)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-5}$	90	47 BROWDER	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

47 BROWDER 97 uses the hermiticity of the CLEO II detector to reconstruct the neutrino energy and momentum.

 $\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.098 ± 0.009 ± 0.006	48 AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$

48 Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties. $\Gamma(\bar{D}^0 X)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.793 ± 0.025 ± 0.045	49 AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$

49 Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties. $\Gamma(D^0 X)/[\Gamma(D^0 X) + \Gamma(\bar{D}^0 X)]$ $\Gamma_{20}/(\Gamma_{20} + \Gamma_{21})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.110 ± 0.010 ± 0.003	AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(D^+ X)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{22}/Γ
0.038±0.009±0.005	50 AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$	

50 Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(D^- X)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{23}/Γ
0.098±0.012±0.014	51 AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$	

51 Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(D^+ X)/[\Gamma(D^+ X) + \Gamma(D^- X)]$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{22}/(\Gamma_{22}+\Gamma_{23})$
0.278±0.052±0.009	AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$	

 $\Gamma(D_s^+ X)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{24}/Γ
0.143±0.016^{+0.051}_{-0.034}	52 AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$	

52 Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(D_s^- X)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{25}/Γ
<0.022	90	53 AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$	

53 Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(D_s^+ X)/[\Gamma(D_s^+ X) + \Gamma(D_s^- X)]$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{24}/(\Gamma_{24}+\Gamma_{25})$
0.966±0.039±0.012	AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$	

 $\Gamma(D_s^- X)/[\Gamma(D_s^+ X) + \Gamma(D_s^- X)]$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{25}/(\Gamma_{24}+\Gamma_{25})$
<0.126	90	AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$	

 $\Gamma(\Lambda_c^+ X)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{26}/Γ
0.029±0.008^{+0.011}_{-0.007}	54 AUBERT,BE	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$	

54 Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$\Gamma(\bar{\Lambda}_c^- X)/\Gamma_{\text{total}}$ VALUE **$0.035 \pm 0.008^{+0.013}_{-0.009}$** DOCUMENT ID55 AUBERT,BE 04B BABR $e^+ e^- \rightarrow \gamma(4S)$ Γ_{27}/Γ

55 Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(\Lambda_c^+ X)/[\Gamma(\Lambda_c^+ X) + \Gamma(\bar{\Lambda}_c^- X)]$ VALUE **$0.452 \pm 0.090 \pm 0.003$** DOCUMENT IDAUBERT,BE 04B BABR $e^+ e^- \rightarrow \gamma(4S)$ $\Gamma_{26}/(\Gamma_{26}+\Gamma_{27})$ $\Gamma(\bar{c}X)/\Gamma_{\text{total}}$ VALUE **$0.983 \pm 0.030^{+0.054}_{-0.051}$** DOCUMENT ID56 AUBERT,BE 04B BABR $e^+ e^- \rightarrow \gamma(4S)$ Γ_{28}/Γ $\Gamma(cX)/\Gamma_{\text{total}}$ VALUE **$0.330 \pm 0.022^{+0.055}_{-0.037}$** DOCUMENT ID57 AUBERT,BE 04B BABR $e^+ e^- \rightarrow \gamma(4S)$ Γ_{29}/Γ

57 Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(\bar{c}cX)/\Gamma_{\text{total}}$ VALUE **$1.313 \pm 0.037^{+0.088}_{-0.075}$** DOCUMENT ID58 AUBERT,BE 04B BABR $e^+ e^- \rightarrow \gamma(4S)$ Γ_{30}/Γ

58 Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

 $\Gamma(\bar{D}^0 \pi^+)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})**4.92 ± 0.20 OUR AVERAGE**EVTSDOCUMENT IDTECNCOMMENT

4.86 $\pm 0.27 \pm 0.09$	59	AUBERT,B	04P BABR	$e^+ e^- \rightarrow \gamma(4S)$
4.97 $\pm 0.12 \pm 0.29$	60,61	AHMED	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
5.0 $\pm 0.7 \pm 0.6$	54	62 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
5.4 $\pm 1.8 \pm 1.2$	14	63 BEBEK	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

 Γ_{31}/Γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.5 $\pm 0.4 \pm 0.5$	304	64 ALAM	94 CLE2	Repl. by AHMED 02B
2.0 $\pm 0.8 \pm 0.6$	12	62 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$
1.9 $\pm 1.0 \pm 0.6$	7	65 ALBRECHT	88K ARG	$e^+ e^- \rightarrow \gamma(4S)$

⁵⁹ AUBERT,B 04P reports $[B(B^+ \rightarrow \bar{D}^0 \pi^+) \times B(\bar{D}^0 \rightarrow K^- \pi^+)] = (1.846 \pm 0.032 \pm 0.097) \times 10^{-4}$. We divide by our best value $B(\bar{D}^0 \rightarrow K^- \pi^+) = (3.80 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶⁰ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁶¹ AHMED 02B reports an additional uncertainty on the branching ratios to account for 4.5% uncertainty on relative production of B^0 and B^+ , which is not included here.

⁶² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D .

⁶³ BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

⁶⁴ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(\bar{D}^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(\bar{D}^0 \rightarrow K^- \pi^+ \pi^0)/B(\bar{D}^0 \rightarrow K^- \pi^+)$ and $B(\bar{D}^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(\bar{D}^0 \rightarrow K^- \pi^+)$.

⁶⁵ ALBRECHT 88K assumes $B^0 \bar{B}^0 : B^+ B^-$ ratio is 45:55. Superseded by ALBRECHT 90J.

$\Gamma(\bar{D}^0 \rho^+)/\Gamma_{\text{total}}$

Γ_{34}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0134 ± 0.0018 OUR AVERAGE				
0.0135 ± 0.0012 ± 0.0015	212	66 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.013 ± 0.004 ± 0.004	19	67 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.021 ± 0.008 ± 0.009	10	68 ALBRECHT	88K ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
66 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(\bar{D}^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(\bar{D}^0 \rightarrow K^- \pi^+ \pi^0)/B(\bar{D}^0 \rightarrow K^- \pi^+)$ and $B(\bar{D}^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(\bar{D}^0 \rightarrow K^- \pi^+)$.				
67 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D .				
68 ALBRECHT 88K assumes $B^0 \bar{B}^0 : B^+ B^-$ ratio is 45:55.				

$\Gamma(\bar{D}^0 K^+)/\Gamma_{\text{total}}$

Γ_{35}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.08 ± 0.24 OUR AVERAGE			
4.09 ± 0.20 ± 0.17	69 AUBERT	04N BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
4.9 ± 0.8 ± 0.2	70 BORNHEIM	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
3.8 ± 0.4 ± 0.2	71,72 SWAIN	03 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.6 ± 0.6 ± 0.2	71,73 ABE	03D BELL	Repl. by SWAIN 03
4.19 ± 0.57 ± 0.40	74 ABE	01I BELL	Repl. by ABE 03D
2.92 ± 0.80 ± 0.28	75 ATHANAS	98 CLE2	Repl. by BORNHEIM 03
69 AUBERT 04N reports $[B(B^+ \rightarrow \bar{D}^0 K^+) / B(B^+ \rightarrow \bar{D}^0 \pi^+)] = (831 \pm 35 \pm 20) \times 10^{-4}$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 \pi^+) = (4.92 \pm 0.20) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
70 BORNHEIM 03 reports $[B(B^+ \rightarrow \bar{D}^0 K^+) / B(B^+ \rightarrow \bar{D}^0 \pi^+)] = (990^{+140+70}_{-120-60}) \times 10^{-4}$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 \pi^+) = (4.92 \pm 0.20) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
71 Flavor specific D^0 meson is reconstructed via $D^0 \rightarrow K^- \pi^+$.			

⁷² SWAIN 03 reports $[B(B^+ \rightarrow \bar{D}^0 K^+) / B(B^+ \rightarrow \bar{D}^0 \pi^+)] = (770 \pm 50 \pm 60) \times 10^{-4}$.

We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 \pi^+) = (4.92 \pm 0.20) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷³ ABE 03D reports $[B(B^+ \rightarrow \bar{D}^0 K^+) / B(B^+ \rightarrow \bar{D}^0 \pi^+)] = (940 \pm 90 \pm 70) \times 10^{-4}$.

We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 \pi^+) = (4.92 \pm 0.20) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷⁴ ABE 01I reports $B(B^+ \rightarrow \bar{D}^0 K^+)/B(B^+ \rightarrow \bar{D}^0 \pi^+) = 0.079 \pm 0.009 \pm 0.006$. We

multiply by our best value $B(B^+ \rightarrow \bar{D}^0 \pi^+) = (5.3 \pm 0.5) \times 10^{-3}$. Our first error is their experiment's error and the second error is systematic error from using our best value.

⁷⁵ ATHANAS 98 reports $[B(B^+ \rightarrow \bar{D}^0 K^+)]/[B(B^+ \rightarrow \bar{D}^0 \pi^+)] = 0.055 \pm 0.014 \pm 0.005$.

We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 \pi^+) = (5.3 \pm 0.5) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{CP(+1)} K^+)/\Gamma_{\text{total}}$

Γ_{36}/Γ

VALUE (units 10^{-4})

3.7±0.5±0.2

DOCUMENT ID

⁷⁶ AUBERT

TECN

BABR

COMMENT

$e^+ e^- \rightarrow \gamma(4S)$

⁷⁶ AUBERT 06J reports $[B(B^+ \rightarrow D_{CP(+1)} K^+) / B(B^+ \rightarrow \bar{D}^0 K^+)] = 0.90 \pm 0.12 \pm$

0.04. We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 K^+) = (4.08 \pm 0.24) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{CP(-1)} K^+)/\Gamma_{\text{total}}$

Γ_{37}/Γ

VALUE (units 10^{-4})

3.5±0.5±0.2

DOCUMENT ID

⁷⁷ AUBERT

TECN

BABR

COMMENT

$e^+ e^- \rightarrow \gamma(4S)$

⁷⁷ AUBERT 06J reports $[B(B^+ \rightarrow D_{CP(-1)} K^+) / B(B^+ \rightarrow \bar{D}^0 K^+)] = 0.86 \pm 0.10 \pm$

0.05. We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 K^+) = (4.08 \pm 0.24) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{CP(+1)} K^+)/\Gamma(D_{CP(+1)} \pi^+)$

Γ_{36}/Γ_{32}

VALUE

0.091±0.012 OUR AVERAGE

DOCUMENT ID

TECN

COMMENT

$0.094 \pm 0.015 \pm 0.007$

⁷⁸ ABE

06 BELL

$e^+ e^- \rightarrow \gamma(4S)$

$0.088 \pm 0.016 \pm 0.005$

⁷⁹ AUBERT

04N BABR

$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.125 \pm 0.036 \pm 0.010$

⁷⁹ ABE

03D BELL

Repl. by SWAIN 03

$0.093 \pm 0.018 \pm 0.008$

⁷⁹ SWAIN

03 BELL

Repl. by ABE 06

⁷⁸ Reports a double ratio of $B(B^+ \rightarrow D_{CP(+1)} K^+)/B(B^+ \rightarrow D_{CP(+1)} \pi^+)$ and

$B(B^+ \rightarrow \bar{D}^0 K^+)/B(B^+ \rightarrow \bar{D}^0 \pi^+)$, $1.13 \pm 0.16 \pm 0.08$. We multiply by our best value of $B(B^+ \rightarrow \bar{D}^0 K^+)/B(B^+ \rightarrow \bar{D}^0 \pi^+) = 0.083 \pm 0.006$. Our first error is their experiment's error and the second error is systematic error from using our best value.

⁷⁹ CP=+1 eigenstate of $D^0 \bar{D}^0$ system is reconstructed via $K^+ K^-$ and $\pi^+ \pi^-$.

$\Gamma(D_{CP(-1)}K^+)/\Gamma(D_{CP(-1)}\pi^+)$ Γ_{37}/Γ_{33}

VALUE	DOCUMENT ID	TECN	COMMENT
0.097±0.016±0.007	80 ABE	06 BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.119±0.028±0.006	81 ABE	03D BELL	Repl. by SWAIN 03
0.108±0.019±0.007	81 SWAIN	03 BELL	Repl. by ABE 06
80 Reports a double ratio of $B(B^+ \rightarrow D_{CP(-1)}K^+)/B(B^+ \rightarrow D_{CP(-1)}\pi^+)$ and $B(B^+ \rightarrow \bar{D}^0 K^+)/B(B^+ \rightarrow \bar{D}^0 \pi^+)$, $1.17 \pm 0.14 \pm 0.14$. We multiply by our best value of $B(B^+ \rightarrow \bar{D}^0 K^+)/B(B^+ \rightarrow \bar{D}^0 \pi^+) = 0.083 \pm 0.006$. Our first error is their experiment's error and the second error is systematic error from using our best value.			
81 $CP=-1$ eigenstate of $D^0 \bar{D}^0$ system is reconstructed via $K_S^0 \pi^0$, $K_S^0 \omega$, $K_S^0 \phi$, $K_S^0 \eta$, and $K_S^0 \eta'$.			

 $\Gamma([K^-\pi^+]_D K^+)/\Gamma([K^+\pi^-]_D K^+)$ Γ_{38}/Γ_{39}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.029	90	82 AUBERT	05G BABR	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.044	90	83 SAIGO	05 BELL	$e^+e^- \rightarrow \gamma(4S)$
<0.026	90	84 AUBERT,B	04L BABR	Repl. by AUBERT 05G
82 AUBERT 05G extract a constraint on the magnitude of the ratio of amplitudes $ A(B^+ \rightarrow D^0 K^+)/A(B^+ \rightarrow \bar{D}^0 K^+) < 0.23$ at 90% CL (Bayesian). Similar measurements from $B^+ \rightarrow D^{*0} K^+$ are also reported.				
83 SAIGO 05 extract a constraint on the magnitude of the ratio of amplitudes $ A(B^+ \rightarrow D^0 K^+)/A(B^+ \rightarrow \bar{D}^0 K^+) < 0.27$ at 90% CL.				
84 AUBERT,B 04L extract a constraint on the magnitude of the ratio of amplitudes $ A(B^+ \rightarrow D^0 K^+)/A(B^+ \rightarrow \bar{D}^0 K^+) < 0.22$ at 90% CL.				

 $\Gamma([K^-\pi^+]_D K^*(892)^+)/\Gamma([K^+\pi^-]_D K^*(892)^+)$ Γ_{40}/Γ_{41}

VALUE	DOCUMENT ID	TECN	COMMENT
0.046±0.031±0.008	AUBERT,B	05v BABR	$e^+e^- \rightarrow \gamma(4S)$

 $\Gamma([K^-\pi^+]_D \pi^+)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.74^{+0.52}_{-0.47}±0.03	85 SAIGO	05 BELL	$e^+e^- \rightarrow \gamma(4S)$

85 SAIGO 05 reports $[B(B^+ \rightarrow [K^-\pi^+]_D \pi^+) \times B(D^0 \rightarrow K^-\pi^+)] = (6.6^{+1.9}_{-1.7} \pm 0.5) \times 10^{-7}$. We divide by our best value $B(D^0 \rightarrow K^-\pi^+) = (3.80 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma([\pi^+\pi^-\pi^0]_D K^-)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
5.5^{+1.0}_{-1.0}±0.7	86 AUBERT,B	05T BABR	$e^+e^- \rightarrow \gamma(4S)$

86 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma([K^-\pi^+]_D\pi^+)/\Gamma(\bar{D}^0\pi^+)$ Γ_{42}/Γ_{31} VALUE (units 10^{-3}) **$3.5^{+1.0}_{-0.9}\pm0.2$** DOCUMENT ID

SAIGO

TECN

BELL

COMMENT $e^+e^- \rightarrow \gamma(4S)$ $\Gamma(\bar{D}^0 K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{44}/Γ VALUE (units 10^{-4}) **6.3 ± 0.8 OUR AVERAGE** $6.3\pm0.7\pm0.5$ $6.1\pm1.6\pm1.7$ DOCUMENT ID

87 AUBERT

TECN

04Q BABR

COMMENT

87 MAHAPATRA 02 CLE2

 $e^+e^- \rightarrow \gamma(4S)$ 87 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D_{CP(-1)}K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{45}/Γ VALUE (units 10^{-4}) **$2.0\pm0.9\pm0.3$** DOCUMENT ID

88 AUBERT,B

TECN

05U BABR

COMMENT $e^+e^- \rightarrow \gamma(4S)$ 88 AUBERT,B 05U reports $[B(B^+ \rightarrow D_{CP(-1)}K^*(892)^+) / B(B^+ \rightarrow \bar{D}^0 K^*(892)^+)] = 0.325 \pm 0.13 \pm 0.04$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 K^*(892)^+) = (6.3 \pm 0.8) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma(D_{CP(+1)}K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{46}/Γ VALUE (units 10^{-4}) **$6.2\pm1.3\pm0.8$** DOCUMENT ID

89 AUBERT,B

TECN

05U BABR

COMMENT $e^+e^- \rightarrow \gamma(4S)$ 89 AUBERT,B 05U reports $[B(B^+ \rightarrow D_{CP(+1)}K^*(892)^+) / B(B^+ \rightarrow \bar{D}^0 K^*(892)^+)] = 0.98 \pm 0.20 \pm 0.055$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^0 K^*(892)^+) = (6.3 \pm 0.8) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma(\bar{D}^0 K^+\bar{K}^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ VALUE (units 10^{-4}) **$5.5\pm1.4\pm0.8$** DOCUMENT ID

90 DRUTSKOY

TECN

02 BELL

COMMENT $e^+e^- \rightarrow \gamma(4S)$ 90 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\bar{D}^0 K^+\bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{48}/Γ VALUE (units 10^{-4}) **$7.5\pm1.3\pm1.1$** DOCUMENT ID

91 DRUTSKOY

TECN

02 BELL

COMMENT $e^+e^- \rightarrow \gamma(4S)$ 91 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\bar{D}^0\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{49}/Γ VALUE **$0.0115\pm0.0029\pm0.0021$** DOCUMENT ID

92 BORTOLETTO92

TECN

CLEO

COMMENT $e^+e^- \rightarrow \gamma(4S)$ 92 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\overline{D}^0 \pi^+ \pi^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0051±0.0034±0.0023	93 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

93 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

 $\Gamma(\overline{D}^0 \pi^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0042±0.0023±0.0020	94 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

94 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

 $\Gamma(\overline{D}^0 a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0045±0.0019±0.0031	95 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$

95 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

 $\Gamma(\overline{D}^0 \omega \pi^+)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0041±0.0007±0.0006	96 ALEXANDER 01B	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

96 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

 $\Gamma(D^*(2010)^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.35±0.22 OUR AVERAGE					
1.25±0.08±0.22			97 ABE	04D BELL	$e^+ e^- \rightarrow \gamma(4S)$
1.9 ± 0.7 ± 0.3	14		98 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
2.6 ± 1.4 ± 0.7	11		99 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$
2.4 +1.7 +1.0 -1.6 -0.6	3		100 BEBEK	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.	90	101 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
5. ± 2. ± 3.	7	102 ALBRECHT	87C ARG	$e^+ e^- \rightarrow \gamma(4S)$

97 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

98 ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

99 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses the Mark III branching fractions for the D .

100 BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

101 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$. The authors also find the product branching fraction into $D^{**} \pi$ followed by $D^{**} \rightarrow D^*(2010) \pi$ to be $0.0014^{+0.0008}_{-0.0006} \pm 0.0003$ where D^{**} represents all orbitally excited D mesons.

¹⁰² ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

$\Gamma(D^- \pi^+ \pi^+)/\Gamma_{\text{total}}$					Γ_{55}/Γ
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$1.02 \pm 0.04 \pm 0.15$		103	ABE	04D BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.4	90	104	ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<7	90	105	BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
$2.5 \begin{array}{l} +4.1 \\ -2.3 \end{array} \begin{array}{l} +2.4 \\ -0.8 \end{array}$	1	106	BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

¹⁰³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹⁰⁴ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the Mark III $B(D^+ \rightarrow K^- \pi^+ \pi^+)$.

¹⁰⁵ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D . The product branching fraction into $D_0^*(2340)\pi$ followed by $D_0^*(2340) \rightarrow D\pi$ is < 0.005 at 90%CL and into $D_2^*(2460)$ followed by $D_2^*(2460) \rightarrow D\pi$ is < 0.004 at 90%CL.

¹⁰⁶ BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. $B(D^- \rightarrow K^+ \pi^- \pi^-) = (9.1 \pm 1.3 \pm 0.4)\%$ is assumed.

$\Gamma(D^+ K^0)/\Gamma_{\text{total}}$					Γ_{56}/Γ
VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<5.0	90	107	AUBERT,B	05E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

¹⁰⁷ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 \pi^+)/\Gamma_{\text{total}}$					Γ_{57}/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.0046 ± 0.0004 OUR AVERAGE					
0.00434 $\pm 0.00047 \pm 0.00018$		108 BRANDENB...	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	
0.0052 $\pm 0.0007 \pm 0.0007$	71	109 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	
0.0072 $\pm 0.0018 \pm 0.0016$		110 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	
0.0040 $\pm 0.0014 \pm 0.0012$	9	110 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0027 ± 0.0044		111 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

¹⁰⁸ BRANDENBURG 98 assume equal production of B^+ and B^0 at $\Upsilon(4S)$ and use the D^* reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of $B(D^* \rightarrow D\pi)$.

¹⁰⁹ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

¹¹⁰ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

¹¹¹ This is a derived branching ratio, using the inclusive pion spectrum and other two-body B decays. BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$.

$\Gamma(\bar{D}^*(2007)^0 \omega \pi^+)/\Gamma_{\text{total}}$	Γ_{60}/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.0045 ± 0.0010 ± 0.0007	112 ALEXANDER	01B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
112 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.			

$\Gamma(\bar{D}^*(2007)^0 \rho^+)/\Gamma_{\text{total}}$	Γ_{61}/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0098 ± 0.0017 OUR AVERAGE				
0.0098 ± 0.0006 ± 0.0017		113 CSORNA	03 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.010 ± 0.006 ± 0.004	7	114 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0168 ± 0.0021 ± 0.0028	86	115 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
113 Assumes equal production of B^0 and B^+ at the $\gamma(4S)$ resonance. The second error combines the systematic and theoretical uncertainties in quadrature. CSORNA 03 includes data used in ALAM 94. A full angular fit to three complex helicity amplitudes is performed.				
114 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.				
115 ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$. The nonresonant $\pi^+ \pi^0$ contribution under the ρ^+ is negligible.				

$\Gamma(\bar{D}^*(2007)^0 K^+)/\Gamma_{\text{total}}$	Γ_{62}/Γ		
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.7 ± 0.4 OUR AVERAGE			
3.72 ± 0.27 ± 0.35		116 AUBERT	$e^+ e^- \rightarrow \gamma(4S)$
3.59 ± 0.97 ± 0.31		117 ABE	$e^+ e^- \rightarrow \gamma(4S)$
116 AUBERT 05N reports $[B(B^+ \rightarrow \bar{D}^*(2007)^0 K^+) / B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+)] = 0.0813 \pm 0.0040^{+0.0042}_{-0.0031}$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+) = (4.6 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
117 ABE 01I reports $B(B^+ \rightarrow \bar{D}^*(2007)^0 K^+)/B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+) = 0.078 \pm 0.019 \pm 0.009$. We multiply by our best value $B(B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+) = (4.6 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and the second error is systematic error from using our best value.			

$\Gamma(\bar{D}_{CP(+1)}^{*0} K^+)/\Gamma(\bar{D}_{CP(+1)}^{*0} \pi^+)$	Γ_{63}/Γ_{58}		
VALUE	DOCUMENT ID	TECN	COMMENT
0.095 ± 0.017 OUR AVERAGE			
0.11 ± 0.02 ± 0.02		118 ABE	$e^+ e^- \rightarrow \gamma(4S)$
0.086 ± 0.021 ± 0.007		119 AUBERT	$e^+ e^- \rightarrow \gamma(4S)$

- 118 Reports a double ratio of $B(B^+ \rightarrow (D_{CP(+1)}^*)^0 K^+)/B(B^+ \rightarrow (D_{CP(+1)}^*)^0 \pi^+)$ and $B(B^+ \rightarrow \bar{D}^{*0} K^+)/B(B^+ \rightarrow \bar{D}^{*0} \pi^+)$, $1.41 \pm 0.25 \pm 0.06$. We multiply by our best value of $B(B^+ \rightarrow \bar{D}^{*0} K^+)/B(B^+ \rightarrow \bar{D}^{*0} \pi^+) = 0.080 \pm 0.011$. Our first error is their experiment's error and the second error is systematic error from using our best value.
 119 Uses $D^{*0} \rightarrow D^0 \pi^0$ with D^0 reconstructed in the CP -even eigenstates $K^+ K^-$ and $\pi^+ \pi^-$.

$$\Gamma(\bar{D}_{CP(-1)}^{*0} K^+)/\Gamma(D_{CP(-1)}^{*0} \pi^+) \quad \Gamma_{64}/\Gamma_{59}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.09±0.03 ±0.01	120 ABE	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$

- 120 Reports a double ratio of $B(B^+ \rightarrow (D_{CP(-1)}^*)^0 K^+)/B(B^+ \rightarrow (D_{CP(-1)}^*)^0 \pi^+)$ and $B(B^+ \rightarrow \bar{D}^{*0} K^+)/B(B^+ \rightarrow \bar{D}^{*0} \pi^+)$, $1.15 \pm 0.31 \pm 0.12$. We multiply by our best value of $B(B^+ \rightarrow \bar{D}^{*0} K^+)/B(B^+ \rightarrow \bar{D}^{*0} \pi^+) = 0.080 \pm 0.011$. Our first error is their experiment's error and the second error is systematic error from using our best value.

$$\Gamma(\bar{D}^*(2007)^0 K^*(892)^+)/\Gamma_{\text{total}} \quad \Gamma_{65}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.1±1.4 OUR AVERAGE			

- 8.3±1.1±1.0
 7.2±2.2±2.6

121 AUBERT	04K BABR	$e^+ e^- \rightarrow \gamma(4S)$
122 MAHAPATRA	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

121 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

122 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and an unpolarized final state.

$$\Gamma(\bar{D}^*(2007)^0 K^+ \bar{K}^0)/\Gamma_{\text{total}} \quad \Gamma_{66}/\Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<10.6	90	123 DRUTSKOY	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$

123 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(\bar{D}^*(2007)^0 K^+ K^*(892)^0)/\Gamma_{\text{total}} \quad \Gamma_{67}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
15.3±3.1±2.9	124 DRUTSKOY	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$

124 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(\bar{D}^*(2007)^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{68}/\Gamma$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.03 ±0.12 OUR AVERAGE				

- 1.055±0.047±0.129
 0.94 ±0.20 ±0.17

125 MAJUMDER	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$
126,127 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

125 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

126 ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

127 The three pion mass is required to be between 1.0 and 1.6 GeV consistent with an a_1 meson. (If this channel is dominated by a_1^+ , the branching ratio for $\bar{D}^{*0} a_1^+$ is twice that for $\bar{D}^{*0} \pi^+ \pi^+ \pi^-$.)

$\Gamma(\overline{D}^*(2007)^0 a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0188±0.0040±0.0034	128,129 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

128 ALAM 94 value is twice their $\Gamma(\overline{D}^*(2007)^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ value based on their observation that the three pions are dominantly in the $a_1(1260)$ mass range 1.0 to 1.6 GeV.

129 ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

 $\Gamma(\overline{D}^*(2007)^0 \pi^- \pi^+ \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0180±0.0024±0.0027	130 ALEXANDER	01B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

130 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

 $\Gamma(\overline{D}^{*0} 3\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
5.67±0.91±0.85	131 MAJUMDER	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$

131 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^*(2010)^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00017	90	132 BRANDENB...	98 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

132 BRANDENBURG 98 assume equal production of B^+ and B^0 at $\gamma(4S)$ and use the D^* partial reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of $B(D^* \rightarrow D\pi)$.

 $\Gamma(D^*(2010)^+ K^0)/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.0 \times 10^{-6}$	90	133 AUBERT,B	05E BABR	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<9.5 \times 10^{-5}$	90	133 GRITSAN	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
-----------------------	----	-------------	---------	----------------------------------

133 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0152±0.0071±0.0001	26	134 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.043 ± 0.013 ± 0.026	24	135 ALBRECHT	87C ARG	$e^+ e^- \rightarrow \gamma(4S)$
-----------------------	----	--------------	---------	----------------------------------

134 ALBRECHT 90J reports $0.018 \pm 0.007 \pm 0.005$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D .

135 ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\gamma(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\gamma(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

$\Gamma(D^*(2010)^-\pi^+\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{75}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.56±0.26±0.33		136 MAJUMDER	04 BELL	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10 90 137 ALBRECHT 90J ARG $e^+e^- \rightarrow \gamma(4S)$

136 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

137 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

 $\Gamma(\bar{D}_1^*(2420)^0\pi^+)/\Gamma_{\text{total}}$ Γ_{76}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0015±0.0006 OUR AVERAGE				Error includes scale factor of 1.3.
0.0011±0.0005±0.0002	8	138 ALAM	94 CLE2	$e^+e^- \rightarrow \gamma(4S)$
0.0025±0.0007±0.0006		139 ALBRECHT	94D ARG	$e^+e^- \rightarrow \gamma(4S)$

138 ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0\pi^+)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and assuming $B(D_1(2420)^0 \rightarrow D^*(2010)^+\pi^-) = 67\%$.

139 ALBRECHT 94D assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0\pi^+)$ assuming $B(D_1(2420)^0 \rightarrow D^*(2010)^+\pi^-) = 67\%$.

 $\Gamma(\bar{D}_1(2420)^0\pi^+ \times B(\bar{D}_1^0 \rightarrow \bar{D}^0\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{77}/Γ

<u>VALUE</u> (units 10^{-4})		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.85±0.29^{+0.35}_{-0.55}		140 ABE	05A BELL	$e^+e^- \rightarrow \gamma(4S)$

140 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}_2^*(2462)^0\pi^+ \times B(\bar{D}_2^*(2462)^0 \rightarrow D^-\pi^+))/\Gamma_{\text{total}}$ Γ_{78}/Γ

<u>VALUE</u> (units 10^{-4})		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.4±0.3±0.72		141 ABE	04D BELL	$e^+e^- \rightarrow \gamma(4S)$

141 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}_0^*(2308)^0\pi^+ \times B(\bar{D}_0^*(2308)^0 \rightarrow D^-\pi^+))/\Gamma_{\text{total}}$ Γ_{79}/Γ

<u>VALUE</u> (units 10^{-4})		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.1±0.6±1.8		142 ABE	04D BELL	$e^+e^- \rightarrow \gamma(4S)$

142 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}_1(2421)^0\pi^+ \times B(\bar{D}_1(2421)^0 \rightarrow D^{*-}\pi^+))/\Gamma_{\text{total}}$ Γ_{80}/Γ

<u>VALUE</u> (units 10^{-4})		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.8±0.7±1.3		143 ABE	04D BELL	$e^+e^- \rightarrow \gamma(4S)$

143 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}_2^*(2462)^0\pi^+ \times B(\bar{D}_2^*(2462)^0 \rightarrow D^{*-}\pi^+))/\Gamma_{\text{total}}$ Γ_{81}/Γ

<u>VALUE</u> (units 10^{-4})		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.8±0.3±0.4		144 ABE	04D BELL	$e^+e^- \rightarrow \gamma(4S)$

144 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(\overline{D}'_1(2427)^0 \pi^+ \times B(\overline{D}'_1(2427)^0 \rightarrow D^{*-} \pi^+)) / \Gamma_{\text{total}} \quad \Gamma_{82}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
5.0 ± 0.4 ± 1.1	145 ABE	04D BELL	$e^+ e^- \rightarrow \gamma(4S)$

145 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(\overline{D}_1(2420)^0 \pi^+ \times B(\overline{D}_1^0 \rightarrow \overline{D}^{*0} \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{83}/\Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.06	90	146 ABE	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$

146 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(\overline{D}_1^*(2420)^0 \rho^+)/\Gamma_{\text{total}} \quad \Gamma_{84}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0014	90	147 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

147 ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ assuming $B(D_1(2420)^0 \rightarrow D^*(2010)^+ \pi^-) = 67\%$.

$$\Gamma(\overline{D}_2^*(2460)^0 \pi^+)/\Gamma_{\text{total}} \quad \Gamma_{85}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0013	90	148 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0028 90 149 ALAM 94 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

<0.0023 90 150 ALBRECHT 94D ARG $e^+ e^- \rightarrow \gamma(4S)$

148 ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the Mark III $B(D^+ \rightarrow K^- \pi^+ \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^+ \pi^-) = 30\%$.

149 ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the Mark III $B(D^+ \rightarrow K^- \pi^+ \pi^+)$, the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+ \pi^-) = 20\%$.

150 ALBRECHT 94D assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+ \pi^-) = 30\%$.

$$\Gamma(\overline{D}_2^*(2460)^0 \pi^+ \times B(\overline{D}_2^{*0} \rightarrow \overline{D}^{*0} \pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_{86}/\Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.22	90	151 ABE	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$

151 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(\overline{D}_2^*(2460)^0 \rho^+)/\Gamma_{\text{total}} \quad \Gamma_{87}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0047	90	152 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
<0.005	90	153 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

152 ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the Mark III $B(D^+ \rightarrow K^- \pi^+ \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^+ \pi^-) = 30\%$.

153 ALAM 94 assume equal production of B^+ and B^0 at the $\gamma(4S)$ and use the Mark III $B(D^+ \rightarrow K^- \pi^+ \pi^+)$, the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and $B(D_2^*(2460)^0 \rightarrow D^*(2010)^+ \pi^-) = 20\%$.

$\Gamma(\bar{D}^0 D_s^+)/\Gamma_{\text{total}}$ Γ_{88}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0109±0.0027 OUR AVERAGE				
0.0100±0.0026±0.0013	154	GIBAUT	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.015 ± 0.008 ± 0.002	155	ALBRECHT	92G ARG	$e^+ e^- \rightarrow \gamma(4S)$
0.013 ± 0.006 ± 0.002	5	156 BORTOLETTO90	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
154 GIBAUT 96 reports $0.0126 \pm 0.0022 \pm 0.0025$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
155 ALBRECHT 92G reports $0.024 \pm 0.012 \pm 0.004$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 branching ratios, e.g., $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$.				
156 BORTOLETTO 90 reports 0.029 ± 0.013 for $B(D_s^+ \rightarrow \phi\pi^+) = 0.02$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(D_{s0}(2317)^+\bar{D}^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^+\pi^0))/\Gamma_{\text{total}}$ Γ_{89}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.74^{+0.23}_{-0.19} OUR AVERAGE			
0.82 ^{+0.35} _{-0.21} ± 0.11	157,158 AUBERT,B	04S BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.66 ^{+0.27} _{-0.24} ± 0.09	157,159 KROKOVNY	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
157 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.			
158 AUBERT,B 04S reports $(1.0 \pm 0.3 \pm 0.4) \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
159 KROKOVNY 03B reports $(0.81 \pm 0.30 \pm 0.27) \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.036 \pm 0.009$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

 $\Gamma(D_{s0}(2317)^+\bar{D}^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^{*+}\gamma))/\Gamma_{\text{total}}$ Γ_{90}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.76	90	160 KROKOVNY	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
160 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

 $\Gamma(D_{s0}(2317)^+\bar{D}^*(2010)^0 \times B(D_{s0}(2317)^+ \rightarrow D_s^+\pi^0))/\Gamma_{\text{total}}$ Γ_{91}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.9 ^{+0.6} _{-0.3}	161 AUBERT,B	04S BABR	$e^+ e^- \rightarrow \gamma(4S)$
161 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.			

$\Gamma(D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0)) / \Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.4^{+0.6}_{-0.5}$ OUR AVERAGE	Error includes scale factor of 1.3.		
$2.2^{+0.8}_{-0.7} \pm 0.3$	162,163 AUBERT,B	04S BABR	$e^+ e^- \rightarrow \gamma(4S)$
$1.0^{+0.5}_{-0.4} \pm 0.1$	162,164 KROKOVNY	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
162 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.			
163 AUBERT,B 04S reports $(2.7 \pm 0.7)^{+1.0}_{-0.8} \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.036 \pm 0.009$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
164 KROKOVNY 03B reports $(1.19^{+0.61}_{-0.49} \pm 0.36) \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.036 \pm 0.009$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

 $\Gamma(D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)) / \Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$0.47^{+0.14}_{-0.12}$ OUR AVERAGE			
$0.49^{+0.20}_{-0.14} \pm 0.06$	165,166 AUBERT,B	04S BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.46 \pm 0.15 \pm 0.06$	165,167 KROKOVNY	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
165 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.			
166 AUBERT,B 04S reports $(0.6 \pm 0.2)^{+0.2}_{-0.1} \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.036 \pm 0.009$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
167 KROKOVNY 03B reports $(0.56^{+0.16}_{-0.15} \pm 0.17) \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.036 \pm 0.009$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

 $\Gamma(D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.22	90	168 KROKOVNY	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
168 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

 $\Gamma(D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \pi^0)) / \Gamma_{\text{total}}$ Γ_{95}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.27	90	169 KROKOVNY	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
169 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

$$\Gamma(D_{sJ}(2457)^+ \bar{D}^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \gamma)) / \Gamma_{\text{total}} \quad \Gamma_{96}/\Gamma$$

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.98	90	170 KROKOVNY	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$

170 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(D_{sJ}(2457)^+ \bar{D}^*(2010)^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^{*+} \pi^0)) / \Gamma_{\text{total}} \quad \Gamma_{97}/\Gamma$$

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.6 \pm 1.7^{+3.2}_{-2.4}	171 AUBERT,B	04S BABR	$e^+ e^- \rightarrow \gamma(4S)$

171 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(D_{sJ}(2457)^+ \bar{D}^*(2010)^0 \times B(D_{sJ}(2457)^+ \rightarrow D_s^+ \gamma)) / \Gamma_{\text{total}} \quad \Gamma_{98}/\Gamma$$

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.4 \pm 0.4^{+0.6}_{-0.4}	172 AUBERT,B	04S BABR	$e^+ e^- \rightarrow \gamma(4S)$

172 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(\bar{D}^0 D_{sJ}(2536)^+ \times B(D_{sJ}(2536)^+ \rightarrow D^*(2007)^0 K^+)) / \Gamma_{\text{total}} \quad \Gamma_{99}/\Gamma$$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

$$\Gamma(\bar{D}^*(2007)^0 D_{sJ}(2536)^+ \times B(D_{sJ}(2536)^+ \rightarrow D^*(2007)^0 K^+)) / \Gamma_{\text{total}} \quad \Gamma_{100}/\Gamma$$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<7	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

$$\Gamma(\bar{D}^0 D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)) / \Gamma_{\text{total}} \quad \Gamma_{101}/\Gamma$$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

$$\Gamma(\bar{D}^*(2007)^0 D_{sJ}(2573)^+ \times B(D_{sJ}(2573)^+ \rightarrow D^0 K^+)) / \Gamma_{\text{total}} \quad \Gamma_{102}/\Gamma$$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$

$$\Gamma(\bar{D}^0 D_s^{*+}) / \Gamma_{\text{total}} \quad \Gamma_{103}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0072 \pm 0.0026 OUR AVERAGE			

0.0069 $\pm 0.0025 \pm 0.0009$ 173 GIBAUT 96 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

0.010 $\pm 0.008 \pm 0.001$ 174 ALBRECHT 92G ARG $e^+ e^- \rightarrow \gamma(4S)$

173 GIBAUT 96 reports $0.0087 \pm 0.0027 \pm 0.0017$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

174 ALBRECHT 92G reports $0.016 \pm 0.012 \pm 0.003$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 branching ratios, e.g., $B(D^0 \rightarrow K^- \pi^+) = 3.71 \pm 0.25\%$.

$\Gamma(\bar{D}^*(2007)^0 D_s^+)/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.010±0.004 OUR AVERAGE			
0.011±0.004±0.001	175 GIBAUT	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.008±0.006±0.001	176 ALBRECHT	92G ARG	$e^+ e^- \rightarrow \gamma(4S)$
175 GIBAUT 96 reports $0.0140 \pm 0.0043 \pm 0.0035$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
176 ALBRECHT 92G reports $0.013 \pm 0.009 \pm 0.002$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 and $D^*(2007)^0$ branching ratios, e.g., $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$ and $B(D^*(2007)^0 \rightarrow D^0\pi^0) = 55 \pm 6\%$.			

 $\Gamma(\bar{D}^*(2007)^0 D_s^{*+})/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.022±0.007 OUR AVERAGE			
0.025±0.009±0.003	177 GIBAUT	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
0.019±0.010±0.002	178 ALBRECHT	92G ARG	$e^+ e^- \rightarrow \gamma(4S)$
177 GIBAUT 96 reports $0.0310 \pm 0.0088 \pm 0.0065$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
178 ALBRECHT 92G reports $0.031 \pm 0.016 \pm 0.005$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 and $D^*(2007)^0$ branching ratios, e.g., $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$ and $B(D^*(2007)^0 \rightarrow D^0\pi^0) = 55 \pm 6\%$.			

 $\Gamma(D_s^{(*)+}\bar{D}^{**0})/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
(2.73 ±0.93 ±0.68) × 10⁻²			
179 AHMED	00B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
179 AHMED 00B reports their experiment's uncertainties ($\pm 0.78 \pm 0.48 \pm 0.68\%$), where the first error is statistical, the second is systematic, and the third is the uncertainty in the $D_s \rightarrow \phi\pi$ branching fraction. We combine the first two in quadrature.			

 $\Gamma(\bar{D}^*(2007)^0 D^*(2010)^+)/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.011	90	BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

 $[\Gamma(\bar{D}^0 D^*(2010)^+) + \Gamma(\bar{D}^*(2007)^0 D^+)]/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.013	90	BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$

 $\Gamma(\bar{D}^0 D^*(2010)^+)/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE (units 10 ⁻⁴)	DOCUMENT ID	TECN	COMMENT
4.57±0.71±0.56	180 MAJUMDER	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$

180 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\bar{D}^0 D^+)/\Gamma_{\text{total}}$ Γ_{110}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.83±0.78±0.58		181 MAJUMDER	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$	■

• • • We do not use the following data for averages, fits, limits, etc. • • •

<67 90 BARATE 98Q ALEP $e^+ e^- \rightarrow Z$

181 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}^0 D^+ K^0)/\Gamma_{\text{total}}$ Γ_{111}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2.8	90	182 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$	

182 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}^*(2007)^0 D^+ K^0)/\Gamma_{\text{total}}$ Γ_{112}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<6.1	90	183 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$	

183 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}^0 \bar{D}^*(2010)^+ K^0)/\Gamma_{\text{total}}$ Γ_{113}/Γ

<u>VALUE (units 10^{-3})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
5.2^{+1.0}_{-0.9}±0.7		184 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$	

184 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}^*(2007)^0 D^*(2010)^+ K^0)/\Gamma_{\text{total}}$ Γ_{114}/Γ

<u>VALUE (units 10^{-3})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
7.8^{+2.3}_{-2.1}±1.4		185 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$	

185 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}^0 D^0 K^+)/\Gamma_{\text{total}}$ Γ_{115}/Γ

<u>VALUE (units 10^{-3})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.37±0.32 OUR AVERAGE		Error includes scale factor of 1.5.			
1.17±0.21±0.15		186 CHISTOV	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$	■
1.9 ± 0.3 ± 0.3		186 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$	

186 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}^*(2010)^0 D^0 K^+)/\Gamma_{\text{total}}$ Γ_{116}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<3.8	90	187 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$	

187 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\bar{D}^0 D^*(2007)^0 K^+)/\Gamma_{\text{total}}$ Γ_{117}/Γ

<u>VALUE (units 10^{-3})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.7±0.7±0.7		188 AUBERT	03X BABR	$e^+ e^- \rightarrow \gamma(4S)$	

188 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\overline{D}^*(2007)^0 D^*(2007)^0 K^+)/\Gamma_{\text{total}}$ Γ_{118}/Γ

<u>VALUE</u> (units 10^{-3})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.3^{+1.1}_{-1.0} \pm 1.2$	189 AUBERT	03x BABR	$e^+ e^- \rightarrow \gamma(4S)$

189 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D^- D^+ K^+)/\Gamma_{\text{total}}$ Γ_{119}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.4	90	190 AUBERT	03x BABR	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.90	90	190 CHISTOV	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$
-------	----	-------------	---------	----------------------------------

190 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D^- D^*(2010)^+ K^+)/\Gamma_{\text{total}}$ Γ_{120}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.7	90	191 AUBERT	03x BABR	$e^+ e^- \rightarrow \gamma(4S)$

191 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D^*(2010)^- D^+ K^+)/\Gamma_{\text{total}}$ Γ_{121}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.5^{+0.3 \pm 0.2}$	90	192 AUBERT	03x BABR	$e^+ e^- \rightarrow \gamma(4S)$

192 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D^*(2010)^- D^*(2010)^+ K^+)/\Gamma_{\text{total}}$ Γ_{122}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.8	90	193 AUBERT	03x BABR	$e^+ e^- \rightarrow \gamma(4S)$

193 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma((\overline{D}+\overline{D}^*)(D+D^*)K)/\Gamma_{\text{total}}$ Γ_{123}/Γ

<u>VALUE</u> (units 10^{-2})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.5^{+0.3 \pm 0.5}$	194 AUBERT	03x BABR	$e^+ e^- \rightarrow \gamma(4S)$

194 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(D_s^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{124}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.00017	90	195 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

195 ALEXANDER 93B reports $< 2.0 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$. $[\Gamma(D_s^+ \pi^0) + \Gamma(D_s^{*+} \pi^0)]/\Gamma_{\text{total}}$ $(\Gamma_{124} + \Gamma_{125})/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0006	90	196 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$

196 ALBRECHT 93E reports $< 0.9 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.

$\Gamma(D_s^{*+}\pi^0)/\Gamma_{\text{total}}$ Γ_{125}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00027	90	197 ALEXANDER 93B CLE2	e ⁺ e ⁻ →	$\gamma(4S)$

197 ALEXANDER 93B reports $< 3.2 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.044$.

 $\Gamma(D_s^+\eta)/\Gamma_{\text{total}}$ Γ_{126}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0004	90	198 ALEXANDER 93B CLE2	e ⁺ e ⁻ →	$\gamma(4S)$

198 ALEXANDER 93B reports $< 4.6 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.

 $\Gamma(D_s^{*+}\eta)/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0006	90	199 ALEXANDER 93B CLE2	e ⁺ e ⁻ →	$\gamma(4S)$

199 ALEXANDER 93B reports $< 7.5 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.

 $\Gamma(D_s^+\rho^0)/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00031	90	200 ALEXANDER 93B CLE2	e ⁺ e ⁻ →	$\gamma(4S)$

200 ALEXANDER 93B reports $< 3.7 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.

 $[\Gamma(D_s^+\rho^0) + \Gamma(D_s^+\bar{K}^*(892)^0)]/\Gamma_{\text{total}}$ $(\Gamma_{128} + \Gamma_{138})/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0021	90	201 ALBRECHT 93E ARG	e ⁺ e ⁻ →	$\gamma(4S)$

201 ALBRECHT 93E reports $< 3.4 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.

 $\Gamma(D_s^{*+}\rho^0)/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0004	90	202 ALEXANDER 93B CLE2	e ⁺ e ⁻ →	$\gamma(4S)$

202 ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.

 $[\Gamma(D_s^{*+}\rho^0) + \Gamma(D_s^{*+}\bar{K}^*(892)^0)]/\Gamma_{\text{total}}$ $(\Gamma_{129} + \Gamma_{139})/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0012	90	203 ALBRECHT 93E ARG	e ⁺ e ⁻ →	$\gamma(4S)$

203 ALBRECHT 93E reports $< 2.0 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.

$\Gamma(D_s^+ \omega)/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0004	90	204 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0021	90	205 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$
204 ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.				
205 ALBRECHT 93E reports $< 3.4 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.				

 $\Gamma(D_s^{*+} \omega)/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0006	90	206 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0012	90	207 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$
206 ALEXANDER 93B reports $< 6.8 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.				
207 ALBRECHT 93E reports $< 1.9 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.				

 $\Gamma(D_s^+ a_1(1260)^0)/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0018	90	208 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$
208 ALBRECHT 93E reports $< 3.0 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.				

 $\Gamma(D_s^{*+} a_1(1260)^0)/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0013	90	209 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$
209 ALBRECHT 93E reports $< 2.2 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.				

 $\Gamma(D_s^+ \phi)/\Gamma_{\text{total}}$ Γ_{134}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.9 $\times 10^{-6}$	90	210 AUBERT	06F BABR	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0010	90	211 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$
<0.00026	90	212 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
210 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				
211 ALBRECHT 93E reports $< 1.7 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.				
212 ALEXANDER 93B reports $< 3.1 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.				

$\Gamma(D_s^{*+} \phi)/\Gamma_{\text{total}}$					Γ_{135}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.2 \times 10^{-5}$	90	213 AUBERT	06F BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.0013	90	214 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$	
<0.00035	90	215 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
213 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.					
214 ALBRECHT 93E reports $< 2.1 \times 10^{-3}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.044$.					
215 ALEXANDER 93B reports $< 4.2 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.044$.					

$\Gamma(D_s^{*+} \bar{K}^0)/\Gamma_{\text{total}}$					Γ_{136}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0009	90	216 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.0015	90	217 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$	
216 ALEXANDER 93B reports $< 10.3 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.044$.					
217 ALBRECHT 93E reports $< 2.5 \times 10^{-3}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.044$.					

$\Gamma(D_s^{*+} \bar{K}^0)/\Gamma_{\text{total}}$					Γ_{137}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0009	90	218 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.0019	90	219 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$	
218 ALEXANDER 93B reports $< 10.9 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.044$.					
219 ALBRECHT 93E reports $< 3.1 \times 10^{-3}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.044$.					

$\Gamma(D_s^{*+} \bar{K}^*(892)^0)/\Gamma_{\text{total}}$					Γ_{138}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0004	90	220 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
220 ALEXANDER 93B reports $< 4.4 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.044$.					

$\Gamma(D_s^{*+} \bar{K}^*(892)^0)/\Gamma_{\text{total}}$					Γ_{139}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0004	90	221 ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
221 ALEXANDER 93B reports $< 4.3 \times 10^{-4}$ for $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^{*+} \rightarrow \phi\pi^+) = 0.044$.					

$\Gamma(D_s^- \pi^+ K^+)/\Gamma_{\text{total}}$ Γ_{140}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0007	90	222 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$

222 ALBRECHT 93E reports $< 1.1 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.

 $\Gamma(D_s^{*-} \pi^+ K^+)/\Gamma_{\text{total}}$ Γ_{141}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0010	90	223 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$

223 ALBRECHT 93E reports $< 1.6 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.

 $\Gamma(D_s^- \pi^+ K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{142}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.005	90	224 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$

224 ALBRECHT 93E reports $< 8.6 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.

 $\Gamma(D_s^{*-} \pi^+ K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{143}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.007	90	225 ALBRECHT	93E ARG	$e^+ e^- \rightarrow \gamma(4S)$

225 ALBRECHT 93E reports $< 1.1 \times 10^{-2}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.044$.

 $\Gamma(\eta_c K^+)/\Gamma_{\text{total}}$ Γ_{144}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.91 ± 0.13 OUR AVERAGE

0.87 ± 0.15	226,227 AUBERT	06E BABR	$e^+ e^- \rightarrow \gamma(4S)$
1.4 $^{+0.3}_{-0.2}$ ± 0.4	228 AUBERT,B	05L BABR	$e^+ e^- \rightarrow \gamma(4S)$
1.25 ± 0.14 $^{+0.39}_{-0.40}$	229 FANG	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.69 $^{+0.26}_{-0.21}$ ± 0.22	230 EDWARDS	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.06 $\pm 0.12 \pm 0.18$	227,231 AUBERT,B	04B BABR	$e^+ e^- \rightarrow \gamma(4S)$
--------------------------	------------------	----------	----------------------------------

226 Perform measurements of absolute branching fractions using a missing mass technique.

227 The ratio of $B(B^\pm \rightarrow K^\pm \eta_c)$ $B(\eta_c \rightarrow K\bar{K}\pi) = (7.4 \pm 0.5 \pm 0.7) \times 10^{-5}$ reported in AUBERT,B 04B and $B(B^\pm \rightarrow K^\pm \eta_c) = (8.7 \pm 1.5) \times 10^{-3}$ reported in AUBERT 06E contribute to the determination of $B(\eta_c \rightarrow K\bar{K}\pi)$, which is used by others for normalization.

228 AUBERT,B 05L reports $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c(1S) \rightarrow p\bar{p})] = (1.8 $^{+0.3}_{-0.2} \pm 0.2) \times 10^{-6}$. We divide by our best value $B(\eta_c(1S) \rightarrow p\bar{p}) = (1.3 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.$

229 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

- 230 EDWARDS 01 assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \rightarrow \gamma\eta_c)$ in those modes have been accounted for.
 231 AUBERT,B 04B reports $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c(1S) \rightarrow K\bar{K}\pi)] = (0.074 \pm 0.005 \pm 0.007) \times 10^{-3}$. We divide by our best value $B(\eta_c(1S) \rightarrow K\bar{K}\pi) = (7.0 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta'_c K^+)/\Gamma_{\text{total}}$	Γ_{145}/Γ		
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.4 \pm 1.8 \pm 0.3$	232 AUBERT	06E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

232 Perform measurements of absolute branching fractions using a missing mass technique.

$\Gamma(\eta_c K^+)/\Gamma(J/\psi(1S) K^+)$	$\Gamma_{144}/\Gamma_{146}$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.33 \pm 0.10 \pm 0.43$	233 AUBERT,B	04B BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

233 Uses BABAR measurement of $B(B^+ \rightarrow J/\psi K^+) = (10.1 \pm 0.3 \pm 0.5) \times 10^{-4}$.

$\Gamma(J/\psi(1S) K^+)/\Gamma_{\text{total}}$	Γ_{146}/Γ			
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.08 ± 0.35 OUR FIT				
10.22 ± 0.35 OUR AVERAGE				
8.1 \pm 1.3 \pm 0.7	234	AUBERT	06E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
10.61 \pm 0.15 \pm 0.48	235	AUBERT	05J BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
10.2 \pm 1.0 \pm 0.4	236	AUBERT,B	05L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
10.1 \pm 0.2 \pm 0.7	235	ABE	03B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
10.2 \pm 0.8 \pm 0.7	235	JESSOP	97 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
9.3 \pm 3.1 \pm 0.1	237	BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
8.1 \pm 3.5 \pm 0.1	6	238 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10.1 \pm 0.3 \pm 0.5	235	AUBERT	02 BABR	Repl. by AUBERT 05J
11.0 \pm 1.5 \pm 0.9	59	235 ALAM	94 CLE2	Repl. by JESSOP 97
22 \pm 10 \pm 2		BUSKULIC	92G ALEP	$e^+ e^- \rightarrow Z$
7 \pm 4	3	239 ALBRECHT	87D ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
10 \pm 7 \pm 2	3	240 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
9 \pm 5	3	241 ALAM	86 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

234 Perform measurements of absolute branching fractions using a missing mass technique.

235 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

236 AUBERT,B 05L reports $[B(B^+ \rightarrow J/\psi(1S) K^+) \times B(J/\psi(1S) \rightarrow p\bar{p})] = (2.2 \pm 0.2 \pm 0.1) \times 10^{-6}$. We divide by our best value $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.17 \pm 0.08) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

237 BORTOLETTO 92 reports $(8 \pm 2 \pm 2) \times 10^{-4}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

238 ALBRECHT 90J reports $(7 \pm 3 \pm 1) \times 10^{-4}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first

error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

239 ALBRECHT 87D assume $B^+ B^- / B^0 \bar{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.

240 BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

241 ALAM 86 assumes B^\pm / B^0 ratio is 60/40.

$\Gamma(J/\psi(1S) K^+ \pi^+ \pi^-) / \Gamma_{\text{total}}$

Γ_{147} / Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.07 ± 0.19 OUR AVERAGE			Error includes scale factor of 1.9. See the ideogram below.		
1.16 ± 0.07 ± 0.09			242 AUBERT	05R BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
0.69 ± 0.18 ± 0.12			243 ACOSTA	02F CDF	$p\bar{p}$ 1.8 TeV
1.39 ± 0.82 ± 0.01			244 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
1.39 ± 0.91 ± 0.01	6	245 ALBRECHT	87D ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.9	90		246 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$


242 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

243 ACOSTA 02F uses as reference of $B(B \rightarrow J/\psi(1S) K^+) = (10.1 \pm 0.6) \times 10^{-4}$. The second error includes the systematic error and the uncertainties of the branching ratio.

244 BORTOLETTO 92 reports $(1.2 \pm 0.6 \pm 0.4) \times 10^{-3}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

245 ALBRECHT 87D reports $(1.2 \pm 0.8) \times 10^{-3}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. They actually report 0.0011 ± 0.0007 assuming $B^+ B^- / B^0 \bar{B}^0$ ratio is 55/45. We rescale to 50/50. Analysis explicitly removes $B^+ \rightarrow \psi(2S) K^+$.

246 ALBRECHT 90J reports $< 1.6 \times 10^{-3}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0594$. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.



$$\Gamma(h_c(1P)K^+ \times B(h_c(1P) \rightarrow J/\psi\pi^+\pi^-))/\Gamma_{\text{total}} \quad \Gamma_{147}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.4 \times 10^{-6}$	90	247 AUBERT	05R BABR	$e^+e^- \rightarrow \gamma(4S)$

247 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(X(3872)K^+)/\Gamma_{\text{total}} \quad \Gamma_{149}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2 \times 10^{-4}$	90	248 AUBERT	06E BABR	$e^+e^- \rightarrow \gamma(4S)$

248 Perform measurements of absolute branching fractions using a missing mass technique.

$$\Gamma(X(3872)K^+ \times B(X \rightarrow J/\psi\pi^+\pi^-))/\Gamma_{\text{total}} \quad \Gamma_{150}/\Gamma$$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
11.4±2.0 OUR AVERAGE			

$10.1 \pm 2.5 \pm 1.0$ 249 AUBERT 06 BABR $e^+e^- \rightarrow \gamma(4S)$

$13.0 \pm 2.9 \pm 0.7$ 250 CHOI 03 BELL $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.8 ± 4.1 249 AUBERT 05R BABR Repl. by AUBERT 06

249 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

250 CHOI 03 reports $[B(B^+ \rightarrow X(3872)K^+) \times B(X \rightarrow J/\psi\pi^+\pi^-)] / B(B^+ \rightarrow \psi(2S)K^+) = 0.0200 \pm 0.0038 \pm 0.0023$. We multiply by our best value $B(B^+ \rightarrow \psi(2S)K^+) = (6.48 \pm 0.35) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(X(3872)K^+ \times B(X(3872) \rightarrow D^0\bar{D}^0))/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.0 \times 10^{-5}$	90	251 CHISTOV	04 BELL	$e^+e^- \rightarrow \gamma(4S)$

251 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(X(3872)K^+ \times B(X(3872) \rightarrow D^+D^-))/\Gamma_{\text{total}}$ Γ_{152}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.0 \times 10^{-5}$	90	252 CHISTOV	04 BELL	$e^+e^- \rightarrow \gamma(4S)$

252 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(X(3872)K^+ \times B(X(3872) \rightarrow D^0\bar{D}^0\pi^0))/\Gamma_{\text{total}}$ Γ_{153}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.0 \times 10^{-5}$	90	253 CHISTOV	04 BELL	$e^+e^- \rightarrow \gamma(4S)$

253 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(X(3872)K^+ \times B(X(3872) \rightarrow J/\psi(1S)\eta))/\Gamma_{\text{total}}$ Γ_{154}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.7 \times 10^{-6}$	90	254 AUBERT	04Y BABR	$e^+e^- \rightarrow \gamma(4S)$

254 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(X(3872)^+K^0 \times B(X(3872)^+ \rightarrow J/\psi(1S)\pi^+\pi^0))/\Gamma_{\text{total}}$ Γ_{155}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<22	90	255 AUBERT	05B BABR	$e^+e^- \rightarrow \gamma(4S)$

255 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. The isovector-X hypothesis is excluded with a likelihood test at 1×10^{-4} level. $\Gamma(Y(4260)^0 K^+ \times B(Y^0 \rightarrow J/\psi\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{156}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<29	95	256 AUBERT	06 BABR	$e^+e^- \rightarrow \gamma(4S)$

256 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(J/\psi(1S)K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{157}/Γ For polarization information see the Listings at the end of the “ B^0 Branching Ratios” section.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.41 ± 0.08 OUR AVERAGE				

1.454 ± 0.047 ± 0.097 257 AUBERT 05J BABR $e^+e^- \rightarrow \gamma(4S)$ 1.28 ± 0.07 ± 0.14 257 ABE 02N BELL $e^+e^- \rightarrow \gamma(4S)$ 1.41 ± 0.23 ± 0.24 257 JESSOP 97 CLE2 $e^+e^- \rightarrow \gamma(4S)$ 1.58 ± 0.47 ± 0.27 258 ABE 96H CDF $p\bar{p}$ at 1.8 TeV1.51 ± 1.08 ± 0.02 259 BORTOLETTO92 CLEO $e^+e^- \rightarrow \gamma(4S)$ 1.86 ± 1.30 ± 0.02 260 ALBRECHT 90J ARG $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.37 ± 0.09 ± 0.11 257 AUBERT 02 BABR Repl. by AUBERT 05J

1.78 ± 0.51 ± 0.23 13 257 ALAM 94 CLE2 Sup. by JESSOP 97

- 257 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
 258 ABE 96H assumes that $B(B^+ \rightarrow J/\psi K^+) = (1.02 \pm 0.14) \times 10^{-3}$.
 259 BORTOLETTO 92 reports $(1.3 \pm 0.9 \pm 0.3) \times 10^{-3}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
 260 ALBRECHT 90J reports $(1.6 \pm 1.1 \pm 0.3) \times 10^{-3}$ for $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.94 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(J/\psi(1S)K^*(892)^+)/\Gamma(J/\psi(1S)K^+)$ $\Gamma_{157}/\Gamma_{146}$

VALUE	DOCUMENT ID	TECN	COMMENT
1.39±0.09 OUR AVERAGE			
1.37±0.05±0.08	AUBERT	05J BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
1.45±0.20±0.17	261 JESSOP	97 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
1.92±0.60±0.17	ABE	96Q CDF	$p\bar{p}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.37±0.10±0.08	262 AUBERT	02 BABR	Repl. by AUBERT 05J
261 JESSOP 97 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The measurement is actually measured as an average over kaon charged and neutral states.			
262 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.			

 $\Gamma(J/\psi(1S)K(1270)^+)/\Gamma_{\text{total}}$ Γ_{158}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.80±0.34±0.39			
263 ABE	01L BELL	$e^+ e^- \rightarrow \Upsilon(4S)$	

263 Uses the PDG value of $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.00 \pm 0.10) \times 10^{-3}$. $\Gamma(J/\psi(1S)K(1400)^+)/\Gamma(J/\psi(1S)K(1270)^+)$ $\Gamma_{159}/\Gamma_{158}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.30	90	ABE	01L BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

 $\Gamma(J/\psi(1S)\eta K^+)/\Gamma_{\text{total}}$ Γ_{160}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
10.8±2.3±2.4			
264 AUBERT	04Y BABR	$e^+ e^- \rightarrow \Upsilon(4S)$	

264 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(J/\psi(1S)\phi K^+)/\Gamma_{\text{total}}$ Γ_{161}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
(5.2±1.7) × 10⁻⁵ OUR AVERAGE Error includes scale factor of 1.2.			
$(4.4 \pm 1.4 \pm 0.5) \times 10^{-5}$	265 AUBERT	030 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$(8.8^{+3.5}_{-3.0} \pm 1.3) \times 10^{-5}$	266 ANASTASSOV	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

265 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.266 ANASTASSOV 00 finds 10 events on a background of 0.5 ± 0.2 . Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$, a uniform Dalitz plot distribution, isotropic $J/\psi(1S)$ and ϕ decays, and $B(B^+ \rightarrow J/\psi(1S)\phi K^+) = B(B^0 \rightarrow J/\psi(1S)\phi K^0)$.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma_{\text{total}}$ Γ_{162}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$(4.9 \pm 0.6) \times 10^{-5}$ OUR FIT			Error includes scale factor of 1.5.
$(3.8 \pm 0.6 \pm 0.3) \times 10^{-5}$	267 ABE	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$

267 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma(J/\psi(1S)K^+)$ $\Gamma_{162}/\Gamma_{146}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.049 ± 0.006 OUR FIT				Error includes scale factor of 1.5.

0.053 ± 0.004 OUR AVERAGE

$0.0537 \pm 0.0045 \pm 0.0011$	AUBERT	04P BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$0.05 \pm 0.019 \pm 0.001$	ABE	96R CDF	$p\bar{p} 1.8 \text{ TeV}$	
0.052 ± 0.024	BISHAI	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.0391 \pm 0.0078 \pm 0.0019$	AUBERT	02F BABR	Repl. by AUBERT 04P	
0.043 ± 0.023	5 268 ALEXANDER	95 CLE2	Sup. by BISHAI 96	

268 Assumes equal production of $B^+ B^-$ and $B^0 \bar{B}^0$ on $\gamma(4S)$.

$\Gamma(J/\psi(1S)\rho^+)/\Gamma_{\text{total}}$ Γ_{163}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.7 \times 10^{-4}$	90	BISHAI	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(J/\psi(1S)a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{164}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-3}$	90	BISHAI	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(J/\psi(1S)p\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{165}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
11.8 ± 3.1 OUR AVERAGE				

$11.7 \pm 2.8 \pm 1.8$	269 XIE	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$	
$12 \pm 9 \pm 6$	269 AUBERT	03K BABR	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<41	90	ZANG	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$
-------	----	------	---------	----------------------------------

269 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)\bar{\Sigma}^0 p)/\Gamma_{\text{total}}$ Γ_{166}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-5}$	90	270 XIE	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$

270 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)D^+)/\Gamma_{\text{total}}$ Γ_{167}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<12	90	271 AUBERT	05U BABR	$e^+ e^- \rightarrow \gamma(4S)$

271 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(J/\psi(1S)\bar{D}^0\pi^+)/\Gamma_{\text{total}}$		Γ_{168}/Γ		
<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.5	90	272 ZHANG	05B BELL	$e^+e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<5.2	90	272 AUBERT	05R BABR	$e^+e^- \rightarrow \gamma(4S)$
272 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

$\Gamma(\psi(2S)K^+)/\Gamma_{\text{total}}$		Γ_{169}/Γ			
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.48 ± 0.35 OUR AVERAGE					
4.9 \pm 1.6 \pm 0.4			273 AUBERT	06E BABR	$e^+e^- \rightarrow \gamma(4S)$
6.17 \pm 0.32 \pm 0.44			274 AUBERT	05J BABR	$e^+e^- \rightarrow \gamma(4S)$
6.9 \pm 0.6			274 ABE	03B BELL	$e^+e^- \rightarrow \gamma(4S)$
7.8 \pm 0.7 \pm 0.9			274 RICHICHI	01 CLE2	$e^+e^- \rightarrow \gamma(4S)$
5.5 \pm 1.0 \pm 0.6			275 ABE	980 CDF	$p\bar{p}$ 1.8 TeV
18 \pm 8 \pm 4	5		274 ALBRECHT	90J ARG	$e^+e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
6.4 \pm 0.5 \pm 0.8			274 AUBERT	02 BABR	Repl. by AUBERT 05J
6.1 \pm 2.3 \pm 0.9	7		274 ALAM	94 CLE2	Repl. by RICHICHI 01
<5	90		274 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
22 \pm 17	3		276 ALBRECHT	87D ARG	$e^+e^- \rightarrow \gamma(4S)$

273 Perform measurements of absolute branching fractions using a missing mass technique.
 274 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.
 275 ABE 980 reports $[B(B^+ \rightarrow \psi(2S)K^+)]/[B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.558 \pm 0.082 \pm 0.056$. We multiply by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (9.9 \pm 1.0) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
 276 ALBRECHT 87D assume $B^+ B^- / B^0 \bar{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.

$\Gamma(\psi(2S)K^+)/\Gamma(J/\psi(1S)K^+)$		$\Gamma_{169}/\Gamma_{146}$		
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.64 \pm 0.06 \pm 0.07$		277 AUBERT	02 BABR	$e^+e^- \rightarrow \gamma(4S)$
277 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

$\Gamma(\psi(2S)K^{*(892)}+)/\Gamma_{\text{total}}$		Γ_{170}/Γ		
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.7 ± 1.4 OUR AVERAGE				
5.92 \pm 0.85 \pm 0.89		278 AUBERT	05J BABR	$e^+e^- \rightarrow \gamma(4S)$
9.2 \pm 1.9 \pm 1.2		278 RICHICHI	01 CLE2	$e^+e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<30	90	278 ALAM	94 CLE2	Repl. by RICHICHI 01
<35	90	278 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
<49	90	278 ALBRECHT	90J ARG	$e^+e^- \rightarrow \gamma(4S)$
278 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

$\Gamma(\psi(2S)K^{*(892)}+)/\Gamma(\psi(2S)K^+)$		$\Gamma_{170}/\Gamma_{169}$		
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.96 \pm 0.15 \pm 0.09$		AUBERT	05J BABR	$e^+e^- \rightarrow \gamma(4S)$

$\Gamma(\psi(2S) K^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{171}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0019±0.0011±0.0004	3	279 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \gamma(4S)$

279 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\psi(3770) K^+)/\Gamma_{\text{total}}$ Γ_{172}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.49±0.13 OUR AVERAGE			

3.5 ± 2.5 ± 0.3	280 AUBERT	06E BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.48±0.11±0.07	281 CHISTOV	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$

280 Perform measurements of absolute branching fractions using a missing mass technique.

281 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\psi(3770) K^+ \times B(\psi(3770) \rightarrow D^0 \bar{D}^0))/\Gamma_{\text{total}}$ Γ_{173}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.34±0.08±0.05	282 CHISTOV	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$

282 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\psi(3770) K^+ \times B(\psi(3770) \rightarrow D^+ D^- K^+))/\Gamma_{\text{total}}$ Γ_{174}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.14±0.08±0.02	283 CHISTOV	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$

283 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\chi_{c0} \pi^+ \times B(\chi_{c0} \rightarrow \pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{175}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.3	90	284 AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$

284 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\chi_{c0}(1P) K^+)/\Gamma_{\text{total}}$ Γ_{176}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.6 +0.5 -0.4 OUR AVERAGE				

1.39±0.49±0.11 285 AUBERT,B 05N BABR $e^+ e^- \rightarrow \gamma(4S)$ 1.96±0.35^{+2.00}_{-0.42} 286 GARMASH 05 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.8 90 287 AUBERT 06E BABR $e^+ e^- \rightarrow \gamma(4S)$ <8.9 90 286 AUBERT 05K BABR $e^+ e^- \rightarrow \gamma(4S)$

2.7 ± 0.7 288 AUBERT 04T BABR Repl. by AUBERT,B 04P

3.0 ± 0.8 ± 0.3 289 AUBERT,B 04P BABR Repl. by AUBERT,B 05N

6.0 +2.1 ± 1.1 290 ABE 02B BELL Repl. by GARMASH 05

<4.8 90 291 EDWARDS 01 CLE2 $e^+ e^- \rightarrow \gamma(4S)$

- 285 AUBERT,B 05N reports $(0.66 \pm 0.22 \pm 0.08) \times 10^{-6}$ for $B(B^+ \rightarrow \chi_c^0 K^+) \times B(\chi_c^0 \rightarrow \pi^+ \pi^-)$. We compute $B(B^+ \rightarrow \chi_c^0 K^+)$ using the PDG value $B(\chi_c^0 \rightarrow \pi^+ \pi^-) = (7.1 \pm 0.6) \times 10^{-3}$ and 2/3 for the $\pi^+ \pi^-$ fraction.
- 286 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.
- 287 Perform measurements of absolute branching fractions using a missing mass technique.
- 288 The measurement performed using decay channels $\chi_c^0 \rightarrow \pi^+ \pi^-$ and $\chi_c^0 \rightarrow K^+ K^-$. The ratio of the branching ratios for these channels is found to be consistent with world average.
- 289 AUBERT 04P reports $B(B^+ \rightarrow \chi_c^0 K^+) \times B(\chi_c^0 \rightarrow \pi^+ \pi^-) = (1.5 \pm 0.4 \pm 0.1) \times 10^{-6}$ and used PDG value of $B(\chi_c^0 \rightarrow \pi\pi) = (7.4 \pm 0.8) \times 10^{-3}$ and Clebsh-Gordan coefficient to compute $B(B^\pm \rightarrow \chi_c^0 K^\pm)$.
- 290 ABE 02B measures the ratio of $B(B^+ \rightarrow \chi_c^0 K^+)/B(B^+ \rightarrow J/\psi(1S) K^+) = 0.60 + 0.21 - 0.18 \pm 0.05 \pm 0.08$, where the third error is due to the uncertainty in the $B(\chi_c^0 \rightarrow \pi^+ \pi^-)$, and uses $B(B^+ \rightarrow J/\psi(1S) K^+) = (10.0 \pm 1.0) \times 10^{-4}$ to obtain the result.
- 291 EDWARDS 01 assumes equal production of B^0 and B^+ at the $\gamma(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \rightarrow \gamma \eta_c)$ in those modes have been accounted for.

$\Gamma(\chi_{c0} K^*(892)^+)/\Gamma_{\text{total}}$					Γ_{177}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.86 \times 10^{-3}$	90	292 AUBERT	05K BABR	$e^+ e^- \rightarrow \gamma(4S)$	

292 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\chi_{c2} K^+)/\Gamma_{\text{total}}$					Γ_{178}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.9 \times 10^{-5}$	90	293 SONI	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.0 \times 10^{-4}$	90	294 AUBERT	06E BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$<3.0 \times 10^{-5}$	90	293 AUBERT	05K BABR	Repl. by AUBERT 06E	

293 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

294 Perform measurements of absolute branching fractions using a missing mass technique.

$\Gamma(\chi_{c2} K^*(892)^+)/\Gamma_{\text{total}}$					Γ_{179}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.2 \times 10^{-5}$	90	295 AUBERT	05K BABR	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.27 \times 10^{-4}$ 90 295 SONI 06 BELL $e^+ e^- \rightarrow \gamma(4S)$

295 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\chi_{c1}(1P) K^+)/\Gamma_{\text{total}}$					Γ_{180}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
5.3 ± 0.7 OUR AVERAGE				Error includes scale factor of 1.7. See the ideogram below.	
8.1 ± 1.4 ± 0.7		296 AUBERT	06E BABR	$e^+ e^- \rightarrow \gamma(4S)$	
4.49 ± 0.19 ± 0.53		297 SONI	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$	
5.79 ± 0.26 ± 0.65		297 AUBERT	05J BABR	$e^+ e^- \rightarrow \gamma(4S)$	
15.5 ± 5.4 ± 2.0		298 ACOSTA	02F CDF	$p\bar{p} 1.8 \text{ TeV}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.8 ± 0.9 ± 0.3	299 AUBERT	02 BABR	Repl. by AUBERT 05J
9.7 ± 4.0 ± 0.9	6 297 ALAM	94 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
19 ± 13 ± 6	300 ALBRECHT	92E ARG	$e^+ e^- \rightarrow \gamma(4S)$


296 Perform measurements of absolute branching fractions using a missing mass technique.

297 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

298 ACOSTA 02F uses as reference of $B(B \rightarrow J/\psi(1S) K^+) = (10.1 \pm 0.6) \times 10^{-4}$. The second error includes the systematic error and the uncertainties of the branching ratio.

299 AUBERT 02 reports $(7.5 \pm 0.9 \pm 0.8) \times 10^{-4}$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$. We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (35.6 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

300 ALBRECHT 92E assumes no $\chi_{c2}(1P)$ production and $B(\gamma(4S) \rightarrow B^+ B^-) = 50\%$.



301 AUBERT 02 reports $0.75 \pm 0.08 \pm 0.05$ for $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$.

We rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (35.6 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\chi_{c1}(1P) K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{181}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

3.6 ± 0.9 OUR AVERAGE

$4.05 \pm 0.59 \pm 0.95$	302	SONI	06	BELL $e^+ e^- \rightarrow \gamma(4S)$
$2.94 \pm 0.95 \pm 0.98$	302	AUBERT	05J	BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	302 ALAM	94	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
-----	----	----------	----	---------------------------------------

302 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\chi_{c1}(1P) K^*(892)^+)/\Gamma(\chi_{c1}(1P) K^+)$ $\Gamma_{181}/\Gamma_{180}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.51 ± 0.17 ± 0.16	AUBERT	05J	BABR $e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(K^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{182}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

2.41 ± 0.17 OUR AVERAGE

Error includes scale factor of 1.4. See the ideogram below.

$2.60 \pm 0.13 \pm 0.10$	303	AUBERT,BE	05E	BABR $e^+ e^- \rightarrow \gamma(4S)$
$2.20 \pm 0.19 \pm 0.11$	303	CHAO	04	BELL $e^+ e^- \rightarrow \gamma(4S)$
$1.88^{+0.37}_{-0.33}{}^{+0.21}_{-0.18}$	303	BORNHEIM	03	CLE2 $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.23 \pm 0.17 \pm 0.11$	303	AUBERT	04M	BABR Repl. by AUBERT,BE 05E
--------------------------	-----	--------	-----	-----------------------------

$1.94^{+0.31}_{-0.30} \pm 0.16$	303	CASEY	02	BELL Repl. by CHAO 04
---------------------------------	-----	-------	----	-----------------------

$1.37^{+0.57}_{-0.48}{}^{+0.19}_{-0.18}$	303	ABE	01H	BELL Repl. by CASEY 02
--	-----	-----	-----	------------------------

$1.82^{+0.33}_{-0.30} \pm 0.20$	303	AUBERT	01E	BABR Repl. by AUBERT 04M
---------------------------------	-----	--------	-----	--------------------------

$1.82^{+0.46}_{-0.40} \pm 0.16$	303	CRONIN-HEN..00	CLE2	Repl. by BORNHEIM 03
---------------------------------	-----	----------------	------	----------------------

$2.3^{+1.1}_{-1.0} \pm 0.36$	GODANG	98	CLE2	Repl. by CRONIN-HENNESSY 00
------------------------------	--------	----	------	-----------------------------

< 4.8	90	ASNER	96	CLE2 Repl. by GODANG 98
-------	----	-------	----	-------------------------


<19	90	ALBRECHT	91B	ARG $e^+ e^- \rightarrow \gamma(4S)$
-----	----	----------	-----	--------------------------------------

<10	90	304 Avery	89B	CLEO $e^+ e^- \rightarrow \gamma(4S)$
-----	----	-----------	-----	---------------------------------------

<68	90	AVERY	87	CLEO $e^+ e^- \rightarrow \gamma(4S)$
-----	----	-------	----	---------------------------------------

303 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

304 Avery 89B reports $< 9 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.



$\Gamma(K^+\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
1.21±0.08 OUR AVERAGE				
1.20±0.07±0.06		305 AUBERT	05L	BABR $e^+e^- \rightarrow \gamma(4S)$
1.20±0.13 ^{+0.13} _{-0.09}		305 CHAO	04	BELL $e^+e^- \rightarrow \gamma(4S)$
1.29 ^{+0.24} _{-0.22} ^{+0.12} _{-0.11}		305 BORNHEIM	03	CLE2 $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.28 ^{+0.12} _{-0.11} ^{±0.10}		305 AUBERT	03L	BABR Repl. by AUBERT 05L
1.3 ^{+0.25} _{-0.24} ^{±0.13}		305 CASEY	02	BELL Repl. by CHAO 04
1.63 ^{+0.35} _{-0.33} ^{+0.16} _{-0.18}		305 ABE	01H	BELL Repl. by CASEY 02
1.08 ^{+0.21} _{-0.19} ^{±0.10}		305 AUBERT	01E	BABR Repl. by AUBERT 03L
1.16 ^{+0.30} _{-0.27} ^{+0.14} _{-0.13}		305 CRONIN-HEN..00	CLE2	Repl. by BORNHEIM 03
<1.6	90	GODANG	98	CLE2 Repl. by CRONIN-HENNESSY 00
<1.4	90	ASNER	96	CLE2 Repl. by GODANG 98

305 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^+\pi^0)/\Gamma(K^0\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
2.38^{+0.98}_{-1.10}^{+0.39}_{-0.26}	ABE	01H	BELL $e^+e^- \rightarrow \gamma(4S)$

$\Gamma_{183}/\Gamma_{182}$

$\Gamma(\eta' K^+)/\Gamma_{\text{total}}$ Γ_{184}/Γ VALUE (units 10^{-5}) **7.05 ± 0.35 OUR AVERAGE**

		DOCUMENT ID	TECN	COMMENT	
6.89 $\pm 0.20 \pm 0.32$	306	AUBERT	05M BABR	$e^+ e^- \rightarrow \gamma(4S)$	
7.9 $^{+1.2}_{-1.1} \pm 0.9$	306	ABE	01M BELL	$e^+ e^- \rightarrow \gamma(4S)$	
8.0 $^{+1.0}_{-0.9} \pm 0.7$	306	RICHICHI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
7.69 $\pm 0.35 \pm 0.44$	306	AUBERT	03W BABR	Repl. by AUBERT 05M	
7.0 $\pm 0.8 \pm 0.5$	306	AUBERT	01G BABR	Repl. by AUBERT 03W	
6.5 $^{+1.5}_{-1.4} \pm 0.9$	BEHRENS	98 CLE2	Repl. by RICHICHI 00		

306 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\eta' K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{185}/Γ VALUECL%

		DOCUMENT ID	TECN	COMMENT	
$<1.4 \times 10^{-5}$	90	307 AUBERT,B	04D BABR	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<3.5 \times 10^{-5}$	90	307 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	

$<1.3 \times 10^{-4}$	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00	
-----------------------	----	---------	---------	----------------------	--


307 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\eta K^+)/\Gamma_{\text{total}}$ Γ_{186}/Γ VALUE (units 10^{-6})CL%

		DOCUMENT ID	TECN	COMMENT	
2.6 ± 0.6 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.	
3.3 $\pm 0.6 \pm 0.3$	308	AUBERT,B	05K BABR	$e^+ e^- \rightarrow \gamma(4S)$	
2.1 $\pm 0.6 \pm 0.2$	308	CHANG	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$	
2.2 $^{+2.8}_{-2.2}$	308	RICHICHI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.4 $\pm 0.8 \pm 0.2$	308	AUBERT	04H BABR	Repl. by AUBERT,B 05K	
<14	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00	

308 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.



$\Gamma(\eta K^*(892)^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
2.6 ±0.4 OUR AVERAGE				
2.56±0.40±0.24	309	AUBERT,B	04D BABR	$e^+ e^- \rightarrow \gamma(4S)$
2.64 ^{+0.96} _{-0.82} ±0.33	309	RICHICHI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.0 90 BEHRENS 98 CLE2 Repl. by RICHICHI 00

309 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\omega K^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
0.51±0.07 OUR AVERAGE				
0.48±0.08±0.04	310	AUBERT	04H BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.65 ^{+0.13} _{-0.12} ±0.06	310	WANG	04A BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.32 ^{+0.24} _{-0.19} ±0.08	310	JESSOP	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.92^{+0.26}_{-0.23}±0.10 310 LU 02 BELL Repl. by WANG 04A
 <0.4 90 310 AUBERT 01G BABR $e^+ e^- \rightarrow \gamma(4S)$
 1.5^{+0.7}_{-0.6}±0.2 310 BERGFELD 98 CLE2 Repl. by JESSOP 00

310 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\omega K^*(892)^+)/\Gamma_{\text{total}}$					Γ_{189}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 7.4	90	311 AUBERT	050 BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<87	90	311 BERGFELD	98 CLE2		
311 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.					

$\Gamma(a_0^0 K^+)/\Gamma_{\text{total}}$					Γ_{191}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2.5	90	312 AUBERT,BE	04 BABR	$e^+ e^- \rightarrow \gamma(4S)$	
312 Assumes equal production of charged and neutral B mesons from $\gamma(4S)$ decays.					

$\Gamma(a_0^+ K^0)/\Gamma_{\text{total}}$					Γ_{190}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<3.9	90	313 AUBERT,BE	04 BABR	$e^+ e^- \rightarrow \gamma(4S)$	
313 Assumes equal production of charged and neutral B mesons from $\gamma(4S)$ decays.					

$\Gamma(K^*(892)^0 \pi^+)/\Gamma_{\text{total}}$					Γ_{192}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.16±0.19 OUR AVERAGE				Error includes scale factor of 1.8.	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
1.35±0.12 ^{+0.08} _{-0.09}		314 AUBERT,B	05N BABR	$e^+ e^- \rightarrow \gamma(4S)$	
0.98±0.09 ^{+0.11} _{-0.12}		314 GARMASH	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$	
1.55±0.18 ^{+0.15} _{-0.40}		314,315 AUBERT,B	04P BABR	Repl. by AUBERT,B 05N	
1.94 ^{+0.42} _{-0.39} ^{+0.41} _{-0.71}		316 GARMASH	02 BELL	Repl. by GARMASH 05	
<11.9	90	317 ABE	00C SLD	$e^+ e^- \rightarrow Z$	
< 1.6	90	314 JESSOP	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
<39	90	318 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$	
< 4.1	90	ASNER	96 CLE2	Repl. by JESSOP 00	
<48	90	319 ABREU	95N DLPH	Sup. by ADAM 96D	
<17	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \gamma(4S)$	
<15	90	320 Avery	89B CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
<26	90	AVERY	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	

314 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.	
315 AUBERT 04P also report a branching ratio for $B^+ \rightarrow$ "higher K^* resonances" π^+ , $K^* \rightarrow K^+ \pi^-$, $(25.1 \pm 2.0^{+11.0}_{-5.7}) \times 10^{-6}$.	
316 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.	
317 ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.	
318 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.	
319 Assumes a B^0, B^- production fraction of 0.39 and a B_s production fraction of 0.12.	
320 Avery 89B reports $< 1.3 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.	

$\Gamma(K^*(892)^+\pi^0)/\Gamma_{\text{total}}$				Γ_{193}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$6.9 \pm 2.0 \pm 1.3$		321 AUBERT	05X BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<31	90	321 JESSOP	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<99	90	ASNER	96 CLE2	Repl. by JESSOP 00

321 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+\pi^-\pi^+)/\Gamma_{\text{total}}$				Γ_{194}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
5.6 ± 0.9 OUR AVERAGE			Error includes scale factor of 2.6.	
$6.41 \pm 0.24 \pm 0.40$	322 AUBERT,B	05N BABR	$e^+e^- \rightarrow \Upsilon(4S)$	
$4.66 \pm 0.21 \pm 0.43$	322 GARMASH	05 BELL	$e^+e^- \rightarrow \Upsilon(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5.36 \pm 0.31 \pm 0.51$	322 GARMASH	04 BELL	Repl. by GARMASH 05	
$5.91 \pm 0.38 \pm 0.32$	323 AUBERT	03M BABR	Repl. by AUBERT,B 05N	
$5.56 \pm 0.58 \pm 0.77$	324 GARMASH	02 BELL	Repl. by GARMASH 04	
322 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				
323 Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.				
324 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0\pi^+$ and $\bar{D}^0 \rightarrow K^+\pi^-$ with $B(B^+ \rightarrow \bar{D}^0\pi^+) \cdot B(\bar{D}^0 \rightarrow K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.				

$\Gamma(K^+\pi^-\pi^+\text{nonresonant})/\Gamma_{\text{total}}$				Γ_{195}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.31^{+0.10}_{-0.08}$ OUR AVERAGE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.29 \pm 0.06^{+0.08}_{-0.05}$	325 AUBERT,B	05N BABR	$e^+e^- \rightarrow \Upsilon(4S)$	
$1.73 \pm 0.17^{+1.72}_{-0.80}$	325 GARMASH	05 BELL	$e^+e^- \rightarrow \Upsilon(4S)$	
325 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				
326 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.				
327 Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.				
328 Avery 89B reports $< 1.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.				

$\Gamma(K^+ f_0(980) \times B(f_0 \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{196}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
8.9 ± 1.0 OUR AVERAGE				
9.47 ± 0.97 $^{+0.62}_{-0.88}$	329	AUBERT,B	05N BABR	$e^+ e^- \rightarrow \gamma(4S)$
7.55 ± 1.24 $^{+1.63}_{-1.18}$	329	GARMASH	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9.2 ± 1.2 $^{+2.1}_{-2.6}$	330	AUBERT,B	04P BABR	Repl. by AUBERT,B 05N
9.6 $^{+2.5}_{-2.3}$ $^{+3.7}_{-1.7}$	331	GARMASH	02 BELL	Repl. by GARMASH 05
<80	90	332	AVERY	89B CLEO $e^+ e^- \rightarrow \gamma(4S)$
329 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				
330 AUBERT,B 04P also reports $B(B^+ \rightarrow \text{"higher } f^0 \text{ resonances"} \pi^+, f(980)^0 \rightarrow \pi^+ \pi^-) = (3.2 \pm 1.2)^{+6.0}_{-2.9} \times 10^{-6}$.				
331 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \times B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$. Only charged pions from the $f_0(980)$ are used.				
332 Avery 89B reports $< 7 \times 10^{-5}$ assuming the $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.				

 $\Gamma(f_2(1270)^0 K^+)/\Gamma_{\text{total}}$ Γ_{197}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.3 × 10⁻⁶	90	333	GARMASH 05 BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.6 × 10 ⁻⁵	90	334	AUBERT,B 05N BABR	$e^+ e^- \rightarrow \gamma(4S)$
333 GARMASH 05 reports 1.3×10^{-6} at 90% CL for $B(B^+ \rightarrow f_2(1270) \pi^+) \times B(f_2(1270) \rightarrow \pi^+ \pi^-)$. We rescaled it using the PDG value $B(f_2(1270) \rightarrow \pi\pi) = 84.7\%$ and 2/3 for the $\pi^+ \pi^-$ mode.				
334 AUBERT,B 05N reports 8.9×10^{-6} at 90% CL for $B(B^+ \rightarrow f_2(1270) \pi^+) \times B(f_2(1270) \rightarrow \pi^+ \pi^-)$. We rescaled it using the PDG value $B(f_2(1270) \rightarrow \pi\pi) = 84.7\%$ and 2/3 for the $K^+ \pi^-$ fraction.				

 $\Gamma(f_0^*(1370)^0 K^+ \times B(f_0^*(1370)^0 \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{198}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<10.7 × 10⁻⁶	90	335	AUBERT,B 05N BABR	$e^+ e^- \rightarrow \gamma(4S)$
335 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

 $\Gamma(\rho^0(1450) K^+ \times B(\rho^0(1450) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{199}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<11.7 × 10⁻⁶	90	336	AUBERT,B 05N BABR	$e^+ e^- \rightarrow \gamma(4S)$
336 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

 $\Gamma(f_0(1500) K^+ \times B(f_0(1500) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{200}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.4 × 10⁻⁶	90	337	AUBERT,B 05N BABR	$e^+ e^- \rightarrow \gamma(4S)$
337 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

$\Gamma(f'_2(1525) K^+ \times B(f'_2(1525) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{201}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.4 \times 10^{-6}$	90	338 AUBERT,B	05N BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

338 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K^+ \rho^0) / \Gamma_{\text{total}}$ Γ_{202}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
5.0 ± 0.7 OUR AVERAGE				

$5.07 \pm 0.75^{+0.55}_{-0.88}$ 339 AUBERT,B 05N BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$4.78 \pm 0.75^{+1.01}_{-0.97}$ 339 GARMASH 05 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.2	90	340 AUBERT,B	04P BABR	Repl. by AUBERT,B 05N
< 12	90	341 GARMASH	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
< 86	90	342 ABE	00C SLD	$e^+ e^- \rightarrow Z$
< 17	90	339 JESSOP	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
< 120	90	343 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
< 19	90	ASNER	96 CLE2	Repl. by JESSOP 00
< 190	90	344 ABREU	95N DLPH	Sup. by ADAM 96D
< 180	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
< 80	90	345 AVERY	89B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
< 260	90	AVERY	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

339 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

340 AUBERT 04P reports a central value of $(3.9 \pm 1.2^{+1.3}_{-3.5}) \times 10^{-6}$ for this branching ratio.

341 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

342 ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

343 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

344 Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

345 AVERY 89B reports $< 7 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

 $\Gamma(K_0^*(1430)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{203}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
38 ± 5 OUR AVERAGE			

$36.6 \pm 1.8 \pm 4.7$ 346 AUBERT,B 05N BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$45.0 \pm 2.9^{+15.0}_{-10.7}$ 346 GARMASH 05 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

346 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K_2^*(1430)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{204}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.9 \times 10^{-6}$	90	347 GARMASH	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<2.3 \times 10^{-5}$	90	348 AUBERT,B	05N BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$<6.8 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
347 GARMASH 05 reports 2.3×10^{-6} at 90% CL for $B(B^+ \rightarrow K_2^*(1430)^0 \pi^+) \times B(K_2^*(1430)^0 \rightarrow K^+ \pi^-)$. We rescaled it using the PDG value $B(K_2^*(1430)^0 \rightarrow K\pi) = 49.9\%$ and 2/3 for the $K^+ \pi^-$ mode.				
348 AUBERT,B 05N reports 7.7×10^{-6} at 90% CL for $B(B^+ \rightarrow K_2^*(1430)^0 \pi^+) \times B(K_2^*(1430)^0 \rightarrow K^+ \pi^-)$. We rescaled it using the PDG value $B(K_2^*(1430)^0 \rightarrow K\pi) = 49.9\%$ and 2/3 for the $K^+ \pi^-$ fraction.				

 $\Gamma(K^*(1410)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{205}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.5 \times 10^{-5}$	90	349 GARMASH	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
349 GARMASH 05 reports 2.0×10^{-6} at 90% CL for $B(B^+ \rightarrow K^*(1410)^0 \pi^+) \times B(K^*(1410)^0 \rightarrow K^+ \pi^-)$. We rescaled it using the PDG value $B(K^*(1410)^0 \rightarrow K\pi) = 6.6\%$ and 2/3 for the $K^+ \pi^-$ mode.				

 $\Gamma(K^*(1680)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{206}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-5}$	90	350 GARMASH	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<1.5 \times 10^{-5}$	90	351 AUBERT,B	05N BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
350 GARMASH 05 reports 3.1×10^{-6} at 90% CL for $B(B^+ \rightarrow K^*(1680)^0 \pi^+) \times B(K^*(1680)^0 \rightarrow K^+ \pi^-)$. We rescaled it using the PDG value $B(K^*(1680)^0 \rightarrow K\pi) = 38.7\%$ and 2/3 for the $K^+ \pi^-$ mode.				
351 AUBERT,B 05N reports 3.8×10^{-6} at 90% CL for $B(B^+ \rightarrow K^*(1680)^0 \pi^+) \times B(K^*(1680)^0 \rightarrow K^+ \pi^-)$. We rescaled it using the PDG value $B(K^*(1680)^0 \rightarrow K\pi) = 38.7\%$ and 2/3 for the $K^+ \pi^-$ fraction.				

 $\Gamma(K^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_{207}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	90	352 AUBERT	03M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<4.5 \times 10^{-6}$	90	353 GARMASH	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$<7.0 \times 10^{-6}$	90	354 GARMASH	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
352 Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.				
353 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				
354 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.				

$\Gamma(K^-\pi^+\pi^+ \text{nonresonant})/\Gamma_{\text{total}}$				Γ_{208}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.6 \times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$\Gamma(K_1(1400)^0\pi^+)/\Gamma_{\text{total}}$				Γ_{209}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.6 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$\Gamma(K^0\pi^+\pi^0)/\Gamma_{\text{total}}$				Γ_{210}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<66 \times 10^{-6}$	90	355 ECKHART	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
355 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				
$\Gamma(K^0\rho^+)/\Gamma_{\text{total}}$				Γ_{211}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<4.8 \times 10^{-5}$	90	ASNER	96 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$\Gamma(K^*(892)^+\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_{212}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.1 \times 10^{-3}$	90	ALBRECHT	91E ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$\Gamma(K^*(892)^+\rho^0)/\Gamma_{\text{total}}$				Γ_{213}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$10.6^{+3.0}_{-2.6} \pm 2.4$	356 AUBERT	03V BABR	$e^+e^- \rightarrow \Upsilon(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 74	90	357 GODANG	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<900	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
356 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				
357 Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 4.9×10^{-5} .				
$\Gamma(K^*(892)^0\rho^+)/\Gamma_{\text{total}}$				Γ_{214}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.9 \pm 1.7 \pm 1.2$	358 ZHANG	05D BELL	$e^+e^- \rightarrow \Upsilon(4S)$	
358 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				
$\Gamma(K^*(892)^+K^*(892)^0)/\Gamma_{\text{total}}$				Γ_{215}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.1 \times 10^{-5}$	90	359 GODANG	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
359 Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 4.8×10^{-5} .				
$\Gamma(K_1(1400)^+\rho^0)/\Gamma_{\text{total}}$				Γ_{216}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.8 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_2^*(1430)^+\rho^0)/\Gamma_{\text{total}}$		Γ_{217}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(K^+\bar{K}^0)/\Gamma_{\text{total}}$		Γ_{218}/Γ		
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
1.20 ± 0.32 OUR AVERAGE				

$1.0 \pm 0.4 \pm 0.1$	360	ABE	05G BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.5 \pm 0.5 \pm 0.1$	360	AUBERT,BE	05E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 2.5	90	360 AUBERT	04M BABR	Repl. by AUBERT,BE 05E
< 3.3	90	360 CHAO	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
< 3.3	90	360 BORNHEIM	03 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
< 2.0	90	360 CASEY	02 BELL	Repl. by CHAO 04
< 5.0	90	360 ABE	01H BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
< 2.4	90	360 AUBERT	01E BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
< 5.1	90	360 CRONIN-HEN..00	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
< 21	90	GODANG	98 CLE2	Repl. by CRONIN-HENNESSY 00

360 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{K}^0 K^+ \pi^0)/\Gamma_{\text{total}}$		Γ_{219}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<24 \times 10^{-6}$	90	361 ECKHART	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

361 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ K_S^0 K_S^0)/\Gamma_{\text{total}}$		Γ_{220}/Γ		
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
11.5 ± 1.3 OUR AVERAGE				
$10.7 \pm 1.2 \pm 1.0$	362	AUBERT,B	04V BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$13.4 \pm 1.9 \pm 1.5$	362	GARMASH	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

362 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K_S^0 K_S^0 \pi^+)/\Gamma_{\text{total}}$		Γ_{221}/Γ		
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.2	90	363 GARMASH	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

363 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$		Γ_{222}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.3 \times 10^{-6}$	90	364 AUBERT	03M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

$< 13 \times 10^{-6}$	90	365 GARMASH	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$< 1.2 \times 10^{-5}$	90	366 GARMASH	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
364 Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.				
365 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				
366 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.				

$\Gamma(K^+ K^- \pi^+ \text{nonresonant})/\Gamma_{\text{total}}$ Γ_{223}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.5 \times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

 $\Gamma(K^+ K^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{224}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.3 \times 10^{-6}$	90	367 AUBERT	03M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<2.4 \times 10^{-6}$	90	368 GARMASH	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$<3.2 \times 10^{-6}$	90	369 GARMASH	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

367 Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

368 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

369 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0 \pi^+$ and $\bar{D}^0 \rightarrow K^+ \pi^-$ with $B(B^+ \rightarrow \bar{D}^0 \pi^+) \cdot B(\bar{D}^0 \rightarrow K^+ \pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

 $\Gamma(K^+ K^+ \pi^- \text{nonresonant})/\Gamma_{\text{total}}$ Γ_{225}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<8.79 \times 10^{-5}$	90	ABBIENDI	00B OPAL	$e^+ e^- \rightarrow Z$

 $\Gamma(K^+ K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{226}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.3 \times 10^{-6}$	90	370 JESSOP	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<1.29 \times 10^{-4}$	90	ABBIENDI	00B OPAL	$e^+ e^- \rightarrow Z$
$<1.38 \times 10^{-4}$	90	371 ABE	00C SLD	$e^+ e^- \rightarrow Z$

370 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

371 ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

 $\Gamma(K^+ f_J(2220))/\Gamma_{\text{total}}$ Γ_{227}/Γ

<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	372 HUANG	03 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
372 No evidence is found for such decay and set a limit on $B(B^+ \rightarrow f_J(2220)) \times B(f_J(2220) \rightarrow \phi\phi) < 1.2 \times 10^{-6}$ at 90%CL where the $f_J(2220)$ is a possible glueball state.			
3.06 $\pm 0.12 \pm 0.23$	373 GARMASH	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

$3.01 \pm 0.19 \text{ OUR AVERAGE}$

$2.96 \pm 0.21 \pm 0.16$	374 AUBERT	03M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
3.28 $\pm 0.18 \pm 0.28$	373 GARMASH	04 BELL	Repl. by GARMASH 05
3.53 $\pm 0.37 \pm 0.45$	375 GARMASH	02 BELL	Repl. by GARMASH 04
<20	376 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$
<32	377 ABREU	95N DLPH	Sup. by ADAM 96D
<35	ALBRECHT	91E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

- 373 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
 374 Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.
 375 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0\pi^+$ and $\bar{D}^0 \rightarrow K^+\pi^-$ with $B(B^+ \rightarrow \bar{D}^0\pi^+) \cdot B(\bar{D}^0 \rightarrow K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.
 376 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.
 377 Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

$\Gamma(K^+\phi)/\Gamma_{\text{total}}$ Γ_{229}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
9.0 ±0.8 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
7.6 ±1.3 ±0.6	378	ACOSTA	05J CDF	$p\bar{p}$ at 1.96 TeV
9.60±0.92 ^{+1.05} _{-0.85}	379	GARMASH	05 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
10.0 ±0.9 ±0.5	379	AUBERT	04A BABR	$e^+e^- \rightarrow \Upsilon(4S)$
5.5 ±2.1 ±0.6	379	BRIERE	01 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9.4 ±1.1 ±0.7	379	CHEN	03B BELL	Repl. by GARMASH 05
14.6 ±3.0 ±2.0	380	GARMASH	02 BELL	Repl. by CHEN 03B
7.7 ±1.6 ±0.8	379	AUBERT	01D BABR	$e^+e^- \rightarrow \Upsilon(4S)$
<144	90	381 ABE	00C SLD	$e^+e^- \rightarrow Z$
< 5	90	379 BERGFELD	98 CLE2	
<280	90	382 ADAM	96D DLPH	$e^+e^- \rightarrow Z$
< 12	90	ASNER	96 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<440	90	383 ABREU	95N DLPH	Sup. by ADAM 96D
<180	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 90	90	384 Avery	89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<210	90	AVERY	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

378 Uses $B(B^+ \rightarrow J/\psi K^+) = (1.00 \pm 0.04) \times 10^{-3}$ and $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

379 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.


380 Uses a reference decay mode $B^+ \rightarrow \bar{D}^0\pi^+$ and $\bar{D}^0 \rightarrow K^+\pi^-$ with $B(B^+ \rightarrow \bar{D}^0\pi^+) \cdot B(\bar{D}^0 \rightarrow K^+\pi^-) = (20.3 \pm 2.0) \times 10^{-5}$.

381 ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7 \pm 1.8)\%$ and $f_{B_s} = (10.5 \pm 1.8)\%$.

382 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

383 Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

384 Avery 89B reports $< 8 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.



$$\Gamma(f_0(980)K^+ \times B(f_0(980) \rightarrow K^+ K^-))/\Gamma_{\text{total}} \quad \Gamma_{230}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.9 \times 10^{-6}$	90	385 GARMASH	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$

385 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(a_2(1320)K^+ \times B(a_2(1320) \rightarrow K^+ K^-))/\Gamma_{\text{total}} \quad \Gamma_{231}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-6}$	90	386 GARMASH	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$

386 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(f'_2(1525)K^+ \times B(f'_2(1525) \rightarrow K^+ K^-))/\Gamma_{\text{total}} \quad \Gamma_{232}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.9 \times 10^{-6}$	90	387 GARMASH	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$

387 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(\phi(1680)K^+ \times B(\phi(1680) \rightarrow K^+ K^-))/\Gamma_{\text{total}} \quad \Gamma_{233}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.8 \times 10^{-6}$	90	388 GARMASH	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$

388 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^+ K^- K^+ \text{nonresonant})/\Gamma_{\text{total}}$ Γ_{234}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.40 \pm 0.15^{+0.26}_{-0.60}$		389 GARMASH	05 BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.8	90	BERGFELD	96B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
------	----	----------	----------	----------------------------------

389 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K^*(892)^+ K^+ K^-)/\Gamma_{\text{total}}$ Γ_{235}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.6 \times 10^{-3}$	90	ALBRECHT	91E ARG	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(K^*(892)^+ \phi)/\Gamma_{\text{total}}$ Γ_{236}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$9.6 \pm 3.0 \text{ OUR AVERAGE}$				Error includes scale factor of 1.9.

$12.7^{+2.2}_{-2.0} \pm 1.1$		390 AUBERT	03V BABR	$e^+ e^- \rightarrow \gamma(4S)$
------------------------------	--	------------	----------	----------------------------------

$6.7^{+2.1}_{-1.9} + 0.7_{-1.0}$		390 CHEN	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
----------------------------------	--	----------	----------	----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.7^{+4.2}_{-3.4} \pm 1.7$		390 AUBERT	01D BABR	Repl. by AUBERT 03v
-----------------------------	--	------------	----------	---------------------

< 22.5	90	390 BRIERE	01 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
--------	----	------------	---------	----------------------------------

< 41	90	390 BERGFELD	98 CLE2	
------	----	--------------	---------	--

< 70	90	ASNER	96 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
------	----	-------	---------	----------------------------------

<1300	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \gamma(4S)$
-------	----	----------	---------	----------------------------------

390 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(K_1(1400)^+ \phi)/\Gamma_{\text{total}}$ Γ_{237}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.1 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(K_2^*(1430)^+ \phi)/\Gamma_{\text{total}}$ Γ_{238}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.4 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma(K^+ \phi \phi)/\Gamma_{\text{total}}$ Γ_{239}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.6^{+1.1}_{-0.9} \pm 0.3$		391 HUANG	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$

391 Assumes equal production of B^0 and B^+ at the $\gamma(4S)$ and for a $\phi\phi$ invariant mass below $2.85 \text{ GeV}/c^2$.

 $\Gamma(K^*(892)^+ \gamma)/\Gamma_{\text{total}}$ Γ_{240}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.03 \pm 0.26 \text{ OUR AVERAGE}$				

3.87 $\pm 0.28 \pm 0.26$		392 AUBERT,BE	04A BABR	$e^+ e^- \rightarrow \gamma(4S)$
--------------------------	--	---------------	----------	----------------------------------

4.25 $\pm 0.31 \pm 0.24$		393 NAKAO	04 BELL	$e^+ e^- \rightarrow \gamma(4S)$
--------------------------	--	-----------	---------	----------------------------------

$3.76^{+0.89}_{-0.83} \pm 0.28$		393 COAN	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
---------------------------------	--	----------	---------	----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.83 \pm 0.62 \pm 0.22$	393 AUBERT	02c BABR	Repl. by AUBERT,BE 04A
$5.7 \pm 3.1 \pm 1.1$	394 AMMAR	93 CLE2	Repl. by COAN 00
< 55	90 395 ALBRECHT	89G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
< 55	90 395 AVERY	89B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
< 180	90 AVERY	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

392 Uses the production ratio of charged and neutral B from $\Upsilon(4S)$ decays $R^{+}/0 = 1.006 \pm 0.048$.

393 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

394 AMMAR 93 observed 4.1 ± 2.3 events above background.

395 Assumes the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$.

$\Gamma(K_1(1270)^+\gamma)/\Gamma_{\text{total}}$

Γ_{241}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$4.3 \pm 0.9 \pm 0.9$		396 YANG	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9.9	90 396 NISHIDA	02 BELL	Repl. by YANG 05
< 730	90 397 ALBRECHT	89G ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

396 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

397 ALBRECHT 89G reports < 0.0066 assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\eta K^+\gamma)/\Gamma_{\text{total}}$

Γ_{242}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$8.4 \pm 1.5 \pm 1.2$	398,399 NISHIDA	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

398 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

399 $m_{\eta K} < 2.4 \text{ GeV}/c^2$

$\Gamma(\phi K^+\gamma)/\Gamma_{\text{total}}$

Γ_{243}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$3.4 \pm 0.9 \pm 0.4$	400 DRUTSKOY	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

400 Assumes equal production of B^+ and B^0 at $\Upsilon(4S)$.

$\Gamma(K^+\pi^-\pi^+\gamma)/\Gamma_{\text{total}}$

Γ_{244}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$2.50 \pm 0.18 \pm 0.22$	401 YANG	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.4 \pm 0.5 \pm 0.4$	401,402 NISHIDA	02 BELL	Repl. by YANG 05
-----------------------	-----------------	---------	------------------

401 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

402 $M_{K\pi\pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K^*(892)^0\pi^+\gamma)/\Gamma_{\text{total}}$	Γ_{245}/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$(2.0^{+0.7}_{-0.6} \pm 0.2) \times 10^{-5}$	403,404	NISHIDA	02 BELL $e^+e^- \rightarrow \Upsilon(4S)$

403 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.404 $M_{K\pi\pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K^+\rho^0\gamma)/\Gamma_{\text{total}}$	Γ_{246}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.0 \times 10^{-5}$	90	405,406	NISHIDA	02 BELL $e^+e^- \rightarrow \Upsilon(4S)$

405 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.406 $M_{K\pi\pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K^+\pi^-\pi^+\gamma \text{ nonresonant})/\Gamma_{\text{total}}$	Γ_{247}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9.2 \times 10^{-6}$	90	407,408	NISHIDA	02 BELL $e^+e^- \rightarrow \Upsilon(4S)$

407 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.408 $M_{K\pi\pi} < 2.4 \text{ GeV}/c^2$.

$\Gamma(K_1(1400)^+\gamma)/\Gamma_{\text{total}}$	Γ_{248}/Γ				
<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.5 \times 10^{-5}$	90	409	YANG	05 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $<5.0 \times 10^{-5}$ 90 409 NISHIDA 02 BELL Repl. by YANG 05 <0.0022 90 410 ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$ 409 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.410 ALBRECHT 89G reports < 0.0020 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_2^*(1430)^+\gamma)/\Gamma_{\text{total}}$	Γ_{249}/Γ			
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.45^{+0.40 \pm 0.15}$	411	AUBERT,B	04U BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 <140 90 412 ALBRECHT 89G ARG $e^+e^- \rightarrow \Upsilon(4S)$ 411 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.412 ALBRECHT 89G reports < 0.0013 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(1680)^+\gamma)/\Gamma_{\text{total}}$	Γ_{250}/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0019	90	413	ALBRECHT 89G ARG	$e^+e^- \rightarrow \Upsilon(4S)$

413 ALBRECHT 89G reports < 0.0017 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_3^*(1780)^+\gamma)/\Gamma_{\text{total}}$				Γ_{251}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.9 \times 10^{-5}$	90	414,415 NISHIDA	05 BELL	$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0055	90	416 ALBRECHT	89G ARG	$e^+e^- \rightarrow \Upsilon(4S)$
---------	----	--------------	---------	-----------------------------------

414 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

415 Uses $B(K_3^*(1780) \rightarrow \eta K) = 0.11^{+0.05}_{-0.04}$.

416 ALBRECHT 89G reports < 0.005 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_4^*(2045)^+\gamma)/\Gamma_{\text{total}}$				Γ_{252}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0099	90	417 ALBRECHT	89G ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

417 ALBRECHT 89G reports < 0.0090 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\rho^+\gamma)/\Gamma_{\text{total}}$				Γ_{253}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.8 \times 10^{-6}$	90	418 AUBERT	05 BABR	$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.2 $\times 10^{-6}$	90	418 MOHAPATRA	05 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------	----	---------------	---------	-----------------------------------

<2.1 $\times 10^{-6}$	90	418 AUBERT	04C BABR	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------	----	------------	----------	-----------------------------------

<1.3 $\times 10^{-5}$	90	418,419 COAN	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------	----	--------------	---------	-----------------------------------

418 Assumes equal production of B^+ and B^0 at $\Upsilon(4S)$.

419 No evidence for a nonresonant $K\pi\gamma$ contamination was seen; the central value assumes no contamination.

$\Gamma(\pi^+\pi^0)/\Gamma_{\text{total}}$				Γ_{254}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
0.55 \pm 0.06 OUR AVERAGE					

0.58 \pm 0.06 \pm 0.04		420 AUBERT	05L BABR	$e^+e^- \rightarrow \Upsilon(4S)$
----------------------------	--	------------	----------	-----------------------------------

0.50 \pm 0.12 \pm 0.05		420 CHAO	04 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
----------------------------	--	----------	---------	-----------------------------------

0.46 $^{+0.18}_{-0.16}$ $^{+0.06}_{-0.07}$		420 BORNHEIM	03 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
--	--	--------------	---------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.55^{+0.10}_{-0.19} \pm 0.06$		420 AUBERT	03L BABR	Repl. by AUBERT 05L
---------------------------------	--	------------	----------	---------------------

$0.74^{+0.23}_{-0.22} \pm 0.09$		420 CASEY	02 BELL	Repl. by CHAO 04
---------------------------------	--	-----------	---------	------------------

< 1.34	90	420 ABE	01H BELL	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	---------	----------	-----------------------------------

< 0.96	90	420 AUBERT	01E BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	------------	----------	-----------------------------------

< 1.27	90	420 CRONIN-HEN..00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	--------------------	------	-----------------------------------

< 2.0	90	GODANG	98 CLE2	Repl. by CRONIN-HENNESSY 00
-------	----	--------	---------	-----------------------------

< 1.7	90	ASNER	96 CLE2	Repl. by GODANG 98
-------	----	-------	---------	--------------------

< 24	90	420 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
------	----	--------------	---------	-----------------------------------

<230	90	421 BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
------	----	-----------	---------	-----------------------------------

420 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

421 BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$.

$\Gamma(\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{255}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$16.2 \pm 1.2 \pm 0.9$		422 AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.9 \pm 3.3 \pm 1.6$	422	AUBERT	03M BABR	Repl. by AUBERT 05G	
< 130	90	423 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$	
< 220	90	424 ABREU	95N DLPH	Sup. by ADAM 96D	
< 450	90	425 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$	
< 190	90	426 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	

422 Assumes equal production of B^0 and B^+ at the $\gamma(4S)$; charm and charmonium contributions are subtracted, otherwise no assumptions about intermediate resonances.

423 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

424 Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

425 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$.

426 BORTOLETTO 89 reports $< 1.7 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(\rho^0\pi^+)/\Gamma_{\text{total}}$ Γ_{256}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.87 ± 0.11 OUR AVERAGE					
$0.88 \pm 0.10 \pm 0.09$	427	AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$	
$0.80 \pm 0.23 \pm 0.07$	427	GORDON	02 BELL	$e^+ e^- \rightarrow \gamma(rS)$	
$1.04 \pm 0.33 \pm 0.21$	427	JESSOP	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.95 \pm 0.11 \pm 0.09$	427	AUBERT	04Z BABR	Repl. by AUBERT 05G	
< 8.3	90	428 ABE	00C SLD	$e^+ e^- \rightarrow Z$	
< 16	90	429 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$	
< 4.3	90	ASNER	96 CLE2	Repl. by JESSOP 00	
< 26	90	430 ABREU	95N DLPH	Sup. by ADAM 96D	
< 15	90	427 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$	
< 17	90	431 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
< 23	90	431 BEBEK	87 CLEO	$e^+ e^- \rightarrow \gamma(4S)$	
< 60	90	GILES	84 CLEO	Repl. by BEBEK 87	

427 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

428 ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7 \pm 1.8)\%$ and $f_{B_s} = (10.5 \pm 1.8)\%$.

429 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

430 Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

431 Papers assume the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

 $[\Gamma(K^*(892)^0\pi^+) + \Gamma(\rho^0\pi^+)/\Gamma_{\text{total}}$ $(\Gamma_{192} + \Gamma_{256})/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
--------------	--------------------	-------------	----------------	--

$(17 \pm 12 \pm 2) \times 10^{-5}$	432 ADAM	96D DLPH	$e^+ e^- \rightarrow Z$	
--	----------	----------	-------------------------	--

432 ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

$\Gamma(\pi^+ f_0(980) \times B(f_0(980) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{257}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-6}$	90	433 AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<1.4 \times 10^{-4}$	90	434 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
433 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				
434 BORTOLETTO 89 reports $< 1.2 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.				

 $\Gamma(\pi^+ f_2(1270)) / \Gamma_{\text{total}}$ Γ_{258}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$8.2 \pm 2.1 \pm 1.4$	435,436	AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<240	90	437 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
435 Reported $B(B^+ \rightarrow f_2(1270)\pi^+) \times B(f_2(1270) \rightarrow \pi^+ \pi^-) = (2.3 \pm 0.6 \pm 0.4) \times 10^{-6}$ and rescaled using $B(f_2(1270) \rightarrow \pi^+ \pi^-) = 0.28$.				
436 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				
437 BORTOLETTO 89 reports $< 2.1 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.				

 $\Gamma(\rho(1450)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{259}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.3	90	438 AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$
438 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

 $\Gamma(f_0(1370)\pi^+ \times B(f_0(1370) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{260}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.0	90	439 AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$
439 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

 $\Gamma(f_0(600)\pi^+ \times B(f_0(600) \rightarrow \pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{261}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.1	90	440 AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$
440 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

 $\Gamma(\pi^+ \pi^- \pi^+ \text{nonresonant}) / \Gamma_{\text{total}}$ Γ_{262}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.6 \times 10^{-6}$	90	441 AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<4.1 \times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+ e^- \rightarrow \gamma(4S)$
441 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

 $\Gamma(\pi^+ \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{263}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.9 \times 10^{-4}$	90	442 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$
442 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\gamma(4S)$.				

$\Gamma(\rho^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{264}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
12.0 ± 1.9 OUR AVERAGE				
$13.2 \pm 2.3^{+1.4}_{-1.9}$	443	ZHANG	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$
$10.9 \pm 1.9 \pm 1.9$	443	AUBERT	04Z BABR	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 43	90	443,444 JESSOP	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
< 77	90	ASNER	96 CLE2	Repl. by JESSOP 00
< 550	90	443 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$

443 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.444 Assumes no nonresonant contributions of $B^+ \rightarrow \pi^+ \pi^0 \pi^0$. $\Gamma(\pi^+ \pi^- \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{265}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 4.0 \times 10^{-3}$	90	445 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$

445 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\gamma(4S)$. $\Gamma(\rho^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{266}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.6 ± 0.6 OUR AVERAGE				
$2.25^{+0.57}_{-0.54} \pm 0.58$	446	AUBERT	03V BABR	$e^+ e^- \rightarrow \gamma(4S)$
$3.17 \pm 0.71^{+0.38}_{-0.67}$	447	ZHANG	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<100 90 448 ALBRECHT 90B ARG $e^+ e^- \rightarrow \gamma(4S)$ 446 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.447 Assumes equal production of B^0 and B^+ at the $\gamma(4S)$ and the systematic error includes the error associated with the helicity-mix uncertainty.448 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\gamma(4S)$. $\Gamma(a_1(1260)^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{267}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.7 \times 10^{-3}$	90	449 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$

449 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\gamma(4S)$. $\Gamma(a_1(1260)^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{268}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 9.0 \times 10^{-4}$	90	450 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$

450 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\gamma(4S)$. $\Gamma(\omega \pi^+)/\Gamma_{\text{total}}$ Γ_{269}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.59 ± 0.10 OUR AVERAGE				
Error includes scale factor of 1.2.				
$0.55 \pm 0.09 \pm 0.05$	451	AUBERT	04H BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.57^{+0.14}_{-0.13} \pm 0.06$	451	WANG	04A BELL	$e^+ e^- \rightarrow \gamma(4S)$
$1.13^{+0.33}_{-0.29} \pm 0.14$	451	JESSOP	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.42^{+0.20}_{-0.18} \pm 0.05$	451 LU	02 BELL	Repl. by WANG 04A
$0.66^{+0.21}_{-0.18} \pm 0.07$	451 AUBERT	01G BABR	Repl. by AUBERT 04H
< 2.3	90	451 BERGFELD	98 CLE2 Repl. by JESSOP 00
<40	90	452 ALBRECHT	90B ARG $e^+ e^- \rightarrow \Upsilon(4S)$

451 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

452 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\omega\rho^+)/\Gamma_{\text{total}}$

Γ_{270}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$12.6^{+3.7}_{-3.3} \pm 1.6$		453 AUBERT	050 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<61	90	453 BERGFELD	98 CLE2
-----	----	--------------	---------

453 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta\pi^+)/\Gamma_{\text{total}}$

Γ_{271}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
4.9 ± 0.5 OUR AVERAGE				
$5.1 \pm 0.6 \pm 0.3$		454 AUBERT,B	05K BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$4.8 \pm 0.7 \pm 0.3$		454 CHANG	05A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.2^{+2.8}_{-1.2}$		454 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.3 \pm 1.0 \pm 0.3$		454 AUBERT	04H BABR	Repl. by AUBERT,B 05K
< 15	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00
<700	90	454 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

454 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta'\pi^+)/\Gamma_{\text{total}}$

Γ_{272}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$4.0 \pm 0.8 \pm 0.4$		455 AUBERT,B	05K BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.5	90	455 AUBERT	04H BABR	Repl. by AUBERT,B 05K
< 7.0	90	455 ABE	01M BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
<12	90	455 AUBERT	01G BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
<12	90	455 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<31	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

455 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta'\rho^+)/\Gamma_{\text{total}}$

Γ_{273}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	90	456 AUBERT,B	04D BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.3 \times 10^{-5}$	90	456 RICHICHI	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$<4.7 \times 10^{-5}$	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

456 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta\rho^+)/\Gamma_{\text{total}}$ Γ_{274}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.4 \pm 1.9 \pm 1.1$		457 AUBERT,B	05K BABR	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<14	90	457 AUBERT,B	04D BABR	Repl. by AUBERT,B 05K
<15	90	457 RICHICHI	00 CLE2	$e^+e^- \rightarrow \gamma(4S)$
<32	90	BEHRENS	98 CLE2	Repl. by RICHICHI 00

457 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\phi\pi^+)/\Gamma_{\text{total}}$ Γ_{275}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<4.1 \times 10^{-7}$	90	458 AUBERT	04A BABR	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1.4 \times 10^{-6}$	90	458 AUBERT	01D BABR	$e^+e^- \rightarrow \gamma(4S)$
$<1.53 \times 10^{-4}$	90	459 ABE	00C SLD	$e^+e^- \rightarrow Z$
$<0.5 \times 10^{-5}$	90	458 BERGFELD	98 CLE2	

458 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.459 ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$. $\Gamma(\phi\rho^+)/\Gamma_{\text{total}}$ Γ_{276}/Γ

<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>
$<1.6 \times 10^{-5}$		460 BERGFELD	98 CLE2

460 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(a_0^0\pi^+)/\Gamma_{\text{total}}$ Γ_{277}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.8	90	461 AUBERT,BE	04 BABR	$e^+e^- \rightarrow \gamma(4S)$

461 Assumes equal production of charged and neutral B mesons from $\gamma(4S)$ decays. $\Gamma(\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$ Γ_{278}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<8.6 \times 10^{-4}$	90	462 ALBRECHT	90B ARG	$e^+e^- \rightarrow \gamma(4S)$

462 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$. $\Gamma(\rho^0 a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{279}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.2 \times 10^{-4}$	90	463 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.0 × 10⁻⁴ 90 ALBRECHT 90B ARG $e^+e^- \rightarrow \gamma(4S)$ <3.2 × 10⁻³ 90 BEBEK 87 CLEO $e^+e^- \rightarrow \gamma(4S)$ 463 BORTOLETTO 89 reports $< 5.4 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0\bar{B}^0$.
We rescale to 50%.464 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\gamma(4S)$.

$\Gamma(\rho^0 a_2(1320)^+)/\Gamma_{\text{total}}$					Γ_{280}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.2 \times 10^{-4}$	90	465 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.6 \times 10^{-3}$	90	466 BEBEK	87	CLEO	$e^+ e^- \rightarrow \gamma(4S)$
465 BORTOLETTO 89 reports $< 6.3 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.					
466 BEBEK 87 reports $< 2.3 \times 10^{-3}$ assuming the $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.					

$\Gamma(\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0)/\Gamma_{\text{total}}$					Γ_{281}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.3 \times 10^{-3}$	90	467 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$	

467 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\gamma(4S)$.

$\Gamma(a_1(1260)^+ a_1(1260)^0)/\Gamma_{\text{total}}$					Γ_{282}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-2}$	90	468 ALBRECHT	90B ARG	$e^+ e^- \rightarrow \gamma(4S)$	

468 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\gamma(4S)$.

$\Gamma(h^+ \pi^0)/\Gamma_{\text{total}}$					Γ_{283}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$(1.6^{+0.6}_{-0.5} \pm 0.36) \times 10^{-5}$		GODANG	98	CLE2	$e^+ e^- \rightarrow \gamma(4S)$

$\Gamma(\omega h^+)/\Gamma_{\text{total}}$					Γ_{284}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$h^+ = K^+$ or π^+					

$1.38^{+0.27}_{-0.24}$ OUR AVERAGE

$1.34^{+0.33}_{-0.29} \pm 0.11$	469 LU	02	BELL	$e^+ e^- \rightarrow \gamma(4S)$	
$1.43^{+0.36}_{-0.32} \pm 0.20$	469 JESSOP	00	CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$2.5^{+0.8}_{-0.7} \pm 0.3$	469 BERGFELD	98	CLE2	Repl. by JESSOP 00	
469 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.					

$\Gamma(h^+ X^0 (\text{Familon}))/\Gamma_{\text{total}}$					Γ_{285}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<4.9 \times 10^{-5}$	90	470 AMMAR	01B CLE2	$e^+ e^- \rightarrow \gamma(4S)$	

470 AMMAR 01B searched for the two-body decay of the B meson to a massless neutral feebly-interacting particle X^0 such as the famililon, the Nambu-Goldstone boson associated with a spontaneously broken global family symmetry.

$\Gamma(p\bar{p}\pi^+)/\Gamma_{\text{total}}$ Γ_{286}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.06^{+0.73}_{-0.62} \pm 0.37$	471,472	WANG	04	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.7	90	471,473	ABE	02K	BELL	Repl. by WANG 04
<500	90	474	ABREU	95N	DLPH	Sup. by ADAM 96D
<160	90	475	BEBEK	89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
570 ± 150	± 210	476	ALBRECHT	88F	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

471 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

472 The branching fraction for $M_{p\bar{p}} < 2.85$ is also reported.

473 Explicitly vetoes resonant production of $p\bar{p}$ from Charmonium states.

474 Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12.

475 BEBEK 89 reports $< 1.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

476 ALBRECHT 88F reports $(5.2 \pm 1.4 \pm 1.9) \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(p\bar{p}\pi^+ \text{nonresonant})/\Gamma_{\text{total}}$ Γ_{287}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.3 \times 10^{-5}$	90	BERGFELD	96B	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

 $\Gamma(p\bar{p}\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{288}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.2 \times 10^{-4}$	90	477	ALBRECHT	88F ARG $e^+ e^- \rightarrow \Upsilon(4S)$

477 ALBRECHT 88F reports $< 4.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(p\bar{p}K^+)/\Gamma_{\text{total}}$ Γ_{289}/Γ

<u>VALUE</u> (units 10^{-6})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.6 ± 1.0 OUR AVERAGE	Error includes scale factor of 2.4.		
6.7 $\pm 0.5 \pm 0.4$	478,479,480	AUBERT,B	05L BABR $e^+ e^- \rightarrow \Upsilon(4S)$
$4.59^{+0.38}_{-0.34} \pm 0.50$	478,479,480	WANG	05A BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.66^{+0.67}_{-0.57} \pm 0.62$	478,479,481	WANG	04	BELL	Repl. by WANG 05A
$4.3^{+1.1}_{-0.9} \pm 0.5$	478,479	ABE	02K	BELL	Repl. by WANG 04

478 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

479 Explicitly vetoes resonant production of $p\bar{p}$ from Charmonium states.

480 Provides also results with $M_{p\bar{p}} < 2.85 \text{ GeV}/c^2$ and angular asymmetry of $p\bar{p}$ system.

481 The branching fraction for $M_{p\bar{p}} < 2.85$ is also reported.

$\Gamma(\Theta(1710)^{++} \bar{p} \times B(\Theta(1710)^{++} \rightarrow p K^+)) / \Gamma_{\text{total}}$ Γ_{290}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<0.91	90	482 WANG	05A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.0	90	482,483 AUBERT,B	05L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
482 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				
483 Provides upper limits depending on the pentaquark masses between 1.43 to 2.0 GeV/c^2 .				

 $\Gamma(f_J(2220)K^+ \times B(f_J(2220) \rightarrow p \bar{p})) / \Gamma_{\text{total}}$ Γ_{291}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<4.1	90	484 WANG	05A BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
484 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				

 $\Gamma(p\bar{\Lambda}(1520)) / \Gamma_{\text{total}}$ Γ_{292}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	485 AUBERT,B	05L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
485 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				

 $\Gamma(p\bar{p}K^+ \text{ nonresonant}) / \Gamma_{\text{total}}$ Γ_{293}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.9 \times 10^{-5}$	90	BERGFELD	96B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

 $\Gamma(p\bar{p}K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{294}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$10.3^{+3.6+1.3}_{-2.8-1.7}$	486,487 WANG	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

486 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.487 The branching fraction for $M_{p\bar{p}} < 2.85$ is also reported. $\Gamma(p\bar{\Lambda}) / \Gamma_{\text{total}}$ Γ_{295}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.9 \times 10^{-7}$	90	488 CHANG	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5 $\times 10^{-6}$ 90 488 BORNHEIM 03 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$ <2.2 $\times 10^{-6}$ 90 488 ABE 020 BELL $e^+ e^- \rightarrow \Upsilon(4S)$ <2.6 $\times 10^{-6}$ 90 488 COAN 99 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$ <6 $\times 10^{-5}$ 90 489 AVERY 89B CLEO $e^+ e^- \rightarrow \Upsilon(4S)$ <9.3 $\times 10^{-5}$ 90 490 ALBRECHT 88F ARG $e^+ e^- \rightarrow \Upsilon(4S)$ 488 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.489 AVERY 89B reports $< 5 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.490 ALBRECHT 88F reports $< 8.5 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(p\bar{\Lambda}\gamma)/\Gamma_{\text{total}}$ Γ_{296}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.16^{+0.58}_{-0.53} \pm 0.20$	491	LEE	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.9 90 492 EDWARDS 03 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

491 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

492 Corresponds to $E_\gamma > 1.5$ GeV. The limit changes to 3.3×10^{-6} for $E_\gamma > 2.0$ GeV.

 $\Gamma(p\bar{\Sigma}\gamma)/\Gamma_{\text{total}}$ Γ_{297}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.6	90	493 LEE	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7.9 90 494 EDWARDS 03 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

493 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

494 Corresponds to $E_\gamma > 1.5$ GeV. The limit changes to 6.4×10^{-6} for $E_\gamma > 2.0$ GeV.

 $\Gamma(p\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{298}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.0 \times 10^{-4}$	90	495 ALBRECHT	88F ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

495 ALBRECHT 88F reports $< 1.8 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(\Lambda\bar{\Lambda}\pi^+)/\Gamma_{\text{total}}$ Γ_{299}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.8	90	496 LEE	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

496 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\Lambda\bar{\Lambda}K^+)/\Gamma_{\text{total}}$ Γ_{300}/Γ

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.91^{+0.9}_{-0.70} \pm 0.38$	497	LEE	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

497 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\Delta^0 p)/\Gamma_{\text{total}}$ Γ_{301}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.8 \times 10^{-4}$	90	498 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

498 BORTOLETTO 89 reports $< 3.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(\Delta^{++}\bar{p})/\Gamma_{\text{total}}$ Γ_{302}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.5 \times 10^{-4}$	90	499 BORTOLETTO89	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

499 BORTOLETTO 89 reports $< 1.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(D^+ p\bar{p})/\Gamma_{\text{total}}$				Γ_{303}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-5}$	90	500 ABE	02W BELL	$e^+ e^- \rightarrow \gamma(4S)$
500 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

$\Gamma(D^*(2010)^+ p\bar{p})/\Gamma_{\text{total}}$				Γ_{304}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-5}$	90	501 ABE	02W BELL	$e^+ e^- \rightarrow \gamma(4S)$
501 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				

$\Gamma(\Lambda_c^- p\pi^+)/\Gamma_{\text{total}}$				Γ_{305}/Γ
VALUE (units 10^{-4})		DOCUMENT ID	TECN	COMMENT
2.1 ± 0.7 OUR AVERAGE				
$2.4 \pm 0.6 \pm 0.6$		502 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$1.9 \pm 0.5 \pm 0.5$		503 GABYSHEV	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$6.2^{+2.3}_{-2.0} \pm 1.6$		504 FU	97 CLE2	Repl. by DYTMAN 02
502 DYTMAN 02 reports $(2.4^{+0.63}_{-0.62}) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
503 GABYSHEV 02 reports $(1.87^{+0.51}_{-0.49}) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
504 FU 97 uses PDG 96 values of Λ_c branching fraction.				

$\Gamma(\Lambda_c^- p\pi^+\pi^0)/\Gamma_{\text{total}}$				Γ_{306}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
$1.81 \pm 0.29^{+0.52}_{-0.50}$		505, 506 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<3.12	90	507 FU	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
505 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				
506 DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.				
507 FU 97 uses PDG 96 values of Λ_c branching ratio.				

$\Gamma(\Lambda_c^- p\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_{307}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
$2.25 \pm 0.25^{+0.63}_{-0.61}$		508, 509 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<1.46	90	510 FU	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
508 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.				
509 DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.				
510 FU 97 uses PDG 96 values of Λ_c branching ratio.				

$\Gamma(\bar{\Lambda}_c^- p \pi^+ \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$	Γ_{308}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.34 \times 10^{-2}$	90	511 FU	97 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

511 FU 97 uses PDG 96 values of Λ_c branching ratio.

$\Gamma(\Sigma_c(2455)^0 p)/\Gamma_{\text{total}}$	Γ_{309}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.8 \times 10^{-4}$	90	512,513 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<9.3 \times 10^{-5}$	90	512,514 GABYSHEV	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$
-----------------------	----	------------------	---------	----------------------------------

512 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

513 DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio.

514 Uses the value for $\Lambda_c \rightarrow p K^- \pi^+$ branching ratio ($5.0 \pm 1.3\%$).

$\Gamma(\bar{\Sigma}_c(2520)^0 p)/\Gamma_{\text{total}}$	Γ_{310}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.6 \times 10^{-5}$	90	515,516 GABYSHEV	02 BELL	$e^+ e^- \rightarrow \gamma(4S)$

515 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

516 Uses the value for $\Lambda_c \rightarrow p K^- \pi^+$ branching ratio ($5.0 \pm 1.3\%$).

$\Gamma(\bar{\Sigma}_c(2455)^0 p \pi^0)/\Gamma_{\text{total}}$	Γ_{311}/Γ		
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.4 \pm 1.4 \pm 1.1$	517,518 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

517 DYTMAN 02 reports $(4.4 \pm 1.4) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

518 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\bar{\Sigma}_c(2455)^0 p \pi^- \pi^+)/\Gamma_{\text{total}}$	Γ_{312}/Γ		
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.4 \pm 1.3 \pm 1.1$	519,520 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

519 DYTMAN 02 reports $(4.4 \pm 1.3) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

520 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\bar{\Sigma}_c(2455)^{--} p \pi^+ \pi^+)/\Gamma_{\text{total}}$	Γ_{313}/Γ		
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 1.0 \pm 0.7$	521,522 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

521 DYTMAN 02 reports $(2.8 \pm 1.0) \times 10^{-4}$ for $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 0.05$. We rescale to our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

522 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(\bar{\Lambda}_c(2593)^-/\bar{\Lambda}_c(2625)^- p\pi^+)/\Gamma_{\text{total}}$ Γ_{314}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.9 \times 10^{-4}$	90	523,524 DYTMAN	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

523 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.524 DYTMAN 02 measurement uses $B(\Lambda_c^- \rightarrow \bar{p} K^+ \pi^-) = 5.0 \pm 1.3\%$. The second error includes the systematic and the uncertainty of the branching ratio. $\Gamma(\pi^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{315}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0039	90	525 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$

525 WEIR 90B assumes B^+ production cross section from LUND. $\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{316}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0091	90	526 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$

526 WEIR 90B assumes B^+ production cross section from LUND. $\Gamma(\pi^+ \nu \bar{\nu})/\Gamma_{\text{total}}$ Γ_{317}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-4}$	90	527 AUBERT	05H BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

527 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{318}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
$8.0^{+2.2}_{-1.9}$ OUR AVERAGE				Error includes scale factor of 1.4.

10.5 $^{+2.5}_{-2.2} \pm 0.7$ 528 AUBERT 03U BABR $e^+ e^- \rightarrow \Upsilon(4S)$ 6.3 $^{+1.9}_{-1.7} \pm 0.3$ 529 ISHIKAWA 03 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<	14	90	528 ABE	02 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
<	9	90	528 AUBERT	02L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
<	24	90	530 ANDERSON	01B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<	990	90	531 ALBRECHT	91E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
<	68000	90	532 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$
<	600	90	533 AVERY	89B CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
<	2500	90	534 AVERY	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

528 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.529 Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$. The second error is a total of systematic uncertainties including model dependence.

530 The result is for di-lepton masses above 0.5 GeV.

531 ALBRECHT 91E reports $< 9.0 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.532 WEIR 90B assumes B^+ production cross section from LUND.533 Avery 89B reports $< 5 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.534 Avery 87 reports $< 2.1 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K^+\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{319}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

0.34 $^{+0.19}_{-0.14}$ OUR AVERAGE Error includes scale factor of 1.7.0.07 $^{+0.19}_{-0.11}$ ± 0.02 535 AUBERT 03U BABR $e^+ e^- \rightarrow \gamma(4S)$ 0.45 $^{+0.14}_{-0.12}$ ± 0.03 536 ISHIKAWA 03 BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.98 $^{+0.46}_{-0.36}$ ± 0.16 535 ABE 02 BELL Repl. by ISHIKAWA 03< 1.2 90 535 AUBERT 02L BABR $e^+ e^- \rightarrow \gamma(4S)$ < 3.68 90 537 ANDERSON 01B CLE2 $e^+ e^- \rightarrow \gamma(4S)$ < 5.2 90 538 AFFOLDER 99B CDF $p\bar{p}$ at 1.8 TeV

< 10 90 539 ABE 96L CDF Repl. by AF-FOLDER 99B

< 240 90 540 ALBRECHT 91E ARG $e^+ e^- \rightarrow \gamma(4S)$ < 6400 90 541 WEIR 90B MRK2 $e^+ e^-$ 29 GeV< 170 90 542 AVERY 89B CLEO $e^+ e^- \rightarrow \gamma(4S)$ < 380 90 543 AVERY 87 CLEO $e^+ e^- \rightarrow \gamma(4S)$ 535 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.536 Assumes equal production of B^0 and B^+ at $\gamma(4S)$. The second error is a total of systematic uncertainties including model dependence.

537 The result is for di-lepton masses above 0.5 GeV.

538 AFFOLDER 99B measured relative to $B^+ \rightarrow J/\psi(1S) K^+$.539 ABE 96L measured relative to $B^+ \rightarrow J/\psi(1S) K^+$ using PDG 94 branching ratios.540 ALBRECHT 91E reports $< 2.2 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.541 WEIR 90B assumes B^+ production cross section from LUND.542 AVERY 89B reports $< 1.5 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.543 AVERY 87 reports $< 3.2 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%. $\Gamma(K^+\ell^+\ell^-)/\Gamma_{\text{total}}$ Γ_{320}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

5.3 $^{+1.1}_{-1.0}$ ± 0.3 544 ISHIKAWA 03 BELL $e^+ e^- \rightarrow \gamma(4S)$ 544 Assumes equal production of B^0 and B^+ at $\gamma(4S)$. $\Gamma(K^+\bar{\nu}\nu)/\Gamma_{\text{total}}$ Γ_{321}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<5.2 $\times 10^{-5}$ 90 545 AUBERT 05H BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.4 $\times 10^{-4}$ 90 545 BROWDER 01 CLE2 $e^+ e^- \rightarrow \gamma(4S)$ 545 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(K^*(892)^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{322}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.6	90	546 ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.20 $^{+1.34}_{-0.87} \pm 0.28$		547 AUBERT	03U BABR	$e^+ e^- \rightarrow \gamma(4S)$
< 8.9	90	547 ABE	02 BELL	Repl. by ISHIKAWA 03
< 9.5	90	547 AUBERT	02L BABR	$e^+ e^- \rightarrow \gamma(4S)$
< 690	90	548 ALBRECHT	91E ARG	$e^+ e^- \rightarrow \gamma(4S)$

546 Assumes equal production of B^0 and B^+ at $\gamma(4S)$. The second error is a total of systematic uncertainties including model dependence.547 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.548 ALBRECHT 91E reports $< 6.3 \times 10^{-4}$ assuming the $\gamma(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%. $\Gamma(K^*(892)^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{323}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.2	90	549 ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
3.07 $^{+2.58}_{-1.78} \pm 0.42$		550 AUBERT	03U BABR	$e^+ e^- \rightarrow \gamma(4S)$
< 3.9	90	550 ABE	02 BELL	Repl. by ISHIKAWA 03
< 17.0	90	550 AUBERT	02L BABR	$e^+ e^- \rightarrow \gamma(4S)$
< 1200	90	551 ALBRECHT	91E ARG	$e^+ e^- \rightarrow \gamma(4S)$

549 Assumes equal production of B^0 and B^+ at $\gamma(4S)$. The second error is a total of systematic uncertainties including model dependence.550 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.551 ALBRECHT 91E reports $< 1.1 \times 10^{-3}$ assuming the $\gamma(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%. $\Gamma(K^*(892)^+ \ell^+ \ell^-)/\Gamma_{\text{total}}$ Γ_{324}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 22	90	552 ISHIKAWA	03 BELL	$e^+ e^- \rightarrow \gamma(4S)$

552 Assumes equal production of B^0 and B^+ at $\gamma(4S)$. $\Gamma(\pi^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{325}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	553 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$

553 WEIR 90B assumes B^+ production cross section from LUND. $\Gamma(\pi^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{326}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	554 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$

554 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{327}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.8 \times 10^{-6}$	90	555 AUBERT	02L BABR	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0064	90	556 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$
---------	----	----------	----------	--------------------------

555 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.556 WEIR 90B assumes B^+ production cross section from LUND. $\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{328}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	557 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$

557 WEIR 90B assumes B^+ production cross section from LUND. $\Gamma(K^*(892)^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{329}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.9 \times 10^{-6}$	90	558 AUBERT	02L BABR	$e^+ e^- \rightarrow \gamma(4S)$

558 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\pi^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{330}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.6 \times 10^{-6}$	90	559 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0039	90	560 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$
---------	----	----------	----------	--------------------------

559 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.560 WEIR 90B assumes B^+ production cross section from LUND. $\Gamma(\pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{331}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-6}$	90	561 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0091	90	562 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$
---------	----	----------	----------	--------------------------

561 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.562 WEIR 90B assumes B^+ production cross section from LUND. $\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{332}/Γ

Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-6}$	90	563 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0064	90	564 WEIR	90B MRK2	$e^+ e^- 29 \text{ GeV}$
---------	----	----------	----------	--------------------------

563 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.564 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\rho^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{333}/Γ

Test of total lepton number conservation.

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.6	90	565 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

565 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\rho^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{334}/Γ

Test of total lepton number conservation.

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.0	90	566 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

566 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(\rho^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{335}/Γ

Test of total lepton number conservation.

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.3	90	567 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

567 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{336}/Γ

Test of total lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.0 $\times 10^{-6}$	90	568 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0039 90 569 WEIR 90B MRK2 $e^+ e^-$ 29 GeV568 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.569 WEIR 90B assumes B^+ production cross section from LUND. $\Gamma(K^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{337}/Γ

Test of total lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.8 $\times 10^{-6}$	90	570 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0091 90 571 WEIR 90B MRK2 $e^+ e^-$ 29 GeV570 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.571 WEIR 90B assumes B^+ production cross section from LUND. $\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{338}/Γ

Test of total lepton number conservation.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.0 $\times 10^{-6}$	90	572 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0064 90 573 WEIR 90B MRK2 $e^+ e^-$ 29 GeV572 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.573 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^*(892)^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{339}/Γ

Test of total lepton number conservation.

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.8	90	574 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

574 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K^*(892)^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{340}/Γ

Test of total lepton number conservation.

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8.3	90	575 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

575 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$. $\Gamma(K^*(892)^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{341}/Γ

Test of total lepton number conservation.

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.4	90	576 EDWARDS	02B CLE2	$e^+ e^- \rightarrow \gamma(4S)$

576 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.**POLARIZATION IN B^+ DECAY**

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal (L) or both are transverse and parallel (\parallel) or perpendicular (\perp) to each other with the parameters Γ_L/Γ , Γ_\perp/Γ , and the relative phases ϕ_\parallel and ϕ_\perp . See the definitions in the note on "Polarization in B Decays" review in the B^0 Particle Listings.

 Γ_L/Γ in $B^+ \rightarrow \bar{D}^{*0} \rho^+$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.892 ± 0.018 ± 0.016	CSORNA	03	$e^+ e^- \rightarrow \gamma(4S)$

 Γ_L/Γ in $B^+ \rightarrow \bar{D}^{*0} K^{*+}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.86 ± 0.06 ± 0.03	AUBERT	04K BABR	$e^+ e^- \rightarrow \gamma(4S)$

 Γ_L/Γ in $B^+ \rightarrow J/\psi K^{*+}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.604 ± 0.015 ± 0.018	ITOH	05	BELL

 Γ_\perp/Γ in $B^+ \rightarrow J/\psi K^{*+}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.180 ± 0.014 ± 0.010	ITOH	05	BELL

 Γ_L/Γ in $B^+ \rightarrow \phi K^*(892)^+$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.50 ± 0.07 OUR AVERAGE	CHEN	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$

0.52 ± 0.08 ± 0.03

0.46 ± 0.12 ± 0.03

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
AUBERT	03V BABR	$e^+ e^- \rightarrow \gamma(4S)$

Γ_{\perp}/Γ in $B^+ \rightarrow \phi K^{*+}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.19±0.08±0.02	CHEN	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$

 $\phi_{||}$ in $B^+ \rightarrow \phi K^{*+}$

VALUE (°)	DOCUMENT ID	TECN	COMMENT
2.10±0.28±0.04	CHEN	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$

 ϕ_{\perp} in $B^+ \rightarrow \phi K^{*+}$

VALUE (°)	DOCUMENT ID	TECN	COMMENT
2.31±0.30±0.07	CHEN	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$

 Γ_L/Γ in $B^+ \rightarrow \rho^0 K^*(892)^+$

VALUE	DOCUMENT ID	TECN	COMMENT
0.96^{+0.04}_{-0.15}±0.04	AUBERT	03V BABR	$e^+ e^- \rightarrow \gamma(4S)$

 $\Gamma_L/\Gamma(B^+ \rightarrow K^*(892)^0 \rho^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.43±0.11^{+0.05}_{-0.02}	ZHANG	05D BELL	$e^+ e^- \rightarrow \gamma(4S)$

 Γ_L/Γ in $B^+ \rightarrow \rho^+ \rho^0$

VALUE	DOCUMENT ID	TECN	COMMENT
0.96^{+0.05}_{-0.06} OUR AVERAGE	AUBERT	03V BABR	$e^+ e^- \rightarrow \gamma(4S)$

0.97^{+0.03}_{-0.07} ± 0.04

0.948 ± 0.106 ± 0.021

 Γ_L/Γ in $B^+ \rightarrow \omega \rho^+$

VALUE	DOCUMENT ID	TECN	COMMENT
0.88^{+0.12}_{-0.15}±0.03	AUBERT	05O BABR	$e^+ e^- \rightarrow \gamma(4S)$

CP VIOLATION A_{CP} is defined as

$$\frac{B(B^- \rightarrow \bar{f}) - B(B^+ \rightarrow f)}{B(B^- \rightarrow \bar{f}) + B(B^+ \rightarrow f)},$$

the CP -violation charge asymmetry of exclusive B^- and B^+ decay. $A_{CP}(B^+ \rightarrow J/\psi(1S) K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.024±0.014 OUR AVERAGE			

-0.030 ± 0.014 ± 0.010

-0.026 ± 0.022 ± 0.017

0.018 ± 0.043 ± 0.004

577	DOCUMENT ID	TECN	COMMENT
	AUBERT	05J BABR	$e^+ e^- \rightarrow \gamma(4S)$
	ABE	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
	BONVICINI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.03 ± 0.015 ± 0.006

0.003 ± 0.030 ± 0.004

AUBERT 04P BABR Repl. by AUBERT 05J

AUBERT 02F BABR Repl. by AUBERT 04P

$577\text{A} + 0.3\%$ correction is applied due to a slightly higher reconstruction efficiency for the positive kaons.

$A_{CP}(B^+ \rightarrow J/\psi(1S)\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.09 ± 0.08 OUR AVERAGE			

0.123 ± 0.085 ± 0.004	AUBERT	04P BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.023 ± 0.164 ± 0.015	ABE	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.01 ± 0.22 ± 0.01	AUBERT	02F BABR	Repl. by AUBERT 04P

$A_{CP}(B^+ \rightarrow J/\psi K^*(892)^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.048 ± 0.029 ± 0.016	AUBERT	05J BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \psi(2S)K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.025 ± 0.024 OUR AVERAGE			

0.052 ± 0.059 ± 0.020	AUBERT	05J BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.042 ± 0.020 ± 0.017	ABE	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.02 ± 0.091 ± 0.01	578 BONVICINI	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

$578\text{A} + 0.3\%$ correction is applied due to a slightly higher reconstruction efficiency for the positive kaons.

$A_{CP}(B^+ \rightarrow \psi(2S)K^*(892)^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.077 ± 0.207 ± 0.051	AUBERT	05J BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \chi_{c1} K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.003 ± 0.076 ± 0.017	AUBERT	05J BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \chi_{c1} K^*(892)^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.471 ± 0.378 ± 0.268	AUBERT	05J BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \bar{D}^0\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.008 ± 0.008	ABE	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D_{CP(+1)}\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.035 ± 0.024	ABE	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D_{CP(-1)}\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.017 ± 0.026	ABE	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \bar{D}^0 K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.066±0.036	ABE	06	BELL $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.003±0.080±0.037	579 ABE	03D	BELL Repl. by SWAIN 03
0.04 ± 0.06 ± 0.03	580 SWAIN	03	BELL Repl. by ABE 06
579	Corresponds to 90% confidence range $-0.15 < A_{CP} < 0.16$.		
580	Corresponds to 90% confidence range $-0.07 < A_{CP} < 0.15$.		

 $r_B(B^+ \rightarrow D^0 K^+)$

$r_B^{(*)}$ and $\delta_B^{(*)}$ are the amplitude ratios and relative strong phases between the amplitudes of $A_{CP}(B^+ \rightarrow D^{(*)0} K^+)$ and $A_{CP}(B^+ \rightarrow \bar{D}^{(*)0} K^+)$,

VALUE	DOCUMENT ID	TECN	COMMENT
0.12±0.08±0.05	581 AUBERT,B	05Y	BABR $e^+ e^- \rightarrow \gamma(4S)$
581 Uses a Dalitz analysis of neutral D decays to $K_S^0 \pi^+ \pi^-$ in the processes $B^\pm \rightarrow D^{(*)} K^\pm$, $D^* \rightarrow D \pi^0$, $D \gamma$.			

 $\delta_B(B^+ \rightarrow D^0 K^+)$

VALUE (degrees)	DOCUMENT ID	TECN	COMMENT
104±45⁺²³₋₃₂	582 AUBERT,B	05Y	BABR $e^+ e^- \rightarrow \gamma(4S)$
582 Uses a Dalitz analysis of neutral D decays to $K_S^0 \pi^+ \pi^-$ in the processes $B^\pm \rightarrow D^{(*)} K^\pm$, $D^* \rightarrow D \pi^0$, $D \gamma$.			

 $A_{CP}(B^+ \rightarrow [K^- \pi^+]_D K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.88^{+0.77}_{-0.62}±0.06	SAIGO	05	BELL $e^+ e^- \rightarrow \gamma(4S)$

 $A_{CP}(B^+ \rightarrow [K^- \pi^+]_{\bar{D}} K^*(892)^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.22±0.61±0.17	AUBERT,B	05V	BABR $e^+ e^- \rightarrow \gamma(4S)$

 $A_{CP}(B^+ \rightarrow [K^- \pi^+]_D \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.30^{+0.29}_{-0.25}±0.06	SAIGO	05	BELL $e^+ e^- \rightarrow \gamma(4S)$

 $A_{CP}(B^+ \rightarrow [\pi^+ \pi^- \pi^0]_D K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.02±0.16±0.03	AUBERT,B	05T	BABR $e^+ e^- \rightarrow \gamma(4S)$

 $A_{CP}(B^+ \rightarrow D_{CP(+1)} K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.22±0.14 OUR AVERAGE	Error includes scale factor of 1.4.		
0.06±0.14±0.05	ABE	06	BELL $e^+ e^- \rightarrow \gamma(4S)$
0.35±0.13±0.04	AUBERT	06J	BABR $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.07 \pm 0.17 \pm 0.06$	AUBERT	04N BABR	Repl. by AUBERT 06J
$0.29 \pm 0.26 \pm 0.05$	583 ABE	03D BELL	Repl. by SWAIN 03
$0.06 \pm 0.19 \pm 0.04$	584 SWAIN	03 BELL	Repl. by ABE 06

583 Corresponds to 90% confidence range $-0.14 < A_{CP} < 0.73$.

584 Corresponds to 90% confidence range $-0.26 < A_{CP} < 0.38$.

$A_{CP}(B^+ \rightarrow D_{CP(-1)} K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.09 ± 0.10 OUR AVERAGE			

$-0.12 \pm 0.14 \pm 0.05$	ABE	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$
$-0.06 \pm 0.13 \pm 0.04$	AUBERT	06J BABR	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.22 \pm 0.24 \pm 0.04$	585 ABE	03D BELL	Repl. by SWAIN 03
$-0.19 \pm 0.17 \pm 0.05$	586 SWAIN	03 BELL	Repl. by ABE 06

585 Corresponds to 90% confidence range $-0.62 < A_{CP} < 0.18$.

586 Corresponds to 90% confidence range $-0.47 < A_{CP} < 0.11$.

$A_{CP}(B^+ \rightarrow \bar{D}^{*0} \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.014 ± 0.015			
ABE	06	BELL	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow (D_{CP(+1)}^*)^0 \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.021 ± 0.045			
ABE	06	BELL	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow (D_{CP(-1)}^*)^0 \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.090 ± 0.051			
ABE	06	BELL	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D^{*0} K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.089 ± 0.086			
ABE	06	BELL	$e^+ e^- \rightarrow \gamma(4S)$

$r_B^*(B^+ \rightarrow D^{*0} K^+)$

$r_B^{(*)}$ and $\delta_B^{(*)}$ are the amplitude ratios and relative strong phases between the amplitudes of $A_{CP}(B^+ \rightarrow D^{(*)0} K^+)$ and $A_{CP}(B^+ \rightarrow \bar{D}^{(*)0} K^+)$,

VALUE	DOCUMENT ID	TECN	COMMENT
$0.17 \pm 0.10 \pm 0.04$			
587	AUBERT,B	05Y BABR	$e^+ e^- \rightarrow \gamma(4S)$

587 Uses a Dalitz analysis of neutral D decays to $K_S^0 \pi^+ \pi^-$ in the processes $B^\pm \rightarrow D^{(*)} K^\pm$, $D^* \rightarrow D \pi^0$, $D \gamma$.

$\delta_B^*(B^+ \rightarrow D^{*0} K^+)$

VALUE (degrees)	DOCUMENT ID	TECN	COMMENT
-64±41⁺²⁰₋₁₉	588 AUBERT,B	05Y BABR	$e^+ e^- \rightarrow \gamma(4S)$

588 Uses a Dalitz analysis of neutral D decays to $K_S^0 \pi^+ \pi^-$ in the processes $B^\pm \rightarrow D^{(*)} K^\pm$, $D^* \rightarrow D\pi^0$, $D\gamma$.

$A_{CP}(B^+ \rightarrow D_{CP(+1)}^{*0} K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.15±0.16 OUR AVERAGE			
-0.20±0.22±0.04	ABE	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$
-0.10±0.23 ^{+0.03} _{-0.04}	AUBERT	05N BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D_{CP(-1)}^{*0} K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.13±0.30±0.08	ABE	06 BELL	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D_{CP(+1)} K^*(892)^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.08±0.19±0.08	AUBERT,B	05U BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow D_{CP(-1)} K^*(892)^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.26±0.40±0.12	AUBERT,B	05U BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow K_S^0 \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.02 ±0.07 OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.		

-0.09 ±0.05 ±0.01	589 AUBERT,BE	05E BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.05 ±0.05 ±0.01	590 CHAO	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.18 ±0.24	591 CHEN	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.05 ±0.08 ±0.01	592 AUBERT	04M BABR	Repl. by AUBERT,BE 05E
0.07 ^{+0.09} _{-0.08} ^{+0.01} _{-0.03}	593 UNNO	03 BELL	Repl. by CHAO 05A
0.46 ±0.15 ±0.02	594 CASEY	02 BELL	Repl. by UNNO 03
0.098 ^{+0.430} _{-0.343} ^{+0.020} _{-0.063}	595 ABE	01K BELL	Repl. by CASEY 02
-0.21 ±0.18 ±0.03	596 AUBERT	01E BABR	Repl. by AUBERT 04M

589 Corresponds to 90% confidence range $-0.16 < A_{CP} < -0.02$.

590 Corresponds to a 90% CL interval of $-0.04 < A_{CP} < 0.13$.

591 Corresponds to 90% confidence range $-0.22 < A_{CP} < 0.56$.


592 90% CL interval $-0.18 < A_{CP} < 0.08$

593 Corresponds to 90% confidence range $-0.10 < A_{CP} < +0.22$.

594 Corresponds to 90% confidence range $+0.19 < A_{CP} < +0.72$.

595 Corresponds to 90% confidence range $-0.53 < A_{CP} < 0.82$.

596 Corresponds to 90% confidence range $-0.51 < A_{CP} < 0.09$.



$A_{CP}(B^+ \rightarrow K^+ \pi^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.04 ±0.04 OUR AVERAGE			
0.06 ± 0.06 ± 0.01	597 AUBERT	05L BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.04 ± 0.05 ± 0.02	598 CHAO	04B BELL	$e^+ e^- \rightarrow \gamma(4S)$
-0.29 ± 0.23	599 CHEN	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.06 ± 0.06 ± 0.02	597 CHAO	05A BELL	Repl. by CHAO 04B
-0.09 ± 0.09 ± 0.01	600 AUBERT	03L BABR	Repl. by AUBERT 05L
-0.02 ± 0.19 ± 0.02	601 CASEY	02 BELL	Repl. by CHAO 04B
-0.059 ^{+0.222} _{-0.196} ^{+0.055} _{-0.017}	602 ABE	01K BELL	Repl. by CASEY 02
0.00 ± 0.18 ± 0.04	603 AUBERT	01E BABR	Repl. by AUBERT 03L
597 Corresponds to a 90% CL interval of $-0.06 < A_{CP} < 0.18$.			
598 Corresponds to 90% CL interval of $-0.05 < A_{CP} < 0.13$.			
599 Corresponds to 90% confidence range $-0.67 < A_{CP} < 0.09$.			
600 Corresponds to 90% confidence range $-0.24 < A_{CP} < 0.06$.			
601 Corresponds to 90% confidence range $-0.35 < A_{CP} < +0.30$.			
602 Corresponds to 90% confidence range $-0.40 < A_{CP} < 0.36$.			
603 Corresponds to 90% confidence range $-0.30 < A_{CP} < +0.30$.			

$A_{CP}(B^+ \rightarrow K^+\eta')$

VALUE	DOCUMENT ID	TECN	COMMENT
0.020 ± 0.025 OUR AVERAGE			
0.033 $\pm 0.028 \pm 0.005$	604 AUBERT	05M BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.11 $\pm 0.11 \pm 0.02$	605 AUBERT	02E BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.015 $\pm 0.070 \pm 0.009$	606 CHEN	02B BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.03 ± 0.12	607 CHEN	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.037 $\pm 0.045 \pm 0.011$	608 AUBERT	03W BABR	Repl. by AUBERT 05M
0.06 $\pm 0.15 \pm 0.01$	609 ABE	01M BELL	Repl. by CHEN 02B
604	Corresponds to 90% confidence range $-0.012 < A_{CP} < 0.078$.		
605	Corresponds to 90% confidence range $-0.28 < A_{CP} < 0.07$.		
606	Corresponds to 90% confidence range $-0.13 < A_{CP} < 0.10$.		
607	Corresponds to 90% confidence range $-0.17 < A_{CP} < 0.23$.		
608	Corresponds to 90% confidence range $-0.04 < A_{CP} < 0.11$.		
609	Corresponds to 90% confidence range $-0.20 < A_{CP} < 0.32$.		

 $A_{CP}(B^+ \rightarrow \eta K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.25 ± 0.14 OUR AVERAGE			
-0.20 $\pm 0.15 \pm 0.01$	AUBERT,B	05K BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.49 $\pm 0.31 \pm 0.07$	CHANG	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.52 $\pm 0.24 \pm 0.01$	AUBERT	04H BABR	Repl. by AUBERT,B 05K

 $A_{CP}(B^+ \rightarrow \eta K^*(892)^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.13 \pm 0.14 \pm 0.02$			
AUBERT,B	04D BABR	$e^+ e^- \rightarrow \gamma(4S)$	

 $A_{CP}(B^+ \rightarrow \omega K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.02 ± 0.13 OUR AVERAGE			
-0.09 $\pm 0.17 \pm 0.01$	AUBERT	04H BABR	$e^+ e^- \rightarrow \gamma(4S)$
0.06 $\pm 0.21 \pm 0.01$	610 WANG	04A BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.21 $\pm 0.28 \pm 0.03$	611 LU	02 BELL	Repl. by WANG 04A
610	Corresponds to 90% CL interval $0.15 < A_{CP} < 0.90$		
611	Corresponds to 90% confidence range $-0.70 < A_{CP} < +0.38$.		

 $A_{CP}(B^+ \rightarrow K^{*0}\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.068 \pm 0.078 \pm 0.070$			
AUBERT,B	05N BABR	$e^+ e^- \rightarrow \gamma(4S)$	

 $A_{CP}(B^+ \rightarrow K^+\pi^-\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.013 \pm 0.037 \pm 0.011$			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.01 $\pm 0.07 \pm 0.03$	AUBERT	03M BABR	Repl. by AUBERT,B 05N

$A_{CP}(B^+ \rightarrow f_0(980)K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.088 \pm 0.095^{+0.097}_{-0.056}$	AUBERT,B	05N BABR	$e^+ e^- \rightarrow \gamma(4S)$

 $A_{CP}(B^+ \rightarrow \rho^0 K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.32 \pm 0.13^{+0.10}_{-0.08}$	AUBERT,B	05N BABR	$e^+ e^- \rightarrow \gamma(4S)$

 $A_{CP}(B^+ \rightarrow K_0^*(1430)^0 \pi^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.064 \pm 0.032^{+0.023}_{-0.026}$	AUBERT,B	05N BABR	$e^+ e^- \rightarrow \gamma(4S)$

 $A_{CP}(B^+ \rightarrow K^*(892)^+ \pi^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.04 \pm 0.29 \pm 0.05$	AUBERT	05X BABR	$e^+ e^- \rightarrow \gamma(4S)$

 $A_{CP}(B^+ \rightarrow \rho^0 K^*(892)^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.20^{+0.32}_{-0.29} \pm 0.04$	AUBERT	03V BABR	$e^+ e^- \rightarrow \gamma(4S)$

 $A_{CP}(B^+ \rightarrow K^0 K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.15 \pm 0.33 \pm 0.03$	612 AUBERT,BE	05E BABR	$e^+ e^- \rightarrow \gamma(4S)$

612 Corresponds to 90% confidence range $-0.43 < A_{CP} < 0.68$. $A_{CP}(B^+ \rightarrow K^+ K_S^0 K_S^0)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$-0.04 \pm 0.11 \pm 0.02$	613 AUBERT,B	04V BABR	$e^+ e^- \rightarrow \gamma(4S)$

613 Corresponds to 90% confidence range $-0.23 < A_{CP} < 0.15$. $A_{CP}(B^+ \rightarrow K^+ K^- K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.02 \pm 0.07 \pm 0.03$	AUBERT	03M BABR	$e^+ e^- \rightarrow \gamma(4S)$

 $A_{CP}(B^+ \rightarrow \phi K^+)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.01 \pm 0.07 \text{ OUR AVERAGE}$			

 $-0.07 \pm 0.17^{+0.03}_{-0.02}$ ACOSTA 05J CDF $p\bar{p}$ at 1.96 TeV $0.04 \pm 0.09 \pm 0.01$ 614 AUBERT 04A BABR $e^+ e^- \rightarrow \gamma(4S)$ $0.01 \pm 0.12 \pm 0.05$ 615 CHEN 03B BELL $e^+ e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $-0.05 \pm 0.20 \pm 0.03$ 616 AUBERT 02E BABR $e^+ e^- \rightarrow \gamma(4S)$ 614 Corresponds to 90% confidence range $-0.10 < A_{CP} < 0.18$.615 Corresponds to 90% confidence range $-0.20 < A_{CP} < 0.22$.616 Corresponds to 90% confidence range $-0.37 < A_{CP} < 0.28$.

$A_{CP}(B^+ \rightarrow \phi K^*(892)^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.05±0.11 OUR AVERAGE			
-0.02±0.14±0.03	617 CHEN	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.16±0.17±0.03	AUBERT	03V BABR	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.13±0.29 ^{+0.08} _{-0.11}	618 CHEN	03B BELL	Repl. by CHEN 05A
-0.43 ^{+0.36} _{-0.30} ±0.06	619 AUBERT	02E BABR	Repl. by AUBERT 03V
617 Corresponds to 90% confidence range $-0.25 < A_{CP} < 0.22$.			
618 Corresponds to 90% confidence range $-0.64 < A_{CP} < 0.36$.			
619 Corresponds to 90% confidence range $-0.88 < A_{CP} < 0.18$.			

 $A_{CP}(B^+ \rightarrow \eta K^+ \gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.16±0.09±0.06	620 NISHIDA	05	BELL
620 $m_{\eta K^+} < 2.4 \text{ GeV}/c^2$			

 $A_{CP}(B^+ \rightarrow \pi^+ \pi^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.02±0.07 OUR AVERAGE			
-0.01±0.10±0.02	621 AUBERT	05L BABR	$e^+ e^- \rightarrow \gamma(4S)$
-0.02±0.10±0.01	622 CHAO	04B BELL	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.00±0.10±0.02	623 CHAO	05A BELL	Repl. by CHAO 04B
-0.03 ^{+0.18} _{-0.17} ±0.02	624 AUBERT	03L BABR	Repl. by AUBERT 05L
0.30±0.30 ^{+0.06} _{-0.04}	625 CASEY	02	BELL Repl. by CHAO 04B
621 Corresponds to a 90% CL interval of $-0.19 < A_{CP} < 0.21$.			
622 This corresponds to 90% CL interval of $-0.18 < A_{CP} < 0.14$.			
623 Corresponds to a 90% CL interval of $-0.17 < A_{CP} < 0.16$.			
624 Corresponds to 90% confidence range $-0.32 < A_{CP} < 0.27$.			
625 Corresponds to 90% confidence range $-0.23 < A_{CP} < +0.86$.			

 $A_{CP}(B^+ \rightarrow \pi^+ \pi^- \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.007±0.077±0.025	AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.39 ± 0.33 ± 0.12	AUBERT	03M BABR	Repl. by AUBERT 05G

 $A_{CP}(B^+ \rightarrow \rho^0 \pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.074±0.120^{+0.035}_{-0.055}	AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.19 ± 0.11 ± 0.02	AUBERT	04Z BABR	Repl. by AUBERT,B 05G

$A_{CP}(B^+ \rightarrow f_2(1270)\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.004 \pm 0.247^{+0.028}_{-0.032}$	AUBERT,B	05G BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \rho^+ \pi^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.15 ± 0.12 OUR AVERAGE	ZHANG	05A BELL	$e^+ e^- \rightarrow \gamma(4S)$
$0.06 \pm 0.17^{+0.04}_{-0.05}$	AUBERT	04Z BABR	$e^+ e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \rho^+ \rho^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.09 ± 0.16 OUR AVERAGE	AUBERT	03V BABR	$e^+ e^- \rightarrow \gamma(4S)$
$-0.19 \pm 0.23 \pm 0.03$	ZHANG	03B BELL	$e^+ e^- \rightarrow \gamma(4S)$


$A_{CP}(B^+ \rightarrow \omega\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.10 ± 0.22 OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.		
$0.03 \pm 0.16 \pm 0.01$	AUBERT	04H BABR	$e^+ e^- \rightarrow \gamma(4S)$
$0.50^{+0.23}_{-0.20} \pm 0.02$	626 WANG	04A BELL	$e^+ e^- \rightarrow \gamma(4S)$
-0.34 ± 0.25	627 CHEN	00 CLE2	$e^+ e^- \rightarrow \gamma(4S)$
$-0.01^{+0.29}_{-0.31} \pm 0.03$	628 AUBERT	02E BABR	Repl. by AUBERT 04H

626 Corresponds to 90% CL interval $-0.25 < A_{CP} < 0.41$

627 Corresponds to 90% confidence range $-0.75 < A_{CP} < 0.07$.

628 Corresponds to 90% confidence range $-0.50 < A_{CP} < 0.46$.



$A_{CP}(B^+ \rightarrow \omega\rho^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.05±0.26±0.02	AUBERT	050 BABR	$e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \eta\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.05±0.10 OUR AVERAGE			
-0.13±0.12±0.01	AUBERT,B	05K BABR	$e^+e^- \rightarrow \gamma(4S)$
0.07±0.15±0.03	CHANG	05A BELL	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.44±0.18±0.01	AUBERT	04H BABR	Repl. by AUBERT,B 05K

$A_{CP}(B^+ \rightarrow \eta'\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.14±0.16±0.01	AUBERT,B	05K BABR	$e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow \eta\rho^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.02±0.18±0.02	AUBERT,B	05K BABR	$e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow p\bar{p}\pi^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.16±0.22±0.01	WANG	04 BELL	$e^+e^- \rightarrow \gamma(4S)$

$A_{CP}(B^+ \rightarrow p\bar{p}K^+)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.05±0.11±0.01	WANG	04 BELL	$e^+e^- \rightarrow \gamma(4S)$

$\gamma(B^+ \rightarrow D^*(\star) K^+)$

For angle $\gamma(\phi_3)$ of the CKM unitarity triangle, see the review on “CP Violation” in the Reviews section.

VALUE (°)	DOCUMENT ID	TECN	COMMENT
75±20 OUR AVERAGE			
$70 \pm 31^{+18}_{-15}$	629 AUBERT,B	05Y BABR	$e^+ e^- \rightarrow \gamma(4S)$
$77^{+17}_{-19} \pm 17$	630 POLUEKTOV 04	BELL	$e^+ e^- \rightarrow \gamma(4S)$
629 Uses a Dalitz plot analysis of neutral $D \rightarrow K_S^0 \pi^+ \pi^-$ decays coming from $B^\pm \rightarrow D K^\pm$ and $B^\pm \rightarrow D^{*0} K^\pm$ followed by $D^{*0} \rightarrow D \pi^0$, $D \gamma$. The corresponding two standard deviations interval for gamma is $12^\circ < \gamma < 137^\circ$.			
630 Uses a Dalitz plot analysis of the 3-body $D \rightarrow K_S^0 \pi^+ \pi^-$ decays coming from $B^\pm \rightarrow D K^\pm$ and $B^\pm \rightarrow D^* K^\pm$ followed by $D^* \rightarrow D \pi^0$; here we use D to denote that the neutral D meson produced in the decay is an admixture of D^0 and \bar{D}^0 . The corresponding two standard deviations interval for γ is $26^\circ < \gamma < 126^\circ$. POLUEKTOV 04 also reports the amplitude ratios and the strong phases.			

 B^\pm REFERENCES

ABE	06	PR D73 051106	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06	PR D73 011101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06F	PR D73 011103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06J	PR D73 051105R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06K	PR D73 057101	B. Aubert <i>et al.</i>	(BABAR Collab.)
SONI	06	PL B634 155	N. Soni <i>et al.</i>	(BELLE Collab.)
ABE	05A	PRL 94 221805	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	05B	PR D71 072003	K. Abe <i>et al.</i>	(BELLE Collab.)
Also		PR D71 079903(Erratum)	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	05G	PRL 95 231802	K. Abe <i>et al.</i>	(BELLE Collab.)
ACOSTA	05J	PRL 95 031801	D. Acosta <i>et al.</i>	(CDF Collab.)
AUBERT	05	PRL 94 011801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05B	PR D71 031501R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05G	PR D72 032004	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05H	PRL 94 101801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05J	PRL 94 141801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05K	PRL 94 171801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05L	PRL 94 181802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05M	PRL 94 191802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05N	PR D71 031102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05O	PR D71 031103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05R	PR D71 071103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05U	PR D71 091103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05X	PR D71 111101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05B	PRL 95 041804	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05E	PR D72 011102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05G	PR D72 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05K	PRL 95 131803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05L	PR D72 051101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05N	PR D72 072003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05O	PR D72 051102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05T	PR D72 071102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05U	PR D72 071103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05V	PR D72 071104R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	05Y	PRL 95 121802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	05E	PRL 95 221801	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHANG	05	PR D71 072007	M.-C. Chang <i>et al.</i>	(BELLE Collab.)
CHANG	05A	PR D71 091106R	P. Chang <i>et al.</i>	(BELLE Collab.)
CHAO	05A	PR D71 031502R	Y. Chao <i>et al.</i>	(BELLE Collab.)
CHEN	05A	PRL 94 221804	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)
ITOH	05	PRL 95 091601	R. Itoh <i>et al.</i>	(BELLE Collab.)

LEE	05	PRL 95 061802	Y.-J. Lee <i>et al.</i>	(BELLE Collab.)
LIVENTSEV	05	PR D72 051109R	D. Liventsev <i>et al.</i>	(BELLE Collab.)
MAJUMDER	05	PRL 95 041803	G. Majumder <i>et al.</i>	(BELLE Collab.)
MOHAPATRA	05	PR D72 011101R	D. Mohapatra <i>et al.</i>	(BELLE Collab.)
NISHIDA	05	PL B610 23	S. Nishida <i>et al.</i>	(BELLE Collab.)
OKABE	05	PL B614 27	T. Okabe <i>et al.</i>	(BELLE Collab.)
SAIGO	05	PRL 94 091601	M. Saigo <i>et al.</i>	(BELLE Collab.)
WANG	05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
XIE	05	PR D72 051105R	Q.L. Xie <i>et al.</i>	(BELLE Collab.)
YANG	05	PRL 94 111802	H. Yang <i>et al.</i>	(BELLE Collab.)
ZHANG	05A	PRL 94 031801	J. Zhang <i>et al.</i>	(BELLE Collab.)
ZHANG	05B	PR D71 091107R	L.M. Zhang <i>et al.</i>	(BELLE Collab.)
ZHANG	05D	PRL 95 141801	J. Zhang <i>et al.</i>	(BELLE Collab.)
ABDALLAH	04E	EPJ C33 307	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	04A	PR D69 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04C	PRL 92 111801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	04H	PRL 92 061801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04K	PRL 92 141801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04M	PRL 92 201802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04N	PRL 92 202002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04O	PRL 92 221803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04P	PRL 92 241802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04Q	PR D69 051101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04T	PR D69 071103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04Y	PRL 93 041801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	04Z	PRL 93 051802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04B	PR D70 011101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04D	PR D70 032006	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04L	PRL 93 131804	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04P	PR D70 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04U	PR D70 091105R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04V	PRL 93 181805	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	04	PR D70 111102R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	04A	PR D70 112006	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	04B	PR D70 091106	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHAO	04	PR D69 111102R	Y. Chao <i>et al.</i>	(BELLE Collab.)
CHAO	04B	PRL 93 191802	Y. Chao <i>et al.</i>	(BELLE Collab.)
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	04	PRL 92 051801	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
GARMASH	04	PR D69 012001	A. Garmash <i>et al.</i>	(BELLE Collab.)
LEE	04	PRL 93 211801	Y.-J. Lee <i>et al.</i>	(BELLE Collab.)
MAJUMDER	04	PR D70 111103R	G. Majumder <i>et al.</i>	(BELLE Collab.)
NAKAO	04	PR D69 112001	M. Nakao <i>et al.</i>	(BELLE Collab.)
POLUEKTOV	04	PR D70 072003	A. Poluektov <i>et al.</i>	(BELLE Collab.)
SCHWANDA	04	PRL 93 131803	C. Schwanda <i>et al.</i>	(BELLE Collab.)
WANG	04	PRL 92 131801	M.Z. Wang <i>et al.</i>	(BELLE Collab.)
WANG	04A	PR D70 012001	C.H. Wang <i>et al.</i>	(BELLE Collab.)
ZANG	04	PR D69 017101	S.L. Zang <i>et al.</i>	(BELLE Collab.)
ABE	03B	PR D67 032003	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	03D	PRL 90 131803	K. Abe <i>et al.</i>	(BELLE Collab.)
ADAM	03	PR D67 032001	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ADAM	03B	PR D68 012004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ATHAR	03	PR D68 072003	S.B. Athar <i>et al.</i>	(CLEO Collab.)
AUBERT	03K	PRL 90 231801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03L	PRL 91 021801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03M	PRL 91 051801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03O	PRL 91 071801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03U	PRL 91 221802	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03V	PRL 91 171802	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03W	PRL 91 161801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	03X	PR D68 092001	B. Aubert <i>et al.</i>	(BaBar Collab.)
BORNHEIM	03	PR D68 052002	A. Bornheim <i>et al.</i>	(CLEO Collab.)
CHEN	03B	PRL 91 201801	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
CHOI	03	PRL 91 262001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
CSORNA	03	PR D67 112002	S.E. Csorna <i>et al.</i>	(CLEO Collab.)
EDWARDS	03	PR D68 011102R	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
FANG	03	PRL 90 071801	F. Fang <i>et al.</i>	(BELLE Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
ISHIKAWA	03	PRL 91 261601	A. Ishikawa <i>et al.</i>	(BELLE Collab.)

KROKOVNY	03B	PRL 91 262002	P. Krovovny <i>et al.</i>	(BELLE Collab.)
SWAIN	03	PR D68 051101R	S.K. Swain <i>et al.</i>	(BELLE Collab.)
UNNO	03	PR D68 011103R	Y. Unno <i>et al.</i>	(BELLE Collab.)
ZHANG	03B	PRL 91 221801	J. Zhang <i>et al.</i>	(BELLE Collab.)
ABE	02	PRL 88 021801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02B	PRL 88 031802	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02H	PRL 88 171801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02K	PRL 88 181803	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02N	PL B538 11	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02O	PR D65 091103R	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02W	PRL 89 151802	K. Abe <i>et al.</i>	(BELLE Collab.)
ACOSTA	02C	PR D65 092009	D. Acosta <i>et al.</i>	(CDF Collab.)
ACOSTA	02F	PR D66 052005	D. Acosta <i>et al.</i>	(CDF Collab.)
AHMED	02B	PR D66 031101R	S. Ahmed <i>et al.</i>	(CLEO Collab.)
AUBERT	02	PR D65 032001	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02C	PRL 88 101805	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02E	PR D65 051101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02F	PR D65 091101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	02L	PRL 88 241801	B. Aubert <i>et al.</i>	(BaBar Collab.)
BRIERE	02	PRL 89 081803	R. Briere <i>et al.</i>	(CLEO Collab.)
CASEY	02	PR D66 092002	B.C.K. Casey <i>et al.</i>	(BELLE Collab.)
CHEN	02B	PL B546 196	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	02	PL B542 171	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
DYTMAN	02	PR D66 091101R	S.A. Dytman <i>et al.</i>	(CLEO Collab.)
ECKHART	02	PRL 89 251801	E. Eckhart <i>et al.</i>	(CLEO Collab.)
EDWARDS	02B	PR D65 111102R	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
GABYSHEV	02	PR D66 091102R	N. Gabyshev <i>et al.</i>	(BELLE Collab.)
GARMASH	02	PR D65 092005	A. Garmash <i>et al.</i>	(BELLE Collab.)
GODANG	02	PRL 88 021802	R. Godang <i>et al.</i>	(CLEO Collab.)
GORDON	02	PL B542 183	A. Gordon <i>et al.</i>	(BELLE Collab.)
LU	02	PRL 89 191801	R.-S. Lu <i>et al.</i>	(BELLE Collab.)
MAHAPATRA	02	PRL 88 101803	R. Mahapatra <i>et al.</i>	(CLEO Collab.)
NISHIDA	02	PRL 89 231801	S. Nishida <i>et al.</i>	(BELLE Collab.)
ABE	01H	PRL 87 101801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01I	PRL 87 111801	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01K	PR D64 071101	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01L	PRL 87 161601	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	01M	PL B517 309	K. Abe <i>et al.</i>	(BELLE Collab.)
ALEXANDER	01B	PR D64 092001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AMMAR	01B	PRL 87 271801	R. Ammar <i>et al.</i>	(CLEO Collab.)
ANDERSON	01B	PRL 87 181803	S. Anderson <i>et al.</i>	(CLEO Collab.)
AUBERT	01D	PRL 87 151801	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01E	PRL 87 151802	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01F	PRL 87 201803	B. Aubert <i>et al.</i>	(BaBar Collab.)
AUBERT	01G	PRL 87 221802	B. Aubert <i>et al.</i>	(BaBar Collab.)
BARATE	01E	EPJ C19 213	R. Barate <i>et al.</i>	(ALEPH Collab.)
BRIERE	01	PRL 86 3718	R.A. Biere <i>et al.</i>	(CLEO Collab.)
BROWDER	01	PRL 86 2950	T.E. Browder <i>et al.</i>	(CLEO Collab.)
EDWARDS	01	PRL 86 30	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
GRITSAN	01	PR D64 077501	A. Gritsan <i>et al.</i>	(CLEO Collab.)
RICHICHI	01	PR D63 031103R	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ABBIENDI	00B	PL B476 233	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABE	00C	PR D62 071101R	K. Abe <i>et al.</i>	(SLD Collab.)
AHMED	00B	PR D62 112003	S. Ahmed <i>et al.</i>	(CLEO Collab.)
ANASTASSOV	00	PRL 84 1393	A. Anastassov <i>et al.</i>	(CLEO Collab.)
BARATE	00R	PL B492 275	R. Barate <i>et al.</i>	(ALEPH Collab.)
BEHRENS	00	PR D61 052001	B.H. Behrens <i>et al.</i>	(CLEO Collab.)
BONVICINI	00	PRL 84 5940	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
CHEN	00	PRL 85 525	S. Chen <i>et al.</i>	(CLEO Collab.)
COAN	00	PRL 84 5283	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN... CSORNA	00	PRL 85 515 PR D61 111101	D. Cronin-Hennessy <i>et al.</i> S.E. Csorna <i>et al.</i>	(CLEO Collab.)
JESSOP	00	PRL 85 2881	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
RICHICHI	00	PRL 85 520	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ABBIENDI	99J	EPJ C12 609	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AFFOLDER	99B	PRL 83 3378	T. Affolder <i>et al.</i>	(CDF Collab.)
BARTELT	99	PRL 82 3746	J. Bartelt <i>et al.</i>	(CLEO Collab.)
COAN	99	PR D59 111101	T.E. Coan <i>et al.</i>	(CLEO Collab.)
ABE	98B	PR D57 5382	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98O	PR D58 072001	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98Q	PR D58 092002	F. Abe <i>et al.</i>	(CDF Collab.)

ACCIARRI	98S	PL B438 417	M. Acciari <i>et al.</i>	(L3 Collab.)
ANASTASSOV	98	PRL 80 4127	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ATHANAS	98	PRL 80 5493	M. Athanas <i>et al.</i>	(CLEO Collab.)
BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
BEHRENS	98	PRL 80 3710	B.H. Behrens <i>et al.</i>	(CLEO Collab.)
BERGFELD	98	PRL 81 272	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BRANDENB...	98	PRL 80 2762	G. Brandenbrug <i>et al.</i>	(CLEO Collab.)
GODANG	98	PRL 80 3456	R. Godang <i>et al.</i>	(CLEO Collab.)
ABE	97J	PRL 79 590	K. Abe <i>et al.</i>	(SLD Collab.)
ACCIARRI	97F	PL B396 327	M. Acciari <i>et al.</i>	(L3 Collab.)
ARTUSO	97	PL B399 321	M. Artuso <i>et al.</i>	(CLEO Collab.)
ATHANAS	97	PRL 79 2208	M. Athanas <i>et al.</i>	(CLEO Collab.)
BROWDER	97	PR D56 11	T. Browder <i>et al.</i>	(CLEO Collab.)
FU	97	PRL 79 3125	X. Fu <i>et al.</i>	(CLEO Collab.)
JESSOP	97	PRL 79 4533	C.P. Jessop <i>et al.</i>	(CLEO Collab.)
ABE	96B	PR D53 3496	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96C	PRL 76 4462	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96H	PRL 76 2015	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96L	PRL 76 4675	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96Q	PR D54 6596	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96R	PRL 77 5176	F. Abe <i>et al.</i>	(CDF Collab.)
ADAM	96D	ZPHY C72 207	W. Adam <i>et al.</i>	(DELPHI Collab.)
ALEXANDER	96T	PRL 77 5000	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ASNER	96	PR D53 1039	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BERGFELD	96B	PRL 77 4503	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
BISHAI	96	PL B369 186	M. Bishai <i>et al.</i>	(CLEO Collab.)
BUSKULIC	96J	ZPHY C71 31	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
GIBAUT	96	PR D53 4734	D. Gibaut <i>et al.</i>	(CLEO Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	
ABREU	95N	PL B357 255	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95Q	ZPHY C68 13	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ADAM	95	ZPHY C68 363	W. Adam <i>et al.</i>	(DELPHI Collab.)
AKERS	95T	ZPHY C67 379	R. Akers <i>et al.</i>	(OPAL Collab.)
ALBRECHT	95D	PL B353 554	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	95	PL B341 435	J. Alexander <i>et al.</i>	(CLEO Collab.)
Also		PL B347 469 (erratum)	J. Alexander <i>et al.</i>	(CLEO Collab.)
ARTUSO	95	PRL 75 785	M. Artuso <i>et al.</i>	(CLEO Collab.)
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)
BUSKULIC	95	PL B343 444	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABE	94D	PRL 72 3456	F. Abe <i>et al.</i>	(CDF Collab.)
ALAM	94	PR D50 43	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	94D	PL B335 526	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ATHANAS	94	PRL 73 3503	M. Athanas <i>et al.</i>	(CLEO Collab.)
Also		PRL 74 3090 (erratum)	M. Athanas <i>et al.</i>	(CLEO Collab.)
PDG	94	PR D50 1173	L. Montanet <i>et al.</i>	(CERN, LBL, BOST+)
STONE	94	HEPSY 93-11	S. Stone	
Published in B Decays, 2nd Edition, World Scientific, Singapore				
ABREU	93D	ZPHY C57 181	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	93G	PL B312 253	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	93C	PL B307 247	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ALBRECHT	93E	ZPHY C60 11	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	93B	PL B319 365	J. Alexander <i>et al.</i>	(CLEO Collab.)
AMMAR	93	PRL 71 674	R. Ammar <i>et al.</i>	(CLEO Collab.)
BEAN	93B	PRL 70 2681	A. Bean <i>et al.</i>	(CLEO Collab.)
BUSKULIC	93D	PL B307 194	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
Also		PL B325 537 (erratum)	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
SANGHERA	93	PR D47 791	S. Sanghera <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92C	PL B275 195	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92E	PL B277 209	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	92G	ZPHY C54 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	92	PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
BUSKULIC	92G	PL B295 396	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ALBRECHT	91B	PL B254 288	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91C	PL B255 297	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	91E	PL B262 148	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BERKELMAN	91	ARNPS 41 1	K. Berkelman, S. Stone	(CORN, SYRA)
"Decays of B Mesons"				
FULTON	91	PR D43 651	R. Fulton <i>et al.</i>	(CLEO Collab.)
ALBRECHT	90B	PL B241 278	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	90J	ZPHY C48 543	H. Albrecht <i>et al.</i>	(ARGUS Collab.)

ANTREASYAN	90B	ZPHY C48 553	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
BORTOLETTO	90	PRL 64 2117	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
Also		PR D45 21	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
WEIR	90B	PR D41 1384	A.J. Weir <i>et al.</i>	(Mark II Collab.)
ALBRECHT	89G	PL B229 304	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AVERY	89B	PL B223 470	P. Avery <i>et al.</i>	(CLEO Collab.)
BEBEK	89	PRL 62 8	C. Bebek <i>et al.</i>	(CLEO Collab.)
BORTOLETTO	89	PRL 62 2436	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
ALBRECHT	88F	PL B209 119	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	88K	PL B215 424	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87C	PL B185 218	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	87D	PL B199 451	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AVERY	87	PL B183 429	P. Avery <i>et al.</i>	(CLEO Collab.)
BEBEK	87	PR D36 1289	C. Bebek <i>et al.</i>	(CLEO Collab.)
ALAM	86	PR D34 3279	M.S. Alam <i>et al.</i>	(CLEO Collab.)
PDG	86	PL 170B	M. Aguilar-Benitez <i>et al.</i>	(CERN, CIT+)
GILES	84	PR D30 2279	R. Giles <i>et al.</i>	(CLEO Collab.)