

THE $\rho(1450)$ AND THE $\rho(1700)$

Updated April 2006 by S. Eidelman (Novosibirsk) and J.J Hernandez-Rey (Valencia).

In our 1988 edition, we replaced the $\rho(1600)$ entry with two new ones, the $\rho(1450)$ and the $\rho(1700)$, because there was emerging evidence that the 1600-MeV region actually contains two ρ -like resonances. ERKAL 86 had pointed out this possibility with a theoretical analysis on the consistency of 2π and 4π electromagnetic form factors and the $\pi\pi$ scattering length. DONNACHIE 87, with a full analysis of data on the 2π and 4π final states in e^+e^- annihilation and photoproduction reactions, had also argued that in order to obtain a consistent picture, two resonances were necessary. The existence of $\rho(1450)$ was supported by the analysis of $\eta\rho^0$ mass spectra obtained in photoproduction and e^+e^- annihilation (DONNACHIE 87B), as well as that of $e^+e^- \rightarrow \omega\pi$ (DONNACHIE 91).

The analysis of DONNACHIE 87 was further extended by CLEGG 88, 94 to include new data on 4π systems produced in e^+e^- annihilation, and in τ decays (τ decays to 4π and e^+e^- annihilation to 4π can be related by the Conserved Vector Current assumption). These systems were successfully analyzed using interfering contributions from two ρ -like states, and from the tail of the $\rho(770)$ decaying into two-body states. While specific conclusions on $\rho(1450) \rightarrow 4\pi$ were obtained, little could be said about the $\rho(1700)$.

Independent evidence for two 1^- states is provided by KILLIAN 80 in 4π electroproduction at $\langle Q^2 \rangle = 1$ (GeV/c) 2 , and by FUKUI 88 in a high-statistics sample of the $\eta\pi\pi$ system in π^-p charge exchange.

This scenario with two overlapping resonances is supported by other data. BISELLO 89 measured the pion form factor in the interval 1.35–2.4 GeV and observed a deep minimum around 1.6 GeV. The best fit was obtained with the hypothesis of ρ -like resonances at 1420 and 1770 MeV, with widths of about 250 MeV. ANTONELLI 88 found that the $e^+e^- \rightarrow \eta\pi^+\pi^-$ cross section is better fitted with two fully interfering Breit-Wigners, with parameters in fair agreement with those of

DONNACHIE 87 and BISELLO 89. These results can be considered as a confirmation of the $\rho(1450)$.

Decisive evidence for the $\pi\pi$ decay mode of both $\rho(1450)$ and $\rho(1700)$ came from recent results in $\bar{p}p$ annihilation at rest (ABELE 97). It was shown that these resonances also possess a $K\bar{K}$ decay mode (ABELE 98, BERTIN 98B, ABELE 99D). High statistics studies of the decays $\tau \rightarrow \pi\pi\nu_\tau$ (BARATE 97M, URHEIM 97), and $\tau \rightarrow 4\pi\nu_\tau$ (EDWARDS 00A), also require the $\rho(1450)$, but are not sensitive to the $\rho(1700)$, because it is too close to the τ mass. Recently in a very high statistics study of the $\tau \rightarrow \pi\pi\nu_\tau$ decay performed at Belle (ABE 05H) both $\rho(1450)$ and $\rho(1700)$ were observed for the first time in τ decays.

The structure of these ρ states is not yet completely clear. BARNES 97 and CLOSE 97C claim that $\rho(1450)$ has a mass consistent with radial $2S$, but its decays show characteristics of hybrids, and suggest that this state may be a $2S$ -hybrid mixture. DONNACHIE 99 argues that hybrid states could have a 4π decay mode dominated by the $a_1\pi$. Such behavior has recently been observed by AKHMETSHIN 99E in $e^+e^- \rightarrow 4\pi$ in the energy range 1.05–1.38 GeV, and by EDWARDS 00A in $\tau \rightarrow 4\pi$ decays. ALEXANDER 01B observed the $\rho(1450) \rightarrow \omega\pi$ decay mode in B-meson decays, however, didn't find $\rho(1700) \rightarrow \omega\pi^0$. A similar conclusion is made by AKHMETSHIN 03B who studied the process $e^+e^- \rightarrow \omega\pi^0$. Various decay modes of the $\rho(1450)$ and $\rho(1700)$ were observed in $\bar{p}n$ and $\bar{p}p$ annihilation (ABELE 01B, BARGIOTTI 03B), but no definite conclusions could be drawn. More data should be collected to clarify the nature of the ρ states, particularly in the energy range above 1.6 GeV.

We also list under the $\rho(1450)$ the $\phi\pi$ state with $J^{PC} = 1^{--}$ or $C(1480)$ observed by BITYUKOV 87. While ACHASOV 96B shows that it may be a threshold effect, CLEGG 88 and LANDSBERG 92 suggest two independent vector states with this decay mode. Note, however, that $C(1480)$ in its $\phi\pi$ decay mode was not confirmed by e^+e^- (DOLINSKY 91, BISELLO 91C) and $\bar{p}p$ (ABELE 97H) experiments.

Several observations on the $\omega\pi$ system in the 1200-MeV region (FRENKIEL 72, COSME 76, BARBER 80C, ASTON 80C, ATKINSON 84C, BRAU 88, AMSLER 93B) may be interpreted in

terms of either $J^P = 1^- \rho(770) \rightarrow \omega\pi$ production (**LAYSSAC 71**), or $J^P = 1^+ b_1(1235)$ production (**BRAU 88**, **AMSLER 93B**). We argue that no special entry for a $\rho(1250)$ is needed. The LASS amplitude analysis (**ASTON 91B**) showing evidence for $\rho(1270)$ is preliminary and needs confirmation. For completeness, the relevant observations are listed under the $\rho(1450)$.

Evidence for ρ -like mesons decaying into 6π states was first noted by **CLEGG 90** in the analysis of 6π mass spectra from e^+e^- annihilation (**BISELLO 81**, **CASTRO 88**) and diffractive photoproduction (**ATKINSON 85**). **CLEGG 90** argued that two states at about 2.1 and 1.8 GeV exist: while the former is a candidate for a new resonance ($\rho(2150)$), the latter could be a manifestation of the $\rho(1700)$ distorted by threshold effects. Recently, the E687 Collaboration at Fermilab reported an observation of a narrow dip structure at 1.9 GeV in the $3\pi^+3\pi^-$ diffractive photoproduction (**FRABETTI 01**). A similar effect of the dip in the cross section of $e^+e^- \rightarrow 6\pi$ around 1.9 GeV has been earlier reported by DM2 (**CASTRO 88**), where 6π included both $3\pi^+3\pi^-$ and $2\pi^+2\pi^-2\pi^0$. Later the dip in the R value (the total cross section of $e^+e^- \rightarrow$ hadrons divided by the cross section of $e^+e^- \rightarrow \mu^+\mu^-$) was observed by **ANTONELLI 96**, again around 1.9 GeV. This energy is close to the $N\bar{N}$ threshold which hints to the possible relation between the dip and $N\bar{N}$, e.g., the frequently discussed narrow $N\bar{N}$ resonance or just a threshold effect. Such behaviour is also characteristic of exotic objects like vector $q\bar{q}$ hybrids. Note that **AGNELLO 02** failed to find this state in the reaction $\bar{n}p \rightarrow 3\pi^+2\pi^-\pi^0$. A reanalysis of the E687 data by **FRABETTI 04** shows that a dip may arise due to interference of a narrow object with a broad $\rho(1700)$ independently of the nature of the former. Recently BaBar studied the processes $e^+e^- \rightarrow 3\pi^+3\pi^-$ and $e^+e^- \rightarrow 2\pi^+2\pi^-2\pi^0$ using the radiative return and observed a structure around 1.9 GeV in both final states (**AUBERT 06D**). The data are not well described by a single Breit-Wigner state, and a good fit is achieved while taking into account the interference of such a structure with a Jacob-Slansky amplitude for continuum. The mass of this state obtained by BaBar is consistent with **ANTONELLI 96** and **FRABETTI 01**, but the width is substantially larger. We list

these observations under a separate particle $\rho(1900)$, which needs confirmation.