

CHARMED BARYONS ($C = +1$)

$$\begin{aligned}\Lambda_c^+ &= u d c, & \Sigma_c^{++} &= u u c, & \Sigma_c^+ &= u d c, & \Sigma_c^0 &= d d c, \\ \Xi_c^+ &= u s c, & \Xi_c^0 &= d s c, & \Omega_c^0 &= s s c\end{aligned}$$

Λ_c^+

$$I(J^P) = 0(\frac{1}{2}^+)$$

J is not well measured; $\frac{1}{2}$ is the quark-model prediction.

Mass $m = 2286.46 \pm 0.14$ MeV

Mean life $\tau = (200 \pm 6) \times 10^{-15}$ s ($S = 1.6$)

$$c\tau = 59.9 \mu\text{m}$$

Decay asymmetry parameters

$$\Lambda\pi^+ \quad \alpha = -0.91 \pm 0.15$$

$$\Sigma^+\pi^0 \quad \alpha = -0.45 \pm 0.32$$

$$\Lambda\ell^+\nu_\ell \quad \alpha = -0.86 \pm 0.04$$

$$[\alpha(\Lambda_c^+) + \alpha(\bar{\Lambda}_c^-)] / [\alpha(\Lambda_c^+) - \alpha(\bar{\Lambda}_c^-)] \text{ in } \Lambda_c^+ \rightarrow \Lambda\pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda}\pi^- = -0.07 \pm 0.31$$

$$[\alpha(\Lambda_c^+) + \alpha(\bar{\Lambda}_c^-)] / [\alpha(\Lambda_c^+) - \alpha(\bar{\Lambda}_c^-)] \text{ in } \Lambda_c^+ \rightarrow \Lambda e^+ \nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} e^- \bar{\nu}_e = 0.00 \pm 0.04$$

Nearly all branching fractions of the Λ_c^+ are measured relative to the $pK^-\pi^+$ mode, but there are no model-independent measurements of this branching fraction. We explain how we arrive at our value of $B(\Lambda_c^+ \rightarrow pK^-\pi^+)$ in a Note at the beginning of the branching-ratio measurements in the Listings. When this branching fraction is eventually well determined, all the other branching fractions will slide up or down proportionally as the true value differs from the value we use here.

| Λ_c^+ DECAY MODES | Fraction (Γ_i/Γ) | Scale factor/ Confidence level | p (MeV/c) |
|---|------------------------------------|-----------------------------------|----------------|
| Hadronic modes with a p: $S = -1$ final states | | | |
| $p\bar{K}^0$ | (2.3 \pm 0.6) % | | 873 |
| $pK^-\pi^+$ | [a] (5.0 \pm 1.3) % | | 823 |
| $p\bar{K}^*(892)^0$ | [b] (1.6 \pm 0.5) % | | 685 |
| $\Delta(1232)^{++}K^-$ | (8.6 \pm 3.0) $\times 10^{-3}$ | | 710 |
| $\Lambda(1520)\pi^+$ | [b] (1.8 \pm 0.6) % | | 627 |
| $pK^-\pi^+$ nonresonant | (2.8 \pm 0.8) % | | 823 |
| $p\bar{K}^0\pi^0$ | (3.3 \pm 1.0) % | | 823 |

| | | |
|---|----------------------------------|-----|
| $p\bar{K}^0\eta$ | (1.2 ± 0.4) % | 568 |
| $p\bar{K}^0\pi^+\pi^-$ | (2.6 ± 0.7) % | 754 |
| $pK^-\pi^+\pi^0$ | (3.4 ± 1.0) % | 759 |
| $pK^*(892)^-\pi^+$ | [b] (1.1 ± 0.5) % | 580 |
| $p(K^-\pi^+)_{\text{nonresonant}}\pi^0$ | (3.6 ± 1.2) % | 759 |
| $\Delta(1232)\bar{K}^*(892)$ | seen | 419 |
| $pK^-\pi^+\pi^+\pi^-$ | (1.1 ± 0.8) × 10 ⁻³ | 671 |
| $pK^-\pi^+\pi^0\pi^0$ | (8 ± 4) × 10 ⁻³ | 678 |

Hadronic modes with a p : $S = 0$ final states

| | | |
|----------------------------|--------------------------------------|-----|
| $p\pi^+\pi^-$ | (3.5 ± 2.0) × 10 ⁻³ | 927 |
| $p f_0(980)$ | [b] (2.8 ± 1.9) × 10 ⁻³ | 622 |
| $p\pi^+\pi^+\pi^-\pi^-$ | (1.8 ± 1.2) × 10 ⁻³ | 852 |
| pK^+K^- | (7.7 ± 3.5) × 10 ⁻⁴ | 616 |
| $p\phi$ | [b] (8.2 ± 2.7) × 10 ⁻⁴ | 590 |
| $pK^+K^- \text{ non-}\phi$ | (3.5 ± 1.7) × 10 ⁻⁴ | 616 |

Hadronic modes with a hyperon: $S = -1$ final states

| | | |
|---|--------------------------------------|------------|
| $\Lambda\pi^+$ | (1.01 ± 0.28) % | 864 |
| $\Lambda\pi^+\pi^0$ | (3.6 ± 1.3) % | 844 |
| $\Lambda\rho^+$ | < 5 % | CL=95% 635 |
| $\Lambda\pi^+\pi^+\pi^-$ | (2.6 ± 0.7) % | 807 |
| $\Sigma(1385)^+\pi^+\pi^-$, $\Sigma^{*+} \rightarrow$ | (7 ± 4) × 10 ⁻³ | 688 |
| $\Lambda\pi^+$ | | |
| $\Sigma(1385)^-\pi^+\pi^+$, $\Sigma^{*-} \rightarrow$ | (5.5 ± 1.7) × 10 ⁻³ | 688 |
| $\Lambda\pi^-$ | | |
| $\Lambda\pi^+\rho^0$ | (1.1 ± 0.5) % | 523 |
| $\Sigma(1385)^+\rho^0$, $\Sigma^{*+} \rightarrow \Lambda\pi^+$ | (3.7 ± 3.1) × 10 ⁻³ | 363 |
| $\Lambda\pi^+\pi^+\pi^- \text{ nonresonant}$ | < 8 × 10 ⁻³ | CL=90% 807 |
| $\Lambda\pi^+\pi^+\pi^-\pi^0 \text{ total}$ | (1.8 ± 0.8) % | 757 |
| $\Lambda\pi^+\eta$ | [b] (1.8 ± 0.6) % | 691 |
| $\Sigma(1385)^+\eta$ | [b] (8.5 ± 3.3) × 10 ⁻³ | 570 |
| $\Lambda\pi^+\omega$ | [b] (1.2 ± 0.5) % | 517 |
| $\Lambda\pi^+\pi^+\pi^-\pi^0$, no η or ω | < 7 × 10 ⁻³ | CL=90% 757 |
| $\Lambda K^+\bar{K}^0$ | (6.5 ± 2.0) × 10 ⁻³ | 443 |
| $\Xi(1690)^0K^+$, $\Xi^{*0} \rightarrow \Lambda\bar{K}^0$ | (1.9 ± 0.7) × 10 ⁻³ | 286 |
| $\Sigma^0\pi^+$ | (1.04 ± 0.31) % | 825 |
| $\Sigma^+\pi^0$ | (1.00 ± 0.34) % | 827 |
| $\Sigma^+\eta$ | (5.5 ± 2.3) × 10 ⁻³ | 714 |
| $\Sigma^+\pi^+\pi^-$ | (3.6 ± 1.0) % | 804 |
| $\Sigma^+\rho^0$ | < 1.4 % | CL=95% 575 |
| $\Sigma^-\pi^+\pi^+$ | (1.9 ± 0.8) % | 799 |
| $\Sigma^0\pi^+\pi^0$ | (1.8 ± 0.8) % | 803 |
| $\Sigma^0\pi^+\pi^+\pi^-$ | (8.3 ± 3.1) × 10 ⁻³ | 763 |
| $\Sigma^+\pi^+\pi^-\pi^0$ | — | 767 |

| | | | |
|---|-----|------------------------------------|-----|
| $\Sigma^+ \omega$ | [b] | (2.7 \pm 1.0) % | 569 |
| $\Sigma^+ K^+ K^-$ | | (2.8 \pm 0.8) $\times 10^{-3}$ | 349 |
| $\Sigma^+ \phi$ | [b] | (3.2 \pm 1.0) $\times 10^{-3}$ | 295 |
| $\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow$ | | (8.2 \pm 3.1) $\times 10^{-4}$ | 286 |
| $\Sigma^+ K^-$ | | | |
| $\Sigma^+ K^+ K^-$ nonresonant | | < 7 $\times 10^{-4}$ CL=90% | 349 |
| $\Xi^0 K^+$ | | (3.9 \pm 1.4) $\times 10^{-3}$ | 653 |
| $\Xi^- K^+ \pi^+$ | | (4.9 \pm 1.7) $\times 10^{-3}$ | 566 |
| $\Xi(1530)^0 K^+$ | [b] | (2.6 \pm 1.0) $\times 10^{-3}$ | 473 |

Hadronic modes with a hyperon: $S = 0$ final states

| | | | |
|-----------------------|-----|------------------------------------|-----|
| ΛK^+ | | (7.5 \pm 2.6) $\times 10^{-4}$ | 781 |
| $\Sigma^0 K^+$ | | (5.8 \pm 2.4) $\times 10^{-4}$ | 735 |
| $\Sigma^+ K^+ \pi^-$ | | (1.7 \pm 0.7) $\times 10^{-3}$ | 670 |
| $\Sigma^+ K^*(892)^0$ | [b] | (2.8 \pm 1.1) $\times 10^{-3}$ | 470 |
| $\Sigma^- K^+ \pi^+$ | | < 1.0 $\times 10^{-3}$ CL=90% | 664 |

Doubly Cabibbo-suppressed modes

| | | | |
|---------------|--|-------------------------------|-----|
| $p K^+ \pi^-$ | | < 2.3 $\times 10^{-4}$ CL=90% | 823 |
|---------------|--|-------------------------------|-----|

Semileptonic modes

| | | | |
|---------------------------|-----|---------------------|-----|
| $\Lambda \ell^+ \nu_\ell$ | [c] | (2.0 \pm 0.6) % | 871 |
| $\Lambda e^+ \nu_e$ | | (2.1 \pm 0.6) % | 871 |
| $\Lambda \mu^+ \nu_\mu$ | | (2.0 \pm 0.7) % | 867 |

Inclusive modes

| | | | |
|------------------------------|-----|---------------------|-------|
| e^+ anything | | (4.5 \pm 1.7) % | — |
| $p e^+$ anything | | (1.8 \pm 0.9) % | — |
| p anything | | (50 \pm 16) % | — |
| p anything (no Λ) | | (12 \pm 19) % | — |
| n anything | | (50 \pm 16) % | — |
| n anything (no Λ) | | (29 \pm 17) % | — |
| Λ anything | | (35 \pm 11) % | S=1.4 |
| Σ^\pm anything | [d] | (10 \pm 5) % | — |
| 3prongs | | (24 \pm 8) % | — |

$\Delta C = 1$ weak neutral current ($C1$) modes, or Lepton number (L) violating modes

| | | | |
|------------------------|------|-------------------------------|-----|
| $p \mu^+ \mu^-$ | $C1$ | < 3.4 $\times 10^{-4}$ CL=90% | 937 |
| $\Sigma^- \mu^+ \mu^+$ | L | < 7.0 $\times 10^{-4}$ CL=90% | 812 |

$\Lambda_c(2593)^+$

$$I(J^P) = 0(\frac{1}{2}^-)$$

The spin-parity follows from the fact that $\Sigma_c(2455)\pi$ decays, with little available phase space, are dominant. This assumes that $J^P = 1/2^+$ for the $\Sigma_c(2455)$.

$$\begin{aligned} \text{Mass } m &= 2595.4 \pm 0.6 \text{ MeV } (S = 1.1) \\ m - m_{\Lambda_c^+} &= 308.9 \pm 0.6 \text{ MeV } (S = 1.1) \\ \text{Full width } \Gamma &= 3.6^{+2.0}_{-1.3} \text{ MeV} \end{aligned}$$

$\Lambda_c^+\pi\pi$ and its submode $\Sigma_c(2455)\pi$ — the latter just barely — are the only strong decays allowed to an excited Λ_c^+ having this mass; and the submode seems to dominate.

| $\Lambda_c(2593)^+$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $\Lambda_c^+\pi^+\pi^-$ | [e] $\approx 67\%$ | 124 |
| $\Sigma_c(2455)^{++}\pi^-$ | $24 \pm 7\%$ | 28 |
| $\Sigma_c(2455)^0\pi^+$ | $24 \pm 7\%$ | 28 |
| $\Lambda_c^+\pi^+\pi^-$ 3-body | $18 \pm 10\%$ | 124 |
| $\Lambda_c^+\pi^0$ | [f] not seen | 261 |
| $\Lambda_c^+\gamma$ | not seen | 291 |

 $\Lambda_c(2625)^+$

$$I(J^P) = 0(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

$$\begin{aligned} \text{Mass } m &= 2628.1 \pm 0.6 \text{ MeV } (S = 1.5) \\ m - m_{\Lambda_c^+} &= 341.7 \pm 0.6 \text{ MeV } (S = 1.6) \\ \text{Full width } \Gamma &< 1.9 \text{ MeV, CL} = 90\% \end{aligned}$$

$\Lambda_c^+\pi\pi$ and its submode $\Sigma(2455)\pi$ are the only strong decays allowed to an excited Λ_c^+ having this mass.

| $\Lambda_c(2625)^+$ DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | p (MeV/c) |
|---|--------------------------------|------------------|-------------|
| $\Lambda_c^+\pi^+\pi^-$ | [e] $\approx 67\%$ | | 184 |
| $\Sigma_c(2455)^{++}\pi^-$ | <5 | 90% | 102 |
| $\Sigma_c(2455)^0\pi^+$ | <5 | 90% | 102 |
| $\Lambda_c^+\pi^+\pi^-$ 3-body | large | | 184 |
| $\Lambda_c^+\pi^0$ | [f] not seen | | 293 |
| $\Lambda_c^+\gamma$ | not seen | | 319 |

$\Sigma_c(2455)$

$$I(J^P) = 1(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

$\Sigma_c(2455)^{++}$ mass $m = 2454.02 \pm 0.18$ MeV

$\Sigma_c(2455)^+$ mass $m = 2452.9 \pm 0.4$ MeV

$\Sigma_c(2455)^0$ mass $m = 2453.76 \pm 0.18$ MeV

$m_{\Sigma_c^{++}} - m_{\Lambda_c^+} = 167.56 \pm 0.11$ MeV

$m_{\Sigma_c^+} - m_{\Lambda_c^+} = 166.4 \pm 0.4$ MeV

$m_{\Sigma_c^0} - m_{\Lambda_c^+} = 167.30 \pm 0.11$ MeV

$m_{\Sigma_c^{++}} - m_{\Sigma_c^0} = 0.27 \pm 0.11$ MeV ($S = 1.1$)

$m_{\Sigma_c^+} - m_{\Sigma_c^0} = -0.9 \pm 0.4$ MeV

$\Sigma_c(2455)^{++}$ full width $\Gamma = 2.23 \pm 0.30$ MeV

$\Sigma_c(2455)^+$ full width $\Gamma < 4.6$ MeV, CL = 90%

$\Sigma_c(2455)^0$ full width $\Gamma = 2.2 \pm 0.4$ MeV ($S = 1.4$)

$\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

| $\Sigma_c(2455)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|--|--------------------------------|-------------|
| $\Lambda_c^+ \pi$ | ≈ 100 % | 94 |

$\Sigma_c(2520)$

$$I(J^P) = 1(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$\Sigma_c(2520)^{++}$ mass $m = 2518.4 \pm 0.6$ MeV ($S = 1.4$)

$\Sigma_c(2520)^+$ mass $m = 2517.5 \pm 2.3$ MeV

$\Sigma_c(2520)^0$ mass $m = 2518.0 \pm 0.5$ MeV

$m_{\Sigma_c(2520)^{++}} - m_{\Lambda_c^+} = 231.9 \pm 0.6$ MeV ($S = 1.5$)

$m_{\Sigma_c(2520)^+} - m_{\Lambda_c^+} = 231.0 \pm 2.3$ MeV

$m_{\Sigma_c(2520)^0} - m_{\Lambda_c^+} = 231.6 \pm 0.5$ MeV ($S = 1.1$)

$m_{\Sigma_c(2520)^{++}} - m_{\Sigma_c(2520)^0} = 0.3 \pm 0.6$ MeV ($S = 1.2$)

$\Sigma_c(2520)^{++}$ full width $\Gamma = 14.9 \pm 1.9$ MeV

$\Sigma_c(2520)^+$ full width $\Gamma < 17$ MeV, CL = 90%

$\Sigma_c(2520)^0$ full width $\Gamma = 16.1 \pm 2.1$ MeV

$\Lambda_c^+ \pi$ is the only strong decay allowed to a Σ_c having this mass.

| $\Sigma_c(2520)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|------------------------------|--------------------------------|-------------|
| $\Lambda_c^+ \pi$ | $\approx 100\%$ | 180 |

$\Sigma_c(2800)$

$$I(J^P) = 1(?)$$

$\Sigma_c(2800)^{++}$ mass $m = 2801^{+4}_{-6}$

$\Sigma_c(2800)^+$ mass $m = 2792^{+14}_{-5}$

$\Sigma_c(2800)^0$ mass $m = 2802^{+4}_{-7}$

$$m_{\Sigma_c(2800)^{++}} - m_{\Lambda_c^+} = 514^{+4}_{-6}$$

$$m_{\Sigma_c(2800)^+} - m_{\Lambda_c^+} = 505^{+14}_{-5}$$

$$m_{\Sigma_c(2800)^0} - m_{\Lambda_c^+} = 515^{+4}_{-7}$$

$\Sigma_c(2800)^{++}$ full width $\Gamma = 75^{+22}_{-17}$

$\Sigma_c(2800)^+$ full width $\Gamma = 62^{+60}_{-40}$

$\Sigma_c(2800)^0$ full width $\Gamma = 61^{+28}_{-18}$

| $\Sigma_c(2800)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|------------------------------|--------------------------------|-------------|
| $\Lambda_c^+ \pi$ | seen | 443 |

Ξ_c^+

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2467.9 \pm 0.4$ MeV

Mean life $\tau = (442 \pm 26) \times 10^{-15}$ s (S = 1.3)

$$c\tau = 132 \mu\text{m}$$

| Ξ_c^+ DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | p (MeV/c) |
|-----------------------|--------------------------------|------------------|-------------|
|-----------------------|--------------------------------|------------------|-------------|

No absolute branching fractions have been measured.

The following are branching ratios relative to $\Xi^- \pi^+ \pi^+$.


Cabibbo-favored ($S = -2$) decays

| | | | |
|----------------------------------|-------|-------------------|---------|
| $p K_S^0 K_S^0$ | [g] | 0.087 ± 0.022 | 767 |
| $\Lambda \bar{K}^0 \pi^+$ | | — | 852 |
| $\Sigma(1385)^+ \bar{K}^0$ | [b,g] | 1.0 ± 0.5 | 746 |
| $\Lambda K^- \pi^+ \pi^+$ | [g] | 0.323 ± 0.033 | 787 |
| $\Lambda \bar{K}^*(892)^0 \pi^+$ | [b,g] | <0.2 | 90% 608 |

| | | | |
|-----------------------------|-----------------------|-----|-----|
| $\Sigma(1385)^+ K^- \pi^+$ | [b,g] <0.3 | 90% | 678 |
| $\Sigma^+ K^- \pi^+$ | [g] 0.94 ± 0.11 | | 811 |
| $\Sigma^+ \bar{K}^*(892)^0$ | [b,g] 0.81 ± 0.15 | | 658 |
| $\Sigma^0 K^- \pi^+ \pi^+$ | [g] 0.29 ± 0.16 | | 735 |
| $\Xi^0 \pi^+$ | [g] 0.55 ± 0.16 | | 877 |
| $\Xi^- \pi^+ \pi^+$ | [g] DEFINED AS 1 | | 851 |
| $\Xi(1530)^0 \pi^+$ | [b,g] <0.1 | 90% | 750 |
| $\Xi^0 \pi^+ \pi^0$ | [g] 2.34 ± 0.68 | | 856 |
| $\Xi^0 \pi^+ \pi^+ \pi^-$ | [g] 1.74 ± 0.50 | | 818 |
| $\Xi^0 e^+ \nu_e$ | [g] 2.3 ± 0.7 | | 884 |
| $\Omega^- K^+ \pi^+$ | [g] 0.07 ± 0.04 | | 399 |

Cabibbo-suppressed decays

| | | |
|--|-----------------------|-----|
| $p K^- \pi^+$ | [g] 0.21 ± 0.03 | 944 |
| $p \bar{K}^*(892)^0$ | [b,g] 0.12 ± 0.02 | 828 |
| $\Sigma^+ K^+ K^-$ | [g] 0.15 ± 0.07 | 580 |
| $\Sigma^+ \phi$ | [b,g] <0.11 | 90% |
| $\Xi(1690)^0 K^+, \Xi(1690)^0 \rightarrow$ | [g] <0.05 | 90% |
| $\Sigma^+ K^-$ | | 501 |



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2471.0 \pm 0.4$ MeV

$$m_{\Xi_c^0} - m_{\Xi_c^+} = 3.1 \pm 0.5$$
 MeV

$$\text{Mean life } \tau = (112^{+13}_{-10}) \times 10^{-15}$$
 s

$$c\tau = 33.6$$
 μm

Decay asymmetry parameters

$$\Xi^- \pi^+ \quad \alpha = -0.6 \pm 0.4$$

No absolute branching fractions have been measured. Several measurements of ratios of fractions may be found in the Listings that follow.

| Ξ_c^0 DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|--|--------------------------------|-------------|
| $p K^- K^- \pi^+$ | seen | 676 |
| $p K^- \bar{K}^*(892)^0$ | seen | 413 |
| $p K^- K^- \pi^+ \text{ no } \bar{K}^*(892)^0$ | seen | 676 |
| ΛK_S^0 | seen | 906 |
| $\Lambda \bar{K}^0 \pi^+ \pi^-$ | seen | 787 |

| | | |
|---------------------------------|------|-----|
| $\Lambda K^- \pi^+ \pi^+ \pi^-$ | seen | 703 |
| $\Xi^- \pi^+$ | seen | 875 |
| $\Xi^- \pi^+ \pi^+ \pi^-$ | seen | 817 |
| $\Omega^- K^+$ | seen | 523 |
| $\Xi^- e^+ \nu_e$ | seen | 882 |
| $\Xi^- \ell^+ \text{anything}$ | seen | — |

$\Xi_c'^+$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2575.7 \pm 3.1$ MeV

$$m_{\Xi_c'^+} - m_{\Xi_c^+} = 107.8 \pm 3.0 \text{ MeV}$$

The $\Xi_c'^+ - \Xi_c^+$ mass difference is too small for any strong decay to occur.

$\Xi_c'^+$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$\Xi_c^+ \gamma$

seen

106

$\Xi_c'^0$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2578.0 \pm 2.9$ MeV

$$m_{\Xi_c'^0} - m_{\Xi_c^0} = 107.0 \pm 2.9 \text{ MeV}$$

The $\Xi_c'^0 - \Xi_c^0$ mass difference is too small for any strong decay to occur.

Ξ_c^0 DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$\Xi_c^0 \gamma$

seen

105

$\Xi_c(2645)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

J^P has not been measured; $\frac{3}{2}^+$ is the quark-model prediction.

$\Xi_c(2645)^+$ mass $m = 2646.6 \pm 1.4$ MeV ($S = 1.6$)

$\Xi_c(2645)^0$ mass $m = 2646.1 \pm 1.2$ MeV

$$m_{\Xi_c(2645)^+} - m_{\Xi_c^0} = 175.6 \pm 1.4 \text{ MeV} \quad (S = 1.7)$$

$$m_{\Xi_c(2645)^0} - m_{\Xi_c^+} = 178.2 \pm 1.1 \text{ MeV}$$

$\Xi_c(2645)^+$ full width $\Gamma < 3.1$ MeV, CL = 90%
 $\Xi_c(2645)^0$ full width $\Gamma < 5.5$ MeV, CL = 90%

$\Xi_c \pi$ is the only strong decay allowed to a Ξ_c resonance having this mass.

| $\Xi_c(2645)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---------------------------|--------------------------------|-------------|
| $\Xi_c^0 \pi^+$ | seen | 102 |
| $\Xi_c^+ \pi^-$ | seen | 107 |

$\Xi_c(2790)$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

J^P has not been measured; $\frac{1}{2}^-$ is the quark-model prediction.

$\Xi_c(2790)^+$ mass = 2789.2 ± 3.2 MeV
 $\Xi_c(2790)^0$ mass = 2791.9 ± 3.3 MeV
 $m_{\Xi_c(2790)^+} - m_{\Xi_c^0} = 318.2 \pm 3.2$ MeV
 $m_{\Xi_c(2790)^0} - m_{\Xi_c^+} = 324.0 \pm 3.3$ MeV
 $\Xi_c(2790)^+$ width < 15 MeV, CL = 90%
 $\Xi_c(2790)^0$ width < 12 MeV, CL = 90%

| $\Xi_c(2790)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---------------------------|--------------------------------|-------------|
| $\Xi_c' \pi$ | seen | 159 |

$\Xi_c(2815)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

J^P has not been measured; $\frac{3}{2}^-$ is the quark-model prediction.

$\Xi_c(2815)^+$ mass $m = 2816.5 \pm 1.2$ MeV
 $\Xi_c(2815)^0$ mass $m = 2818.2 \pm 2.1$ MeV
 $m_{\Xi_c(2815)^+} - m_{\Xi_c^+} = 348.6 \pm 1.2$ MeV
 $m_{\Xi_c(2815)^0} - m_{\Xi_c^0} = 347.2 \pm 2.1$ MeV
 $\Xi_c(2815)^+$ full width $\Gamma < 3.5$ MeV, CL = 90%
 $\Xi_c(2815)^0$ full width $\Gamma < 6.5$ MeV, CL = 90%

The $\Xi_c \pi \pi$ modes are consistent with being entirely via $\Xi_c(2645) \pi$.

| $\Xi_c(2815)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---------------------------|--------------------------------|-------------|
| $\Xi_c^+ \pi^+ \pi^-$ | seen | 196 |
| $\Xi_c^0 \pi^+ \pi^-$ | seen | 191 |



$$I(J^P) = 0(\frac{1}{2}^+)$$

J^P has not been measured; $\frac{1}{2}^+$ is the quark-model prediction.

Mass $m = 2697.5 \pm 2.6$ MeV (S = 1.2)

Mean life $\tau = (69 \pm 12) \times 10^{-15}$ s

$$c\tau = 21 \text{ }\mu\text{m}$$

No absolute branching fractions have been measured.

| Ω_c^0 DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|--|--------------------------------|-------------|
| $\Sigma^+ K^- K^- \pi^+$ | seen | 691 |
| $\Xi^0 K^- \pi^+$ | seen | 903 |
| $\Xi^- K^- \pi^+ \pi^+$ | seen | 832 |
| $\Omega^- e^+ \nu_e$ | seen | 830 |
| $\Omega^- \pi^+$ | seen | 822 |
| $\Omega^- \pi^+ \pi^0$ | seen | 798 |
| $\Omega^- \pi^- \pi^+ \pi^+$ | seen | 754 |

NOTES

- [a] See the note on “ Λ_c^+ Branching Fractions” in the Λ_c^+ Particle Listings.
- [b] This branching fraction includes all the decay modes of the final-state resonance.
- [c] An ℓ indicates an e or a μ mode, not a sum over these modes.
- [d] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [e] Assuming isospin conservation, so that the other third is $\Lambda_c^+ \pi^0 \pi^0$.
- [f] A test that the isospin is indeed 0, so that the particle is indeed a Λ_c^+ .
- [g] No absolute branching fractions have been measured. The value here is the branching *ratio* relative to $\Xi^- \pi^+ \pi^+$.