

CHARMED MESONS ($C = \pm 1$)

$D^+ = c\bar{d}$, $D^0 = c\bar{u}$, $\bar{D}^0 = \bar{c}u$, $D^- = \bar{c}d$, similarly for D^* 's

D^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

Mass $m = 1869.3 \pm 0.4$ MeV ($S = 1.1$)

Mean life $\tau = (1040 \pm 7) \times 10^{-15}$ s

$$c\tau = 311.8 \mu\text{m}$$

c-quark decays

$$\Gamma(c \rightarrow \ell^+ \text{anything})/\Gamma(c \rightarrow \text{anything}) = 0.096 \pm 0.004 \text{ [a]}$$

$$\Gamma(c \rightarrow D^*(2010)^+ \text{anything})/\Gamma(c \rightarrow \text{anything}) = 0.255 \pm 0.017$$

CP-violation decay-rate asymmetries

$$A_{CP}(K_S^0 \pi^\pm) = -0.016 \pm 0.017$$

$$A_{CP}(K_S^0 K^\pm) = 0.07 \pm 0.06$$

$$A_{CP}(K^+ K^- \pi^\pm) = 0.007 \pm 0.008$$

$$A_{CP}(K^\pm K^{*0}) = 0.005 \pm 0.017$$

$$A_{CP}(\phi \pi^\pm) = -0.001 \pm 0.015$$

$$A_{CP}(\pi^+ \pi^- \pi^\pm) = -0.02 \pm 0.04$$

$$A_{CP}(K_S^0 K^\pm \pi^+ \pi^-) = -0.04 \pm 0.07$$

T-violation decay-rate asymmetry

$$A_T(K_S^0 K^\pm \pi^+ \pi^-) = 0.02 \pm 0.07$$

$D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$ form factors

$$r_v = 1.62 \pm 0.08 \quad (S = 1.5)$$

$$r_2 = 0.83 \pm 0.05$$

$$r_3 = 0.0 \pm 0.4$$

$$\Gamma_L/\Gamma_T = 1.13 \pm 0.08$$

$$\Gamma_+/\Gamma_- = 0.22 \pm 0.06 \quad (S = 1.6)$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as \bar{K}^0 modes. Nearly always

it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$.

D⁺ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Inclusive modes			
e^+ anything	(17.2 \pm 1.9) %		—
K^- anything	(27.5 \pm 2.4) %		—
\bar{K}^0 anything + K^0 anything	(61 \pm 8) %		—
K^+ anything	(5.5 \pm 1.6) %		—
$\bar{K}^*(892)^0$ anything	(23 \pm 5) %		—
$K^*(892)^0$ anything	< 6.6 %	CL=90%	—
η anything	[b] < 13 %	CL=90%	—
ϕ anything	< 1.8 %	CL=90%	—
ϕe^+ anything	< 1.6 %	CL=90%	—
Leptonic and semileptonic modes			
$e^+ \nu_e$	< 2.4 $\times 10^{-5}$	CL=90%	935
$\mu^+ \nu_\mu$	(4.4 \pm 0.7) $\times 10^{-4}$		932
$\bar{K}^0 e^+ \nu_e$	(8.6 \pm 0.5) %		868
$\bar{K}^0 \mu^+ \nu_\mu$	(9.5 \pm 0.8) %		865
$K^- \pi^+ e^+ \nu_e$	(4.5 \pm 1.0) %	S=1.1	863
$\bar{K}^*(892)^0 e^+ \nu_e$, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.74 \pm 0.21) %		722
$K^- \pi^+ e^+ \nu_e$ nonresonant	< 7 $\times 10^{-3}$	CL=90%	863
$K^- \pi^+ \mu^+ \nu_\mu$	(4.0 \pm 0.5) %		851
$\bar{K}^*(892)^0 \mu^+ \nu_\mu$, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(3.7 \pm 0.3) %		717
$K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	(2.1 \pm 0.6) $\times 10^{-3}$		851
$(\bar{K}^*(892)\pi)^0 e^+ \nu_e$	< 1.2 %	CL=90%	712
$(\bar{K}\pi\pi)^0 e^+ \nu_e$ non- $\bar{K}^*(892)$	< 9 $\times 10^{-3}$	CL=90%	846
$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 1.7 $\times 10^{-3}$	CL=90%	825
$\pi^0 e^+ \nu_e$	(4.4 \pm 0.7) $\times 10^{-3}$		930

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\bar{K}^*(892)^0 e^+ \nu_e$	(5.61 \pm 0.31) %	S=1.1	722
$\bar{K}^*(892)^0 \mu^+ \nu_\mu$	(5.5 \pm 0.5) %	S=1.1	717
$\bar{K}_1(1270)^0 \mu^+ \nu_\mu$	< 4 %	CL=95%	493
$\bar{K}_0^*(1430)^0 \mu^+ \nu_\mu$	< 2.5 $\times 10^{-4}$		388
$\bar{K}_2^*(1430)^0 \mu^+ \nu_\mu$	< 1.1 %	CL=95%	380
$\bar{K}^*(1680)^0 \mu^+ \nu_\mu$	< 1.6 $\times 10^{-3}$		105

$\rho^0 e^+ \nu_e$	(2.2 \pm 0.4) $\times 10^{-3}$	774
$\rho^0 \mu^+ \nu_\mu$	(3.4 \pm 0.8) $\times 10^{-3}$	770
$\omega e^+ \nu_e$	(1.6 $^{+0.7}_{-0.6}$) $\times 10^{-3}$	771
$\phi e^+ \nu_e$	< 2.09 %	CL=90% 657
$\phi \mu^+ \nu_\mu$	< 3.72 %	CL=90% 651
$\eta \ell^+ \nu_\ell$	< 7 $\times 10^{-3}$	CL=90% 854
$\eta'(958) \mu^+ \nu_\mu$	< 1.1 %	CL=90% 684

Hadronic modes with a \bar{K} or $\bar{K} K \bar{K}$

$K_S^0 \pi^+$	(1.47 \pm 0.06) %	S=1.1	862
$K^- \pi^+ \pi^+$	[c] (9.51 \pm 0.34) %	S=1.1	845
$\bar{K}^*(892)^0 \pi^+$, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	[d] (1.33 \pm 0.11) %		714
$\bar{K}_0^*(1430)^0 \pi^+$, $\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$	[d] (2.41 \pm 0.24) %		382
$\bar{K}^*(1680)^0 \pi^+$, $\bar{K}^*(1680)^0 \rightarrow K^- \pi^+$	[d] (4.0 \pm 0.8) $\times 10^{-3}$		58
$K^- \pi^+ \pi^+$ nonresonant	[d] (9.0 \pm 0.7) %		845
$K_S^0 \pi^+ \pi^0$	[c] (7.0 \pm 0.5) %	S=1.2	845
$K_S^0 \rho^+$	(4.8 \pm 1.1) %		677
$\bar{K}^*(892)^0 \pi^+$, $\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$	(1.3 \pm 0.6) %		714
$K_S^0 \pi^+ \pi^0$ nonresonant	(9 \pm 7) $\times 10^{-3}$		845
$K^- \pi^+ \pi^+ \pi^0$	[c] (6.00 \pm 0.28) %	S=1.1	816
$\bar{K}^*(892)^0 \rho^+$ total, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(1.3 \pm 0.8) %		422
$\bar{K}_1(1400)^0 \pi^+$, $\bar{K}_1(1400)^0 \rightarrow K^- \pi^+ \pi^0$	(1.8 \pm 0.7) %		390
$K^- \rho^+ \pi^+$ total	(2.6 \pm 1.6) %		613
$K^- \rho^+ \pi^+$ 3-body	(9 \pm 6) $\times 10^{-3}$		613
$\bar{K}^*(892)^0 \pi^+ \pi^0$ total, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(4.2 \pm 0.6) %		690
$\bar{K}^*(892)^0 \pi^+ \pi^0$ 3-body, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$	(2.7 \pm 0.8) %		690
$K^*(892)^- \pi^+ \pi^+$ 3-body, $K^*(892)^- \rightarrow K^- \pi^0$	(6 \pm 3) $\times 10^{-3}$		688
$K^- \pi^+ \pi^+ \pi^0$ nonresonant	[e] (1.0 \pm 0.7) %		816
$K_S^0 \pi^+ \pi^+ \pi^-$	[c] (3.11 \pm 0.21) %	S=1.1	814
$K_S^0 a_1(1260)^+$, $a_1(1260)^+ \rightarrow \pi^+ \pi^+ \pi^-$	(1.8 \pm 0.3) %		328
$\bar{K}_1(1400)^0 \pi^+$, $\bar{K}_1(1400)^0 \rightarrow K_S^0 \pi^+ \pi^-$	(1.8 \pm 0.7) %		390

$K^*(892)^-\pi^+\pi^+$ 3-body,	(1.3 ± 0.6) %		688
$K^*(892)^-\rightarrow K_S^0\pi^-$			
$K_S^0\rho^0\pi^+$ total	(1.86 ± 0.34) %	CL=90%	610
$K_S^0\rho^0\pi^+$ 3-body	(2.2 ± 2.2) × 10 ⁻³		610
$K_S^0\pi^+\pi^+\pi^-$ nonresonant	(3.7 ± 1.9) × 10 ⁻³		814
$K^-3\pi^+\pi^-$	[c] (5.8 ± 0.6) × 10 ⁻³	S=1.1	772
$\bar{K}^*(892)^0\pi^+\pi^+\pi^-$,	(1.2 ± 0.4) × 10 ⁻³		645
$\bar{K}^*(892)^0\rightarrow K^-\pi^+$			
$\bar{K}^*(892)^0\rho^0\pi^+$,	(2.3 ± 0.4) × 10 ⁻³		239
$\bar{K}^*(892)^0\rightarrow K^-\pi^+$			
$K^-\rho^0\pi^+\pi^+$	(1.75 ± 0.29) × 10 ⁻³		524
$K^-3\pi^+\pi^-$ nonresonant	(4.1 ± 3.0) × 10 ⁻⁴		772
$K^+2K_S^0$	(4.7 ± 2.1) × 10 ⁻³		545
$K^+K^-K_S^0\pi^+$	(2.4 ± 0.6) × 10 ⁻⁴		435

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$K_S^0 a_1(1260)^+$	(3.6 ± 0.6) %		328
$K_S^0 a_2(1320)^+$	< 1.5 × 10 ⁻³	CL=90%	199
$\bar{K}^*(892)^0\rho^+$ total	[e] (1.8 ± 1.4) %		422
$\bar{K}^*(892)^0\rho^+$ S-wave	[e] (1.4 ± 1.5) %		422
$\bar{K}^*(892)^0\rho^+$ P-wave	< 1 × 10 ⁻³	CL=90%	422
$\bar{K}^*(892)^0\rho^+$ D-wave	(8 ± 7) × 10 ⁻³		422
$\bar{K}^*(892)^0\rho^+$ D-wave longitudinal	< 7 × 10 ⁻³	CL=90%	422
$\bar{K}_1(1270)^0\pi^+$	< 7 × 10 ⁻³	CL=90%	487
$\bar{K}_1(1400)^0\pi^+$	(4.3 ± 1.5) %	S=1.2	390
$\bar{K}^*(892)^0\pi^+\pi^0$ total	(5.8 ± 2.9) %		690
$\bar{K}^*(892)^0\pi^+\pi^0$ 3-body	[e] (3.6 ± 2.1) %		690
$K^*(892)^-\pi^+\pi^+$ total	—		688
$K^*(892)^-\pi^+\pi^+$ 3-body	(1.8 ± 1.1) %	S=1.2	688
$\bar{K}^*(892)^0a_1(1260)^+$	(9.4 ± 1.9) × 10 ⁻³		†

Pionic modes

$\pi^+\pi^0$	(1.28 ± 0.09) × 10 ⁻³	925
$\pi^+\pi^+\pi^-$	(3.31 ± 0.21) × 10 ⁻³	908
$\rho^0\pi^+$	(1.07 ± 0.11) × 10 ⁻³	766
$\pi^+(\pi^+\pi^-)_{S\text{-wave}}$	(1.86 ± 0.18) × 10 ⁻³	908
$\sigma\pi^+$, $\sigma \rightarrow \pi^+\pi^-$	(1.53 ± 0.32) × 10 ⁻³	—
$f_0(980)\pi^+$,	(2.1 ± 0.5) × 10 ⁻⁴	669
$f_0(980) \rightarrow \pi^+\pi^-$		
$f_0(1370)\pi^+$,	(8 ± 6) × 10 ⁻⁵	—
$f_0(1370) \rightarrow \pi^+\pi^-$		

$f_2(1270)\pi^+$,	$(4.8 \pm 1.3) \times 10^{-4}$	485
$f_2(1270) \rightarrow \pi^+ \pi^-$		
$\pi^+ 2\pi^0$	$(4.8 \pm 0.4) \times 10^{-3}$	910
$\pi^+ \pi^+ \pi^- \pi^0$	$(1.18 \pm 0.09) \%$	883
$\eta \pi^+, \eta \rightarrow \pi^+ \pi^- \pi^0$	$(7.9 \pm 0.7) \times 10^{-4}$	848
$\omega \pi^+, \omega \rightarrow \pi^+ \pi^- \pi^0$	$< 3 \times 10^{-4}$ CL=90%	763
$3\pi^+ 2\pi^-$	$(1.68 \pm 0.17) \times 10^{-3}$ S=1.1	845

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\eta \pi^+$	$(3.50 \pm 0.32) \times 10^{-3}$	848
$\omega \pi^+$	$< 3.4 \times 10^{-4}$ CL=90%	763
$\eta \rho^+$	$< 7 \times 10^{-3}$ CL=90%	655
$\eta'(958)\pi^+$	$(5.3 \pm 1.1) \times 10^{-3}$	680
$\eta'(958)\rho^+$	$< 6 \times 10^{-3}$ CL=90%	348

Hadronic modes with a $K\bar{K}$ pair

$K^+ K_S^0$	$(2.96 \pm 0.19) \times 10^{-3}$	792
$K^+ K^- \pi^+$	[c] $(1.00 \pm 0.04) \%$	S=1.2 744
$\phi \pi^+, \phi \rightarrow K^+ K^-$	$(3.2 \pm 0.4) \times 10^{-3}$	647
$K^+ \bar{K}^*(892)^0,$	$(3.02 \pm 0.35) \times 10^{-3}$	613
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$		
$K^+ \bar{K}_0^*(1430)^0,$	$(3.7 \pm 0.4) \times 10^{-3}$	-
$\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$		
$K_S^0 K_S^0 \pi^+$	—	741
$K^*(892)^+ K_S^0,$	$(5.3 \pm 2.3) \times 10^{-3}$	611
$K^*(892)^+ \rightarrow K_S^0 \pi^+$		
$K^+ K^- \pi^+ \pi^0$	—	682
$\phi \pi^+ \pi^0, \phi \rightarrow K^+ K^-$	$(1.1 \pm 0.5) \%$	619
$\phi \rho^+, \phi \rightarrow K^+ K^-$	$< 7 \times 10^{-3}$ CL=90%	258
$K^+ K^- \pi^+ \pi^0$ non- ϕ	$(1.5 \pm 0.7) \%$	682
$K^+ K_S^0 \pi^+ \pi^-$	$(1.75 \pm 0.21) \times 10^{-3}$	678
$K_S^0 K^- \pi^+ \pi^+$	$(2.39 \pm 0.23) \times 10^{-3}$	678
$K^*(892)^+ \bar{K}^*(892)^0,$	$(5.8 \pm 2.4) \times 10^{-3}$	280
$K^{*+} \rightarrow K_S^0 \pi^+, \bar{K}^{*0} \rightarrow K^- \pi^+$		
$K_S^0 K^- \pi^+ \pi^+ (\text{non-}K^{*+} \bar{K}^{*0})$	$< 4 \times 10^{-3}$ CL=90%	678
$K^+ K^- \pi^+ \pi^+ \pi^-$	$(2.3 \pm 1.2) \times 10^{-4}$	600

Fractions of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\phi \pi^+$	$(6.5 \pm 0.7) \times 10^{-3}$	647
$\phi \pi^+ \pi^0$	$(2.3 \pm 1.0) \%$	619

$\phi\rho^+$	< 1.5	%	CL=90%	259
$K^*(892)^+ K_S^0$	(1.6 ± 0.7)	%		611
$K^*(892)^+ \bar{K}^*(892)^0$	(2.6 ± 1.1)	%		280

Doubly Cabibbo-suppressed modes

$K^+\pi^0$	< 4.2	$\times 10^{-4}$	CL=90%	864
$K^+\pi^+\pi^-$	(6.4 ± 0.8)	$\times 10^{-4}$		845
$K^+\rho^0$	(2.5 ± 0.7)	$\times 10^{-4}$		678
$K^*(892)^0\pi^+, K^*(892)^0 \rightarrow$	(3.0 ± 0.6)	$\times 10^{-4}$		714
$K^+\pi^-$				—
$K^+f_0(980), f_0(980) \rightarrow$	(5.7 ± 3.5)	$\times 10^{-5}$		—
$\pi^+\pi^-$				—
$K_2^*(1430)^0\pi^+, K_2^*(1430)^0 \rightarrow$	(5.2 ± 3.5)	$\times 10^{-5}$		—
$K^+\pi^-$				—
$K^+K^+K^-$	(9.0 ± 2.1)	$\times 10^{-5}$		550

 $\Delta C = 1$ weak neutral current (*C1*) modes, or**Lepton Family number (*LF*) or Lepton number (*L*) violating modes**

$\pi^+e^+e^-$	<i>C1</i>	< 7.4	$\times 10^{-6}$	CL=90%	929
$\pi^+\phi, \phi \rightarrow e^+e^-$	[<i>f</i>]	(2.7 $^{+3.6}_{-1.8}$)	$\times 10^{-6}$		—
$\pi^+\mu^+\mu^-$	<i>C1</i>	< 8.8	$\times 10^{-6}$	CL=90%	917
$\rho^+\mu^+\mu^-$	<i>C1</i>	< 5.6	$\times 10^{-4}$	CL=90%	757
$K^+e^+e^-$	[<i>g</i>]	< 6.2	$\times 10^{-6}$	CL=90%	869
$K^+\mu^+\mu^-$	[<i>g</i>]	< 9.2	$\times 10^{-6}$	CL=90%	856
$\pi^+e^\pm\mu^\mp$	<i>LF</i>	[<i>h</i>] < 3.4	$\times 10^{-5}$	CL=90%	926
$K^+e^\pm\mu^\mp$	<i>LF</i>	[<i>h</i>] < 6.8	$\times 10^{-5}$	CL=90%	866
$\pi^-e^+e^+$	<i>L</i>	< 3.6	$\times 10^{-6}$	CL=90%	929
$\pi^-\mu^+\mu^+$	<i>L</i>	< 4.8	$\times 10^{-6}$	CL=90%	917
$\pi^-e^+\mu^+$	<i>L</i>	< 5.0	$\times 10^{-5}$	CL=90%	926
$\rho^-\mu^+\mu^+$	<i>L</i>	< 5.6	$\times 10^{-4}$	CL=90%	757
$K^-e^+e^+$	<i>L</i>	< 4.5	$\times 10^{-6}$	CL=90%	869
$K^-\mu^+\mu^+$	<i>L</i>	< 1.3	$\times 10^{-5}$	CL=90%	856
$K^-e^+\mu^+$	<i>L</i>	< 1.3	$\times 10^{-4}$	CL=90%	866
$K^*(892)^-\mu^+\mu^+$	<i>L</i>	< 8.5	$\times 10^{-4}$	CL=90%	703

D⁰

$I(J^P) = \frac{1}{2}(0^-)$

Mass $m = 1864.5 \pm 0.4$ MeV ($S = 1.1$) $m_{D^\pm} - m_{D^0} = 4.78 \pm 0.10$ MeV ($S = 1.1$)Mean life $\tau = (410.1 \pm 1.5) \times 10^{-15}$ s $c\tau = 122.9$ μm $|m_{D_1^0} - m_{D_2^0}| < 7 \times 10^{10} \hbar \text{ s}^{-1}$, CL = 95% [*j*] $(\Gamma_{D_1^0} - \Gamma_{D_2^0})/\Gamma = 2y = (1.4 \pm 1.0) \times 10^{-2}$

$$\begin{aligned} \Gamma(K^+ \ell^- \bar{\nu}_\ell \text{ (via } \overline{D^0})) / \Gamma(K^- \ell^+ \nu_\ell) &< 0.005, \text{ CL} = 90\% \\ \Gamma(K^+ \pi^- \text{ (via } \overline{D^0})) / \Gamma(K^- \pi^+) &< 4.0 \times 10^{-4}, \text{ CL} = 95\% \\ \Gamma(K_S^0 \pi^+ \pi^- \text{ (in } D^0 \rightarrow \overline{D^0})) / \Gamma(K_S^0 \pi^+ \pi^-) &\quad \Gamma_0 / \Gamma_0 \\ &< 0.0063, \text{ CL} = 95\% \end{aligned}$$

***CP*-violation decay-rate asymmetries**

$$\begin{aligned} A_{CP}(K^+ K^-) &= 0.014 \pm 0.010 \\ A_{CP}(K_S^0 K_S^0) &= -0.23 \pm 0.19 \\ A_{CP}(\pi^+ \pi^-) &= 0.013 \pm 0.012 \\ A_{CP}(\pi^0 \pi^0) &= 0.00 \pm 0.05 \\ A_{CP}(\pi^+ \pi^- \pi^0) &= 0.01^{+0.10}_{-0.09} \\ A_{CP}(K_S^0 \phi) &= -0.03 \pm 0.09 \\ A_{CP}(K_S^0 \pi^0) &= 0.001 \pm 0.013 \\ A_{CP}(K^\pm \pi^\mp) &= 0.05 \pm 0.04 \\ A_{CP}(K^\mp \pi^\pm \pi^0) &= -0.03 \pm 0.09 \\ A_{CP}(K^\pm \pi^\mp \pi^0) &= 0.00 \pm 0.05 \\ A_{CP}(K_S^0 \pi^+ \pi^-) &= -0.009^{+0.026}_{-0.061} \\ A_{CP}(K^\pm \pi^\mp \pi^+ \pi^-) &= -0.02 \pm 0.04 \\ A_{CP}(K^+ K^- \pi^+ \pi^-) &= -0.08 \pm 0.07 \end{aligned}$$

***T*-violation decay-rate asymmetry**

$$A_T(K^+ K^- \pi^+ \pi^-) = 0.01 \pm 0.07$$

***CPT*-violation decay-rate asymmetry**

$$A_{CPT}(K^\mp \pi^\pm) = 0.008 \pm 0.008$$

Most decay modes (other than the semileptonic modes) that involve a neutral K meson are now given as K_S^0 modes, not as $\overline{K^0}$ modes. Nearly always it is a K_S^0 that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that $2\Gamma(K_S^0) = \Gamma(\overline{K^0})$.

D^0 DECAY MODES	Fraction (Γ_i / Γ)	Scale factor/ Confidence level	p (MeV/c)
Topological modes			
0-prongs	[j] (19 ± 6) %		—
2-prongs	(67 ± 6) %		—
4-prongs	[k] (13.8 ± 0.5) %		—
6-prongs	(1.2 + 1.3 - 0.7) × 10 ⁻³		—

Inclusive modes

e^+ anything	[I]	(6.71 \pm 0.29) %	-
μ^+ anything		(6.5 \pm 0.7) %	-
K^- anything		(53 \pm 4) %	S=1.3
\bar{K}^0 anything + K^0 anything		(42 \pm 5) %	-
K^+ anything		(3.4 \pm 0.6) %	-
$\bar{K}^*(892)^0$ anything		(9 \pm 4) %	-
$K^*(892)^0$ anything		(2.8 \pm 1.3) %	-
η anything	[b]	< 13 %	CL=90%
ϕ anything		(1.7 \pm 0.8) %	-

Semileptonic modes

$K^- e^+ \nu_e$		(3.51 \pm 0.11) %	867
$K^- \mu^+ \nu_\mu$		(3.19 \pm 0.16) %	863
$K^*(892)^- e^+ \nu_e$		(2.17 \pm 0.16) %	719
$K^*(892)^- \mu^+ \nu_\mu$		(1.95 \pm 0.25) %	714
$K^- \pi^+ \pi^- \mu^+ \nu_\mu$	< 1.2	$\times 10^{-3}$ CL=90%	821
$(\bar{K}^*(892)\pi)^- \mu^+ \nu_\mu$	< 1.4	$\times 10^{-3}$ CL=90%	692
$\pi^- e^+ \nu_e$		(2.81 \pm 0.19) $\times 10^{-3}$	927
$\pi^- \mu^+ \nu_\mu$		(2.4 \pm 0.4) $\times 10^{-3}$	924
$\rho^- e^+ \nu_e$		(1.9 \pm 0.4) $\times 10^{-3}$	771

Hadronic modes with one \bar{K}

$K^- \pi^+$		(3.80 \pm 0.07) %	S=1.1	861
$K_S^0 \pi^0$		(1.14 \pm 0.12) %	860	
$K_S^0 \pi^+ \pi^-$	[c]	(2.90 \pm 0.19) %	842	
$K_S^0 \rho^0$		(7.5 \pm 0.6) $\times 10^{-3}$	674	
$K_S^0 \omega, \omega \rightarrow \pi^+ \pi^-$		(2.1 \pm 0.6) $\times 10^{-4}$	670	
$K_S^0 f_0(980),$ $f_0(980) \rightarrow \pi^+ \pi^-$		(1.36 \pm 0.30) $\times 10^{-3}$	549	
$K_S^0 f_2(1270),$ $f_2(1270) \rightarrow \pi^+ \pi^-$		(1.3 \pm 1.1) $\times 10^{-4}$	262	
$K_S^0 f_0(1370),$ $f_0(1370) \rightarrow \pi^+ \pi^-$		(2.5 \pm 0.6) $\times 10^{-3}$		†
$K^*(892)^- \pi^+,$ $K^*(892)^- \rightarrow K_S^0 \pi^-$		(1.91 \pm 0.14) %	711	
$K^*(892)^+ \pi^-,$ $K^*(892)^+ \rightarrow K_S^0 \pi^+$	[m]	(10 \pm 12) $\times 10^{-5}$	711	
$K_0^*(1430)^- \pi^+,$ $K_0^*(1430)^- \rightarrow K_S^0 \pi^-$		(2.8 \pm 0.6) $\times 10^{-3}$	378	

$K_2^*(1430)^-\pi^+$,	(3.2 \pm 2.1) $\times 10^{-4}$	367
$K_2^*(1430)^-\rightarrow K_S^0\pi^-$		
$K^*(1680)^-\pi^+$,	(6 \pm 5) $\times 10^{-4}$	46
$K^*(1680)^-\rightarrow K_S^0\pi^-$		
$K_S^0\pi^+\pi^-$ nonresonant	(2.6 \pm 5.9) $\times 10^{-4}$	842
$K^-\pi^+\pi^0$	[c] (14.1 \pm 0.5) %	S=1.2
$K^-\rho^+$	(11.0 \pm 0.7) %	675
$K^-\rho(1700)^+$,	(8.0 \pm 1.7) $\times 10^{-3}$	†
$\rho(1700)^+\rightarrow\pi^+\pi^0$		
$K^*(892)^-\pi^+$,	(2.25 \pm 0.36) %	711
$K^*(892)^-\rightarrow K^-\pi^0$		
$\bar{K}^*(892)^0\pi^0$,	(1.91 \pm 0.24) %	711
$\bar{K}^*(892)^0\rightarrow K^-\pi^+$		
$K_0^*(1430)^-\pi^+$,	(4.6 \pm 2.2) $\times 10^{-3}$	378
$K_0^*(1430)^-\rightarrow K^-\pi^0$		
$\bar{K}_0^*(1430)^0\pi^0$,	(5.8 \pm 4.6) $\times 10^{-3}$	379
$\bar{K}_0^*(1430)^0\rightarrow K^-\pi^+$		
$K^*(1680)^-\pi^+$,	(1.8 \pm 0.7) $\times 10^{-3}$	46
$K^*(1680)^-\rightarrow K^-\pi^0$		
$K^-\pi^+\pi^0$ nonresonant	(1.13 \pm 0.54) %	844
$K_S^0\pi^0\pi^0$	—	843
$\bar{K}^*(892)^0\pi^0$,	(6.3 \pm 1.8) $\times 10^{-3}$	711
$\bar{K}^*(892)^0\rightarrow K_S^0\pi^0$		
$K_S^0\pi^0\pi^0$ nonresonant	(4.2 \pm 1.1) $\times 10^{-3}$	843
$K^-\pi^+\pi^+\pi^-$	[c] (7.72 \pm 0.28) %	S=1.3
$K^-\pi^+\rho^0$ total	(6.4 \pm 0.4) %	609
$K^-\pi^+\rho^0$ 3-body	(4.9 \pm 2.2) $\times 10^{-3}$	609
$\bar{K}^*(892)^0\rho^0$,	(1.00 \pm 0.22) %	416
$\bar{K}^*(892)^0\rightarrow K^-\pi^+$		
$K^-a_1(1260)^+$,	(3.6 \pm 0.6) %	327
$a_1(1260)^+\rightarrow\pi^+\pi^+\pi^-$		
$\bar{K}^*(892)^0\pi^+\pi^-$ total,	(1.5 \pm 0.4) %	685
$\bar{K}^*(892)^0\rightarrow K^-\pi^+$		
$\bar{K}^*(892)^0\pi^+\pi^-$ 3-body,	(9.7 \pm 2.1) $\times 10^{-3}$	685
$\bar{K}^*(892)^0\rightarrow K^-\pi^+$		
$K_1(1270)^-\pi^+$,	[e] (2.9 \pm 0.3) $\times 10^{-3}$	484
$K_1(1270)^-\rightarrow K^-\pi^+\pi^-$		
$K^-\pi^+\pi^+\pi^-$ nonresonant	(1.80 \pm 0.25) %	812
$K_S^0\pi^+\pi^-\pi^0$	[c] (5.3 \pm 0.6) %	812
$K_S^0\eta, \eta\rightarrow\pi^+\pi^-\pi^0$	(8.6 \pm 1.4) $\times 10^{-4}$	772
$K_S^0\omega, \omega\rightarrow\pi^+\pi^-\pi^0$	(9.8 \pm 1.8) $\times 10^{-3}$	670

$K^*(892)^-\rho^+$,	(2.1 \pm 0.8) %	416
$K^*(892)^-\rightarrow K_S^0\pi^-$		
$K_1(1270)^-\pi^+$,	[e] (2.2 \pm 0.6) $\times 10^{-3}$	484
$K_1(1270)^-\rightarrow K_S^0\pi^-\pi^0$		
$\overline{K}^*(892)^0\pi^+\pi^-$ 3-body,	(2.4 \pm 0.5) $\times 10^{-3}$	685
$\overline{K}^*(892)^0\rightarrow K_S^0\pi^0$		
$K_S^0\pi^+\pi^-\pi^0$ nonresonant	(1.1 \pm 1.1) %	812
$K^-\pi^+\pi^+\pi^-\pi^0$	(4.1 \pm 0.4) %	771
$\overline{K}^*(892)^0\pi^+\pi^-\pi^0$,	(1.2 \pm 0.6) %	643
$\overline{K}^*(892)^0\rightarrow K^-\pi^+$		
$K^-\pi^+\omega$, $\omega \rightarrow \pi^+\pi^-\pi^0$	(2.7 \pm 0.5) %	605
$\overline{K}^*(892)^0\omega$,	(6.5 \pm 2.4) $\times 10^{-3}$	410
$\overline{K}^*(892)^0\rightarrow K^-\pi^+$,		
$K_S^0\eta\pi^0$	(5.2 \pm 1.2) $\times 10^{-3}$	721
$K_S^0a_0(980)$, $a_0(980)\rightarrow\eta\pi^0$	(6.2 \pm 2.0) $\times 10^{-3}$	—
$\overline{K}^*(892)^0\eta$, $\overline{K}^*(892)^0\rightarrow K_S^0\pi^0$	(1.5 \pm 0.5) $\times 10^{-3}$	—
$K_S^02\pi^+2\pi^-$	(2.75 \pm 0.31) $\times 10^{-3}$	768
$K_S^0\rho^0\pi^+\pi^-$, no $K^*(892)^-$	(1.1 \pm 0.7) $\times 10^{-3}$	—
$K^*(892)^-\pi^+\pi^+\pi^-$,	(5 \pm 8) $\times 10^{-4}$	642
$K^*(892)^-\rightarrow K_S^0\pi^-$,		
no ρ^0		
$K^*(892)^-\rho^0\pi^+$,	(1.7 \pm 0.7) $\times 10^{-3}$	230
$K^*(892)^-\rightarrow K_S^0\pi^-$		
$K_S^02\pi^+2\pi^-$ nonresonant	< 1.3 $\times 10^{-3}$ CL=90%	768
$K^-3\pi^+2\pi^-$	(2.1 \pm 0.5) $\times 10^{-4}$	713

Fractions of many of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. (Modes for which there are only upper limits and $\overline{K}^*(892)\rho$ submodes only appear below.)

$K_S^0\eta$	(3.8 \pm 0.6) $\times 10^{-3}$	772
$K_S^0\omega$	(1.10 \pm 0.20) %	670
$K_S^0\eta'(958)$	(9.1 \pm 1.4) $\times 10^{-3}$	565
$K^-a_1(1260)^+$	(7.5 \pm 1.1) %	327
$\overline{K}^0a_1(1260)^0$	< 1.9 %	322
$K^-a_2(1320)^+$	< 2 $\times 10^{-3}$ CL=90%	197
$\overline{K}^*(892)^0\pi^+\pi^-$ total	(2.3 \pm 0.5) %	685
$\overline{K}^*(892)^0\pi^+\pi^-$ 3-body	(1.46 \pm 0.32) %	685
$\overline{K}^*(892)^0\rho^0$	(1.50 \pm 0.33) %	417
$\overline{K}^*(892)^0\rho^0$ transverse	(1.6 \pm 0.5) %	417
$\overline{K}^*(892)^0\rho^0$ S-wave	(2.9 \pm 0.6) %	417

$\bar{K}^*(892)^0 \rho^0$	<i>S</i> -wave long.	< 3	$\times 10^{-3}$	CL=90%	417
$\bar{K}^*(892)^0 \rho^0$	<i>P</i> -wave	< 3	$\times 10^{-3}$	CL=90%	417
$\bar{K}^*(892)^0 \rho^0$	<i>D</i> -wave	(2.0 \pm 0.6) %			417
$K^*(892)^- \rho^+$		(6.4 \pm 2.5) %			417
$K^*(892)^- \rho^+$	longitudinal	(3.1 \pm 1.2) %			417
$K^*(892)^- \rho^+$	transverse	(3.4 \pm 2.0) %			417
$K^*(892)^- \rho^+$	<i>P</i> -wave	< 1.5	%	CL=90%	417
$K_1(1270)^- \pi^+$		[e] (1.12 \pm 0.31) %			484
$K_1(1400)^- \pi^+$		< 1.2	%	CL=90%	386
$\bar{K}_1(1400)^0 \pi^0$		< 3.7	%	CL=90%	387
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$		(1.8 \pm 0.9) %			643
$K^- \pi^+ \omega$		(3.0 \pm 0.6) %			605
$\bar{K}^*(892)^0 \omega$		(1.1 \pm 0.4) %			410
$K^- \pi^+ \eta'(958)$		(7.2 \pm 1.8) $\times 10^{-3}$			479
$\bar{K}^*(892)^0 \eta'(958)$		< 1.1	$\times 10^{-3}$	CL=90%	118

Hadronic modes with three K 's

$K_S^0 K^+ K^-$		(4.58 \pm 0.34) $\times 10^{-3}$		544	
$K_S^0 a_0(980)^0, a_0^0 \rightarrow K^+ K^-$		(3.0 \pm 0.4) $\times 10^{-3}$		—	
$K^- a_0(980)^+, a_0^+ \rightarrow K^+ K_S^0$		(6.1 \pm 1.8) $\times 10^{-4}$		—	
$K^+ a_0(980)^-, a_0^- \rightarrow K^- K_S^0$		< 1.1	$\times 10^{-4}$	CL=95%	—
$K_S^0 f_0(980), f_0 \rightarrow K^+ K^-$		< 1.0	$\times 10^{-4}$	CL=95%	—
$K_S^0 \phi, \phi \rightarrow K^+ K^-$		(2.10 \pm 0.16) $\times 10^{-3}$		520	
$K_S^0 f_0(1400), f_0 \rightarrow K^+ K^-$		(1.7 \pm 1.1) $\times 10^{-4}$		—	
$3K_S^0$		(9.3 \pm 1.3) $\times 10^{-4}$		538	
$K^+ K^- K^- \pi^+$		(2.11 \pm 0.31) $\times 10^{-4}$		434	
$K^+ K^- \bar{K}^*(892)^0,$		(4.2 \pm 1.7) $\times 10^{-5}$		†	
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$					
$K^- \pi^+ \phi, \phi \rightarrow K^+ K^-$		(3.8 \pm 1.6) $\times 10^{-5}$		422	
$\phi \bar{K}^*(892)^0,$		(1.01 \pm 0.20) $\times 10^{-4}$		†	
$\phi \rightarrow K^+ K^-$,					
$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$					
$K^+ K^- K^- \pi^+$ nonresonant		(3.2 \pm 1.4) $\times 10^{-5}$		434	
$K_S^0 K_S^0 K^\pm \pi^\mp$		(6.1 \pm 1.3) $\times 10^{-4}$		427	

Pionic modes

$\pi^+ \pi^-$		(1.364 \pm 0.032) $\times 10^{-3}$		922	
$\pi^0 \pi^0$		(7.9 \pm 0.8) $\times 10^{-4}$		922	
$\pi^+ \pi^- \pi^0$		(1.31 \pm 0.06) %		907	
$\rho^+ \pi^-$		(10.0 \pm 0.6) $\times 10^{-3}$		764	
$\rho^0 \pi^0$		(3.2 \pm 0.4) $\times 10^{-3}$		764	
$\rho^- \pi^+$		(4.5 \pm 0.4) $\times 10^{-3}$		764	
$f_0(980) \pi^0, f_0(980) \rightarrow$		< 3.4	$\times 10^{-6}$	CL=95%	—
$\pi^+ \pi^-$					

$f_0(600)\pi^0$, $f_0(600) \rightarrow$	< 2.7	$\times 10^{-5}$ CL=95%	-
$\pi^+\pi^-$			
$(\pi^+\pi^-)S\text{-wave}\pi^0$	< 2.5	$\times 10^{-4}$ CL=95%	907
$3\pi^0$	< 3.5	$\times 10^{-4}$ CL=90%	908
$2\pi^+2\pi^-$	(7.31 \pm 0.27) $\times 10^{-3}$		879
$\pi^+\pi^-2\pi^0$	(9.8 \pm 0.9) $\times 10^{-3}$		882
$\eta\pi^0$	[n] (5.6 \pm 1.4) $\times 10^{-4}$		846
$\omega\pi^0$	[n] < 2.6 $\times 10^{-4}$ CL=90%		761
$2\pi^+2\pi^-\pi^0$	(4.1 \pm 0.5) $\times 10^{-3}$		844
$\eta\pi^+\pi^-$	[n] < 1.9 $\times 10^{-3}$ CL=90%		827
$\omega\pi^+\pi^-$	[n] (1.6 \pm 0.5) $\times 10^{-3}$		738
$3\pi^+3\pi^-$	(4.0 \pm 1.1) $\times 10^{-4}$		795

Hadronic modes with a $K\bar{K}$ pair

K^+K^-	(3.84 \pm 0.10) $\times 10^{-3}$	791
$2K_S^0$	(3.7 \pm 0.7) $\times 10^{-4}$	788
$K_S^0K^-\pi^+$	(3.4 \pm 0.5) $\times 10^{-3}$ S=1.1	739
$\bar{K}^*(892)^0K_S^0$,	< 6 $\times 10^{-4}$ CL=90%	608
$\bar{K}^*(892)^0 \rightarrow K^-\pi^+$		
$K^*(892)^+K^-$, $K^*(892)^+ \rightarrow K_S^0\pi^+$	(1.2 \pm 0.3) $\times 10^{-3}$	610
$K_S^0K^-\pi^+$ nonresonant	(1.1 \pm 1.1) $\times 10^{-3}$	739
$K_S^0K^+\pi^-$	(2.6 \pm 0.5) $\times 10^{-3}$	739
$K^*(892)^0K_S^0$, $K^*(892)^0 \rightarrow K^+\pi^-$	< 3 $\times 10^{-4}$ CL=90%	608
$K^*(892)^-K^+$, $K^*(892)^- \rightarrow K_S^0\pi^-$	(7 \pm 4) $\times 10^{-4}$	610
$K_S^0K^+\pi^-$ nonresonant	(1.9 \pm 1.1) $\times 10^{-3}$	739
$K^+K^-\pi^0$	(1.3 \pm 0.4) $\times 10^{-3}$	743
$K_S^0K_S^0\pi^0$	< 5.9 $\times 10^{-4}$	740
$K^+K^-\pi^+\pi^-$	[o] (2.32 \pm 0.13) $\times 10^{-3}$	676
$\phi\pi^+\pi^-$ 3-body, $\phi \rightarrow K^+K^-$	(2.3 \pm 2.3) $\times 10^{-5}$	614
$\phi\rho^0$, $\phi \rightarrow K^+\rho^-$	(6.7 \pm 0.6) $\times 10^{-4}$	250
$K^+K^-\rho^0$ 3-body	(5 \pm 7) $\times 10^{-5}$	302
$f_0(980)\pi^+\pi^-$, $f_0 \rightarrow K^+K^-$	(3.5 \pm 0.9) $\times 10^{-4}$	-
$K^*(892)^0K^\mp\pi^\pm$ 3-body,	[p] (2.5 \pm 0.5) $\times 10^{-4}$	531
$K^{*0} \rightarrow K^\pm\pi^\mp$		
$K^*(892)^0\bar{K}^*(892)^0$, $K^{*0} \rightarrow K^\pm\pi^\mp$	(7 \pm 5) $\times 10^{-5}$	272
$K_1(1270)^\pm K^\mp$,	(7.6 \pm 1.7) $\times 10^{-4}$	-
$K_1(1270)^\pm \rightarrow K^\pm\pi^+\pi^-$		
$K_1(1400)^\pm K^\mp$,	(5.1 \pm 1.2) $\times 10^{-4}$	-
$K_1(1400)^\pm \rightarrow K^\pm\pi^+\pi^-$		

$K_S^0 K_S^0 \pi^+ \pi^-$	$(1.26 \pm 0.24) \times 10^{-3}$	673
$K_S^0 K^- \pi^+ \pi^+ \pi^-$	$< 1.5 \times 10^{-4} \text{CL}=90\%$	595
$K^+ K^- \pi^+ \pi^- \pi^0$	$(3.1 \pm 2.0) \times 10^{-3}$	600

Fractions of most of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\bar{K}^*(892)^0 K_S^0$	$< 8 \times 10^{-4} \text{CL}=90\%$	608
$K^*(892)^+ K^-$	$(3.7 \pm 0.8) \times 10^{-3}$	610
$K^*(892)^0 K_S^0$	$< 4 \times 10^{-4} \text{CL}=90\%$	608
$K^*(892)^- K^+$	$(2.0 \pm 1.1) \times 10^{-3}$	610
$\phi \pi^0$	$(7.4 \pm 0.5) \times 10^{-4}$	644
$\phi \eta$	$(1.4 \pm 0.4) \times 10^{-4}$	489
$\phi \omega$	$< 2.1 \times 10^{-3} \text{CL}=90\%$	237

Radiative modes

$\rho^0 \gamma$	$< 2.4 \times 10^{-4} \text{CL}=90\%$	771
$\omega \gamma$	$< 2.4 \times 10^{-4} \text{CL}=90\%$	768
$\phi \gamma$	$(2.4 \pm 0.7) \times 10^{-5}$	654
$\bar{K}^*(892)^0 \gamma$	$< 7.6 \times 10^{-4} \text{CL}=90\%$	719

Doubly Cabibbo suppressed (DC) modes or $\Delta C = 2$ forbidden via mixing (C2M) modes

$K^+ \ell^- \bar{\nu}_\ell$ (via \bar{D}^0)	C2M	$< 1.8 \times 10^{-4} \text{CL}=90\%$	—
$K^+ \text{ or } K^*(892)^+ e^- \bar{\nu}_e$ (via \bar{D}^0)	C2M	$< 6 \times 10^{-5} \text{CL}=90\%$	—
$K^+ \pi^-$	DC	$(1.43 \pm 0.04) \times 10^{-4}$	861
$K^+ \pi^-$ (via \bar{D}^0)	C2M	$< 1.5 \times 10^{-5} \text{CL}=95\%$	861
$K_S^0 \pi^+ \pi^-$ (in $D^0 \rightarrow \bar{D}^0$)	C2M	$< 1.8 \times 10^{-4} \text{CL}=95\%$	—
$K^*(892)^+ \pi^-$, $K^*(892)^+ \rightarrow K_S^0 \pi^+$	DC	$(10 \pm 12) \times 10^{-5}$	711
$K^+ \pi^- \pi^0$	DC	$(3.29 \pm 0.30) \times 10^{-4}$	844
$K^+ \pi^- \pi^+ \pi^-$	DC	$(2.49 \pm 0.21) \times 10^{-4}$	812
$K^+ \pi^- \pi^+ \pi^-$ (via \bar{D}^0)	C2M	$< 4 \times 10^{-4} \text{CL}=90\%$	812
$\mu^- \text{ anything}$ (via \bar{D}^0)	C2M	$< 4 \times 10^{-4} \text{CL}=90\%$	—

$\Delta C = 1$ weak neutral current (C1) modes, Lepton Family number (LF) violating modes, or Lepton number (L) violating modes

$\gamma \gamma$	C1	$< 2.6 \times 10^{-5} \text{CL}=90\%$	932
$e^+ e^-$	C1	$< 1.2 \times 10^{-6} \text{CL}=90\%$	932
$\mu^+ \mu^-$	C1	$< 1.3 \times 10^{-6} \text{CL}=90\%$	926
$\pi^0 e^+ e^-$	C1	$< 4.5 \times 10^{-5} \text{CL}=90\%$	927
$\pi^0 \mu^+ \mu^-$	C1	$< 1.8 \times 10^{-4} \text{CL}=90\%$	915

$\eta e^+ e^-$	<i>C1</i>	< 1.1	$\times 10^{-4} \text{CL}=90\%$	852
$\eta \mu^+ \mu^-$	<i>C1</i>	< 5.3	$\times 10^{-4} \text{CL}=90\%$	838
$\pi^+ \pi^- e^+ e^-$	<i>C1</i>	< 3.73	$\times 10^{-4} \text{CL}=90\%$	922
$\rho^0 e^+ e^-$	<i>C1</i>	< 1.0	$\times 10^{-4} \text{CL}=90\%$	771
$\pi^+ \pi^- \mu^+ \mu^-$	<i>C1</i>	< 3.0	$\times 10^{-5} \text{CL}=90\%$	894
$\rho^0 \mu^+ \mu^-$	<i>C1</i>	< 2.2	$\times 10^{-5} \text{CL}=90\%$	754
$\omega e^+ e^-$	<i>C1</i>	< 1.8	$\times 10^{-4} \text{CL}=90\%$	768
$\omega \mu^+ \mu^-$	<i>C1</i>	< 8.3	$\times 10^{-4} \text{CL}=90\%$	751
$K^- K^+ e^+ e^-$	<i>C1</i>	< 3.15	$\times 10^{-4} \text{CL}=90\%$	791
$\phi e^+ e^-$	<i>C1</i>	< 5.2	$\times 10^{-5} \text{CL}=90\%$	654
$K^- K^+ \mu^+ \mu^-$	<i>C1</i>	< 3.3	$\times 10^{-5} \text{CL}=90\%$	709
$\phi \mu^+ \mu^-$	<i>C1</i>	< 3.1	$\times 10^{-5} \text{CL}=90\%$	631
$\bar{K}^0 e^+ e^-$	[<i>g</i>]	< 1.1	$\times 10^{-4} \text{CL}=90\%$	866
$\bar{K}^0 \mu^+ \mu^-$	[<i>g</i>]	< 2.6	$\times 10^{-4} \text{CL}=90\%$	852
$K^- \pi^+ e^+ e^-$	<i>C1</i>	< 3.85	$\times 10^{-4} \text{CL}=90\%$	861
$\bar{K}^*(892)^0 e^+ e^-$	[<i>g</i>]	< 4.7	$\times 10^{-5} \text{CL}=90\%$	719
$K^- \pi^+ \mu^+ \mu^-$	<i>C1</i>	< 3.59	$\times 10^{-4} \text{CL}=90\%$	829
$\bar{K}^*(892)^0 \mu^+ \mu^-$	[<i>g</i>]	< 2.4	$\times 10^{-5} \text{CL}=90\%$	700
$\pi^+ \pi^- \pi^0 \mu^+ \mu^-$	<i>C1</i>	< 8.1	$\times 10^{-4} \text{CL}=90\%$	863
$\mu^\pm e^\mp$	<i>LF</i>	[<i>h</i>] < 8.1	$\times 10^{-7} \text{CL}=90\%$	929
$\pi^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>h</i>] < 8.6	$\times 10^{-5} \text{CL}=90\%$	924
$\eta e^\pm \mu^\mp$	<i>LF</i>	[<i>h</i>] < 1.0	$\times 10^{-4} \text{CL}=90\%$	848
$\pi^+ \pi^- e^\pm \mu^\mp$	<i>LF</i>	[<i>h</i>] < 1.5	$\times 10^{-5} \text{CL}=90\%$	911
$\rho^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>h</i>] < 4.9	$\times 10^{-5} \text{CL}=90\%$	767
$\omega e^\pm \mu^\mp$	<i>LF</i>	[<i>h</i>] < 1.2	$\times 10^{-4} \text{CL}=90\%$	764
$K^- K^+ e^\pm \mu^\mp$	<i>LF</i>	[<i>h</i>] < 1.8	$\times 10^{-4} \text{CL}=90\%$	754
$\phi e^\pm \mu^\mp$	<i>LF</i>	[<i>h</i>] < 3.4	$\times 10^{-5} \text{CL}=90\%$	648
$\bar{K}^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>h</i>] < 1.0	$\times 10^{-4} \text{CL}=90\%$	862
$K^- \pi^+ e^\pm \mu^\mp$	<i>LF</i>	[<i>h</i>] < 5.53	$\times 10^{-4} \text{CL}=90\%$	848
$\bar{K}^*(892)^0 e^\pm \mu^\mp$	<i>LF</i>	[<i>h</i>] < 8.3	$\times 10^{-5} \text{CL}=90\%$	714
$\pi^- \pi^- e^+ e^+ + \text{c.c.}$	<i>L</i>	< 1.12	$\times 10^{-4} \text{CL}=90\%$	922
$\pi^- \pi^- \mu^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 2.9	$\times 10^{-5} \text{CL}=90\%$	894
$K^- \pi^- e^+ e^+ + \text{c.c.}$	<i>L</i>	< 2.06	$\times 10^{-4} \text{CL}=90\%$	861
$K^- \pi^- \mu^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 3.9	$\times 10^{-4} \text{CL}=90\%$	829
$K^- K^- e^+ e^+ + \text{c.c.}$	<i>L</i>	< 1.52	$\times 10^{-4} \text{CL}=90\%$	791
$K^- K^- \mu^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 9.4	$\times 10^{-5} \text{CL}=90\%$	709
$\pi^- \pi^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 7.9	$\times 10^{-5} \text{CL}=90\%$	911
$K^- \pi^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 2.18	$\times 10^{-4} \text{CL}=90\%$	848
$K^- K^- e^+ \mu^+ + \text{c.c.}$	<i>L</i>	< 5.7	$\times 10^{-5} \text{CL}=90\%$	754

D*(2007)⁰

$I(J^P) = \frac{1}{2}(1^-)$
 I, J, P need confirmation.

Mass $m = 2006.7 \pm 0.4$ MeV ($S = 1.1$)

$$m_{D^{*0}} - m_{D^0} = 142.12 \pm 0.07 \text{ MeV}$$

Full width $\Gamma < 2.1$ MeV, CL = 90%

$\overline{D}^*(2007)^0$ modes are charge conjugates of modes below.

D*(2007)⁰ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$$D^0\pi^0$$

(61.9 ± 2.9) %

43

$$D^0\gamma$$

(38.1 ± 2.9) %

137

D*(2010)[±]

$I(J^P) = \frac{1}{2}(1^-)$
 I, J, P need confirmation.

Mass $m = 2010.0 \pm 0.4$ MeV ($S = 1.1$)

$$m_{D^*(2010)^+} - m_{D^+} = 140.64 \pm 0.10 \text{ MeV} \quad (S = 1.1)$$

$$m_{D^*(2010)^+} - m_{D^0} = 145.421 \pm 0.010 \text{ MeV} \quad (S = 1.1)$$

Full width $\Gamma = 96 \pm 22$ keV

$D^*(2010)^-$ modes are charge conjugates of the modes below.

D*(2010)[±] DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$$D^0\pi^+$$

(67.7 ± 0.5) %

39

$$D^+\pi^0$$

(30.7 ± 0.5) %

38

$$D^+\gamma$$

(1.6 ± 0.4) %

136

D₁(2420)⁰

$I(J^P) = \frac{1}{2}(1^+)$
 I, J, P need confirmation.

Mass $m = 2422.3 \pm 1.3$ MeV ($S = 1.2$)

$$m_{D_1^0} - m_{D^{*+}} = 411.7 \pm 0.8$$

Full width $\Gamma = 20.4 \pm 1.7$ MeV

$\overline{D}_1(2420)^0$ modes are charge conjugates of modes below.

D₁(2420)⁰ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$$D^*(2010)^+\pi^-$$

seen

355

$$D^0\pi^+\pi^-$$

seen

426

$$D^+\pi^-$$

not seen

474

$$D^{*0}\pi^+\pi^-$$

not seen

281

$D_2^*(2460)^0$

$$I(J^P) = \frac{1}{2}(2^+)$$

$J^P = 2^+$ assignment strongly favored.

Mass $m = 2461.1 \pm 1.6$ MeV ($S = 1.3$)

$$m_{D_2^{*0}} - m_{D^+} = 593.9 \pm 0.8$$

Full width $\Gamma = 43 \pm 4$ MeV ($S = 1.8$)

$\overline{D}_2^*(2460)^0$ modes are charge conjugates of modes below.

$D_2^*(2460)^0$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$D^+ \pi^-$	seen	506
$D^*(2010)^+ \pi^-$	seen	389
$D^0 \pi^+ \pi^-$	not seen	462
$D^{*0} \pi^+ \pi^-$	not seen	325

$D_2^*(2460)^{\pm}$

$$I(J^P) = \frac{1}{2}(2^+)$$

$J^P = 2^+$ assignment strongly favored.

Mass $m = 2459 \pm 4$ MeV ($S = 1.7$)

$$m_{D_2^*(2460)^{\pm}} - m_{D_2^*(2460)^0} = 2.4 \pm 1.7$$
 MeV

Full width $\Gamma = 29 \pm 5$ MeV

$D_2^*(2460)^-$ modes are charge conjugates of modes below.

$D_2^*(2460)^{\pm}$ DECAY MODES

Fraction (Γ_i/Γ)

p (MeV/c)

$D^0 \pi^+$	seen	507
$D^{*0} \pi^+$	seen	390
$D^+ \pi^+ \pi^-$	not seen	456
$D^{*+} \pi^+ \pi^-$	not seen	319

NOTES

- [a] This result applies to $Z^0 \rightarrow c\bar{c}$ decays only. Here ℓ^+ is an average (not a sum) of e^+ and μ^+ decays.
- [b] This is a weighted average of D^\pm (44%) and D^0 (56%) branching fractions. See " $D^+ \text{ and } D^0 \rightarrow (\eta \text{ anything}) / (\text{total } D^+ \text{ and } D^0)$ " under " D^+ Branching Ratios" in the Particle Listings.
- [c] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers in the Particle Listings.
- [d] These subfractions of the $K^- \pi^+ \pi^+$ mode are uncertain: see the Particle Listings.
- [e] The two experiments measuring this fraction are in serious disagreement. See the Particle Listings.
- [f] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+ e^+ e^-$ final state.
- [g] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [h] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [i] This $D_1^0 - D_2^0$ limit is inferred from the $D^0 - \overline{D}^0$ mixing ratio $\Gamma(K^+ \pi^- \text{ (via } \overline{D}^0)) / \Gamma(K^- \pi^+)$ near the end of the D^0 Listings.
- [j] This value is obtained by subtracting the branching fractions for 2-, 4- and 6-prongs from unity.
- [k] This is the sum of our $K^- \pi^+ \pi^+ \pi^-$, $K^- \pi^+ \pi^+ \pi^- \pi^0$, $\overline{K}^0 2\pi^+ 2\pi^-$, $2\pi^+ 2\pi^-$, $2\pi^+ 2\pi^- \pi^0$, $K^+ K^- \pi^+ \pi^-$, and $K^+ K^- \pi^+ \pi^- \pi^0$, branching fractions.
- [l] The branching fractions for the $K^- e^+ \nu_e$, $K^*(892)^- e^+ \nu_e$, $\pi^- e^+ \nu_e$, and $\rho^- e^+ \nu_e$ modes add up to 6.14 ± 0.20 %.
- [m] This is a doubly Cabibbo-suppressed mode.
- [n] This branching fraction includes all the decay modes of the resonance in the final state.
- [o] The experiments on the division of this charge mode amongst its submodes disagree, and the submode branching fractions here add up to considerably more than the charged-mode fraction.
- [p] However, these upper limits are in serious disagreement with values obtained in another experiment.