

## $K_0^*(1950)$

$I(J^P) = \frac{1}{2}(0^+)$

### OMMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the  $K^- \pi^+$  system. Needs confirmation.

### $K_0^*(1950)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>1945 \pm 10 \pm 20</math></b>	<sup>1</sup> ASTON 88 LASS	0		$11 K^- p \rightarrow K^- \pi^+ n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1917 $\pm$ 12	<sup>2</sup> ZHOU 06 RVUE			$K^- p \rightarrow K^- \pi^+ n$
1820 $\pm$ 40	<sup>3</sup> ANISOVICH 97C RVUE			$11 K^- p \rightarrow K^- \pi^+ n$

<sup>1</sup> We take the central value of the two solutions and the larger error given.

<sup>2</sup> S-matrix pole. Using ASTON 88 and assuming  $K_0^*(800)$ ,  $K_0^*(1430)$ .

<sup>3</sup> T-matrix pole. Reanalysis of ASTON 88 data.

### $K_0^*(1950)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>201 \pm 34 \pm 79</math></b>	<sup>4</sup> ASTON 88 LASS	0		$11 K^- p \rightarrow K^- \pi^+ n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

145 $\pm$ 38	<sup>5</sup> ZHOU 06 RVUE			$K^- p \rightarrow K^- \pi^+ n$
250 $\pm$ 100	<sup>6</sup> ANISOVICH 97C RVUE			$11 K^- p \rightarrow K^- \pi^+ n$

<sup>4</sup> We take the central value of the two solutions and the larger error given.

<sup>5</sup> S-matrix pole. Using ASTON 88 and assuming  $K_0^*(800)$ ,  $K_0^*(1430)$ .

<sup>6</sup> T-matrix pole. Reanalysis of ASTON 88 data.

### $K_0^*(1950)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 K\pi$	(52 $\pm$ 14) %

### $K_0^*(1950)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$		$\Gamma_1/\Gamma$
<b><math>0.52 \pm 0.08 \pm 0.12</math></b>	<sup>7</sup> ASTON 88 LASS	0

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\sim 0.60$	<sup>8</sup> ZHOU 06 RVUE			$K^- p \rightarrow K^- \pi^+ n$
-------------	---------------------------	--	--	---------------------------------

<sup>7</sup> We take the central value of the two solutions and the larger error given.

<sup>8</sup> S-matrix pole. Using ASTON 88 and assuming  $K_0^*(800)$ ,  $K_0^*(1430)$ .

## **$K_0^*(1950)$ REFERENCES**

ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng
ANISOVICH	97C	PL B413 137	A.V. Anisovich, A.V. Sarantsev
ASTON	88	NP B296 493	D. Aston <i>et al.</i> (SLAC, NAGO, CINC, INUS)

---

## **OTHER RELATED PAPERS**

---

ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
KATAEV	05	PAN 68 567 Translated from YAF 68 597.	A.L. Kataev	
JAMIN	00	NP B587 331	M. Jamin <i>et al.</i>	
SHAKIN	00	PR D62 114014	C.M. Shakin, H. Wang	