

**$a_2(1700)$**

$I^G(J^{PC}) = 1^-(2^{++})$

OMMITTED FROM SUMMARY TABLE

### **$a_2(1700)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>1732 \pm 16</math> OUR AVERAGE</b>		Error includes scale factor of 1.9. See the ideogram below.			
$1737 \pm 5 \pm 7$		ABE	04	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
$1698 \pm 44$		<sup>1</sup> AMSLER	02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta\eta$
$1660 \pm 40$		ABELE	99B	CBAR	$1.94 \bar{p}p \rightarrow \pi^0 \eta\eta$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
$1722 \pm 9 \pm 15$	18k	<sup>2</sup> SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
$1702 \pm 7$	80k	<sup>3</sup> UMAN	06	E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
$1721 \pm 13 \pm 44$	145k	LU	05	B852	$18 \pi^- p \rightarrow \omega\pi^- \pi^0 p$
$\sim 1775$		<sup>4</sup> GRYGOREV	99	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$1752 \pm 21 \pm 4$		ACCIARRI	97T	L3	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$


<sup>1</sup> T-matrix pole.

<sup>2</sup> From analysis of L3 data at 183–209 GeV.

<sup>3</sup> Statistical error only.

<sup>4</sup> Possibly two  $J^P = 2^+$  resonances with isospins 0 and 1.


WEIGHTED AVERAGE  
 $1732 \pm 16$  (Error scaled by 1.9)



## $a_2(1700)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>194± 40 OUR AVERAGE</b>	Error includes scale factor of 1.6. See the ideogram below.				
151± 22±24	ABE	04	BELL	10.6	$e^+e^- \rightarrow e^+e^- K^+K^-$
265± 55	5 AMSLER	02	CBAR	0.9	$\bar{p}p \rightarrow \pi^0\eta\eta$
280± 70	ABELE	99B	CBAR	1.94	$\bar{p}p \rightarrow \pi^0\eta\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
336± 20±20	18k	6 SCHEGELSKY	06	RVUE 0	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
417± 19	80k	7 UMAN	06	E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
279± 49±66	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega\pi^-\pi^0 p$
150±110±34		ACCIARRI	97T	L3	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

<sup>5</sup> T-matrix pole.  
<sup>6</sup> From analysis of L3 data at 183–209 GeV.  
<sup>7</sup> Statistical error only.



## $a_2(1700)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \eta\pi$	seen
$\Gamma_2 \quad \gamma\gamma$	
$\Gamma_3 \quad \rho\pi$	

$\Gamma_4$	$f_2(1270)\pi$			
$\Gamma_5$	$K\bar{K}$		seen	
$\Gamma_6$	$\omega\pi^-\pi^0$		seen	
$\Gamma_7$	$\omega\rho$		seen	

### $a_2(1700)$ PARTIAL WIDTHS

#### $\Gamma(\eta\pi)$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_1$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$9.5 \pm 2.0$	870	<sup>8</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

#### $\Gamma(\gamma\gamma)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$0.30 \pm 0.05$	870	<sup>8</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

#### $\Gamma(K\bar{K})$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_5$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$5.0 \pm 3.0$	870	<sup>8</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

<sup>8</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $a_2(1700)$  mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

### $a_2(1700) \Gamma(i) \Gamma(\gamma\gamma)/\Gamma(\text{total})$

$[\Gamma(\rho\pi) + \Gamma(f_2(1270)\pi)] \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$(\Gamma_3 + \Gamma_4)\Gamma_2/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.29 \pm 0.04 \pm 0.02$		ACCIARRI 97T	L3	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$0.37^{+0.12}_{-0.08} \pm 0.10$	18k	<sup>9</sup> SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

<sup>9</sup> From analysis of L3 data at 183–209 GeV.

#### $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

#### $\Gamma_5\Gamma_2/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$20.6 \pm 4.2 \pm 4.6$	<sup>10</sup> ABE 04	BELL	$10.6 e^+e^- \rightarrow e^+e^-K^+K^-$	
10 Assuming spin 2.				

### $a_2(1700)$ BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$				$\Gamma_3/\Gamma_4$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$3.4 \pm 0.4 \pm 0.1$	18k	<sup>11</sup> SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

<sup>11</sup> From analysis of L3 data at 183–209 GeV.

## **$a_2(1700)$ REFERENCES**

SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>
LU	05	PRL 94 032002	M. Lu <i>et al.</i>
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>
ABELE	99B	EPJ C8 67	A. Abele <i>et al.</i>
GRYGOREV	99	PAN 62 470 Translated from YAF 62 513.	V.K. Grygorev <i>et al.</i>
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>

(FNAL E835)  
(BNL E852 Collab.)  
(BELLE Collab.)

(Crystal Barrel Collab.)

(L3 Collab.)

---

## **OTHER RELATED PAPERS**

---

BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>
BARBERIS	00H	PL B488 225	D. Barberis <i>et al.</i>

(WA 102 Collab.)

---