

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ***$$

According to the quark model, the  $\Xi_c^+$  (quark content *usc*) and  $\Xi_c^0$  form an isospin doublet, and the spin-parity ought to be  $J^P = 1/2^+$ . None of  $I$ ,  $J$ , or  $P$  has actually been measured.

### $\Xi_c^+$ MASS

The fit uses the  $\Xi_c^+$  and  $\Xi_c^0$  mass and mass-difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2467.9 ± 0.4 OUR FIT</b>				
<b>2467.6 ± 0.4 OUR AVERAGE</b>				
2468.1 ± 0.4 ± 0.2	4950 ± 286	<sup>1</sup> LESIAK	05 BELL	$e^+ e^-$ , $\gamma(4S)$
2465.8 ± 1.9 ± 2.5	90	FRABETTI	98 E687	$\gamma$ Be, $\bar{E}_\gamma = 220$ GeV
2467.0 ± 1.6 ± 2.0	147	EDWARDS	96 CLE2	$e^+ e^- \approx \gamma(4S)$
2465.1 ± 3.6 ± 1.9	30	ALBRECHT	90F ARG	$e^+ e^-$ at $\gamma(4S)$
2467 ± 3 ± 4	23	ALAM	89 CLEO	$e^+ e^-$ 10.6 GeV
2466.5 ± 2.7 ± 1.2	5	BARLAG	89C ACCM	$\pi^-$ Cu 230 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2464.4 ± 2.0 ± 1.4	30	FRABETTI	93B E687	See FRABETTI 98
2459 ± 5 ± 30	56	<sup>2</sup> COTEUS	87 SPEC	$nA \simeq 600$ GeV
2460 ± 25	82	BIAGI	83 SPEC	$\Sigma^-$ Be 135 GeV


<sup>1</sup> The systematic error was (wrongly) given the other way round in LESIAK 05; see the erratum.

<sup>2</sup> Although COTEUS 87 claims to agree well with BIAGI 83 on the mass and width, there appears to be a discrepancy between the two experiments. BIAGI 83 sees a single peak (stated significance about 6 standard deviations) in the  $\Lambda K^- \pi^+ \pi^+$  mass spectrum. COTEUS 87 sees two peaks in the same spectrum, one at the  $\Xi_c^+$  mass, the other 75 MeV lower. The latter is attributed to  $\Xi_c^+ \rightarrow \Sigma^0 K^- \pi^+ \pi^+ \rightarrow (\Lambda \gamma) K^- \pi^+ \pi^+$ , with the  $\gamma$  unseen. The combined significance of the double peak is stated to be 5.5 standard deviations. But the absence of any trace of a lower peak in BIAGI 83 seems to us to throw into question the interpretation of the lower peak of COTEUS 87.

### $\Xi_c^+$ MEAN LIFE

VALUE ( $10^{-15}$ s)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>442 ± 26 OUR AVERAGE</b>		Error includes scale factor of 1.3.		See the ideogram below.
503 ± 47 ± 18	250	MAHMOOD	02 CLE2	$e^+ e^- \approx \gamma(4S)$
439 ± 22 ± 9	532	LINK	01D FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV
$340 \pm 70 \pm 20$	56	FRABETTI	98 E687	$\gamma$ Be, $\bar{E}_\gamma = 220$ GeV
$400 \pm 180 \pm 100$	102	COTEUS	87 SPEC	$nA \simeq 600$ GeV
$480 \pm 210 \pm 200$	53	BIAGI	85C SPEC	$\Sigma^-$ Be 135 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				

$410^{+110}_{-80} \pm 20$	30	FRABETTI	93B	E687	See FRABETTI 98
$200^{+110}_{-60}$	6	BARLAG	89C	ACCM	$\pi^- (K^-)$ Cu 230 GeV



### $\Xi_c^+$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
<b>No absolute branching fractions have been measured.</b>		
The following are branching <i>ratios</i> relative to $\Xi^- \pi^+ \pi^+$ .		
<b>Cabibbo-favored (<math>S = -2</math>) decays</b>		
$\Gamma_1 p K_S^0 K_S^0$	[a] $0.087 \pm 0.022$	
$\Gamma_2 \Lambda \bar{K}^0 \pi^+$	—	
$\Gamma_3 \Sigma(1385)^+ \bar{K}^0$	[a,b] $1.0 \pm 0.5$	
$\Gamma_4 \Lambda K^- \pi^+ \pi^+$	[a] $0.323 \pm 0.033$	
$\Gamma_5 \Lambda \bar{K}^*(892)^0 \pi^+$	[a,b] $< 0.2$	90%
$\Gamma_6 \Sigma(1385)^+ K^- \pi^+$	[a,b] $< 0.3$	90%
$\Gamma_7 \Sigma^+ K^- \pi^+$	[a] $0.94 \pm 0.11$	
$\Gamma_8 \Sigma^+ \bar{K}^*(892)^0$	[a,b] $0.81 \pm 0.15$	
$\Gamma_9 \Sigma^0 K^- \pi^+ \pi^+$	[a] $0.29 \pm 0.16$	

$\Gamma_{10}$	$\Xi^0 \pi^+$	[a] 0.55 $\pm$ 0.16	
$\Gamma_{11}$	$\Xi^- \pi^+ \pi^+$	[a] DEFINED AS 1	
$\Gamma_{12}$	$\Xi(1530)^0 \pi^+$	[a,b] <0.1	90%
$\Gamma_{13}$	$\Xi^0 \pi^+ \pi^0$	[a] 2.34 $\pm$ 0.68	
$\Gamma_{14}$	$\Xi^0 \pi^+ \pi^+ \pi^-$	[a] 1.74 $\pm$ 0.50	
$\Gamma_{15}$	$\Xi^0 e^+ \nu_e$	[a] 2.3 $\pm$ 0.7	
$\Gamma_{16}$	$\Omega^- K^+ \pi^+$	[a] 0.07 $\pm$ 0.04	

**Cabibbo-suppressed decays**

$\Gamma_{17}$	$p K^- \pi^+$	[a] 0.21 $\pm$ 0.03	
$\Gamma_{18}$	$p \bar{K}^*(892)^0$	[a,b] 0.12 $\pm$ 0.02	
$\Gamma_{19}$	$\Sigma^+ K^+ K^-$	[a] 0.15 $\pm$ 0.07	
$\Gamma_{20}$	$\Sigma^+ \phi$	[a,b] <0.11	90%
$\Gamma_{21}$	$\Xi(1690)^0 K^+, \Xi(1690)^0 \rightarrow \Sigma^+ K^-$	[a] <0.05	90%

[a] No absolute branching fractions have been measured. The value here is the branching *ratio* relative to  $\Xi^- \pi^+ \pi^+$ .

[b] This branching fraction includes all the decay modes of the final-state resonance.

 **$\Xi_c^+$  BRANCHING RATIOS****— Cabibbo-favored ( $S = -2$ ) decays —** **$\Gamma(p K_S^0 K_S^0)/\Gamma(\Xi^- \pi^+ \pi^+)$**  **$\Gamma_1/\Gamma_{11}$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.087 \pm 0.016 \pm 0.014</math></b>	$168 \pm 27$	LESIAK	05	BELL $e^+ e^-$ , $\gamma(4S)$

 **$\Gamma(\Sigma(1385)^+ \bar{K}^0)/\Gamma(\Xi^- \pi^+ \pi^+)$**  **$\Gamma_3/\Gamma_{11}$** 

Unseen decay modes of the  $\Sigma(1385)^+$  are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.00 \pm 0.49 \pm 0.24</math></b>	20	LINK	03E	FOCS $< 1.72$ , 90% CL

 **$\Gamma(\Lambda K^- \pi^+ \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$**  **$\Gamma_4/\Gamma_{11}$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.323 \pm 0.033</math> OUR AVERAGE</b>				
0.32 $\pm$ 0.03 $\pm$ 0.02	$1177 \pm 55$	LESIAK	05	BELL $e^+ e^-$ , $\gamma(4S)$
0.28 $\pm$ 0.06 $\pm$ 0.06	58	LINK	03E	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV
0.58 $\pm$ 0.16 $\pm$ 0.07	61	BERGFELD	96	CLE2 $e^+ e^- \approx \gamma(4S)$

 **$\Gamma(\Lambda \bar{K}^*(892)^0 \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+)$**  **$\Gamma_5/\Gamma_4$** 

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.5</b>	90	BERGFELD	96	CLE2 $e^+ e^- \approx \gamma(4S)$

### $\Gamma(\Sigma(1385)^+ K^- \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+)$

$\Gamma_6/\Gamma_4$

Unseen decay modes of the  $\Sigma(1385)^+$  are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	BERGFELD	96	CLE2 $e^+ e^- \approx \gamma(4S)$

### $\Gamma(\Sigma^+ K^- \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$

$\Gamma_7/\Gamma_{11}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.94±0.11 OUR AVERAGE</b>				
0.91±0.11±0.04	251	LINK	03E	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV
1.18±0.26±0.17	119	BERGFELD	96	CLE2 $e^+ e^- \approx \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.92±0.20±0.07	3 JUN		00	SELX $\Sigma^-$ nucleus, 600 GeV

<sup>3</sup>This JUN 00 result is redundant with other results given below.

### $\Gamma(\Sigma^+ \bar{K}^*(892)^0)/\Gamma(\Xi^- \pi^+ \pi^+)$

$\Gamma_8/\Gamma_{11}$

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.81±0.15 OUR AVERAGE</b>				
0.78±0.16±0.06	119	LINK	03E	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV
0.92±0.27±0.14	61	BERGFELD	96	CLE2 $e^+ e^- \approx \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	59	AVERY	95	CLE2 $e^+ e^- \approx \gamma(4S)$

### $\Gamma(\Sigma^0 K^- \pi^+ \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+)$

$\Gamma_9/\Gamma_4$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.84±0.36</b>	47	4 COTEUS	87	SPEC $nA \simeq 600$ GeV

<sup>4</sup> See, however, the note on the COTEUS 87  $\Xi_c^+$  mass measurement.

### $\Gamma(\Xi^0 \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$

$\Gamma_{10}/\Gamma_{11}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.55±0.13±0.09</b>	39	EDWARDS	96	CLE2 $e^+ e^- \approx \gamma(4S)$

### $\Gamma(\Xi^- \pi^+ \pi^+)/\Gamma_{\text{total}}$

$\Gamma_{11}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	131	BERGFELD	96	CLE2 $e^+ e^- \approx \gamma(4S)$
seen	160	AVERY	95	CLE2 $e^+ e^- \approx \gamma(4S)$
seen	30	FRABETTI	93B	E687 $\gamma$ Be, $\bar{E}_\gamma = 220$ GeV
seen	30	ALBRECHT	90F	ARG $e^+ e^-$ at $\gamma(4S)$
seen	23	ALAM	89	CLEO $e^+ e^-$ 10.6 GeV

### $\Gamma(\Xi(1530)^0 \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$

$\Gamma_{12}/\Gamma_{11}$

Unseen decay modes of the  $\Xi(1530)^0$  are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.1</b>	90	LINK	03E	FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.2	90	BERGFELD	96	CLE2 $e^+ e^- \approx \gamma(4S)$
------	----	----------	----	-----------------------------------

$\Gamma(\Xi^0 \pi^+ \pi^0)/\Gamma(\Xi^- \pi^+ \pi^+)$   $\Gamma_{13}/\Gamma_{11}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.34±0.57±0.37</b>	81	EDWARDS 96	CLE2	$e^+ e^- \approx \gamma(4S)$

$\Gamma(\Xi(1530)^0 \pi^+)/\Gamma(\Xi^0 \pi^+ \pi^0)$   $\Gamma_{12}/\Gamma_{13}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
<0.3	90	EDWARDS 96	CLE2	$e^+ e^- \approx \gamma(4S)$

$\Gamma(\Xi^0 \pi^+ \pi^+ \pi^-)/\Gamma(\Xi^- \pi^+ \pi^+)$   $\Gamma_{14}/\Gamma_{11}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.74±0.42±0.27</b>	57	EDWARDS 96	CLE2	$e^+ e^- \approx \gamma(4S)$

$\Gamma(\Xi^0 e^+ \nu_e)/\Gamma(\Xi^- \pi^+ \pi^+)$   $\Gamma_{15}/\Gamma_{11}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3±0.6±0.3</b>	41	ALEXANDER 95B	CLE2	$e^+ e^- \approx \gamma(4S)$

$\Gamma(\Omega^- K^+ \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$   $\Gamma_{16}/\Gamma_{11}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.07±0.03±0.03</b>	14	LINK 03E	FOCS	< 0.12, 90% CL

**Cabibbo-suppressed decays**

$\Gamma(p K^- \pi^+)/\Gamma(\Sigma^+ K^- \pi^+)$   $\Gamma_{17}/\Gamma_7$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				

0.22±0.06±0.03 76 JUN 00 SELX  $\Sigma^-$  nucleus, 600 GeV

$\Gamma(p K^- \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$   $\Gamma_{17}/\Gamma_{11}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.214±0.034 OUR AVERAGE</b>				
0.234±0.047±0.022	202	LINK 01B	FOCS	$\gamma$ nucleus
0.20 ± 0.04 ± 0.02	76	JUN 00	SELX	$\Sigma^-$ nucleus, 600 GeV

$\Gamma(p \bar{K}^*(892)^0)/\Gamma(p K^- \pi^+)$   $\Gamma_{18}/\Gamma_{17}$

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.54±0.09±0.05</b>		LINK 01B	FOCS	$\gamma$ nucleus

$\Gamma(\Sigma^+ K^+ K^-)/\Gamma(\Sigma^+ K^- \pi^+)$   $\Gamma_{19}/\Gamma_7$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.16±0.06±0.01</b>	17	LINK 03E	FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(\Sigma^+ \phi)/\Gamma(\Sigma^+ K^- \pi^+)$   $\Gamma_{20}/\Gamma_7$

Unseen decay modes of the  $\phi$  are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.12</b>	90	LINK 03E	FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV

$\Gamma(\Xi(1690)^0 K^+ \times B(\Xi(1690)^0 \rightarrow \Sigma^+ K^-))/\Gamma(\Sigma^+ K^- \pi^+)$					$\Gamma_{21}/\Gamma_7$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.05	90	LINK	03E FOCS	$\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV	

## $\Xi_c^+$ REFERENCES

LESIAK Also	05	PL B605 237 PL B617 198 (erratum)	T. Lesiak <i>et al.</i> T. Lesiak <i>et al.</i>	(BELLE Collab.) (BELLE Collab.)
LINK	03E	PL B571 139	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
MAHMOOD	02	PR D65 031102	A.H. Mahmood <i>et al.</i>	(CLEO Collab.)
LINK	01B	PL B512 277	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	01D	PL B523 53	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
JUN	00	PRL 84 1857	S.Y. Jun <i>et al.</i>	(FNAL SELEX Collab.)
FRAEBETTI	98	PL B427 211	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
BERGFELD	96	PL B365 431	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
EDWARDS	96	PL B373 261	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ALEXANDER Also	95B	PRL 74 3113 PRL 75 4155 (erratum)	J. Alexander <i>et al.</i> J. Alexander <i>et al.</i>	(CLEO Collab.)
AVERY	95	PRL 75 4364	P. Avery <i>et al.</i>	(CLEO Collab.)
FRAEBETTI	93B	PRL 70 1381	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALBRECHT	90F	PL B247 121	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALAM	89	PL B226 401	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BARLAG	89C	PL B233 522	S. Barlag <i>et al.</i>	(ACCMOR Collab.)
COTEUS	87	PRL 59 1530	P. Coteus <i>et al.</i>	(FNAL E400 Collab.)
BIAGI	85C	PL 150B 230	S.F. Biagi <i>et al.</i>	(CERN WA62 Collab.)
BIAGI	83	PL 122B 455	S.F. Biagi <i>et al.</i>	(CERN WA62 Collab.)