

$\Delta(1620)$ S_{31} $I(J^P) = \frac{3}{2}(\frac{1}{2}^-)$ Status: ***

Most of the results published before 1975 were last included in our 1982 edition, Physics Letters **111B** 1 (1982). Some further obsolete results published before 1984 were last included in our 2006 edition, Journal of Physics, G **33** 1 (2006).

 $\Delta(1620)$ BREIT-WIGNER MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|--------------------------|------|-------------------------------------|
| 1600 to 1660 (≈ 1630) OUR ESTIMATE | | | |
| 1615.2 \pm 0.4 | ARNDT 06 | DPWA | $\pi N \rightarrow \pi N, \eta N$ |
| 1672 \pm 7 | MANLEY 92 | IPWA | $\pi N \rightarrow \pi N & N\pi\pi$ |
| 1620 \pm 20 | CUTKOSKY 80 | IPWA | $\pi N \rightarrow \pi N$ |
| 1610 \pm 7 | HOEHLER 79 | IPWA | $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1650 \pm 25 | THOMA 08 | DPWA | Multichannel |
| 1614.1 \pm 1.1 | ARNDT 04 | DPWA | $\pi N \rightarrow \pi N, \eta N$ |
| 1612 \pm 2 | PENNER 02C | DPWA | Multichannel |
| 1617 \pm 15 | VRANA 00 | DPWA | Multichannel |
| 1672 \pm 5 | ARNDT 96 | IPWA | $\gamma N \rightarrow \pi N$ |
| 1617 | ARNDT 95 | DPWA | $\pi N \rightarrow N\pi$ |
| 1669 | LI 93 | IPWA | $\gamma N \rightarrow \pi N$ |
| 1620 | BARNHAM 80 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| 1712.8 \pm 6.0 | ¹ CHEW 80 | BPWA | $\pi^+ p \rightarrow \pi^+ p$ |
| 1786.7 \pm 2.0 | ¹ CHEW 80 | BPWA | $\pi^+ p \rightarrow \pi^+ p$ |
| 1580 | ² LONGACRE 77 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| 1600 | ³ LONGACRE 75 | IPWA | $\pi N \rightarrow N\pi\pi$ |

 $\Delta(1620)$ BREIT-WIGNER WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|-------------------------------------|
| 135 to 150 (≈ 145) OUR ESTIMATE | | | |
| 146.9 \pm 1.9 | ARNDT 06 | DPWA | $\pi N \rightarrow \pi N, \eta N$ |
| 154 \pm 37 | MANLEY 92 | IPWA | $\pi N \rightarrow \pi N & N\pi\pi$ |
| 140 \pm 20 | CUTKOSKY 80 | IPWA | $\pi N \rightarrow \pi N$ |
| 139 \pm 18 | HOEHLER 79 | IPWA | $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 250 \pm 60 | THOMA 08 | DPWA | Multichannel |
| 141.0 \pm 6.0 | ARNDT 04 | DPWA | $\pi N \rightarrow \pi N, \eta N$ |
| 202 \pm 7 | PENNER 02C | DPWA | Multichannel |
| 143 \pm 42 | VRANA 00 | DPWA | Multichannel |
| 147 \pm 8 | ARNDT 96 | IPWA | $\gamma N \rightarrow \pi N$ |
| 108 | ARNDT 95 | DPWA | $\pi N \rightarrow N\pi$ |

| | | | | |
|--------------|-----------------------|----|------|---|
| 184 | LI | 93 | IPWA | $\gamma N \rightarrow \pi N$ |
| 120 | BARNHAM | 80 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| 228.3 ± 18.0 | ¹ CHEW | 80 | BPWA | $\pi^+ p \rightarrow \pi^+ p$ (lower mass) |
| 30.0 ± 6.4 | ¹ CHEW | 80 | BPWA | $\pi^+ p \rightarrow \pi^+ p$ (higher mass) |
| 120 | ² LONGACRE | 77 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| 150 | ³ LONGACRE | 75 | IPWA | $\pi N \rightarrow N\pi\pi$ |

$\Delta(1620)$ POLE POSITION

REAL PART

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|------|--|
| 1590 to 1610 (≈ 1600) OUR ESTIMATE | | | |
| 1595 | ARNDT | 06 | DPWA $\pi N \rightarrow \pi N, \eta N$ |
| 1608 | ⁴ HOEHLER | 93 | SPED $\pi N \rightarrow \pi N$ |
| 1600 ± 15 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1615 ± 25 | THOMA | 08 | DPWA Multichannel |
| 1594 | ARNDT | 04 | DPWA $\pi N \rightarrow \pi N, \eta N$ |
| 1607 | VRANA | 00 | DPWA Multichannel |
| 1585 | ARNDT | 95 | DPWA $\pi N \rightarrow N\pi$ |
| 1587 | ARNDT | 91 | DPWA $\pi N \rightarrow \pi N$ Soln SM90 |
| 1583 or 1583 | ⁵ LONGACRE | 78 | IPWA $\pi N \rightarrow N\pi\pi$ |
| 1575 or 1572 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |

-2×IMAGINARY PART

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-----------------------|------|--|
| 115 to 120 (≈ 118) OUR ESTIMATE | | | |
| 135 | ARNDT | 06 | DPWA $\pi N \rightarrow \pi N, \eta N$ |
| 116 | ⁴ HOEHLER | 93 | SPED $\pi N \rightarrow \pi N$ |
| 120 ± 20 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 180 ± 35 | THOMA | 08 | DPWA Multichannel |
| 118 | ARNDT | 04 | DPWA $\pi N \rightarrow \pi N, \eta N$ |
| 148 | VRANA | 00 | DPWA Multichannel |
| 104 | ARNDT | 95 | DPWA $\pi N \rightarrow N\pi$ |
| 120 | ARNDT | 91 | DPWA $\pi N \rightarrow \pi N$ Soln SM90 |
| 143 or 149 | ⁵ LONGACRE | 78 | IPWA $\pi N \rightarrow N\pi\pi$ |
| 119 or 128 | ² LONGACRE | 77 | IPWA $\pi N \rightarrow N\pi\pi$ |

$\Delta(1620)$ ELASTIC POLE RESIDUE

MODULUS $|r|$

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|--|
| 15 | ARNDT | 06 | DPWA $\pi N \rightarrow \pi N, \eta N$ |
| 19 | HOEHLER | 93 | SPED $\pi N \rightarrow \pi N$ |
| 15 ± 2 | CUTKOSKY | 80 | IPWA $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 17 | ARNDT | 04 | DPWA $\pi N \rightarrow \pi N, \eta N$ |
| 14 | ARNDT | 95 | DPWA $\pi N \rightarrow N\pi$ |
| 15 | ARNDT | 91 | DPWA $\pi N \rightarrow \pi N$ Soln SM90 |

PHASE θ

| VALUE ($^{\circ}$) | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|-------------------------------------|
| - 92 | ARNDT 06 | DPWA | $\pi N \rightarrow \pi N, \eta N$ |
| - 95 | HOEHLER 93 | SPED | $\pi N \rightarrow \pi N$ |
| -110±20 | CUTKOSKY 80 | IPWA | $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -104 | ARNDT 04 | DPWA | $\pi N \rightarrow \pi N, \eta N$ |
| -121 | ARNDT 95 | DPWA | $\pi N \rightarrow N\pi$ |
| -125 | ARNDT 91 | DPWA | $\pi N \rightarrow \pi N$ Soln SM90 |

 $\Delta(1620)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

| Mode | Fraction (Γ_i/Γ) |
|--|--------------------------------|
| $\Gamma_1 N\pi$ | 20–30 % |
| $\Gamma_2 N\pi\pi$ | 70–80 % |
| $\Gamma_3 \Delta\pi$ | 30–60 % |
| $\Gamma_4 \Delta(1232)\pi, D\text{-wave}$ | |
| $\Gamma_5 N\rho$ | 7–25 % |
| $\Gamma_6 N\rho, S=1/2, S\text{-wave}$ | |
| $\Gamma_7 N\rho, S=3/2, D\text{-wave}$ | |
| $\Gamma_8 N(1440)\pi$ | |
| $\Gamma_9 N\gamma$ | 0.004–0.044 % |
| $\Gamma_{10} N\gamma, \text{helicity}=1/2$ | 0.004–0.044 % |

 $\Delta(1620)$ BRANCHING RATIOS

| $\Gamma(N\pi)/\Gamma_{\text{total}}$ | Γ_1/Γ |
|--|---|
| 0.2 to 0.3 OUR ESTIMATE | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | |
| 0.315±0.001 | ARNDT 06 DPWA $\pi N \rightarrow \pi N, \eta N$ |
| 0.09 ± 0.02 | MANLEY 92 IPWA $\pi N \rightarrow \pi N & N\pi\pi$ |
| 0.25 ± 0.03 | CUTKOSKY 80 IPWA $\pi N \rightarrow \pi N$ |
| 0.35 ± 0.06 | HOEHLER 79 IPWA $\pi N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | |
| 0.22 ± 0.12 | THOMA 08 DPWA Multichannel |
| 0.310±0.004 | ARNDT 04 DPWA $\pi N \rightarrow \pi N, \eta N$ |
| 0.34 ± 0.01 | PENNER 02C DPWA Multichannel |
| 0.45 ± 0.05 | VRANA 00 DPWA Multichannel |
| 0.29 | ARNDT 95 DPWA $\pi N \rightarrow N\pi$ |
| 0.60 | ¹ CHEW 80 BPWA $\pi^+ p \rightarrow \pi^+ p$ (lower mass) |
| 0.36 | ¹ CHEW 80 BPWA $\pi^+ p \rightarrow \pi^+ p$ (higher mass) |

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase

ambiguity is resolved by choosing a negative sign for the $\Delta(1620)$ S_{31} coupling to $\Delta(1232)\pi$.

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N\pi \rightarrow \Delta(1620) \rightarrow \Delta(1232)\pi, D\text{-wave} \quad (\Gamma_1 \Gamma_4)^{1/2} / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|----------------------------|------|-------------------------------------|
| -0.36 to -0.28 OUR ESTIMATE | | | |
| -0.24 ± 0.03 | MANLEY 92 | IPWA | $\pi N \rightarrow \pi N & N\pi\pi$ |
| -0.33 ± 0.06 | BARNHAM 80 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| -0.39 | ^{2,6} LONGACRE 77 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| -0.40 | ³ LONGACRE 75 | IPWA | $\pi N \rightarrow N\pi\pi$ |

$$\Gamma(\Delta(1232)\pi, D\text{-wave}) / \Gamma_{\text{total}} \quad \Gamma_4 / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|--------------|
| 0.39 ± 0.02 | VRANA 00 | DPWA | Multichannel |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.48 ± 0.25 | THOMA 08 | DPWA | Multichannel |

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N\pi \rightarrow \Delta(1620) \rightarrow N\rho, S=1/2, S\text{-wave} \quad (\Gamma_1 \Gamma_6)^{1/2} / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|----------------------------|------|-------------------------------------|
| +0.12 to +0.22 OUR ESTIMATE | | | |
| +0.15 ± 0.02 | MANLEY 92 | IPWA | $\pi N \rightarrow \pi N & N\pi\pi$ |
| +0.40 ± 0.10 | BARNHAM 80 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| +0.08 | ^{2,6} LONGACRE 77 | IPWA | $\pi N \rightarrow N\pi\pi$ |
| +0.28 | ³ LONGACRE 75 | IPWA | $\pi N \rightarrow N\pi\pi$ |

$$\Gamma(N\rho, S=1/2, S\text{-wave}) / \Gamma_{\text{total}} \quad \Gamma_6 / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|--------------|
| 0.14 ± 0.03 | VRANA 00 | DPWA | Multichannel |

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N\pi \rightarrow \Delta(1620) \rightarrow N\rho, S=3/2, D\text{-wave} \quad (\Gamma_1 \Gamma_7)^{1/2} / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------------|----------------------------|------|-------------------------------------|
| -0.15 to -0.03 OUR ESTIMATE | | | |
| -0.06 ± 0.02 | MANLEY 92 | IPWA | $\pi N \rightarrow \pi N & N\pi\pi$ |
| -0.13 | ^{2,6} LONGACRE 77 | IPWA | $\pi N \rightarrow N\pi\pi$ |

$$\Gamma(N\rho, S=3/2, D\text{-wave}) / \Gamma_{\text{total}} \quad \Gamma_7 / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|--------------|
| 0.02 ± 0.01 | VRANA 00 | DPWA | Multichannel |

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N\pi \rightarrow \Delta(1620) \rightarrow N(1440)\pi \quad (\Gamma_1 \Gamma_8)^{1/2} / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|-----------------------------|
| 0.11 ± 0.05 | BARNHAM 80 | IPWA | $\pi N \rightarrow N\pi\pi$ |

$$\Gamma(N(1440)\pi) / \Gamma_{\text{total}} \quad \Gamma_8 / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|--------------|
| 0.00 ± 0.01 | VRANA 00 | DPWA | Multichannel |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.19 ± 0.12 | THOMA 08 | DPWA | Multichannel |

$\Delta(1620)$ PHOTON DECAY AMPLITUDES

Papers on γN amplitudes predating 1981 may be found in our 2006 edition,
Journal of Physics, G **33** 1 (2006).

$\Delta(1620) \rightarrow N\gamma$, helicity-1/2 amplitude $A_{1/2}$

| VALUE (GeV $^{-1/2}$) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|------------------------------|
| +0.027±0.011 OUR ESTIMATE | | | |
| 0.050±0.002 | DUGGER 07 | DPWA | $\gamma N \rightarrow \pi N$ |
| 0.035±0.020 | ARNDT 96 | IPWA | $\gamma N \rightarrow \pi N$ |
| 0.035±0.010 | CRAWFORD 83 | IPWA | $\gamma N \rightarrow \pi N$ |
| 0.010±0.015 | AWAJI 81 | DPWA | $\gamma N \rightarrow \pi N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.066 | DRECHSEL 07 | DPWA | $\gamma N \rightarrow \pi N$ |
| -0.050 | PENNER 02D | DPWA | Multichannel |
| 0.042±0.003 | LI 93 | IPWA | $\gamma N \rightarrow \pi N$ |
| 0.066 | WADA 84 | DPWA | Compton scattering |

$\Delta(1620)$ FOOTNOTES

¹ CHEW 80 reports two S_{31} resonances at somewhat higher masses than other analyses.
Problems with this analysis are discussed in section 2.1.11 of HOEHLER 83.

² LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

³ From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

⁴ See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

⁵ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

⁶ LONGACRE 77 considers this coupling to be well determined.

$\Delta(1620)$ REFERENCES

For early references, see Physics Letters **111B** 1 (1982).

| | | | | |
|----------|-----|------------------------|---|------------------------------|
| THOMA | 08 | PL B659 87 | U. Thoma <i>et al.</i> | (CB-ELSA Collab.) |
| DRECHSEL | 07 | EPJ A34 69 | D. Drechsel, S.S. Kamalov, L. Tiator | (MAINZ, JINR) |
| DUGGER | 07 | PR C76 025211 | M. Dugger <i>et al.</i> | (Jefferson Lab CLAS Collab.) |
| ARNDT | 06 | PR C74 045205 | R.A. Arndt <i>et al.</i> | (GWU) |
| PDG | 06 | JPG 33 1 | W.-M. Yao <i>et al.</i> | (PDG Collab.) |
| ARNDT | 04 | PR C69 035213 | R.A. Arndt <i>et al.</i> | (GWU, TRIU) |
| PENNER | 02C | PR C66 055211 | G. Penner, U. Mosel | (GIES) |
| PENNER | 02D | PR C66 055212 | G. Penner, U. Mosel | (GIES) |
| VRANA | 00 | PRPL 328 181 | T.P. Vrana, S.A. Dytman,, T.-S.H. Lee | (PITT+) |
| ARNDT | 96 | PR C53 430 | R.A. Arndt, I.I. Strakovsky, R.L. Workman | (VPI) |
| ARNDT | 95 | PR C52 2120 | R.A. Arndt <i>et al.</i> | (VPI, BRCO) |
| HOEHLER | 93 | πN Newsletter 9 1 | G. Hohler | (KARL) |
| LI | 93 | PR C47 2759 | Z.J. Li <i>et al.</i> | (VPI) |
| MANLEY | 92 | PR D45 4002 | D.M. Manley, E.M. Saleski | (KENT) IJP |
| Also | | PR D30 904 | D.M. Manley <i>et al.</i> | (VPI) |

| | | | | |
|------------------|----|-------------------------|------------------------------|-------------------|
| ARNDT | 91 | PR D43 2131 | R.A. Arndt <i>et al.</i> | (VPI, TELE) IJP |
| WADA | 84 | NP B247 313 | Y. Wada <i>et al.</i> | (INUS) |
| CRAWFORD | 83 | NP B211 1 | R.L. Crawford, W.T. Morton | (GLAS) |
| HOEHLER | 83 | Landolt-Bornstein 1/9B2 | G. Hohler | (KARLT) |
| PDG | 82 | PL 111B 1 | M. Roos <i>et al.</i> | (HELS, CIT, CERN) |
| AWAJI Also | 81 | Bonn Conf. 352 | N. Awaji, R. Kajikawa | (NAGO) |
| BARNHAM | 80 | NP B197 365 | K. Fujii <i>et al.</i> | (NAGO) |
| CHEW | 80 | NP B168 243 | K.W.J. Barnham <i>et al.</i> | (LOIC) |
| CUTKOSKY | 80 | Toronto Conf. 123 | D.M. Chew | (LBL) IJP |
| HOEHLER Also | 79 | Toronto Conf. 19 | R.E. Cutkosky <i>et al.</i> | (CMU, LBL) IJP |
| LONGACRE Also | 78 | PR D20 2839 | R.E. Cutkosky <i>et al.</i> | (CMU, LBL) IJP |
| LONGACRE | 77 | PDAT 12-1 | G. Hohler <i>et al.</i> | (KARLT) IJP |
| LONGACRE | 77 | Toronto Conf. 3 | R. Koch | (KARLT) IJP |
| LONGACRE Also | 75 | PR D17 1795 | R.S. Longacre <i>et al.</i> | (LBL, SLAC) |
| LONGACRE | 75 | NP B122 493 | R.S. Longacre, J. Dolbeau | (SACL) IJP |
| LONGACRE | 75 | NP B108 365 | J. Dolbeau <i>et al.</i> | (SACL) IJP |
| LONGACRE | 75 | PL 55B 415 | R.S. Longacre <i>et al.</i> | (LBL, SLAC) IJP |