

$\psi(2S)$

$I^G(J^{PC}) = 0^-(1^{--})$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

$\psi(2S)$ MASS

OUR FIT includes measurements of $m_{\psi}(2S)$, $m_{\psi}(3770)$, and $m_{\psi}(3770) - m_{\psi}(2S)$.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

3686.09 ± 0.04 OUR FIT Error includes scale factor of 1.6.

3686.093±0.034 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

3686.111±0.025±0.009 AULCHENKO 03 KEDR $e^+ e^- \rightarrow$ hadrons

3685.95 ± 0.10 413 ¹ ARTAMONOV 00 OLYA $e^+ e^- \rightarrow$ hadrons

3685.98 ± 0.09 ± 0.04 ² ARMSTRONG 93B E760 $\bar{p}p \rightarrow e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3686.00 ± 0.10 413 ³ ZHOLENTZ 80 OLYA $e^+ e^-$


¹ Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

² Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $J/\psi(1S)$ mass from AULCHENKO 03.

³ Superseded by ARTAMONOV 00.

WEIGHTED AVERAGE

3686.093±0.034 (Error scaled by 1.4)



$m_{\psi(2S)} - m_{J/\psi(1S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
589.188±0.028 OUR AVERAGE			
589.194±0.027±0.011	⁴ AULCHENKO 03	KEDR	$e^+ e^- \rightarrow$ hadrons
589.7 ± 1.2	LEMOIGNE 82	GOLI	$185 \pi^- Be \rightarrow \gamma \mu^+ \mu^- A$
589.07 ± 0.13	⁴ ZHOLENTZ 80	OLYA	$e^+ e^-$
588.7 ± 0.8	LUTH 75	MRK1	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
588 ± 1	⁵ BAI	98E BES	$e^+ e^-$
⁴ Redundant with data in mass above. ⁵ Systematic errors not evaluated.			

 $\psi(2S)$ WIDTH

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
317± 9 OUR FIT				
286±16 OUR AVERAGE				
358±88± 4		ABLIKIM 08B	BES2	$e^+ e^- \rightarrow$ hadrons
290±25± 4	2.7k	ANDREOTTI 07	E835	$p\bar{p} \rightarrow e^+ e^-, J/\psi X$
331±58± 2		ABLIKIM 06L	BES2	$e^+ e^- \rightarrow$ hadrons
264±27		⁶ BAI 02B	BES2	$e^+ e^-$
287±37±16		⁷ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+ e^-$
⁶ From a simultaneous fit to the hadronic and $\mu^+ \mu^-$ cross section, assuming $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$ and lepton universality. Does not include vacuum polarization correction.				
⁷ The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].				

 $\psi(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(97.85±0.13) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(1.73±0.14) %	S=1.5
Γ_3 light hadrons		
Γ_4 $e^+ e^-$	(7.52±0.17) $\times 10^{-3}$	
Γ_5 $\mu^+ \mu^-$	(7.5 ± 0.8) $\times 10^{-3}$	
Γ_6 $\tau^+ \tau^-$	(3.0 ± 0.4) $\times 10^{-3}$	

Decays into $J/\psi(1S)$ and anything

Γ_7 $J/\psi(1S)$ anything	(57.4 ± 0.9) %	
Γ_8 $J/\psi(1S)$ neutrals	(23.5 ± 0.4) %	
Γ_9 $J/\psi(1S)$ $\pi^+ \pi^-$	(32.6 ± 0.5) %	
Γ_{10} $J/\psi(1S)$ $\pi^0 \pi^0$	(16.84±0.33) %	
Γ_{11} $J/\psi(1S)$ η	(3.16±0.07) %	
Γ_{12} $J/\psi(1S)$ π^0	(1.26±0.13) $\times 10^{-3}$	S=1.3

Hadronic decays

Γ_{13}	$3(\pi^+\pi^-)\pi^0$	$(3.5 \pm 1.6) \times 10^{-3}$	
Γ_{14}	$2(\pi^+\pi^-)\pi^0$	$(2.9 \pm 1.0) \times 10^{-3}$	S=4.6
Γ_{15}	$\rho a_2(1320)$	$(2.6 \pm 0.9) \times 10^{-4}$	
Γ_{16}	$p\bar{p}$	$(2.74 \pm 0.12) \times 10^{-4}$	
Γ_{17}	$\Delta^{++}\bar{\Delta}^{--}$	$(1.28 \pm 0.35) \times 10^{-4}$	
Γ_{18}	$\Lambda\bar{\Lambda}\pi^0$	$< 1.2 \times 10^{-4}$	CL=90%
Γ_{19}	$\Lambda\bar{\Lambda}\eta$	$< 4.9 \times 10^{-5}$	CL=90%
Γ_{20}	$\Lambda\bar{p}K^+$	$(1.00 \pm 0.14) \times 10^{-4}$	
Γ_{21}	$\Lambda\bar{p}K^+\pi^+\pi^-$	$(1.8 \pm 0.4) \times 10^{-4}$	
Γ_{22}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(2.8 \pm 0.6) \times 10^{-4}$	
Γ_{23}	$\Lambda\bar{\Lambda}$	$(2.8 \pm 0.5) \times 10^{-4}$	S=2.6
Γ_{24}	$\Sigma^+\bar{\Sigma}^-$	$(2.6 \pm 0.8) \times 10^{-4}$	
Γ_{25}	$\Sigma^0\bar{\Sigma}^0$	$(2.2 \pm 0.4) \times 10^{-4}$	S=1.5
Γ_{26}	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$(1.1 \pm 0.4) \times 10^{-4}$	
Γ_{27}	$\Xi^-\bar{\Xi}^+$	$(1.8 \pm 0.6) \times 10^{-4}$	S=2.8
Γ_{28}	$\Xi^0\bar{\Xi}^0$	$(2.8 \pm 0.9) \times 10^{-4}$	
Γ_{29}	$\Xi(1530)^0\bar{\Xi}(1530)^0$	$< 8.1 \times 10^{-5}$	CL=90%
Γ_{30}	$\Omega^-\bar{\Omega}^+$	$< 7.3 \times 10^{-5}$	CL=90%
Γ_{31}	$\pi^0 p\bar{p}$	$(1.33 \pm 0.17) \times 10^{-4}$	
Γ_{32}	$\eta p\bar{p}$	$(6.0 \pm 1.2) \times 10^{-5}$	
Γ_{33}	$\omega p\bar{p}$	$(6.9 \pm 2.1) \times 10^{-5}$	
Γ_{34}	$\phi p\bar{p}$	$< 2.4 \times 10^{-5}$	CL=90%
Γ_{35}	$\pi^+\pi^- p\bar{p}$	$(6.0 \pm 0.4) \times 10^{-4}$	
Γ_{36}	$p\bar{n}\pi^-$ or c.c.	$(2.48 \pm 0.17) \times 10^{-4}$	
Γ_{37}	$p\bar{n}\pi^-\pi^0$	$(3.2 \pm 0.7) \times 10^{-4}$	
Γ_{38}	$2(\pi^+\pi^-\pi^0)$	$(4.7 \pm 1.5) \times 10^{-3}$	
Γ_{39}	$\eta\pi^+\pi^-$	$< 1.6 \times 10^{-4}$	CL=90%
Γ_{40}	$\eta\pi^+\pi^-\pi^0$	$(9.5 \pm 1.7) \times 10^{-4}$	
Γ_{41}	$2(\pi^+\pi^-)\eta$	$(1.2 \pm 0.6) \times 10^{-3}$	
Γ_{42}	$\eta'\pi^+\pi^-\pi^0$	$(4.5 \pm 2.1) \times 10^{-4}$	
Γ_{43}	$\omega\pi^+\pi^-$	$(7.3 \pm 1.2) \times 10^{-4}$	S=2.1
Γ_{44}	$b_1^\pm\pi^\mp$	$(4.0 \pm 0.6) \times 10^{-4}$	S=1.1
Γ_{45}	$b_1^0\pi^0$	$(2.4 \pm 0.6) \times 10^{-4}$	
Γ_{46}	$\omega f_2(1270)$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{47}	$\pi^+\pi^- K^+K^-$	$(7.5 \pm 0.9) \times 10^{-4}$	S=1.9
Γ_{48}	$\rho^0 K^+K^-$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{49}	$K^*(892)^0\bar{K}_2^*(1430)^0$	$(1.9 \pm 0.5) \times 10^{-4}$	
Γ_{50}	$K^+K^-\pi^+\pi^-\eta$	$(1.3 \pm 0.7) \times 10^{-3}$	
Γ_{51}	$K^+K^-2(\pi^+\pi^-)\pi^0$	$(1.00 \pm 0.31) \times 10^{-3}$	
Γ_{52}	$K^+K^-2(\pi^+\pi^-)$	$(1.8 \pm 0.9) \times 10^{-3}$	
Γ_{53}	$K_1(1270)^\pm K^\mp$	$(1.00 \pm 0.28) \times 10^{-3}$	
Γ_{54}	$K_S^0 K_S^0 \pi^+\pi^-$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{55}	$\rho^0 p\bar{p}$	$(5.0 \pm 2.2) \times 10^{-5}$	

Γ_{56}	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(6.7 \pm 2.5) \times 10^{-4}$	
Γ_{57}	$2(\pi^+ \pi^-)$	$(2.4 \pm 0.6) \times 10^{-4}$	S=2.2
Γ_{58}	$\rho^0 \pi^+ \pi^-$	$(2.2 \pm 0.6) \times 10^{-4}$	S=1.4
Γ_{59}	$K^+ K^- \pi^+ \pi^- \pi^0$	$(1.26 \pm 0.09) \times 10^{-3}$	
Γ_{60}	$\omega f_0(1710) \rightarrow \omega K^+ K^-$	$(5.9 \pm 2.2) \times 10^{-5}$	
Γ_{61}	$K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}$	$(8.6 \pm 2.2) \times 10^{-4}$	
Γ_{62}	$K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}$	$(9.6 \pm 2.8) \times 10^{-4}$	
Γ_{63}	$K^*(892)^+ K^- \rho^0 + \text{c.c.}$	$(7.3 \pm 2.6) \times 10^{-4}$	
Γ_{64}	$K^*(892)^0 K^- \rho^+ + \text{c.c.}$	$(6.1 \pm 1.8) \times 10^{-4}$	
Γ_{65}	$\eta K^+ K^-$	$< 1.3 \times 10^{-4}$	CL=90%
Γ_{66}	$\omega K^+ K^-$	$(1.85 \pm 0.25) \times 10^{-4}$	S=1.1
Γ_{67}	$3(\pi^+ \pi^-)$	$(3.5 \pm 2.0) \times 10^{-4}$	S=2.8
Γ_{68}	$p\bar{p} \pi^+ \pi^- \pi^0$	$(7.3 \pm 0.7) \times 10^{-4}$	
Γ_{69}	$K^+ K^-$	$(6.3 \pm 0.7) \times 10^{-5}$	
Γ_{70}	$K_S^0 K_L^0$	$(5.4 \pm 0.5) \times 10^{-5}$	
Γ_{71}	$\pi^+ \pi^- \pi^0$	$(1.68 \pm 0.26) \times 10^{-4}$	S=1.4
Γ_{72}	$\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$	$(1.9 \pm 1.2) \times 10^{-4}$	
Γ_{73}	$\rho(770)\pi \rightarrow \pi^+ \pi^- \pi^0$	$(3.2 \pm 1.2) \times 10^{-5}$	S=1.8
Γ_{74}	$\pi^+ \pi^-$	$(8 \pm 5) \times 10^{-5}$	
Γ_{75}	$K_1(1400)^\pm K^\mp$	$< 3.1 \times 10^{-4}$	CL=90%
Γ_{76}	$K^+ K^- \pi^0$	$< 2.96 \times 10^{-5}$	CL=90%
Γ_{77}	$K^+ \bar{K}^*(892)^- + \text{c.c.}$	$(1.7 \pm 0.8) \times 10^{-5}$	
Γ_{78}	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(1.09 \pm 0.20) \times 10^{-4}$	
Γ_{79}	$\phi \pi^+ \pi^-$	$(1.17 \pm 0.29) \times 10^{-4}$	S=1.7
Γ_{80}	$\phi f_0(980) \rightarrow \pi^+ \pi^-$	$(6.8 \pm 2.4) \times 10^{-5}$	S=1.1
Γ_{81}	$2(K^+ K^-)$	$(6.0 \pm 1.4) \times 10^{-5}$	
Γ_{82}	$\phi K^+ K^-$	$(7.0 \pm 1.6) \times 10^{-5}$	
Γ_{83}	$2(K^+ K^-) \pi^0$	$(1.10 \pm 0.28) \times 10^{-4}$	
Γ_{84}	$\phi \eta$	$(2.8 \pm 1.0) \times 10^{-5}$	
Γ_{85}	$\phi \eta'$	$(3.1 \pm 1.6) \times 10^{-5}$	
Γ_{86}	$\omega \eta'$	$(3.2 \pm 2.5) \times 10^{-5}$	
Γ_{87}	$\omega \pi^0$	$(2.1 \pm 0.6) \times 10^{-5}$	
Γ_{88}	$\rho \eta'$	$(1.9 \pm 1.7) \times 10^{-5}$	
Γ_{89}	$\rho \eta$	$(2.2 \pm 0.6) \times 10^{-5}$	S=1.1
Γ_{90}	$\omega \eta$	$< 1.1 \times 10^{-5}$	CL=90%
Γ_{91}	$\phi \pi^0$	$< 4 \times 10^{-6}$	CL=90%
Γ_{92}	$\eta_c \pi^+ \pi^- \pi^0$	$< 1.0 \times 10^{-3}$	CL=90%
Γ_{93}	$p\bar{p} K^+ K^-$	$(2.7 \pm 0.7) \times 10^{-5}$	
Γ_{94}	$\bar{\Lambda} n K_S^0 + \text{c.c.}$	$(8.1 \pm 1.8) \times 10^{-5}$	
Γ_{95}	$\phi f'_2(1525)$	$(4.4 \pm 1.6) \times 10^{-5}$	

Γ_{96}	$\Theta(1540)\overline{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} +$	< 8.8	$\times 10^{-6}$	CL=90%
	c.c.			
Γ_{97}	$\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	< 1.0	$\times 10^{-5}$	CL=90%
Γ_{98}	$\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	< 7.0	$\times 10^{-6}$	CL=90%
Γ_{99}	$\overline{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	< 2.6	$\times 10^{-5}$	CL=90%
Γ_{100}	$\overline{\Theta}(1540)K_S^0 p \rightarrow K_S^0 \bar{p} K^- \bar{n}$	< 6.0	$\times 10^{-6}$	CL=90%
Γ_{101}	$K_S^0 K_S^0$	< 4.6	$\times 10^{-6}$	

Radiative decays

Γ_{102}	$\gamma \chi_{c0}(1P)$	(9.4 \pm 0.4) %		
Γ_{103}	$\gamma \chi_{c1}(1P)$	(8.8 \pm 0.4) %		
Γ_{104}	$\gamma \chi_{c2}(1P)$	(8.3 \pm 0.4) %		
Γ_{105}	$\gamma \eta_c(1S)$	(3.0 \pm 0.5) $\times 10^{-3}$		
Γ_{106}	$\gamma \eta_c(2S)$	< 2.0 $\times 10^{-3}$	CL=90%	
Γ_{107}	$\gamma \pi^0$	< 5.4 $\times 10^{-3}$	CL=95%	
Γ_{108}	$\gamma \eta'(958)$	(1.36 \pm 0.24) $\times 10^{-4}$		
Γ_{109}	$\gamma f_2(1270)$	(2.1 \pm 0.4) $\times 10^{-4}$		
Γ_{110}	$\gamma f_0(1710)$			
Γ_{111}	$\gamma f_0(1710) \rightarrow \gamma \pi \pi$	(3.0 \pm 1.3) $\times 10^{-5}$		
Γ_{112}	$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	(6.0 \pm 1.6) $\times 10^{-5}$		
Γ_{113}	$\gamma \gamma$	< 1.4 $\times 10^{-4}$	CL=90%	
Γ_{114}	$\gamma \eta$	< 9 $\times 10^{-5}$	CL=90%	
Γ_{115}	$\gamma \eta \pi^+ \pi^-$	(8.7 \pm 2.1) $\times 10^{-4}$		
Γ_{116}	$\gamma \eta(1405)$			
Γ_{117}	$\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi$	< 9 $\times 10^{-5}$	CL=90%	
Γ_{118}	$\gamma \eta(1405) \rightarrow \eta \pi^+ \pi^-$	(3.6 \pm 2.5) $\times 10^{-5}$		
Γ_{119}	$\gamma \eta(1475)$			
Γ_{120}	$\gamma \eta(1475) \rightarrow K \bar{K} \pi$	< 1.4 $\times 10^{-4}$	CL=90%	
Γ_{121}	$\gamma \eta(1475) \rightarrow \eta \pi^+ \pi^-$	< 8.8 $\times 10^{-5}$	CL=90%	
Γ_{122}	$\gamma 2(\pi^+ \pi^-)$	(4.0 \pm 0.6) $\times 10^{-4}$		
Γ_{123}	$\gamma K^{*0} K^+ \pi^- + \text{c.c.}$	(3.7 \pm 0.9) $\times 10^{-4}$		
Γ_{124}	$\gamma K^{*0} \bar{K}^{*0}$	(2.4 \pm 0.7) $\times 10^{-4}$		
Γ_{125}	$\gamma K_S^0 K^+ \pi^- + \text{c.c.}$	(2.6 \pm 0.5) $\times 10^{-4}$		
Γ_{126}	$\gamma K^+ K^- \pi^+ \pi^-$	(1.9 \pm 0.5) $\times 10^{-4}$		
Γ_{127}	$\gamma p \bar{p}$	(2.9 \pm 0.6) $\times 10^{-5}$		
Γ_{128}	$\gamma \pi^+ \pi^- p \bar{p}$	(2.8 \pm 1.4) $\times 10^{-5}$		
Γ_{129}	$\gamma 2(\pi^+ \pi^-) K^+ K^-$	< 2.2 $\times 10^{-4}$	CL=90%	
Γ_{130}	$\gamma 3(\pi^+ \pi^-)$	< 1.7 $\times 10^{-4}$	CL=90%	
Γ_{131}	$\gamma K^+ K^- K^+ K^-$	< 4 $\times 10^{-5}$	CL=90%	

$\psi(2S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$

Γ_1

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
258 ± 26	BAI	02B	BES2 $e^+ e^-$
224 ± 56	LUTH	75	MRK1 $e^+ e^-$

$\Gamma(e^+ e^-)$

Γ_4

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.38 ± 0.04 OUR FIT			
2.33 ± 0.07 OUR AVERAGE			
2.338 ± 0.037 ± 0.096	ABLIKIM	08B	BES2 $e^+ e^- \rightarrow \text{hadrons}$
2.330 ± 0.036 ± 0.110	ABLIKIM	06L	BES2 $e^+ e^- \rightarrow \text{hadrons}$
2.44 ± 0.21	⁸ BAI	02B	BES2 $e^+ e^-$
2.14 ± 0.21	ALEXANDER	89	RVUE See γ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.0 ± 0.3	BRANDELIK	79C	DASP $e^+ e^-$
2.1 ± 0.3	⁹ LUTH	75	MRK1 $e^+ e^-$

⁸ From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$, and hadronic channel, assuming $\Gamma_e = \Gamma_\mu = \Gamma_\tau / 0.38847$.

⁹ From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$, and hadronic channels assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$.

$\Gamma(\gamma\gamma)$

Γ_{113}

<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<43	90	BRANDELIK	79C	DASP $e^+ e^-$

$\psi(2S) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$

This combination of a partial width with the partial width into $e^+ e^-$ and with the total width is obtained from the integrated cross section into channel(i) in the $e^+ e^-$ annihilation. We list only data that have not been used to determine the partial width $\Gamma(i)$ or the branching ratio $\Gamma(i)/\text{total}$.

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$

$\Gamma_1 \Gamma_4 / \Gamma$

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.2 ± 0.4	ABRAMS	75	MRK1 $e^+ e^-$

$\Gamma(\tau^+ \tau^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$

$\Gamma_6 \Gamma_4 / \Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9.0 ± 2.6	79	¹⁰ ANASHIN	07	KEDR $e^+ e^- \rightarrow \psi(2S) \rightarrow \tau^+ \tau^-$
¹⁰ Using $\psi(2S)$ total width of 337 ± 13 keV. Systematic errors not evaluated.				


$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_9\Gamma_4/\Gamma$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.777 ± 0.016 OUR FIT				
0.82 ± 0.04 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
0.852 ± 0.010 ± 0.026	19.5k ± 243	ADAM 06	CLEO 3.773 $e^+e^- \rightarrow \gamma\psi(2S)$	
0.76 ± 0.05 ± 0.01	544 11	AUBERT 05D	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$	
0.68 ± 0.09	12	BAI 98E	BES e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.90 ± 0.08 ± 0.06	256	13 AUBERT 07AU	BABR 10.6 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$	

¹¹ AUBERT 05D reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = 0.0450 \pm 0.0018 \pm 0.0022$ keV. We divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.93 \pm 0.06) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹² The value of $\Gamma(e^+e^-)$ quoted in BAI 98E is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6) \times 10^{-2}$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$. Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

¹³ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0186 \pm 0.0012 \pm 0.0011$ keV. We divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.07 \pm 0.13) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

WEIGHTED AVERAGE
0.82 ± 0.04 (Error scaled by 1.6)



$$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} (\text{keV})$$

$\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{10}\Gamma_4/\Gamma$			
<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.401±0.009 OUR FIT				
0.411±0.008±0.018	$3.6k \pm 96$	ADAM	06	CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$
$\Gamma(J/\psi(1S)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{11}\Gamma_4/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
75.2± 2.1 OUR FIT				
87 ± 9 OUR AVERAGE				
83 ± 25 ± 5	14	¹⁴ AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow J/\psi\pi^+\pi^-\pi^0\gamma$
88 ± 6 ± 7	291 ± 24	ADAM	06	CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$
$^{14}\text{AUBERT } 07\text{AU quotes } \Gamma_{ee}^{\psi(2S)} \cdot \mathcal{B}(\psi(2S) \rightarrow J/\psi\eta) \cdot \mathcal{B}(J/\psi \rightarrow \mu^+\mu^-) \cdot \mathcal{B}(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.11 \pm 0.33 \pm 0.07 \text{ eV.}$				
$\Gamma(J/\psi(1S)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{12}\Gamma_4/\Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<8	90	<37	ADAM	06
				CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$
$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{16}\Gamma_4/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.651±0.029 OUR FIT				
0.59 ±0.05 OUR AVERAGE				
0.579 $\pm 0.038 \pm 0.036$	2.7k	ANDREOTTI	07	E835 $p\bar{p} \rightarrow e^+e^-, J/\psi X$
0.70 $\pm 0.17 \pm 0.03$	22	AUBERT	06B	$e^+e^- \rightarrow p\bar{p}\gamma$
$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{23}\Gamma_4/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.5±0.4±0.1		AUBERT	07BD	BABR $10.6 e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{38}\Gamma_4/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.2±3.3±1.3	43	AUBERT	06D	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
$\Gamma(K^+K^-2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{52}\Gamma_4/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.4±2.1±0.3	26	AUBERT	06D	BABR $10.6 e^+e^- \rightarrow K^+K^-2(\pi^+\pi^-)\gamma$
$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{47}\Gamma_4/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.56±0.42±0.16	85	AUBERT	07AK	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
$\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{80}\Gamma_4/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.346±0.168±0.004	6 ± 3	15	AUBERT	07AK BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

¹⁵ AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)] / \Gamma_{\text{total}} \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.17 \pm 0.08 \pm 0.02$ eV. We divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{79} \Gamma_4 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.57 ± 0.23 ± 0.01	10	16	AUBERT,BE 06D	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

¹⁶ AUBERT,BE 06D reports $[\Gamma(\psi(2S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)] / \Gamma_{\text{total}} \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.28 \pm 0.11 \pm 0.02$ eV. We divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(2(\pi^+ \pi^-)\pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{14} \Gamma_4 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
29.7 ± 2.2 ± 1.8	410	AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)\pi^0 \gamma$

$$\Gamma(\omega \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{43} \Gamma_4 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.01 ± 0.84 ± 0.02	37	17	AUBERT	$10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$

¹⁷ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow \omega \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)] / \Gamma_{\text{total}} \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 2.69 \pm 0.73 \pm 0.16$ eV. We divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(2(\pi^+ \pi^-)\eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{41} \Gamma_4 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.87 ± 1.41 ± 0.01	16	18	AUBERT	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)\eta \gamma$

¹⁸ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+ \pi^-)\eta) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)] / \Gamma_{\text{total}} \times [B(\eta \rightarrow 2\gamma)] = 1.13 \pm 0.55 \pm 0.08$ eV. We divide by our best value $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{59} \Gamma_4 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.4 ± 1.3 ± 0.3	32	AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$

$$\Gamma(K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{50} \Gamma_4 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.05 ± 1.80 ± 0.02	7	19	AUBERT 07AU BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$

¹⁹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)] / \Gamma_{\text{total}} \times [B(\eta \rightarrow 2\gamma)] = 1.2 \pm 0.7 \pm 0.1$ eV. We divide by our best value $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\psi(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.9785±0.0013 OUR AVERAGE				
0.9779±0.0015	20 BAI	02B	BES2 $e^+ e^-$	
0.981 ± 0.003	20 LUTH	75	MRK1 $e^+ e^-$	

20 Includes cascade decay into $J/\psi(1S)$.

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.0173±0.0014 OUR AVERAGE	Error includes scale factor of 1.5.			
0.0166±0.0010	21,22 SETH	04	RVUE $e^+ e^-$	
0.0199±0.0019	21 BAI	02B	BES2 $e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.029 ± 0.004	21 LUTH	75	MRK1 $e^+ e^-$	

21 Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.

22 Using $B(\psi(2S) \rightarrow \ell^+ \ell^-) = (0.73 \pm 0.04)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C.

$\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.169±0.026	23 ADAM	05A	CLEO $e^+ e^- \rightarrow \psi(2S)$	
23 Uses $B(J/\psi X)$ from ADAM 05A, $B(\chi_{cJ}\gamma)$, $B(\eta_c\gamma)$ from ATHAR 04 and $B(\ell^+ \ell^-)$ from PDG 04.				

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
75.2± 1.7 OUR FIT				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
88 ± 13	24 FELDMAN	77	RVUE $e^+ e^-$	

24 From an overall fit assuming equal partial widths for $e^+ e^-$ and $\mu^+ \mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID
75±8 OUR FIT	

$\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_4
0.99±0.11 OUR FIT				
• • • We do not use the following data for averages, fits, limits, etc. • • •				

0.89±0.16 BOYARSKI 75C MRK1 $e^+ e^-$

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})

30 ± 4 OUR FIT

30.8±2.1±3.8

DOCUMENT ID

TECN

COMMENT

²⁵ ABLIKIM 06W BES $e^+e^- \rightarrow \psi(2S)$

²⁵ Computed using PDG 02 value of $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$ to estimate the total number of $\psi(2S)$ events.

Γ_6/Γ

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$

VALUE

EVTS

DOCUMENT ID

TECN

COMMENT

0.574 ± 0.009 OUR FIT

0.592 ± 0.018 OUR AVERAGE

$0.5950 \pm 0.0015 \pm 0.0190$ 151k

ADAM

05A

CLEO

$e^+e^- \rightarrow \psi(2S)$

0.51 ± 0.12

BRANDELIK

79C

DASP

$e^+e^- \rightarrow \mu^+\mu^- X$

0.57 ± 0.08

ABRAMS

75B

MRK1

$e^+e^- \rightarrow \mu^+\mu^- X$

Γ_7/Γ

$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\text{anything})$

VALUE (units 10^{-2})

EVTS

DOCUMENT ID

TECN

COMMENT

1.309±0.026 OUR FIT

1.28 ± 0.04 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

$1.22 \pm 0.02 \pm 0.05$ 5097 ± 73 ²⁶ ANDREOTTI 05 E835 $p\bar{p} \rightarrow \psi(2S) \rightarrow e^+e^-$


$1.28 \pm 0.03 \pm 0.02$ ²⁶ AMBROGIANI 00A E835 $p\bar{p} \rightarrow \psi(2S)$

$1.44 \pm 0.08 \pm 0.02$ ²⁶ ARMSTRONG 97 E760 $\bar{p}p \rightarrow \psi(2S)$

²⁶ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

Γ_4/Γ_7

WEIGHTED AVERAGE
 1.28 ± 0.04 (Error scaled by 1.6)



$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\text{anything})$ (units 10^{-2})

$\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\text{anything})$ Γ_5/Γ_7

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0130±0.0014 OUR FIT			
0.014 ±0.003	HILGER	75	SPEC e^+e^-

 $\Gamma(J/\psi(1S)\text{ neutrals})/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>
0.235±0.004 OUR FIT	

 $\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.326 ±0.005 OUR FIT				
0.323 ±0.013 OUR AVERAGE				
0.323 ±0.014	BAI	02B	BES2	e^+e^-
0.32 ±0.04	ABRAMS	75B	MRK1	$e^+e^- \rightarrow J/\psi\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.3354±0.0014±0.0110 60k ²⁷ ADAM 05A CLEO $e^+e^- \rightarrow \psi(2S)$

²⁷ Not independent from other values reported by ADAM 05A.

 $\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_4/Γ_9

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0230±0.0008 OUR FIT			
0.0252±0.0028±0.0011	²⁸ AUBERT	02B	BABR e^+e^-

²⁸ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

 $\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_5/Γ_9

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0229±0.0026 OUR FIT			
0.0224±0.0029 OUR AVERAGE			

0.0216±0.0026±0.0014 ²⁹ AUBERT 02B BABR e^+e^-

0.0327±0.0077±0.0072 ²⁹ GRIBUSHIN 96 FMPS 515 $\pi^-Be \rightarrow 2\mu X$

²⁹ Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

 $\Gamma(\tau^+\tau^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_6/Γ_9

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.2 ±1.1 OUR FIT			
8.73±1.39±1.57	BAI	02	BES e^+e^-

 $\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$ Γ_9/Γ_7


<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.5680±0.0031 OUR FIT				

0.559 ±0.007 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

0.5637±0.0027±0.0046	60k	ADAM	05A	CLEO	$e^+e^- \rightarrow \psi(2S)$
0.525 ±0.009 ±0.022	4090 ± 67	ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
0.536 ±0.007 ±0.016	20k	^{30,31} ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.496 ±0.037		ARMSTRONG	97	E760	$\bar{p}p \rightarrow \psi(2S)$

³⁰ From a fit to the J/ψ recoil mass spectra.

³¹ ABLIKIM 04B quotes $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)$.



$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$	Γ_8/Γ_9
<u>VALUE</u> 0.721 ± 0.008 OUR FIT	<u>DOCUMENT ID</u> TANENBAUM 76
0.73 ± 0.09	<u>TECN</u> MRK1 e^+e^-

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$	Γ_{10}/Γ
<u>VALUE</u> 0.1684 ± 0.0033 OUR FIT	<u>EVTS</u> 13.4k <u>DOCUMENT ID</u> 32 ADAM <u>TECN</u> CLEO $e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •


$0.1652 \pm 0.0014 \pm 0.0058$ 13.4k 32 ADAM 05A CLEO $e^+e^- \rightarrow \psi(2S)$

³² Not independent from other values reported by ADAM 05A.

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{anything})$	Γ_{10}/Γ_7
<u>VALUE</u> 0.2933 ± 0.0032 OUR FIT	<u>EVTS</u> 13.4k <u>DOCUMENT ID</u> ADAM 05A <u>TECN</u> CLEO $e^+e^- \rightarrow \psi(2S)$

0.284 ± 0.010 OUR AVERAGE Error includes scale factor of 2.3. See the ideogram below.

$0.2776 \pm 0.0025 \pm 0.0043$	13.4k	ADAM	05A	CLEO	$e^+e^- \rightarrow \psi(2S)$
$0.300 \pm 0.008 \pm 0.022$	1655 ± 44	ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
$0.328 \pm 0.013 \pm 0.008$		AMBROGIANI	00A	E835	$p\bar{p} \rightarrow \psi(2S)$
0.323 ± 0.033		ARMSTRONG	97	E760	$\bar{p}p \rightarrow \psi(2S)$



$$\Gamma_{10}/\Gamma_7$$

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.516 ± 0.017 OUR FIT				
0.570 ± 0.009 ± 0.026	14k	³³ ABLIKIM	04B BES	$\psi(2S) \rightarrow J/\psi X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.4924 ± 0.0047 ± 0.0086	73k	^{34,35} ADAM	05A CLEO	$e^+e^- \rightarrow \psi(2S)$
0.571 ± 0.018 ± 0.044		³⁶ ANDREOTTI	05 E835	$\psi(2S) \rightarrow J/\psi X$
0.53 ± 0.06		TANENBAUM	76 MRK1	e^+e^-
0.64 ± 0.15		³⁷ HILGER	75 SPEC	e^+e^-

³³ From a fit to the J/ψ recoil mass spectra.

³⁴ Not independent from other values reported by ADAM 05A.

³⁵ Using 13,217 $J/\psi\pi^0\pi^0$ and 60,010 $J/\psi\pi^+\pi^-$ events.

³⁶ Not independent from other values reported by ANDREOTTI 05.


³⁷ Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0316 ± 0.0007 OUR FIT				
0.0296 ± 0.0031 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.		
0.0298 ± 0.0009 ± 0.0023	5.7k	BAI	04I BES2	$\psi(2S) \rightarrow J/\psi\gamma\gamma$
0.0255 ± 0.0029	386	³⁸ OREGLIA	80 CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
0.045 ± 0.012	17	³⁹ BRANDELIK	79B DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
0.042 ± 0.006	164	³⁹ BARTEL	78B CNTR	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0325 ± 0.0006 ± 0.0011	2.8k	⁴⁰ ADAM	05A CLEO	$e^+e^- \rightarrow \psi(2S)$
0.043 ± 0.008	44	TANENBAUM	76 MRK1	e^+e^-

³⁸ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

³⁹ Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

⁴⁰ Not independent from other values reported by ADAM 05A.



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$

VALUE EVTS

0.0550±0.0011 OUR FIT

0.0548±0.0012 OUR AVERAGE

		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0546±0.0010±0.0007	2.8k	ADAM 05A	CLEO	$e^+ e^- \rightarrow \psi(2S)$
0.050 ± 0.006 ± 0.003	298 ± 20	ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$
0.072 ± 0.009		AMBROGIANI 00A	E835	$p\bar{p} \rightarrow \psi(2S)$
0.061 ± 0.015		ARMSTRONG 97	E760	$\bar{p}p \rightarrow \psi(2S)$

Γ_{11}/Γ_7

$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE EVTS

0.0968±0.0033 OUR FIT

0.096 ± 0.010 OUR AVERAGE

		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.098 ± 0.005 ± 0.010	2k	41 ABLIKIM	04B BES	$\psi(2S) \rightarrow J/\psi X$
0.091 ± 0.021		42 HIMEL	80 MRK2	$e^+ e^- \rightarrow \psi(2S) X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0968±0.0019±0.0013	2.8k	43 ADAM	05A CLEO	$e^+ e^- \rightarrow \psi(2S)$
0.095 ± 0.007 ± 0.007		44 ANDREOTTI	05 E835	$\psi(2S) \rightarrow J/\psi X$

Γ_{11}/Γ_9

⁴¹ From a fit to the J/ψ recoil mass spectra.

⁴² The value for $B(\psi(2S) \rightarrow J/\psi(1s)\eta)$ reported in HIMEL 80 is derived using $B(\psi(2S)) \rightarrow J/\psi(1S)\pi^+\pi^-)$ = $(33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-)$ = 0.138 ± 0.018 . Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-)$ = (0.1181 ± 0.0020) .

⁴³ Not independent from other values reported by ADAM 05A.


⁴⁴ Not independent from other values reported by ANDREOTTI 05.

$$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$$

$$\Gamma_{12}/\Gamma$$

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
12.6±1.3 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
13 ± 1 ± 1	88	ADAM	05A	CLEO $e^+ e^- \rightarrow \psi(2S)$
14.3±1.4±1.2	280	BAI	04i	BES2 $\psi(2S) \rightarrow J/\psi \gamma\gamma$
14 ± 6	7	HIMEL	80	MRK2 $e^+ e^-$
9 ± 2 ± 1	23	45 OREGLIA	80	CBAL $\psi(2S) \rightarrow J/\psi 2\gamma$

⁴⁵ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.



$$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\text{anything})$$

$$\Gamma_{12}/\Gamma_7$$

VALUE (units 10^{-2}) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.22 \pm 0.02 \pm 0.01$ 46 ADAM 05A CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma\gamma$

⁴⁶ Not independent from other values reported by ADAM 05A.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{12}/Γ_9

<u>VALUE</u> (units 10^{-2})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.39 \pm 0.04 \pm 0.01$	⁴⁷ ADAM	05A CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma \gamma$
⁴⁷ Not independent from other values reported by ADAM 05A.			

 HADRONIC DECAYS


 $\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
35 ± 16	6	FRANKLIN	83	MRK2 $e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
29 ± 10 OUR AVERAGE				Error includes scale factor of 4.6. See the ideogram below.
24.9 ± 0.7 ± 3.6	2173	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$
125 ± 12 ± 2	410	⁴⁸ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+\pi^-)\pi^0 \gamma$
26.1 ± 0.7 ± 3.0	1703	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
30 ± 8	42	FRANKLIN	83 MRK2	$e^+ e^-$
48 AUBERT 07AU reports $[B(\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0)] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (297 \pm 22 \pm 18) \times 10^{-4} \text{ keV}$. We divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.38 \pm 0.04 \text{ keV}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

WEIGHTED AVERAGE
 29 ± 10 (Error scaled by 4.6)




$\Gamma(\rho a_2(1320))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.55±0.73±0.47		112 ± 31	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.3	90		BAI	98J BES	$e^+ e^-$

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.74±0.12 OUR FIT					
2.95±0.23 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.			
3.36±0.09±0.25	1618	ABLIKIM	07C BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$	
2.87±0.12±0.15	557	PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$	
1.4 ± 0.8	4	BRANDELIK	79C DASP	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$	
2.3 ± 0.7		FELDMAN	77 MRK1	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$	



$\Gamma(p\bar{p})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.4 ± 0.4 OUR FIT			
6.98±0.49±0.97	BAI	01 BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$

$\Gamma(\Delta^{++}\bar{\Delta}^{--})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
12.8±1.0±3.4	157	49 BAI	01 BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

⁴⁹ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

Γ_{15}/Γ

Γ_{16}/Γ

Γ_{16}/Γ_9

Γ_{17}/Γ

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{18}/Γ
<1.2	90	50 ABLIKIM	07H BES2	$e^+ e^- \rightarrow \psi(2S)$	

50 Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

 $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{19}/Γ
<0.49	90	51 ABLIKIM	07H BES2	$e^+ e^- \rightarrow \psi(2S)$	

51 Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

 $\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{20}/Γ
1.0±0.1 ±0.1	74.0	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^-$	

 $\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{21}/Γ
1.8±0.3±0.3	45.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^+\pi^-\pi^-$	

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{22}/Γ
2.8±0.4±0.5	73.4	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}2(\pi^+\pi^-)$	

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$


<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{23}/Γ
2.8 ± 0.5 OUR AVERAGE					Error includes scale factor of 2.6. See the ideogram below.	
3.39±0.20±0.32		337	ABLIKIM	07C BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
6.3 ± 1.7 ± 0.1		52 AUBERT	07BD BABR	10.6 e ⁺ e ⁻	$\rightarrow \Lambda\bar{\Lambda}\gamma$	
3.28±0.23±0.25		208	PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
1.81±0.20±0.27		80	53 BAI	01 BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4 90 FELDMAN 77 MRK1 e⁺ e⁻ $\rightarrow \psi(2S) \rightarrow \text{hadrons}$

52 AUBERT 07BD reports $[B(\psi(2S) \rightarrow \Lambda\bar{\Lambda})] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (15 \pm 4 \pm 1) \times 10^{-4}$ keV. We divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.38 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

53 Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.



$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
25.7±4.4±6.8	35	PEDLAR	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons


Γ_{24}/Γ

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
22 ±4 OUR AVERAGE				Error includes scale factor of 1.5. See the ideogram below.
23.5±3.6±3.2	59	ABLIKIM	07C BES	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
26.3±3.5±2.1	58	PEDLAR	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
12 ±4 ±4	8	54 BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

Γ_{25}/Γ

⁵⁴ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.



VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
11±3±3	14	55 BAI	01 BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons


⁵⁵ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
18 ± 6 OUR AVERAGE			Error includes scale factor of 2.8. See the ideogram below.		
30.3±4.0±3.2	67	ABLIKIM	07C BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons	
23.8±3.0±2.1	63	PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<20 90 FELDMAN 77 MRK1 $e^+ e^- \rightarrow \psi(2S) \rightarrow$
hadrons

⁵⁶ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.



$\Gamma(\Xi^0\Xi^0)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
27.5\pm6.4\pm6.1	19	PEDLAR	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

$\Gamma(\Xi(1530)^0\Xi(1530)^0)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 8.1	90	57 BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<32	90	PEDLAR	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
57	Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.			

$\Gamma(\Omega^-\bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 7.3	90	58 BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<16	90	PEDLAR	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
58	Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.			

$\Gamma(\pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.33\pm0.17 OUR AVERAGE				
1.32 \pm 0.10 \pm 0.15	256 \pm 18	59 ABLIKIM	05E BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$
1.4 \pm 0.5	9	FRANKLIN	83 MRK2	e^+e^-

⁵⁹ Computed using $B(\pi^0 \rightarrow \gamma\gamma) = (98.80 \pm 0.03)\%$.

$\Gamma(\eta p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.60±0.12 OUR AVERAGE				
0.58±0.11±0.07	44.8 ± 8.5	⁶⁰ ABLIKIM	05E BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$
0.8 ± 0.3 ± 0.3	9.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

⁶⁰ Computed using $B(\eta \rightarrow \gamma\gamma) = (39.43 \pm 0.26)\%$.

$\Gamma(\omega p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.69±0.21 OUR AVERAGE				
0.6 ± 0.2 ± 0.2	21.2	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$
0.8 ± 0.3 ± 0.1	14.9 ± 0.1	⁶¹ BAI	03B BES	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

⁶¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(\phi p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.24	90	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.26 90 ⁶² BAI 03B BES $\psi(2S) \rightarrow K^+K^-p\bar{p}$

⁶² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(\pi^+\pi^-p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0±0.4 OUR AVERAGE				
5.9±0.2±0.4	904.5	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-$
8 ± 2		⁶³ TANENBAUM	78 MRK1	$e^+ e^- \rightarrow$

⁶³ Assuming entirely strong decay.

$\Gamma(p\bar{n}\pi^- \text{ or c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.48±0.17 OUR AVERAGE				
2.45±0.11±0.21	851	ABLIKIM	06I BES2	$e^+ e^- \rightarrow p\pi^-X$
2.52±0.12±0.22	849	ABLIKIM	06I BES2	$e^+ e^- \rightarrow \bar{p}\pi^+X$

$\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.18±0.50±0.50	135 ± 21	ABLIKIM	06I BES2	$e^+ e^- \rightarrow p\pi^-\pi^0X$

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(\eta\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{40}/Γ

<i>VALUE</i> (units 10^{-4})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
9.5±0.7±1.5		64 BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadr
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10.3±0.8±1.4	201.7	65 BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $\eta 3\pi (\eta \rightarrow \gamma\gamma)$
8.1±1.4±1.6	50.0	65 BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $\eta 3\pi (\eta \rightarrow 3\pi)$

64 Average of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi$.

65 Not independent from other values reported by BRIERE 05.

 $\Gamma(2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$ Γ_{41}/Γ

<i>VALUE</i> (units 10^{-3})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
1.2±0.6±0.1	16	66 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$
66 AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+\pi^-)\eta) \cdot B(\eta \rightarrow \gamma\gamma) = 1.2 \pm 0.7 \pm 0.1 \text{ eV}$.				


 $\Gamma(\eta'\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{42}/Γ

<i>VALUE</i> (units 10^{-4})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
4.5±1.6±1.3	12.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadr

 $\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{43}/Γ

<i>VALUE</i> (units 10^{-4})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
7.3±1.2 OUR AVERAGE		Error includes scale factor of 2.1. See the ideogram below.		
8.4±0.5±1.2	386	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
12.2±2.2±0.7	37	67 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
8.2±0.5±0.7	391	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $2(\pi^+\pi^-)\pi^0$
4.8±0.6±0.7	100 ± 22	68 BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$

67 AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16 \text{ eV}$.68 Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.



$\Gamma(b_1^\pm\pi^\mp)/\Gamma_{\text{total}}$

Γ_{44}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ±0.6 OUR AVERAGE				Error includes scale factor of 1.1.
5.1 ± 0.6 ± 0.8	202	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
4.18 ^{+0.43} _{-0.42} ± 0.92	170	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
3.2 ± 0.6 ± 0.5	61 ± 11	69,70 BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.2 ± 0.8 ± 1.0	69 BAI		99C BES	Repl. by BAI 03B
69 Assuming $B(b_1 \rightarrow \omega\pi)=1$.				
70 Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.				

$\Gamma(b_1^0\pi^0)/\Gamma_{\text{total}}$

Γ_{45}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.35^{+0.47}_{-0.42} ± 0.40	45	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$

Γ_{46}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ±0.4 OUR AVERAGE					
2.3 ± 0.5 ± 0.4		57	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
2.05 ± 0.41 ± 0.38		62 ± 12	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.5	90	71 BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$	
<1.7	90	BAI	98J BES	Repl. by BAI 03B	
71 Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.					

$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{47}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.5±0.9 OUR AVERAGE		Error includes scale factor of 1.9.		
10.8±1.9±0.2	85	72 AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^- K^+K^- \gamma$
7.1±0.3±0.4	817.2	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow K^+K^-\pi^+\pi^-$
16 ±4		73 TANENBAUM	78 MRK1	e^+e^-
72 AUBERT 07AK reports $[B(\psi(2S) \rightarrow \pi^+\pi^- K^+K^-)] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3} \text{ keV}$. We divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.38 \pm 0.04 \text{ keV}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
73 Assuming entirely strong decay.				

 $\Gamma(\rho^0 K^+K^-)/\Gamma_{\text{total}}$ Γ_{48}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.2±0.2±0.4	223.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow K^+K^-\pi^+\pi^-$

 $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{49}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.86±0.32±0.43		93 ± 16	BAI	04C	$\psi(2S) \rightarrow K^+K^-\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.2		90	BAI	98J BES	e^+e^-

 $\Gamma(K^+K^-\pi^+\pi^-\eta)/\Gamma_{\text{total}}$ Γ_{50}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.3±0.7±0.1	7	74 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\eta\gamma$
74 AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+\pi)\eta) \cdot B(\eta \rightarrow \gamma\gamma) = 1.2 \pm 0.7 \pm 0.1 \text{ eV}$.				

 $\Gamma(K_1(1270)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{53}/Γ

<u>VALUE (units 10^{-4})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.0±1.8±2.1		75 BAI	99C BES	e^+e^-

75 Assuming $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$ $\Gamma(K_S^0 K_S^0 \pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{54}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.20±0.25±0.37	83 ± 9	ABLIKIM	050 BES2	$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\rho^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{55}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.5±0.1 ±0.2	61.1	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-$

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$	Γ_{56}/Γ		
<i>VALUE</i> (units 10^{-4})	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
6.7 \pm 2.5	TANENBAUM 78	MRK1	$e^+ e^-$

$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$	Γ_{57}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±0.6 OUR AVERAGE	Error includes scale factor of 2.2.			
2.2±0.2±0.2	308	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$
4.5±1.0		TANENBAUM	78	MRK1 $e^+ e^-$

$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_{58}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2±0.6 OUR AVERAGE	Error includes scale factor of 1.4.			
2.0±0.2±0.4	285.5	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$
4.2±1.5		TANENBAUM	78	MRK1 $e^+ e^-$

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$					Γ_{59} / Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
12.6 ± 0.9 OUR AVERAGE					
$18.5 \pm 5.6 \pm 0.3$	32	⁷⁶ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$	
$11.7 \pm 1.0 \pm 1.5$	597	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
$12.7 \pm 0.5 \pm 1.0$	711.6	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	

$\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$	Γ_{51}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
10.0 \pm 2.5 \pm 1.8	65	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}$	Γ_{60}/Γ			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
5.9±2.0±0.9	19	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{61}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6±1.3±1.8	238	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{62}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.6 \pm 2.2 \pm 1.7	133	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{63}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.3±2.2±1.4	78	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{64}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.1±1.3±1.2	125	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(\eta K^+ K^-)/\Gamma_{\text{total}}$ Γ_{65}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.3	90	BRIERE	05	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{66}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.85±0.25 OUR AVERAGE				Error includes scale factor of 1.1.
2.38±0.37±0.29	78	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.9 ± 0.3 ± 0.3	76.8	BRIERE	05	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.5 ± 0.3 ± 0.2	23.0 ± 5.2	77 BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

77 Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

 $\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{67}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.5 ± 2.0 OUR AVERAGE				Error includes scale factor of 2.8.
5.45±0.42±0.87	671	ABLIKIM	05H BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+ \pi^-)$
1.5 ± 1.0		78 TANENBAUM	78 MRK1	$e^+ e^-$

78 Assuming entirely strong decay.

 $\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{68}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.3±0.4±0.6	434.9	BRIERE	05	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{69}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.3±0.7 OUR AVERAGE				
6.3±0.6±0.3		DOBBS	06A CLEO	$e^+ e^-$
10 ± 7		BRANDELIK	79C DASP	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 5	90	FELDMAN	77 MRK1	$e^+ e^-$

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5}) EVTS

5.4 ± 0.5 OUR AVERAGE

$5.8 \pm 0.8 \pm 0.4$

$5.24 \pm 0.47 \pm 0.48$ 156 ± 14

DOCUMENT ID

79 DOBBS
BAI

TECN

06A CLEO

04B BES2 $\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$

COMMENT

Γ_{70}/Γ



⁷⁹ Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6860 \pm 0.0027$.

$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4}) EVTS

1.68 ± 0.26 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.


$1.81 \pm 0.18 \pm 0.19$ 260 ± 19 80 ABLIKIM

$1.88^{+0.16}_{-0.15} \pm 0.28$ 194 ADAM

0.85 ± 0.46 4 FRANKLIN

⁸⁰ From a PW analysis of $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$.

Γ_{71}/Γ



$\Gamma(\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

Γ_{72}/Γ

VALUE (units 10^{-4})

DOCUMENT ID

TECN

COMMENT

1.94 ± 0.25 ± 1.15 81 ABLIKIM 05J BES2 $\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$

⁸¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$.

$\Gamma(\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{73}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.32±0.12 OUR AVERAGE	Error includes scale factor of 1.8.				
0.51±0.07±0.11		82	ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$
0.24 $^{+0.08}_{-0.07}$ ±0.02	22	ADAM	05	CLEO	$e^+e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.83	90	1	FRANKLIN	83	MRK2 e^+e^-
<10	90		BARTEL	76	CNTR e^+e^-
<10	90		83 ABRAMS	75	MRK1 e^+e^-

82 From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.83 Final state $\rho^0\pi^0$. $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{74}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
8 ±5		BRANDELIK	79C DASP	e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.1	90	DOBBS	06A CLEO	$e^+e^- \rightarrow \psi(2S)$	
<5	90	FELDMAN	77	MRK1 e^+e^-	

 $\Gamma(K_1(1400)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{75}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.1	90	84 BAI	99C BES	e^+e^-

84 Assuming $B(K_1(1400) \rightarrow K^*\pi) = 0.94 \pm 0.06$ $\Gamma(K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{76}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.96	90	1	FRANKLIN	83	MRK2 $e^+e^- \rightarrow \text{hadrons}$

 $\Gamma(K^+\bar{K}^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{77}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.7$^{+0.8}_{-0.7}$ OUR AVERAGE					
2.9 $^{+1.3}_{-1.7}$ ±0.4		9.6 ± 4.2	ABLIKIM	05I BES2	$e^+e^- \rightarrow \psi(2S)$
1.3 $^{+1.0}_{-0.7}$ ±0.3		7	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<5.4	90		FRANKLIN	83	MRK2 $e^+e^- \rightarrow \text{hadrons}$

 $\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{78}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.9±2.0 OUR AVERAGE				
13.3 $^{+2.4}_{-2.8}$ ±1.7	65.6 ± 9.0	ABLIKIM	05I BES2	$e^+e^- \rightarrow \psi(2S)$
9.2 $^{+2.7}_{-2.2}$ ±0.9	25	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$

$\Gamma(K^+ \bar{K}^*(892)^- + \text{c.c.})/\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})$ Γ_{77}/Γ_{78}

VALUE		DOCUMENT ID	TECN	COMMENT
0.16±0.06 OUR AVERAGE				
0.22 ^{+0.10} _{-0.14}		ABLIKIM	05I	BES2 $e^+ e^- \rightarrow \psi(2S)$
0.14 ^{+0.08} _{-0.06}		ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.17±0.29 OUR AVERAGE Error includes scale factor of 1.7.				
2.39 $\pm 0.94 \pm 0.04$	10 \pm 4	85,86 AUBERT	07AK BABR	$10.6 \frac{e^+ e^-}{\pi^+ \pi^- K^+ K^- \gamma}$
0.9 ± 0.2 ± 0.1	47.6	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
1.5 ± 0.2 ± 0.2	51.5 \pm 8.3	87 BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
85 AUBERT 07AK reports $[B(\psi(2S) \rightarrow \phi \pi^+ \pi^-)] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$ keV. We divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.38 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
86 Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.				
87 Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.				

 $\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.68±0.24 OUR AVERAGE Error includes scale factor of 1.1.				
1.43 $\pm 0.69 \pm 0.02$	6 \pm 3	88,89 AUBERT	07AK BABR	$10.6 \frac{e^+ e^-}{\pi^+ \pi^- K^+ K^- \gamma}$
0.6 ± 0.2 ± 0.1	18.4 \pm 6.4	90 BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
88 AUBERT 07AK reports $[B(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-)] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.34 \pm 0.16 \pm 0.04) \times 10^{-3}$ keV. We divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.38 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
89 Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.				
90 Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.				

 $\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.6±0.1 ±0.1				
59.2		BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.70±0.16 OUR AVERAGE				
0.8 ± 0.2 ± 0.1	36.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$
0.6 ± 0.2 ± 0.1	16.1 \pm 5.0	91 BAI	03B BES	$\psi(2S) \rightarrow 2(K^+ K^-)$
91 Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.				

$\Gamma(2(K^+K^-)\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>
$1.1 \pm 0.2 \pm 0.2$	44.7

 Γ_{83}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(K^+K^-)\pi^0$

 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>
$2.8^{+1.0}_{-0.8}$ OUR AVERAGE	

 Γ_{84}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ADAM 05	CLEO	$e^+e^- \rightarrow \psi(2S)$
ABLIKIM 04K BES		$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\phi\eta')/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>
$3.1 \pm 1.4 \pm 0.7$	8

 Γ_{85}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
92 ABLIKIM 04K BES		$e^+e^- \rightarrow \psi(2S)$

92 Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels. $\Gamma(\omega\eta')/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>
$3.2^{+2.4}_{-2.0} \pm 0.7$	4

 Γ_{86}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
93 ABLIKIM 04K BES		$e^+e^- \rightarrow \psi(2S)$

93 Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels. $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>
2.1 ± 0.6 OUR AVERAGE	

 Γ_{87}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ADAM 05	CLEO	$e^+e^- \rightarrow \psi(2S)$
ABLIKIM 04L BES		$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\rho\eta')/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>
$1.87^{+1.64}_{-1.11} \pm 0.33$	2

 Γ_{88}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ABLIKIM 04L BES		$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\rho\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>
2.2 ± 0.6 OUR AVERAGE	

 Γ_{89}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
Error includes scale factor of 1.1.		

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>
3.0 $^{+1.1}_{-0.9} \pm 0.2$	18
1.78 $^{+0.67}_{-0.62} \pm 0.17$	13

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ADAM 05	CLEO	$e^+e^- \rightarrow \psi(2S)$
ABLIKIM 04L BES		$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>
<1.1	90

 Γ_{90}/Γ

<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
ADAM 05	CLEO	$e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.1	90	ABLIKIM 04K BES	$e^+e^- \rightarrow \psi(2S)$
------	----	-----------------	-------------------------------

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{91}/Γ
<0.4	90	ABLIKIM	04K	BES $e^+ e^- \rightarrow \psi(2S)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.7	90	ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$	

 $\Gamma(\eta_c\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{92}/Γ
<1.0	90	PEDLAR	07	CLEO $e^+ e^- \rightarrow \psi(2S)$	

 $\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{93}/Γ
$2.7 \pm 0.6 \pm 0.4$	30.1	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$	

 $\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{94}/Γ
$0.81 \pm 0.11 \pm 0.14$	50	94 ABLIKIM	08C	BES2 $e^+ e^- \rightarrow J/\psi$	

⁹⁴ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

 $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{95}/Γ
$0.44 \pm 0.12 \pm 0.11$		20 ± 6	BAI	04C	$\psi(2S) \rightarrow 2(K^+K^-)$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
<0.45	90		BAI	98J	BES $e^+ e^- \rightarrow 2(K^+K^-)$	

 $\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{96}/Γ
<0.88	90	BAI	04G	BES2 $e^+ e^-$	

 $\Gamma(\Theta(1540)K^-\bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{97}/Γ
<1.0	90	BAI	04G	BES2 $e^+ e^-$	

 $\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{98}/Γ
<0.70	90	BAI	04G	BES2 $e^+ e^-$	

 $\Gamma(\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{99}/Γ
<2.6	90	BAI	04G	BES2 $e^+ e^-$	

 $\Gamma(\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{100}/Γ
<0.60	90	BAI	04G	BES2 $e^+ e^-$	

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{101}/Γ VALUE (units 10^{-4})**<0.046**DOCUMENT ID

95 BAI

TECN

04D

COMMENT

BES

 $e^+ e^-$

95 Forbidden by CP.

RADIATIVE DECAYS $\Gamma(\gamma \chi_{c0}(1P))/\Gamma_{\text{total}}$ Γ_{102}/Γ VALUE (units 10^{-2})**9.4 ± 0.4 OUR FIT****9.2 ± 0.4 OUR AVERAGE**

9.22 ± 0.11 ± 0.46 72600

9.9 ± 0.5 ± 0.8

7.2 ± 2.3

7.5 ± 2.6

EVTSDOCUMENT IDTECNCOMMENTATHAR 04 CLEO $e^+ e^- \rightarrow \gamma X$ 96 GAISER 86 CBAL $e^+ e^- \rightarrow \gamma X$ 96 BIDDICK 77 CNTR $e^+ e^- \rightarrow \gamma X$ 96 WHITAKER 76 MRK1 $e^+ e^-$ 96 Angular distribution ($1+\cos^2\theta$) assumed. $\Gamma(\gamma \chi_{c1}(1P))/\Gamma_{\text{total}}$ Γ_{103}/Γ VALUE (units 10^{-2})**8.8 ± 0.4 OUR FIT****8.9 ± 0.5 OUR AVERAGE**

9.07 ± 0.11 ± 0.54 76700

9.0 ± 0.5 ± 0.7

7.1 ± 1.9

EVTSDOCUMENT IDTECNCOMMENTATHAR 04 CLEO $e^+ e^- \rightarrow \gamma X$ 97 GAISER 86 CBAL $e^+ e^- \rightarrow \gamma X$ 98 BIDDICK 77 CNTR $e^+ e^- \rightarrow \gamma X$ 97 Angular distribution ($1-0.189 \cos^2\theta$) assumed.

98 Valid for isotropic distribution of the photon.

 $\Gamma(\gamma \chi_{c2}(1P))/\Gamma_{\text{total}}$ Γ_{104}/Γ VALUE (units 10^{-2})**8.3 ± 0.4 OUR FIT****8.8 ± 0.5 OUR AVERAGE** Error includes scale factor of 1.1.

9.33 ± 0.14 ± 0.61 79300

8.0 ± 0.5 ± 0.7

7.0 ± 2.0

EVTSDOCUMENT IDTECNCOMMENTATHAR 04 CLEO $e^+ e^- \rightarrow \gamma X$ 99 GAISER 86 CBAL $e^+ e^- \rightarrow \gamma X$ 100 BIDDICK 77 CNTR $e^+ e^- \rightarrow \gamma X$ 99 Angular distribution ($1-0.052 \cos^2\theta$) assumed.

100 Valid for isotropic distribution of the photon.

 $[\Gamma(\gamma \chi_{c0}(1P)) + \Gamma(\gamma \chi_{c1}(1P)) + \Gamma(\gamma \chi_{c2}(1P))] / \Gamma_{\text{total}} \quad (\Gamma_{102} + \Gamma_{103} + \Gamma_{104}) / \Gamma$ VALUEDOCUMENT IDTECNCOMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

27.6 ± 0.3 ± 2.0 101 ATHAR 04 CLEO $e^+ e^- \rightarrow \gamma X$ 101 Not independent from ATHAR 04 measurements of $B(\gamma \chi_{cJ})$. $\Gamma(\gamma \chi_{c0}(1P))/\Gamma(\gamma \chi_{c1}(1P))$ $\Gamma_{102}/\Gamma_{103}$ VALUEDOCUMENT IDTECNCOMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.02 ± 0.01 ± 0.07 102 ATHAR 04 CLEO $e^+ e^- \rightarrow \gamma X$ 102 Not independent from ATHAR 04 measurements of $B(\gamma \chi_{cJ})$.

$\Gamma(\gamma\chi_{c2}(1P))/\Gamma(\gamma\chi_{c1}(1P))$

$\Gamma_{104}/\Gamma_{103}$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.03 \pm 0.02 \pm 0.03$	103 ATHAR	04 CLEO	$e^+e^- \rightarrow \gamma X$
103 Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.			

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c2}(1P))$

$\Gamma_{102}/\Gamma_{104}$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.99 \pm 0.02 \pm 0.08$	104 ATHAR	04 CLEO	$e^+e^- \rightarrow \gamma X$
104 Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.			

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$

Γ_{105}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.30 ± 0.05 OUR AVERAGE				
$0.32 \pm 0.04 \pm 0.06$	2560	105 ATHAR	04 CLEO	$e^+e^- \rightarrow \gamma X$
0.28 ± 0.06		106 GAISER	86 CBAL	$e^+e^- \rightarrow \gamma X$
105 ATHAR 04 used $\Gamma_{\eta_c(1S)} = 24.8 \pm 4.9$ MeV to obtain this result.				
106 GAISER 86 used $\Gamma_{\eta_c(1S)} = 11.5 \pm 4.5$ MeV to obtain this result.				

$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$

Γ_{106}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<0.20	90	ATHAR	04 CLEO	$e^+e^- \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.2 to 1.3	95	EDWARDS	82C CBAL	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$

Γ_{107}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 54	95	107 LIBERMAN	75 SPEC	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<100	90	WIIK	75 DASP	e^+e^-
107 Restated by us using $B(\psi(2S) \rightarrow \mu^+ \mu^-) = 0.0077$.				

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$

Γ_{108}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.36 ± 0.24 OUR AVERAGE					
1.24 ± 0.27 ± 0.15	23		ABLIKIM	06R BES2	$e^+e^- \rightarrow \psi(2S)$
1.54 ± 0.31 ± 0.20	~ 43		BAI	98F BES	$\psi(2S) \rightarrow \pi^+\pi^- 2\gamma, \pi^+\pi^- 3\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

< 60	90	108 BRAUNSCH...	77 DASP	e^+e^-
< 11	90	109 BARTEL	76 CNTR	e^+e^-

108 Restated by us using total decay width 228 keV.

109 The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{109}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.12±0.19±0.32	110,111	BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.08±0.19±0.33	200.6 ± 18.8	¹¹⁰ BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
2.90±1.08±1.07	29.9 ± 11.1	¹¹⁰ BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$

¹¹⁰ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

¹¹¹ Combining the results from $\pi^+\pi^-$ and $\pi^0\pi^0$ decay modes.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{111}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.301±0.041±0.124	35.6 ± 4.8	¹¹² BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$

¹¹² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{112}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.604±0.090±0.132		39.6 ± 5.9	^{113,114} BAI	03C BES	$\psi(2S) \rightarrow \gamma K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.56	90	6.8 ± 3.1	^{113,114} BAI	03C BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
--------	----	---------------	------------------------	---------	---

¹¹³ Includes unknown branching fractions to K^+K^- or $K_S^0 K_S^0$. We have multiplied the K^+K^- result by a factor of 2 and the $K_S^0 K_S^0$ result by a factor of 4 to obtain the $K\bar{K}$ result.

¹¹⁴ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

 $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$ Γ_{114}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.9	90	BAI	98F	BES	$\psi(2S) \rightarrow \pi^+\pi^- 3\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2	90	YAMADA	77	DASP	$e^+e^- \rightarrow 3\gamma$
----	----	--------	----	------	------------------------------

 $\Gamma(\gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{115}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.71±1.25±1.64	418	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{117}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.9	90	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma K_S^0 K^+\pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3	90	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma K^+K^-\pi^0$
------	----	---------	-----	--

<1.2	90	¹¹⁵ SCHARRE	80	MRK1 $e^+e^- \rightarrow K\bar{K}\pi$
------	----	------------------------	----	---------------------------------------

¹¹⁵ Includes unknown branching fraction $\eta(1405) \rightarrow K\bar{K}\pi$.

 $\Gamma(\gamma\eta(1405) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{118}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.36±0.25±0.05	10	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma\eta(1475) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{120}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.4	90	ABLIKIM	06R	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.5	90	ABLIKIM	06R	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

 $\Gamma(\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{121}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.88	90	ABLIKIM	06R	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

 $\Gamma(\gamma 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{122}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
39.6 ± 2.8 ± 5.0	583	ABLIKIM	07D	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^*0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{123}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
37.0 ± 6.1 ± 7.2	237	ABLIKIM	07D	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^*0 \bar{K}^*)/\Gamma_{\text{total}}$ Γ_{124}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
24.0 ± 4.5 ± 5.0	41	ABLIKIM	07D	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{125}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
25.6 ± 3.6 ± 3.6	115	ABLIKIM	07D	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{126}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
19.1 ± 2.7 ± 4.3	132	ABLIKIM	07D	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{127}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.9 ± 0.4 ± 0.4	142	ABLIKIM	07D	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma\pi^+\pi^- p\bar{p})/\Gamma_{\text{total}}$ Γ_{128}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.8 ± 1.2 ± 0.7	17	ABLIKIM	07D	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma 2(\pi^+\pi^-) K^+ K^-)/\Gamma_{\text{total}}$ Γ_{129}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<22	90	ABLIKIM	07D	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma 3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{130}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<17	90	ABLIKIM	07D	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\gamma K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{131}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4	90	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\psi(2S)$ CROSS-PARTICLE BRANCHING RATIOS

For measurements involving $B(\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$
see the corresponding entries in the $\chi_{cJ}(1P)$ sections.

 $\psi(2S)$ REFERENCES

ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07C	PL B648 149	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07D	PRL 99 011802	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ANASHIN	07	JETPL 85 347	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
		Translated from ZETFP 85 429.		
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Fermilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
PEDLAR	07	PR D75 011102R	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	06G	PR D73 052004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06W	PR D74 112003	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	06A	PR D74 011105R	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05E	PR D71 072006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05I	PL B614 37	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05J	PL B619 247	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05	PRL 94 012005	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05	PR D71 032006	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRIERE	05	PRL 95 062001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
PEDLAR	05	PR D72 051108R	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04K	PR D70 112003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04L	PR D70 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04B	PRL 92 052001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04C	PR D69 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03B	PR D67 052002	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)
AUBERT	02B	PR D65 031101R	B. Aubert <i>et al.</i>	(BaBar Collab.)
BAI	02	PR D65 052004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	02B	PL B550 24	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	
BAI	01	PR D63 032002	J.Z. Bai <i>et al.</i>	(BES Collab.)
AMBROGIANI	00A	PR D62 032004	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)

ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98F	PR D58 097101	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98J	PRL 81 5080	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARMSTRONG	97	PR D55 1153	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 Collab., E706 Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
		Translated from YAF 41 733.		
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
OREGLIA	80	PRL 45 959	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34 1471.		
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
BRAUNSCH...	77	PL 67B 249	W. Braunschweig <i>et al.</i>	(DASP Collab.)
BURMESTER	77	PL 66B 395	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	76	PRL 36 402	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL) IG
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)
ABRAMS	75	Stanford Symp. 25	G.S. Abrams	(LBL)
ABRAMS	75B	PRL 34 1181	G.S. Abrams <i>et al.</i>	(LBL, SLAC)
BOYARSKI	75C	Palermo Conf. 54	A.M. Boyarski <i>et al.</i>	(SLAC, LBL)
HILGER	75	PRL 35 625	E. Hilger <i>et al.</i>	(STAN, PENN)
LIBERMAN	75	Stanford Symp. 55	A.D. Liberman	(STAN)
LUTH	75	PRL 35 1124	V. Luth <i>et al.</i>	(SLAC, LBL) JPC
WIJK	75	Stanford Symp. 69	B.H. Wiik	(DESY)

OTHER RELATED PAPERS

AUBERT,BE	06F	PR D74 111103R	B. Aubert <i>et al.</i>	(BABAR Collab.)
AMBROGIANI	05	PL B610 177	M. Ambrogiani <i>et al.</i>	(FNAL E853 Collab.)
GUO	05	NP A761 269	F.-K. Guo <i>et al.</i>	
VOLOSHIN	05	PR D71 114003	M.B. Voloshin	
ABLIKIM	04I	PR D70 092004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04J	PRL 93 112002	M. Ablikim <i>et al.</i>	(BES Collab.)
LIU	04B	PR D70 094001	K.-Y. Liu, K.-T. Chao	
WANG	04C	PR D70 077505	P. Wang, X.H. Mo, C.Z. Yuan	
BAI	00E	PR D62 032002	J. Bai <i>et al.</i>	(BES Collab.)
CHEN	98	PRL 80 5060	Y.Q. Chen, E. Braaten	
SUZUKI	98	PR D57 5717	M. Suzuki	
BARATE	83	PL 121B 449	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, IND)
AUBERT	75B	PRL 33 1624	J.J. Aubert <i>et al.</i>	(MIT, BNL)
BRAUNSCH...	75B	PL 57B 407	W. Braunschweig <i>et al.</i>	(DASP Collab.)
CAMERINI	75	PRL 35 483	U. Camerini <i>et al.</i>	(WISC, SLAC)
FELDMAN	75B	PRL 35 821	G.J. Feldman <i>et al.</i>	(LBL, SLAC)
GRECO	75	PL 56B 367	M. Greco, G. Pancheri-Srivastava, Y. Srivastava	
JACKSON	75	NIM 128 13	J.D. Jackson, D.L. Scharre	(LBL)
SIMPSON	75	PRL 35 699	J.W. Simpson <i>et al.</i>	(STAN, PENN)
ABRAMS	74	PRL 33 1453	G.S. Abrams <i>et al.</i>	(LBL, SLAC)
