

## THE $\eta(1405)$ , $\eta(1475)$ , $f_1(1420)$ , AND $f_1(1510)$

Revised January 2008 by C. Amsler (Zürich) and A. Masoni (INFN Cagliari).

The first observation of the  $\eta(1440)$  was made in  $p\bar{p}$  annihilation at rest into  $\eta(1440)\pi^+\pi^-$ ,  $\eta(1440) \rightarrow K\bar{K}\pi$  (BAILLON 67). This state was reported to decay through  $a_0(980)\pi$  and  $K^*(892)\bar{K}$  with roughly equal contributions. The  $\eta(1440)$  was also observed in radiative  $J/\psi(1S)$  decay to  $K\bar{K}\pi$  (SCHARRE 80, EDWARDS 82E, AUGUSTIN 90) and  $\gamma\rho$  (BAI 04J). There is now evidence for the existence of two pseudoscalars in this mass region, the  $\eta(1405)$  and  $\eta(1475)$ . The former decays mainly through  $a_0(980)\pi$  (or direct  $K\bar{K}\pi$ ) and the latter mainly to  $K^*(892)\bar{K}$ .

The simultaneous observation of two pseudoscalars is reported in three production mechanisms:  $\pi^-p$  (RATH 89, ADAMS 01); radiative  $J/\psi(1S)$  decay (BAI 90C, AUGUSTIN 92); and  $\bar{p}p$  annihilation at rest (BERTIN 95, BERTIN 97, CICALO 99, NICHTIU 02). All of them give values for the masses, widths, and decay modes in reasonable agreement. However, AUGUSTIN 92 favors a state decaying into  $K^*(892)\bar{K}$  at a lower mass than the state decaying into  $a_0(980)\pi$ , although agreement with MARK-III is not excluded. In  $J/\psi(1S)$  radiative decay, the  $\eta(1405)$  decays into  $K\bar{K}\pi$  through  $a_0(980)\pi$ , and hence a signal is also expected in the  $\eta\pi\pi$  mass spectrum. This was indeed observed by MARK III in  $\eta\pi^+\pi^-$  (BOLTON 92B), which reports a mass of 1400 MeV, in line with the existence of the  $\eta(1405)$  decaying to  $a_0(980)\pi$ . This state is also observed in  $\bar{p}p$  annihilation at rest into  $\eta\pi^+\pi^-\pi^0\pi^0$ , where it decays into  $\eta\pi\pi$  (AMSLER 95F). The intermediate  $a_0(980)\pi$  accounts for roughly half of the  $\eta\pi\pi$  signal, in agreement with MARK III (BOLTON 92B) and DM2 (AUGUSTIN 90).

The existence of the  $\eta(1295)$  is questioned by KLEMPY 05. In KLEMPY 07, the authors also question the existence of the  $\eta(1295)$ , and claim a single pseudoscalar meson in the 1400 MeV region. This conclusion is based on properties of the wave functions in the  ${}^3P_0$  model, and on an unpublished analysis of the annihilation  $\bar{p}p \rightarrow 4\pi\eta$ . The pseudoscalar signal

around 1400 MeV is then attributed to the first radial excitation of the  $\eta$ . However, the  $\eta(1295)$  has been observed by four  $\pi^- p$  experiments (ADAMS 01, FUKUI 91C, ALDE 97B, MANAK 00A), and evidence is reported in  $\bar{p}p$  annihilation (ANISOVICH 01, ABELE 98, AMSLER 04B). In  $J/\psi$  radiative decay, an  $\eta(1295)$  signal is evident in the  $0^{-+}$   $\eta\pi\pi$  wave of DM2 data (AUGUSTIN 92).

Assuming establishment of the  $\eta(1295)$ , the  $\eta(1475)$  could be the first radial excitation of the  $\eta'$ , with the  $\eta(1295)$  being the first radial excitation of the  $\eta$ . Ideal mixing, suggested by the  $\eta(1295)$  and  $\pi(1300)$  mass degeneracy, would then imply that the second isoscalar in the nonet is mainly  $s\bar{s}$ , and hence couples to  $K^*\bar{K}$ , in agreement with the  $\eta(1475)$ . Also its width matches the expected width for the radially excited  $s\bar{s}$  state (CLOSE 97, BARNES 97).

The  $K\bar{K}\pi$  and  $\eta\pi\pi$  channels were studied in  $\gamma\gamma$  collisions by L3 (ACCIARRI 01G). The analysis leads to a clear  $\eta(1475)$  signal in  $K\bar{K}\pi$ , decaying to  $K^*\bar{K}$ , very well identified in the untagged data sample, where contamination from spin 1 resonances is not allowed. At the same time, ACCIARRI 01G did not observe  $\eta(1405)$ , either in  $K\bar{K}\pi$  or  $\eta\pi\pi$ . The observation of the  $\eta(1475)$ , combined with the absence of an  $\eta(1405)$  signal, strengthens the two-resonances hypothesis. Since gluonium production is presumably suppressed in  $\gamma\gamma$  collisions, the ACCIARRI 01G results suggest that  $\eta(1405)$  has a large gluonic content (see also CLOSE 97B, LI 03C).

The ACCIARRI 01G result is somewhat in disagreement with that of CLEO-II, which did not observe any pseudoscalar signal in  $\gamma\gamma \rightarrow \eta(1475) \rightarrow K_S^0 K^\pm \pi^\mp$  (AHOHE 05). However, more data are required. Moreover, after the CLEO-II result, L3 performed a further analysis with full statistics (ACHARD 07), confirming the evidence of the  $\eta(1475)$  observed by ACCIARRI 01G. The CLEO upper limit (AHOHE 05) for  $\Gamma_{\gamma\gamma}(\eta(1475))$ , and the L3 results (ACHARD 07), are consistent with the world average for the  $\eta(1475)$  width.

The gluonium interpretation is not favored by lattice gauge theories which predict the  $0^{-+}$  state above 2 GeV (BALI 93). However, the  $\eta(1405)$  is an excellent candidate for the  $0^{-+}$

glueball in the fluxtube model (FADDEEV 04). In this model, the  $0^{++}$   $f_0(1500)$  glueball is also naturally related to a  $0^{-+}$  glueball with mass degeneracy broken in QCD.

A detailed review of the experimental situation is available in MASONI 06.

Let us now deal with  $1^{++}$  isoscalars. The  $f_1(1420)$ , decaying to  $K^*\bar{K}$ , was first reported in  $\pi^- p$  reactions at 4 GeV/c (DION-ISI 80). However, later analyses found that the 1400–1500 MeV region was far more complex (CHUNG 85, REEVES 86, BIRMAN 88). A reanalysis of the MARK III data in radiative  $J/\psi(1S)$  decay to  $K\bar{K}\pi$  (BAI 90C) shows the  $f_1(1420)$ , decaying into  $K^*\bar{K}$ . Also, a  $C=+1$  state is observed in tagged  $\gamma\gamma$  collisions (*e.g.*, BEHREND 89).

In  $\pi^- p \rightarrow \eta\pi\pi n$  charge-exchange reactions at 8–9 GeV/c, the  $\eta\pi\pi$  mass spectrum is dominated by the  $\eta(1440)$  and  $\eta(1295)$  (ANDO 86, FUKUI 91C), and at 100 GeV/c, ALDE 97B reports the  $\eta(1295)$  and  $\eta(1440)$  decaying to  $\eta\pi^0\pi^0$  with a weak  $f_1(1285)$  signal, and no evidence for the  $f_1(1420)$ .

Axial ( $1^{++}$ ) mesons are not observed in  $\bar{p}p$  annihilation at rest in liquid hydrogen, which proceeds dominantly through  $S$ -wave annihilation. However, in gaseous hydrogen,  $P$ -wave annihilation is enhanced and, indeed, BERTIN 97 reports  $f_1(1420)$  decaying to  $K^*\bar{K}$ .

The  $f_1(1420)$ , decaying into  $K\bar{K}\pi$ , is also seen in  $pp$  central production, together with the  $f_1(1285)$ . The latter decays via  $a_0(980)\pi$ , and the former only via  $K^*\bar{K}$ , while the  $\eta(1440)$  is absent (ARMSTRONG 89, BARBERIS 97C). The  $K_SK_S\pi^0$  decay mode of  $f_1(1420)$  establishes unambiguously  $C=+1$ . On the other hand, there is no evidence for any state decaying to  $\eta\pi\pi$  around 1400 MeV, and hence the  $\eta\pi\pi$  mode of the  $f_1(1420)$  must be suppressed (ARMSTRONG 91B).

We now turn to the experimental evidence for the  $f_1(1510)$ . Two states, the  $f_1(1420)$  and  $f_1(1510)$ , decaying to  $K^*\bar{K}$ , compete for the  $s\bar{s}$  assignment in the  $1^{++}$  nonet. The  $f_1(1510)$  was seen in  $K^- p \rightarrow \Lambda K\bar{K}\pi$  at 4 GeV/c (GAVILLET 82), and at 11 GeV/c (ASTON 88C). Evidence is also reported in  $\pi^- p$  at 8 GeV/c, based on the phase motion of the  $1^{++}$   $K^*\bar{K}$  wave

(BIRMAN 88). A somewhat broader  $1^{++}$  signal is also observed in  $J/\psi$  radiative decay to  $\eta\pi^+\pi^-$  (BAI 99).

The absence of the  $f_1(1420)$  in  $K^-p$  (ASTON 88C) argues against the  $f_1(1420)$  being the  $s\bar{s}$  member of the  $1^{++}$  nonet. However, the  $f_1(1420)$  was reported in  $K^-p$  but not in  $\pi^-p$  (BITYUKOV 84), while two experiments do not observe the  $f_1(1510)$  in  $K^-p$  (BITYUKOV 84, KING 91). It is also not seen in radiative  $J/\psi(1S)$  decay (BAI 90C, AUGUSTIN 92), central collisions (BARBERIS 97C), or  $\gamma\gamma$  collisions (AIHARA 88C), although, surprisingly for an  $s\bar{s}$  state, a signal is reported in  $4\pi$  decays (BAUER 93B). These facts lead to the conclusion that  $f_1(1510)$  is not well established (CLOSE 97D).

Assigning the  $f_1(1420)$  to the  $1^{++}$  nonet, one finds a nonet mixing angle of  $\sim 50^\circ$  (CLOSE 97D). However, arguments favoring the  $f_1(1420)$  being a hybrid  $q\bar{q}g$  meson, or a four-quark state, were put forward by ISHIDA 89 and CALDWELL 90, respectively, while LONGACRE 90 argued for a molecular state formed by the  $\pi$  orbiting in a  $P$ -wave around an  $S$ -wave  $K\bar{K}$  state.

Summarizing, there is convincing evidence for the  $f_1(1420)$  decaying to  $K^*\bar{K}$ , and for two pseudoscalars in the  $\eta(1440)$  region, the  $\eta(1405)$  and  $\eta(1475)$ , decaying to  $a_0(980)\pi$  and  $K^*\bar{K}$ , respectively. The  $f_1(1510)$  is not well established.