

$D_{s2}(2573)^{\pm}$

$I(J^P) = 0(?)$

J^P is natural, width and decay modes consistent with 2^+ .

$D_{s2}(2573)^{\pm}$ MASS

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | CHG | COMMENT |
|---|------|------------------------|------|------|--|
| 2572.6 ± 0.9 OUR AVERAGE | | | | | |
| 2572.2 $\pm 0.3 \pm 1.0$ | | AUBERT,BE | 06E | BABR | $e^+ e^- \rightarrow D K X$ |
| 2574.5 $\pm 3.3 \pm 1.6$ | | ALBRECHT | 96 | ARG | $e^+ e^- \rightarrow D^0 K^+ X$ |
| $2573.2^{+1.7}_{-1.6} \pm 0.9$ | 217 | KUBOTA | 94 | CLE2 | $+ e^+ e^- \sim 10.5$ GeV |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| 2570.0 ± 4.3 | 25 | ¹ EVDOKIMOV | 04 | SELX | $600 \Sigma^- A \rightarrow D^0 K^+ X$ |
| 2568.6 ± 3.2 | 64 | ² HEISTER | 02B | ALEP | $e^+ e^- \rightarrow D^0 K^+ X$ |

¹ Not independent of the mass difference below.

² Calculated using $m_{D^0} = 1864.5 \pm 0.5$ MeV and the mass difference below.


$m_{D_{s2}(2573)^{\pm}} - m_{D^0}$

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|------------------------|------|---|
| 704 $\pm 3 \pm 1$ | 64 | HEISTER | 02B | $ALEP e^+ e^- \rightarrow D^0 K^+ X$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| 705.4 ± 4.3 | 25 | ³ EVDOKIMOV | 04 | SELX $600 \Sigma^- A \rightarrow D^0 K^+ X$ |
| $\bullet \bullet \bullet$ Systematic errors not estimated. | | | | |

$D_{s2}(2573)^{\pm}$ WIDTH

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | CHG | COMMENT |
|---|------|------------------------|------|------|--|
| 20 ± 5 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below. | | | | | |
| 27.1 $\pm 0.6 \pm 5.6$ | | AUBERT,BE | 06E | BABR | $e^+ e^- \rightarrow D K X$ |
| 10.4 $\pm 8.3 \pm 3.0$ | | ALBRECHT | 96 | ARG | $e^+ e^- \rightarrow D^0 K^+ X$ |
| $16^{+5}_{-4} \pm 3$ | 217 | KUBOTA | 94 | CLE2 | $+ e^+ e^- \sim 10.5$ GeV |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| 14 $^{+9}_{-6}$ | 25 | ⁴ EVDOKIMOV | 04 | SELX | $600 \Sigma^- A \rightarrow D^0 K^+ X$ |

⁴ Systematic errors not estimated.



$D_{s2}(2573)^+$ DECAY MODES

$D_{s2}(2573)^-$ modes are charge conjugates of the modes below.

| Mode | Fraction (Γ_i/Γ) |
|----------------------------------|--------------------------------|
| $\Gamma_1 \quad D^0 K^+$ | seen |
| $\Gamma_2 \quad D^*(2007)^0 K^+$ | not seen |

$D_{s2}(2573)^+$ BRANCHING RATIOS

| $\Gamma(D^0 K^+)/\Gamma_{\text{total}}$ | Γ_1/Γ |
|---|---|
| VALUE <u>EVTS</u> seen 217 | <u>DOCUMENT ID</u> KUBOTA <u>TECN</u> 94 <u>CHG</u> ± <u>COMMENT</u> $e^+ e^- \sim 10.5 \text{ GeV}$ |

| $\Gamma(D^*(2007)^0 K^+)/\Gamma(D^0 K^+)$ | Γ_2/Γ_1 |
|---|---|
| VALUE <u>CL%</u> <0.33 90 | <u>DOCUMENT ID</u> KUBOTA <u>TECN</u> 94 <u>CHG</u> + <u>COMMENT</u> $e^+ e^- \sim 10.5 \text{ GeV}$ |

$D_{s2}(2573)^{\pm}$ REFERENCES

| | | | | |
|-----------|-----|---------------|------------------------------|-----------------|
| AUBERT,BE | 06E | PRL 97 222001 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| EVDOKIMOV | 04 | PRL 93 242001 | A.V. Evdokimov <i>et al.</i> | (SELEX Collab.) |
| HEISTER | 02B | PL B526 34 | A. Heister <i>et al.</i> | (ALEPH Collab.) |
| ALBRECHT | 96 | ZPHY C69 405 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| KUBOTA | 94 | PRL 72 1972 | Y. Kubota <i>et al.</i> | (CLEO Collab.) |

———— OTHER RELATED PAPERS ——

| | | | | |
|-----------------------------|-----|---------------|----------------------------|---------|
| COLANGELO | 06 | PL B642 48 | P. Colangelo <i>et al.</i> | |
| CLOSE | 05C | PR D72 094004 | F.E. Close, E.S. Swanson | (OXFTP) |
| SEMENOV | 99 | SPU 42 847 | S.V. Semenov | |
| Translated from UFN 42 937. | | | | |
