

# $\chi_{c2}(1P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

See the Review on “ $\psi(2S)$  and  $\chi_c$  branching ratios” before the  $\chi_{c0}(1P)$  Listings.

## $\chi_{c2}(1P)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>3556.20 ± 0.09 OUR AVERAGE</b>				
3555.3 ± 0.6 ± 2.2	2.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow$ hadrons
3555.70 ± 0.59 ± 0.39		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c2}$
3556.173 ± 0.123 ± 0.020		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
3559.9 ± 2.9		EISENSTEIN	01 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
3556.4 ± 0.7		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3556.22 ± 0.131 ± 0.020	585	<sup>1</sup> ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
3556.9 ± 0.4 ± 0.5	50	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^- X$
3557.8 ± 0.2 ± 4		<sup>2</sup> GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3553.4 ± 2.2	66	<sup>3</sup> LEMOIGNE	82 GOLI	$185 \pi^- \text{Be} \rightarrow \gamma\mu^+\mu^- A$
3555.9 ± 0.7		<sup>4</sup> OREGLIA	82 CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
3557 ± 1.5	69	<sup>5</sup> HIMEL	80 MRK2	$e^+e^- \rightarrow J/\psi 2\gamma$
3551 ± 11	15	BRANDELIK	79B DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
3553 ± 4		<sup>5</sup> BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3553 ± 4 ± 4		<sup>5,6</sup> TANENBAUM	78 MRK1	$e^+e^-$
3563 ± 7	360	<sup>5</sup> BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3543 ± 10	4	WHITAKER	76 MRK1	$e^+e^- \rightarrow J/\psi 2\gamma$

<sup>1</sup> Recalculated by ANDREOTTI 05A, using the value of  $\psi(2S)$  mass from AULCHENKO 03.

<sup>2</sup> Using mass of  $\psi(2S) = 3686.0$  MeV.

<sup>3</sup>  $J/\psi(1S)$  mass constrained to 3097 MeV.

<sup>4</sup> Assuming  $\psi(2S)$  mass = 3686 MeV and  $J/\psi(1S)$  mass = 3097 MeV.

<sup>5</sup> Mass value shifted by us by amount appropriate for  $\psi(2S)$  mass = 3686 MeV and  $J/\psi(1S)$  mass = 3097 MeV.

<sup>6</sup> From a simultaneous fit to radiative and hadronic decay channels.

## $\chi_{c2}(1P)$ WIDTH

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.98 ± 0.11 OUR FIT</b>				
<b>1.95 ± 0.13 OUR AVERAGE</b>				
1.915 ± 0.188 ± 0.013		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
1.96 ± 0.17 ± 0.07	585	<sup>7</sup> ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
2.6 <sup>+1.4</sup> / <sub>-1.0</sub>	50	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^- X$
2.8 <sup>+2.1</sup> / <sub>-2.0</sub>		<sup>8</sup> GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$

<sup>7</sup> Recalculated by ANDREOTTI 05A.

<sup>8</sup> Errors correspond to 90% confidence level; authors give only width range.

## $\chi_{c2}(1P)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level	
<b>Hadronic decays</b>			
$\Gamma_1$	$2(\pi^+ \pi^-)$	( 1.09 ± 0.11 ) %	
$\Gamma_2$	$\rho \rho$		
$\Gamma_3$	$\pi^+ \pi^- \pi^0 \pi^0$	( 2.01 ± 0.26 ) %	
$\Gamma_4$	$\rho^+ \pi^- \pi^0 + \text{c.c.}$	( 2.4 ± 0.4 ) %	
$\Gamma_5$	$K^+ K^- \pi^0 \pi^0$	( 2.3 ± 0.5 ) × 10 <sup>-3</sup>	
$\Gamma_6$	$K^+ \pi^- K^0 \pi^0 + \text{c.c.}$	( 1.51 ± 0.22 ) %	
$\Gamma_7$	$\rho^+ K^- K^0 + \text{c.c.}$	( 4.5 ± 1.4 ) × 10 <sup>-3</sup>	
$\Gamma_8$	$K^*(892)^0 K^+ \pi^- \rightarrow$ $K^+ \pi^- K^0 \pi^0 + \text{c.c.}$	( 3.2 ± 0.9 ) × 10 <sup>-3</sup>	
$\Gamma_9$	$K^*(892)^0 K^0 \pi^0 \rightarrow$ $K^+ \pi^- K^0 \pi^0 + \text{c.c.}$	( 4.2 ± 1.0 ) × 10 <sup>-3</sup>	
$\Gamma_{10}$	$K^*(892)^- K^+ \pi^0 \rightarrow$ $K^+ \pi^- K^0 \pi^0 + \text{c.c.}$	( 4.1 ± 0.9 ) × 10 <sup>-3</sup>	
$\Gamma_{11}$	$K^*(892)^+ K^0 \pi^- \rightarrow$ $K^+ \pi^- K^0 \pi^0 + \text{c.c.}$	( 3.2 ± 0.9 ) × 10 <sup>-3</sup>	
$\Gamma_{12}$	$K^+ K^- \eta \pi^0$	( 1.4 ± 0.5 ) × 10 <sup>-3</sup>	
$\Gamma_{13}$	$\pi^+ \pi^- K^+ K^-$	( 9.0 ± 1.1 ) × 10 <sup>-3</sup>	
$\Gamma_{14}$	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	( 2.3 ± 1.2 ) × 10 <sup>-3</sup>	
$\Gamma_{15}$	$K^*(892)^0 \bar{K}^*(892)^0$	( 2.5 ± 0.5 ) × 10 <sup>-3</sup>	
$\Gamma_{16}$	$3(\pi^+ \pi^-)$	( 8.6 ± 1.8 ) × 10 <sup>-3</sup>	
$\Gamma_{17}$	$\phi \phi$	( 1.47 ± 0.28 ) × 10 <sup>-3</sup>	
$\Gamma_{18}$	$\omega \omega$	( 1.9 ± 0.6 ) × 10 <sup>-3</sup>	
$\Gamma_{19}$	$\pi \pi$	( 2.09 ± 0.23 ) × 10 <sup>-3</sup>	
$\Gamma_{20}$	$\rho^0 \pi^+ \pi^-$	( 3.9 ± 1.7 ) × 10 <sup>-3</sup>	
$\Gamma_{21}$	$\pi^+ \pi^- \eta$	( 5.3 ± 1.4 ) × 10 <sup>-4</sup>	
$\Gamma_{22}$	$\pi^+ \pi^- \eta'$	( 5.5 ± 2.0 ) × 10 <sup>-4</sup>	
$\Gamma_{23}$	$\eta \eta$	< 5 × 10 <sup>-4</sup>	90%
$\Gamma_{24}$	$K^+ K^-$	( 7.6 ± 1.3 ) × 10 <sup>-4</sup>	
$\Gamma_{25}$	$K_S^0 K_S^0$	( 6.2 ± 0.8 ) × 10 <sup>-4</sup>	
$\Gamma_{26}$	$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	( 1.33 ± 0.20 ) × 10 <sup>-3</sup>	
$\Gamma_{27}$	$K^+ K^- \pi^0$	( 3.3 ± 0.8 ) × 10 <sup>-4</sup>	
$\Gamma_{28}$	$K^+ K^- \eta$	< 3.5 × 10 <sup>-4</sup>	90%
$\Gamma_{29}$	$\eta \eta'$	< 2.5 × 10 <sup>-4</sup>	90%
$\Gamma_{30}$	$\eta' \eta'$	< 3.3 × 10 <sup>-4</sup>	90%
$\Gamma_{31}$	$\pi^+ \pi^- K_S^0 K_S^0$	( 2.4 ± 0.6 ) × 10 <sup>-3</sup>	
$\Gamma_{32}$	$K^+ K^- K_S^0 K_S^0$	< 4 × 10 <sup>-4</sup>	90%
$\Gamma_{33}$	$K^+ K^- K^+ K^-$	( 1.77 ± 0.22 ) × 10 <sup>-3</sup>	
$\Gamma_{34}$	$K^+ K^- \phi$	( 1.56 ± 0.33 ) × 10 <sup>-3</sup>	
$\Gamma_{35}$	$K_S^0 K_S^0 \rho \bar{p}$	< 7.9 × 10 <sup>-4</sup>	90%
$\Gamma_{36}$	$\rho \bar{p}$	( 7.2 ± 0.4 ) × 10 <sup>-5</sup>	

$\Gamma_{37}$	$\rho\bar{\rho}\pi^0$	$(4.7 \pm 1.0) \times 10^{-4}$	
$\Gamma_{38}$	$\rho\bar{\rho}\eta$	$(2.0 \pm 0.8) \times 10^{-4}$	
$\Gamma_{39}$	$\pi^+\pi^-\rho\bar{\rho}$	$(1.32 \pm 0.34) \times 10^{-3}$	
$\Gamma_{40}$	$\pi^0\pi^0\rho\bar{\rho}$	$(8.6 \pm 2.6) \times 10^{-4}$	
$\Gamma_{41}$	$\rho\bar{n}\pi^-$	$(1.1 \pm 0.4) \times 10^{-3}$	
$\Gamma_{42}$	$\Lambda\bar{\Lambda}$	$(1.87 \pm 0.27) \times 10^{-4}$	
$\Gamma_{43}$	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$< 3.5 \times 10^{-3}$	90%
$\Gamma_{44}$	$K^+\bar{p}\Lambda + \text{c.c.}$	$(9.1 \pm 1.8) \times 10^{-4}$	
$\Gamma_{45}$	$\Sigma^0\bar{\Sigma}^0$	$< 8 \times 10^{-5}$	90%
$\Gamma_{46}$	$\Sigma^+\bar{\Sigma}^-$	$< 7 \times 10^{-5}$	90%
$\Gamma_{47}$	$\Xi^0\bar{\Xi}^0$	$< 1.1 \times 10^{-4}$	90%
$\Gamma_{48}$	$\Xi^-\bar{\Xi}^+$	$(1.56 \pm 0.35) \times 10^{-4}$	
$\Gamma_{49}$	$J/\psi(1S)\pi^+\pi^-\pi^0$	$< 1.5 \%$	90%

### Radiative decays

$\Gamma_{50}$	$\gamma J/\psi(1S)$	$(19.4 \pm 0.8) \%$	
$\Gamma_{51}$	$\gamma\rho^0$	$< 5 \times 10^{-5}$	90%
$\Gamma_{52}$	$\gamma\omega$	$< 7 \times 10^{-6}$	90%
$\Gamma_{53}$	$\gamma\phi$	$< 1.2 \times 10^{-5}$	90%
$\Gamma_{54}$	$\gamma\gamma$	$(2.60 \pm 0.16) \times 10^{-4}$	

---

### CONSTRAINED FIT INFORMATION

A multiparticle fit to  $\chi_{c1}(1P)$ ,  $\chi_{c0}(1P)$ ,  $\chi_{c2}(1P)$ , and  $\psi(2S)$  with 4 total widths, a partial width, 24 combinations of partial widths obtained from integrated cross section, and 78 branching ratios uses 203 measurements to determine 47 parameters. The overall fit has a  $\chi^2 = 273.3$  for 156 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ .

$x_{13}$	17									
$x_{14}$	4	22								
$x_{15}$	10	9	2							
$x_{17}$	9	7	2	5						
$x_{19}$	13	11	3	7	7					
$x_{20}$	20	4	1	2	2	3				
$x_{24}$	9	8	2	5	5	8	2			
$x_{25}$	14	12	3	7	7	10	3	7		
$x_{33}$	12	10	2	6	6	11	3	7	9	
$x_{36}$	7	6	1	4	2	2	2	2	4	2
$x_{42}$	8	7	2	4	5	9	2	5	7	8
$x_{50}$	27	23	5	14	15	28	6	18	22	25
$x_{54}$	-19	-16	-3	-10	-5	-3	-5	-4	-11	-3
$\Gamma$	-25	-21	-5	-13	-12	-19	-6	-13	-18	-17
	$x_1$	$x_{13}$	$x_{14}$	$x_{15}$	$x_{17}$	$x_{19}$	$x_{20}$	$x_{24}$	$x_{25}$	$x_{33}$
$x_{42}$	0									
$x_{50}$	-12	22								
$x_{54}$	25	2	9							
$\Gamma$	-50	-13	-49	-45						
	$x_{36}$	$x_{42}$	$x_{50}$	$x_{54}$						

### $\chi_{c2}(1P)$ PARTIAL WIDTHS

$$\text{--- } \chi_{c2}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total}) \text{ ---}$$

$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}}$   $\Gamma_{36} \Gamma_{50} / \Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>27.7 ± 1.4 OUR FIT</b>			
<b>27.5 ± 1.5 OUR AVERAGE</b>			
27.0 ± 1.5 ± 1.1	<sup>9</sup> ANDREOTTI 05A	E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
27.7 ± 1.5 ± 2.0	<sup>9,10</sup> ARMSTRONG 92	E760	$\bar{p}p \rightarrow e^+ e^- \gamma$
36 ± 8	<sup>9</sup> BAGLIN 86B	SPEC	$\bar{p}p \rightarrow e^+ e^- X$

<sup>9</sup> Calculated by us using  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$ .

<sup>10</sup> Recalculated by ANDREOTTI 05A.

**$\Gamma(\gamma\gamma) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$   $\Gamma_{54}\Gamma_{50}/\Gamma$**

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>100 ± 6 OUR FIT</b>				
<b>117 ± 10 OUR AVERAGE</b>				
111 ± 12 ± 9	147 ± 15	<sup>11</sup> DOBBS	06 CLE3	10.4 $e^+e^- \rightarrow e^+e^-\chi_{c2}$
114 ± 11 ± 9	136 ± 13.3	<sup>11,12</sup> ABE	02T BELL	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
139 ± 55 ± 21		<sup>11,13</sup> ACCIARRI	99E L3	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
242 ± 65 ± 51		<sup>11,14</sup> ACKER...,K...	98 OPAL	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
150 ± 42 ± 36		<sup>11,15</sup> DOMINICK	94 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
470 ± 240 ± 120		<sup>11,16</sup> BAUER	93 TPC	$e^+e^- \rightarrow e^+e^-\chi_{c2}$

<sup>11</sup> Calculated by us using  $B(J/\psi \rightarrow \ell^+\ell^-) = 0.1187 \pm 0.0008$ .

<sup>12</sup> All systematic errors added in quadrature.

<sup>13</sup> The value for  $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$  reported in ACCIARRI 99E is derived using  $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) \times B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.0162 \pm 0.0014$ .

<sup>14</sup> The value for  $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$  reported in ACKERSTAFF,K 98 is derived using  $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$  and  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$ .

<sup>15</sup> The value for  $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$  reported in DOMINICK 94 is derived using  $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$ ,  $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0627 \pm 0.0020$ , and  $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0597 \pm 0.0025$ .

<sup>16</sup> The value for  $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$  reported in BAUER 93 is derived using  $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$ ,  $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0627 \pm 0.0020$ , and  $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0597 \pm 0.0025$ .

————  $\chi_{c2}(1P) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$  ————

**$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_1\Gamma_{54}/\Gamma$**

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.6 ± 0.5 OUR FIT</b>				
<b>5.2 ± 0.7 OUR AVERAGE</b>				
5.01 ± 0.44 ± 0.55	1597 ± 138	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$
6.4 ± 1.8 ± 0.8		EISENSTEIN	01 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c2}$

**$\Gamma(\rho^0\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{20}\Gamma_{54}/\Gamma$**

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.0 ± 0.9 OUR FIT</b>				
<b>3.2 ± 1.9 ± 0.5</b>	986 ± 578	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$

**$\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_2\Gamma_{54}/\Gamma$**

<u>VALUE (eV)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------	------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7.8	90	<598	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$
------	----	------	--------	---------	--

**$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{13}\Gamma_{54}/\Gamma$**

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------	-------------	--------------------	-------------	----------------

**4.6 ± 0.5 OUR FIT**

<b>4.42 ± 0.42 ± 0.53</b>	780 ± 74	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+K^-\pi^+\pi^-$
---------------------------	----------	--------	---------	---

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{15}\Gamma_{54}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.26±0.24 OUR FIT</b>				
<b>0.8 ±0.17±0.27</b>	151 ± 30	UEHARA 08	BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+ K^- \pi^+ \pi^-$

$$\Gamma(K^+ K^- K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{33}\Gamma_{54}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.91±0.12 OUR FIT</b>				
<b>1.10±0.21±0.15</b>	126 ± 24	UEHARA 08	BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+ K^-)$

$$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{17}\Gamma_{54}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.75±0.14 OUR FIT</b>				
<b>0.58±0.18±0.16</b>	26.5 ± 8.1	UEHARA 08	BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+ K^-)$

$$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{19}\Gamma_{54}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.08±0.13 OUR FIT</b>				
<b>1.14±0.21±0.17</b>	54 ± 10	<sup>17</sup> NAKAZAWA 05	BELL	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$

<sup>17</sup> We have multiplied  $\pi^+ \pi^-$  measurement by 3/2 to obtain  $\pi\pi$ .

$$\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{24}\Gamma_{54}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.39±0.07 OUR FIT</b>				
<b>0.44±0.11±0.07</b>	33 ± 8	NAKAZAWA 05	BELL	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$

$$\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{25}\Gamma_{54}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.32±0.04 OUR FIT</b>				
<b>0.31±0.05±0.03</b>	38 ± 7	CHEN 07B	BELL	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$

## $\chi_{c2}(1P)$ BRANCHING RATIOS

### ———— HADRONIC DECAYS ————

$$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE	DOCUMENT ID
<b>0.0109±0.0011 OUR FIT</b>	

$$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-)) \quad \Gamma_{20}/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.36±0.15 OUR FIT</b>			
<b>0.31±0.17</b>	TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.01±0.25±0.08</b>	903.5	<sup>18</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>18</sup> HE 08B reports  $1.87 \pm 0.07 \pm 0.22 \pm 0.13$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.4±0.4±0.1</b>	1031.9	<sup>19,20</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>19</sup> HE 08B reports  $2.23 \pm 0.11 \pm 0.32 \pm 0.16$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>20</sup> Calculated by us. We have added the values from HE 08B for  $\rho^+\pi^-\pi^0$  and  $\rho^-\pi^+\pi^0$  decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

 $\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.23±0.04±0.01</b>	76.9	<sup>21</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>21</sup> HE 08B reports  $0.21 \pm 0.03 \pm 0.03 \pm 0.01$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+\pi^-K^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.51±0.21±0.06</b>	211.6	<sup>22</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>22</sup> HE 08B reports  $1.41 \pm 0.11 \pm 0.16 \pm 0.10$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\pi^-K^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^+K^-K^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.45±0.13±0.02</b>	62.9	<sup>23</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>23</sup> HE 08B reports  $0.42 \pm 0.11 \pm 0.06 \pm 0.03$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^+K^-K^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0 K^+ \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.32±0.09±0.01</b>	38.7	<sup>24</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>24</sup> HE 08B reports  $0.30 \pm 0.07 \pm 0.04 \pm 0.02$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 K^+ \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0 K^0 \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.42±0.09±0.02</b>	63.0	<sup>25</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>25</sup> HE 08B reports  $0.39 \pm 0.07 \pm 0.05 \pm 0.03$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 K^0 \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.41±0.09±0.02</b>	51.1	<sup>26</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>26</sup> HE 08B reports  $0.38 \pm 0.07 \pm 0.04 \pm 0.03$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^+ K^0 \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.32±0.09±0.01</b>	39.3	<sup>27</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>27</sup> HE 08B reports  $0.30 \pm 0.07 \pm 0.04 \pm 0.02$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+ K^0 \pi^- \rightarrow K^+ \pi^- K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.14±0.05±0.01</b>	22.9	<sup>28</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>28</sup> HE 08B reports  $0.13 \pm 0.04 \pm 0.02 \pm 0.01$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.



$$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}} \qquad \Gamma_{13}/\Gamma$$

VALUE (units  $10^{-3}$ )                      DOCUMENT ID  
**9.0±1.1 OUR FIT**

$$\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.})/\Gamma(\pi^+\pi^-K^+K^-) \qquad \Gamma_{14}/\Gamma_{13}$$

VALUE                      DOCUMENT ID    TECN    COMMENT  
**0.25±0.13 OUR FIT**  
**0.25±0.13**                      TANENBAUM 78    MRK1     $\psi(2S) \rightarrow \gamma\chi_{c2}$

$$\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.})/\Gamma_{\text{total}} \qquad \Gamma_{14}/\Gamma$$

VALUE (units  $10^{-4}$ )                      DOCUMENT ID  
**23±12 OUR FIT**

$$\Gamma(K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}} \qquad \Gamma_{15}/\Gamma$$

VALUE (units  $10^{-3}$ )                      DOCUMENT ID  
**2.5±0.5 OUR FIT**

$$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}} \qquad \Gamma_{16}/\Gamma$$

VALUE (units  $10^{-3}$ )                      DOCUMENT ID    TECN    COMMENT  
**8.6±1.8 OUR EVALUATION**    Treating systematic error as correlated.  
**8.6±1.8 OUR AVERAGE**

8.6±0.9±1.6                      29 BAI                      99B BES     $\psi(2S) \rightarrow \gamma\chi_{c2}$   
8.7±5.9±0.4                      29 TANENBAUM 78    MRK1     $\psi(2S) \rightarrow \gamma\chi_{c2}$

<sup>29</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ . Multiplied by a factor of 2 to convert from  $K_S^0 K^+\pi^-$  to  $K^0 K^+\pi^-$  decay.

$$\Gamma(\phi\phi)/\Gamma_{\text{total}} \qquad \Gamma_{17}/\Gamma$$

VALUE (units  $10^{-3}$ )                      DOCUMENT ID  
**1.47±0.28 OUR FIT**

$$\Gamma(\omega\omega)/\Gamma_{\text{total}} \qquad \Gamma_{18}/\Gamma$$

VALUE (units  $10^{-3}$ )    EVTS                      DOCUMENT ID    TECN    COMMENT  
**1.9±0.6±0.1**    27.7±7.4                      <sup>30</sup> ABLIKIM                      05N BES2     $\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma 6\pi$

<sup>30</sup> ABLIKIM 05N reports  $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.165 \pm 0.044 \pm 0.032) \times 10^{-3}$ . We divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi\pi)/\Gamma_{\text{total}} \qquad \Gamma_{19}/\Gamma$$

VALUE (units  $10^{-3}$ )                      DOCUMENT ID  
**2.09±0.23 OUR FIT**

$$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}} \qquad \Gamma_{20}/\Gamma$$

VALUE (units  $10^{-4}$ )                      DOCUMENT ID  
**39±17 OUR FIT**

$\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<b>0.53±0.14±0.02</b>		<sup>31</sup> ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$
-----------------------	--	---------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6	90	<sup>32</sup> ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \chi_{c2}$
------	----	-----------------------	-----	--

<sup>31</sup> ATHAR 07 reports  $(0.49 \pm 0.12 \pm 0.06) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>32</sup> ABLIKIM 06R reports  $< 1.7 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .

$\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

<b>0.55±0.20±0.02</b>	<sup>33</sup> ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$
-----------------------	---------------------	----	--

<sup>33</sup> ATHAR 07 reports  $(0.51 \pm 0.18 \pm 0.06) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<b>&lt; 5</b>	90	<sup>34</sup> ADAMS	07	CLEO $\psi(2S) \rightarrow \gamma \chi_{c2}$
---------------	----	---------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12	90	<sup>35</sup> BAI	03C	BES $\psi(2S) \rightarrow \gamma \eta \eta \rightarrow 5\gamma$
7.9±4.1±2.4		<sup>36</sup> LEE	85	CBAL $\psi' \rightarrow \text{photons}$

<sup>34</sup> ADAMS 07 reports  $< 4.7 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .

<sup>35</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.3 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ . Multiplied by a factor of 2 to convert from  $K_S^0 K^+ \pi^-$  to  $K^0 K^+ \pi^-$  decay.

<sup>36</sup> Calculated using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.078 \pm 0.008$ .

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>
---	--------------------

**0.76±0.13 OUR FIT**

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>
---	--------------------

**0.62±0.08 OUR FIT**

### $\Gamma(K_S^0 K_S^0)/\Gamma(\pi\pi)$

$\Gamma_{25}/\Gamma_{19}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

#### 0.30±0.05 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.27±0.07±0.04	37,38 CHEN	07B BELL	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
----------------	------------	----------	--------------------------------------

<sup>37</sup> Using  $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  from the  $\pi^+\pi^-$  measurement of NAKAZAWA 05 rescaled by 3/2 to convert to  $\pi\pi$ .

<sup>38</sup> Not independent from other measurements.

### $\Gamma(K_S^0 K_S^0)/\Gamma(K^+K^-)$

$\Gamma_{25}/\Gamma_{24}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

#### 0.82±0.17 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.70±0.21±0.12	39,40 CHEN	07B BELL	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
----------------	------------	----------	--------------------------------------

<sup>39</sup> Using  $\Gamma(K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  from NAKAZAWA 05.

<sup>40</sup> Not independent from other measurements.

### $\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

$\Gamma_{26}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

#### 1.33±0.20 OUR AVERAGE

1.40±0.22±0.06			41 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
----------------	--	--	----------	---------	---

1.12±0.41±0.04	28		42 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c2}$
----------------	----	--	------------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.0	90		43 BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c2}$
------	----	--	--------	---------	--

<sup>41</sup> ATHAR 07 reports  $(1.3 \pm 0.2 \pm 0.1) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>42</sup> We have multiplied the  $K_S^0 K^+ \pi^-$  measurement by a factor of 2 to convert to  $K^0 K^+ \pi^-$ . ABLIKIM 06R reports  $(1.2 \pm 0.4 \pm 0.2) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.6) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>43</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ . Multiplied by a factor of 2 to convert from  $K_S^0 K^+ \pi^-$  to  $K^0 K^+ \pi^-$  decay.

### $\Gamma(K^+K^-\pi^0)/\Gamma_{\text{total}}$

$\Gamma_{27}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.33±0.08±0.01	44 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
----------------	----------	---------	---

<sup>44</sup> ATHAR 07 reports  $(0.31 \pm 0.07 \pm 0.04) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$**   **$\Gamma_{28}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.35</b>	90	<sup>45</sup> ATHAR 07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>45</sup> ATHAR 07 reports  $< 0.33 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .

**$\Gamma(\eta\eta')/\Gamma_{\text{total}}$**   **$\Gamma_{29}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;2.5</b>	90	<sup>46</sup> ADAMS 07	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$

<sup>46</sup> ADAMS 07 reports  $< 2.3 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .

**$\Gamma(\eta'\eta')/\Gamma_{\text{total}}$**   **$\Gamma_{30}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;3.3</b>	90	<sup>47</sup> ADAMS 07	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$

<sup>47</sup> ADAMS 07 reports  $< 3.1 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .

**$\Gamma(\pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}$**   **$\Gamma_{31}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.4 \pm 0.6 \pm 0.1</math></b>	$57 \pm 11$	<sup>48</sup> ABLIKIM 050	BES2	$\psi(2S) \rightarrow \gamma \chi_{c2}$

<sup>48</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (0.207 \pm 0.039 \pm 0.033) \times 10^{-3}$ . We divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$**   **$\Gamma_{32}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;4</b>	90	$2.3 \pm 2.2$	<sup>49</sup> ABLIKIM 050	BES2	$e^+ e^- \rightarrow \chi_{c2} \gamma$

<sup>49</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] < 3.5 \times 10^{-5}$ . We divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .

**$\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$**   **$\Gamma_{33}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>
<b><math>1.77 \pm 0.22</math> OUR FIT</b>	

**$\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$**   **$\Gamma_{34}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.56 \pm 0.32 \pm 0.06</math></b>	52	<sup>50</sup> ABLIKIM 06T	BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>50</sup> ABLIKIM 06T reports  $(1.67 \pm 0.26 \pm 0.24) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \phi) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(K_S^0 K_S^0 \rho \bar{\rho}) / \Gamma_{\text{total}}$**   **$\Gamma_{35} / \Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;7.9</b>	90	<sup>51</sup> ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c2} \gamma$

<sup>51</sup> Using  $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (9.3 \pm 0.6)\%$ .

**$\Gamma(\rho \bar{\rho}) / \Gamma_{\text{total}}$**   **$\Gamma_{36} / \Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>
<b><math>0.72 \pm 0.04</math> OUR FIT</b>	

**$\Gamma(\rho \bar{\rho} \pi^0) / \Gamma_{\text{total}}$**   **$\Gamma_{37} / \Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.47 \pm 0.10 \pm 0.02</math></b>	<sup>52</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>52</sup> ATHAR 07 reports  $(0.44 \pm 0.08 \pm 0.05) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \rho \bar{\rho} \pi^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\rho \bar{\rho} \eta) / \Gamma_{\text{total}}$**   **$\Gamma_{38} / \Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.20 \pm 0.08 \pm 0.01</math></b>	<sup>53</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>53</sup> ATHAR 07 reports  $(0.19 \pm 0.07 \pm 0.02) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \rho \bar{\rho} \eta) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\pi^+ \pi^- \rho \bar{\rho}) / \Gamma_{\text{total}}$**   **$\Gamma_{39} / \Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.32 \pm 0.34</math> OUR EVALUATION</b>	Treating systematic error as correlated.		
<b><math>1.3 \pm 0.4</math> OUR AVERAGE</b>	Error includes scale factor of 1.3.		
$1.17 \pm 0.19 \pm 0.30$	<sup>54</sup> BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c2}$
$2.64 \pm 1.03 \pm 0.14$	<sup>54</sup> TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$

<sup>54</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.3 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$ . Multiplied by a factor of 2 to convert from  $K_S^0 K^+ \pi^-$  to  $K^0 K^+ \pi^-$  decay.

**$\Gamma(\pi^0 \pi^0 \rho \bar{\rho})/\Gamma_{\text{total}}$**   **$\Gamma_{40}/\Gamma$**

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.086 ± 0.026 ± 0.003</b>	29.2	55 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>55</sup> HE 08B reports  $0.08 \pm 0.02 \pm 0.01 \pm 0.01$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^0 \pi^0 \rho \bar{\rho})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\rho \bar{\rho} \pi^-)/\Gamma_{\text{total}}$**   **$\Gamma_{41}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>11.2 ± 3.8 ± 0.4</b>	56 ABLIKIM	06i	BES2 $\psi(2S) \rightarrow \gamma \rho \pi^- X$

<sup>56</sup> ABLIKIM 06i reports  $[\Gamma(\chi_{c2}(1P) \rightarrow \rho \bar{\rho} \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  =  $(0.97 \pm 0.20 \pm 0.26) \times 10^{-4}$ . We divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\Lambda \bar{\Lambda})/\Gamma_{\text{total}}$**   **$\Gamma_{42}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>
<b>1.87 ± 0.27 OUR FIT</b>	

**$\Gamma(\Lambda \bar{\Lambda} \pi^+ \pi^-)/\Gamma_{\text{total}}$**   **$\Gamma_{43}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 3.5</b>	90	57 ABLIKIM	06D	BES2 $\psi(2S) \rightarrow \chi_{c2} \gamma$

<sup>57</sup> Using  $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (9.3 \pm 0.6)\%$ .

**$\Gamma(K^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}$**   **$\Gamma_{44}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.91 ± 0.17 ± 0.04</b>	58 ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>58</sup> ATHAR 07 reports  $(0.85 \pm 0.14 \pm 0.10) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$**   **$\Gamma_{45}/\Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt; 0.8</b>	90	7.5 ± 3.4	59 NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

<sup>59</sup> NAIK 08 reports  $< 0.75 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$   $\Gamma_{46}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<b>&lt;0.7</b>	90	4.0 ± 3.5	<sup>60</sup> NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Sigma^+\bar{\Sigma}^-$
----------------	----	-----------	--------------------	----	---

<sup>60</sup> NAIK 08 reports  $< 0.67 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .

$\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$   $\Gamma_{47}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<b>&lt;1.1</b>	90	2.9 ± 1.7	<sup>61</sup> NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Xi^0\bar{\Xi}^0$
----------------	----	-----------	--------------------	----	---

<sup>61</sup> NAIK 08 reports  $< 1.06 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .

$\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$   $\Gamma_{48}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<b>1.56 ± 0.34 ± 0.06</b>		29 ± 5	<sup>62</sup> NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Xi^+\bar{\Xi}^-$
---------------------------	--	--------	--------------------	----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.7	90		<sup>63</sup> ABLIKIM	06D	BES2 $\psi(2S) \rightarrow \chi_{c2}\gamma$
-------	----	--	-----------------------	-----	---

<sup>62</sup> NAIK 08 reports  $(1.45 \pm 0.30 \pm 0.15) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.69 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>63</sup> Using  $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$ .

$\Gamma(J/\psi(1S)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{49}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b>&lt;0.015</b>	90	BARATE	81	SPEC 190 GeV $\pi^- \text{Be} \rightarrow 2\pi 2\mu$
------------------	----	--------	----	--

————— RADIATIVE DECAYS —————

$\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$   $\Gamma_{50}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.194 ± 0.008 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.199 ± 0.005 ± 0.012	<sup>64</sup> ADAM	05A	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$
-----------------------	--------------------	-----	---

<sup>64</sup> Uses  $B(\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow \gamma \gamma J/\psi)$  from ADAM 05A and  $B(\psi(2S) \rightarrow \gamma \chi_{c2})$  from ATHAR 04.

$\Gamma(\gamma \rho^0)/\Gamma_{\text{total}}$   $\Gamma_{51}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<b>&lt;50</b>	90	17.2 ± 6.8	<sup>65</sup> BENNETT	08A	CLEO $\psi(2S) \rightarrow \gamma \gamma \rho^0$
---------------	----	------------	-----------------------	-----	--

<sup>65</sup> BENNETT 08A reports  $< 50 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$ . We rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .

$\Gamma(\gamma\omega)/\Gamma_{\text{total}}$						$\Gamma_{52}/\Gamma$
VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<7	90	$0.0 \pm 1.8$	<sup>66</sup> BENNETT	08A	CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$
<sup>66</sup> BENNETT 08A reports $< 7.0 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$ . We rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .						

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$						$\Gamma_{53}/\Gamma$
VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<12	90	$1.3 \pm 2.5$	<sup>67</sup> BENNETT	08A	CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$
<sup>67</sup> BENNETT 08A reports $< 13 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$ . We rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 8.69 \times 10^{-2}$ .						

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$		$\Gamma_{54}/\Gamma$
VALUE (units $10^{-4}$ )	DOCUMENT ID	
<b>2.60 ± 0.16 OUR FIT</b>		

$\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$				$\Gamma_{54}/\Gamma_{50}$
VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT	
<b>1.34 ± 0.10 OUR FIT</b>				
<b>0.99 ± 0.18</b>	<sup>68</sup> AMBROGIANI	00B	E835	$\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$
<sup>68</sup> Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$ .				

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}} \times \Gamma(p\bar{p})/\Gamma_{\text{total}}$				$\Gamma_{54}/\Gamma \times \Gamma_{36}/\Gamma$
VALUE (units $10^{-8}$ )	DOCUMENT ID	TECN	COMMENT	
<b>1.87 ± 0.18 OUR FIT</b>				
<b>1.7 ± 0.4 OUR AVERAGE</b>				
1.60 ± 0.42	ARMSTRONG	93	E760	$\bar{p}p \rightarrow \gamma\gamma X$
9.9 ± 4.5	BAGLIN	87B	SPEC	$\bar{p}p \rightarrow \gamma\gamma X$

### $\chi_{c2}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$				$\Gamma_{13}/\Gamma \times \Gamma_{104}^{\psi(2S)}/\Gamma_9^{\psi(2S)}$
VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT	
<b>2.37 ± 0.27 OUR FIT</b>				
<b>2.5 ± 0.9 OUR AVERAGE</b>				Error includes scale factor of 2.3.
1.90 ± 0.14 ± 0.44	BAI	99B	BES	$\psi(2S) \rightarrow \gamma\chi_{c2}$
3.8 ± 0.67	<sup>69</sup> TANENBAUM	78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c2}$
<sup>69</sup> The reported value is derived using $B(\psi(2S) \rightarrow \pi^+\pi^-J/\psi) \times B(J/\psi \rightarrow \ell^+\ell^-) = (4.6 \pm 0.7)\%$ . Calculated by us using $B(J/\psi \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$ .				



$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{\text{total}}} \times \frac{\Gamma_{15} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{\psi(2S)}}{\Gamma_{104}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.1 ± 0.4 OUR FIT</b>			
<b>3.11 ± 0.36 ± 0.48</b>	ABLIKIM	04H BES2	$\psi(2S) \rightarrow \gamma \chi_{c2}$

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow p \bar{p}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{J / \psi(1S) \pi^+ \pi^-} \times \frac{\Gamma_{36} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{\psi(2S)}}{\Gamma_{104}^{\psi(2S)} / \Gamma_9^{\psi(2S)}}$$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.89 ± 0.14 OUR FIT</b>			
<b>1.4 ± 1.1</b>	<sup>70</sup> BAI	98I BES	$\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow \gamma \bar{p} p$

<sup>70</sup> Calculated by us. The value for  $B(\chi_{c2} \rightarrow p \bar{p})$  reported in BAI 98I is derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow p \bar{p}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{36} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.3 ± 0.5 OUR FIT</b>				
<b>6.7 ± 1.1 OUR AVERAGE</b>				Error includes scale factor of 1.5.

7.2 ± 0.7 ± 0.4	121 ± 12	<sup>71</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$
4.4 <sup>+1.6</sup> <sub>-1.4</sub> ± 0.6	14.3 <sup>+5.2</sup> <sub>-4.7</sub>	BAI	04F BES	$\psi(2S) \rightarrow \gamma \chi_{c2}(1P) \rightarrow \gamma \bar{p} p$

<sup>71</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c2} \rightarrow p \bar{p}) = (7.7 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$ .

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Lambda \bar{\Lambda}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{42} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16.2 ± 2.3 OUR FIT</b>				
<b>15.9 ± 2.1 ± 1.0</b>	71 ± 9	<sup>72</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$

<sup>72</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c2} \rightarrow \Lambda \bar{\Lambda}) = (17.0 \pm 2.2 \pm 1.1 \pm 1.1) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$ .

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Lambda \bar{\Lambda}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{J / \psi(1S) \pi^+ \pi^-} \times \frac{\Gamma_{42} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{\psi(2S)}}{\Gamma_{104}^{\psi(2S)} / \Gamma_9^{\psi(2S)}}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.9 ± 0.7 OUR FIT</b>				
<b>7.1<sup>+3.1</sup><sub>-2.9</sub> ± 1.3</b>	8.3 <sup>+3.7</sup> <sub>-3.4</sub>	<sup>73</sup> BAI	03E BES	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$

<sup>73</sup> BAI 03E reports  $[B(\chi_{c2} \rightarrow \Lambda \bar{\Lambda}) B(\psi(2S) \rightarrow \gamma \chi_{c2}) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)] \times [B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p \bar{p})] = (1.33<sup>+0.59</sup><sub>-0.55</sub> \pm 0.25)\%$ . We calculate from this measurement the presented value using  $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$  and  $B(J/\psi \rightarrow p \bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$ .

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{24} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{9}^{\psi(2S)}}$$

VALUE (units 10<sup>-3</sup>)      EVTS      DOCUMENT ID      TECN      COMMENT

**0.199 ± 0.034 OUR FIT**

**0.190 ± 0.034 ± 0.019**      115 ± 13      74 BAI      98I BES       $\psi(2S) \rightarrow \gamma K^+ K^-$

<sup>74</sup> Calculated by us. The value for  $B(\chi_{c2} \rightarrow K^+ K^-)$  reported by BAI 98I is derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{25} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{9}^{\psi(2S)}}$$

VALUE (units 10<sup>-5</sup>)      EVTS      DOCUMENT ID      TECN      COMMENT

**5.4 ± 0.7 OUR FIT**

**5.72 ± 0.76 ± 0.63**      65      ABLIKIM      050 BES2       $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{25} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{9}^{\psi(2S)}}$$

VALUE (units 10<sup>-5</sup>)      DOCUMENT ID      TECN      COMMENT

**16.3 ± 2.0 OUR FIT**

**14.7 ± 4.1 ± 3.3**      75 BAI      99B BES       $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

<sup>75</sup> Calculated by us. The value of  $B(\chi_{c2} \rightarrow K_S^0 K_S^0)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{50} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{9}^{\psi(2S)}}$$

VALUE (units 10<sup>-2</sup>)      EVTS      DOCUMENT ID      TECN      COMMENT

**1.69 ± 0.04 OUR FIT**

**1.34 ± 0.14 OUR AVERAGE**      Error includes scale factor of 1.9. See the ideogram below.

1.62 ± 0.04 ± 0.12      5.8k      BAI      04I BES2       $\psi(2S) \rightarrow J/\psi \gamma \gamma$

0.99 ± 0.10 ± 0.08           GAISER      86 CBAL       $\psi(2S) \rightarrow \gamma X$

1.47 ± 0.17      76 OREGLIA      82 CBAL       $\psi(2S) \rightarrow \gamma \chi_{c2}$

1.8 ± 0.5      77 BRANDELIK      79B DASP       $\psi(2S) \rightarrow \gamma \chi_{c2}$

1.2 ± 0.2      77 BARTEL      78B CNTR       $\psi(2S) \rightarrow \gamma \chi_{c2}$

2.2 ± 1.2      78 BIDDICK      77 CNTR       $e^+ e^- \rightarrow \gamma X$

1.2 ± 0.7      76 WHITAKER      76 MRK1       $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.95 ± 0.02 ± 0.07      12.4k      79 MENDEZ      08 CLEO       $\psi(2S) \rightarrow \gamma \chi_{c2}$

1.85 ± 0.04 ± 0.07      1.9k      80 ADAM      05A CLEO      Repl. by MENDEZ 08


<sup>76</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

<sup>77</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$ .

<sup>78</sup> Assumes isotropic gamma distribution.

<sup>79</sup> Not independent from other measurements of MENDEZ 08.

<sup>80</sup> Not independent from other values reported by ADAM 05A.



$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}} \text{ (units } 10^{-2}\text{)}$$

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{ anything})$$

$$\Gamma_{50}/\Gamma \times \Gamma_{104}^{\psi(2S)}/\Gamma_7^{\psi(2S)} = \Gamma_{50}/\Gamma \times \Gamma_{104}^{\psi(2S)}/(\Gamma_9^{\psi(2S)} + \Gamma_{10}^{\psi(2S)} + \Gamma_{11}^{\psi(2S)}) + 0.341\Gamma_{103}^{\psi(2S)} + 0.194\Gamma_{104}^{\psi(2S)}$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**2.88±0.07 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.12±0.03±0.09	12.4k	<sup>81</sup> MENDEZ	08	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
3.11±0.07±0.07	1.9k	ADAM	05A	CLEO	Repl. by MENDEZ 08

<sup>81</sup> Not independent from other measurements of MENDEZ 08.

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_{50}/\Gamma \times \Gamma_{104}^{\psi(2S)}/\Gamma_9^{\psi(2S)}$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**5.10±0.13 OUR FIT**

**5.53±0.17 OUR AVERAGE**

5.56±0.05±0.16	12.4k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
6.0 ±2.8	1.3k	<sup>82</sup> ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
3.9 ±1.2		<sup>83</sup> HIMEL	80	MRK2	$\psi(2S) \rightarrow \gamma \chi_{c2}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.52±0.13±0.13	1.9k	<sup>84</sup> ADAM	05A	CLEO	Repl. by MENDEZ 08
----------------	------	--------------------	-----	------	--------------------

<sup>82</sup> From a fit to the  $J/\psi$  recoil mass spectra.

<sup>83</sup> The value for  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) \times B(\chi_{c2} \rightarrow \gamma J/\psi(1S))$  reported in HIMEL 80 is derived using  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$  and  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.138 \pm 0.018$ . Calculated by us using  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = (0.1181 \pm 0.0020)$ .

<sup>84</sup> Not independent from other values reported by ADAM 05A.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma\gamma)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))}{\Gamma_{\text{total}}} \quad \frac{\Gamma_{54}/\Gamma \times \Gamma_{104}^{\psi(2S)}/\Gamma_{\psi(2S)}}{\Gamma_{104}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.26 ± 0.16 OUR FIT</b>				
<b>2.73 ± 0.32 OUR AVERAGE</b>				
2.68 ± 0.28 ± 0.15	333 ± 35	ECKLUND	08A CLEO	$\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow 3\gamma$
7.0 ± 2.1 ± 2.0		LEE	85 CBAL	$\psi(2S) \rightarrow \gamma\chi_{c2}$

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)} \quad \frac{\Gamma_{19}/\Gamma \times \Gamma_{104}^{\psi(2S)}/\Gamma_9^{\psi(2S)}}{\Gamma_{104}^{\psi(2S)}/\Gamma_9^{\psi(2S)}}$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.55 ± 0.06 OUR FIT</b>				
<b>0.54 ± 0.06 OUR AVERAGE</b>				
0.66 ± 0.18 ± 0.37	21 ± 6	<sup>85</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$
0.54 ± 0.05 ± 0.04	185 ± 16	<sup>86</sup> BAI	98I BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$

<sup>85</sup> We have multiplied  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$ .

<sup>86</sup> Calculated by us. The value for  $B(\chi_{c2} \rightarrow \pi^+\pi^-)$  reported by BAI 98I is derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D]. We have multiplied  $\pi^+\pi^-$  measurement by 3/2 to obtain  $\pi\pi$ .

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow 2(\pi^+\pi^-))}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)} \quad \frac{\Gamma_1/\Gamma \times \Gamma_{104}^{\psi(2S)}/\Gamma_9^{\psi(2S)}}{\Gamma_{104}^{\psi(2S)}/\Gamma_9^{\psi(2S)}}$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.87 ± 0.27 OUR FIT</b>			
<b>3.1 ± 1.0 OUR AVERAGE</b>			Error includes scale factor of 2.5.
2.3 ± 0.1 ± 0.5	<sup>87</sup> BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c2}$
4.3 ± 0.6	<sup>88</sup> TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c2}$

<sup>87</sup> Calculated by us. The value for  $B(\chi_{c2} \rightarrow 2\pi^+2\pi^-)$  reported in BAI 99B is derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

<sup>88</sup> The value for  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) \times B(\chi_{c2} \rightarrow 2\pi^+\pi^-)$  reported in TANENBAUM 78 is derived using  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times B(J/\psi(1S)\ell^+\ell^-) = (4.6 \pm 0.7)\%$ . Calculated by us using  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$ .

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-K^+K^-)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))}{\Gamma_{\text{total}}} \quad \frac{\Gamma_{33}/\Gamma \times \Gamma_{104}^{\psi(2S)}/\Gamma_{\psi(2S)}}{\Gamma_{104}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.54 ± 0.18 OUR FIT</b>				
<b>1.76 ± 0.16 ± 0.24</b>	160	<sup>89</sup> ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+2K^-$

<sup>89</sup> Calculated by us. The value of  $B(\chi_{c2} \rightarrow 2K^+2K^-)$  reported by ABLIKIM 06T was derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4)\%$ .

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{33} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{9}^{\psi(2S)}}$$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**4.6 ± 0.6 OUR FIT**

<b>3.6 ± 0.6 ± 0.6</b>	90 BAI	99B	BES $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
------------------------	--------	-----	---

<sup>90</sup> Calculated by us. The value of  $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{17} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{9}^{\psi(2S)}}$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.28 ± 0.24 OUR FIT**

<b>1.38 ± 0.24 ± 0.23</b>	41	91 ABLIKIM	06T	BES2 $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
---------------------------	----	------------	-----	--

<sup>91</sup> Calculated by us. The value of  $B(\chi_{c2} \rightarrow \phi\phi)$  reported by ABLIKIM 06T was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$ .

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{17} / \Gamma \times \Gamma_{104}^{\psi(2S)} / \Gamma_{9}^{\psi(2S)}}$$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**3.9 ± 0.7 OUR FIT**

<b>4.8 ± 1.3 ± 1.3</b>	92 BAI	99B	BES $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
------------------------	--------	-----	---

<sup>92</sup> Calculated by us. The value of  $B(\chi_{c2} \rightarrow \phi\phi)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

## MULTIPOLE AMPLITUDES IN $\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$ RADIATIVE DECAY

**$a_2 = M_2 / \sqrt{E_1^2 + M_2^2 + E_3^2}$  Magnetic quadrupole fractional transition amplitude**


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**-0.13 ± 0.05 OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.

$-0.093^{+0.039}_{-0.041} \pm 0.006$	5908	93 AMBROGIANI 02	E835	$\rho\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
--------------------------------------	------	------------------	------	--

$-0.14 \pm 0.06$	1904	93 ARMSTRONG 93E	E760	$\rho\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
------------------	------	------------------	------	--

$-0.333^{+0.116}_{-0.292}$	441	93 OREGLIA 82	CBAL	$\psi(2S) \rightarrow \chi_{c1}\gamma \rightarrow J/\psi\gamma\gamma$
----------------------------	-----	---------------	------	---



**$a_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$  Electric octupole fractional transition amplitude**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.011^{+0.040}_{-0.033}</math> OUR AVERAGE</b>				
$0.020^{+0.055}_{-0.044} \pm 0.009$	5908	AMBROGIANI 02	E835	$\rho\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
$0.00^{+0.06}_{-0.05}$	1904	ARMSTRONG 93E	E760	$\rho\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$

<sup>93</sup> Assuming  $a_3=0$ .

### $\chi_{c2}(1P)$ REFERENCES

BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)
ECKLUND	08A	PR D78 091501R	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102R	H. Mendez <i>et al.</i>	(CLEO Collab.)
NAIK	08	PR D78 031101R	P. Naik <i>et al.</i>	(CLEO Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
ADAMS	07	PR D75 071101R	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
CHEN	07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)
DOBBS	06	PR D73 071101R	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)

NAKAZAWA	05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)
ABE	02T	PL B540 33	K. Abe <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
EISENSTEIN	01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)
AMBROGIANI	00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ACCIARRI	99E	PL B453 73	M. Acciarri <i>et al.</i>	(L3 Collab.)
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
ACKER...,K...	98	PL B439 197	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)
DOMINICK	94	PR D50 4265	J. Dominick <i>et al.</i>	(CLEO Collab.)
ARMSTRONG	93	PRL 70 2988	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
ARMSTRONG	93E	PR D48 3037	T.A. Armstrong <i>et al.</i>	(FNAL-E760 Collab.)
BAUER	93	PL B302 345	D.A. Bauer <i>et al.</i>	(TPC Collab.)
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BAGLIN	87B	PL B187 191	C. Baglin <i>et al.</i>	(R704 Collab.)
BAGLIN	86B	PL B172 455	C. Baglin	(LAPP, CERN, GENO, LYON, OSLO+)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
Also		Private Comm.	M.J. Oreglia	(EFI)
BARATE	81	PR D24 2994	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, CERN+)
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)

### OTHER RELATED PAPERS

ABLIKIM	04I	PR D70 092004	M. Ablikim <i>et al.</i>	(BES Collab.)
BARBERIS	00G	PL B485 357	D. Barberis <i>et al.</i>	(Omega Expt.)
ACCIARRI	99T	PL B461 155	M. Acciarri <i>et al.</i>	(L3 Collab.)
CHEN	90B	PL B243 169	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
AIHARA	88D	PRL 60 2355	H. Aihara <i>et al.</i>	(TPC Collab.)
BARATE	83	PL 121B 449	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, IND)
FELDMAN	75B	PRL 35 821	G.J. Feldman <i>et al.</i>	(LBL, SLAC)
Also		PRL 35 1184 (errat.)	G.J. Feldman <i>et al.</i>	(LBL, SLAC)
TANENBAUM	75	PRL 35 1323	W.M. Tanenbaum <i>et al.</i>	(LBL, SLAC)