

$\eta_c(2S)$

$I^G(J^{PC}) = 0^+(0^{-+})$

Quantum numbers are quark model predictions.


$\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3637 ±4 OUR AVERAGE				Error includes scale factor of 1.7. See the ideogram below.
3626 ±5 ±6	311	1 ABE	07 BELL	$e^+ e^- \rightarrow J/\psi(c\bar{c})$
3645.0 ±5.5 ±4.9	121 ± 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
3642.9 ±3.1 ±1.5	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
3630.8 ±3.4 ±1.0	112 ± 24	AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
3654 ±6 ±8	39 ± 11	CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3639 ±7	98 ± 52	2 AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X c\bar{c}$
3594 ±5		3 EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

¹ From a fit of the J/ψ recoil mass spectrum. Supersedes ABE, K 02 and ABE 04G.

² From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

³ Assuming mass of $\psi(2S) = 3686$ MeV.



$\eta_c(2S)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
14 ± 7 OUR AVERAGE					
6.3±12.4±4.0		61	ASNER	04	CLEO $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
17.0± 8.3±2.5	112 ± 24		AUBERT	04D	BABR $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<23	90	98 ± 52	⁴ AUBERT	06E	BABR $B^\pm \rightarrow K^\pm X_c \bar{c}$
22 ± 14		121 ± 27	AUBERT	05C	BABR $e^+ e^- \rightarrow J/\psi c \bar{c}$
<55	90	39 ± 11	⁵ CHOI	02	BELL $B \rightarrow K K_S K^- \pi^+$
<8.0	95		⁶ EDWARDS	82C	CBAL $e^+ e^- \rightarrow \gamma X$

⁴ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.
⁵ For a mass value of 3654 ± 6 MeV
⁶ For a mass value of 3594 ± 5 MeV

$\eta_c(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	not seen	
Γ_2 $K\bar{K}\pi$	(1.9 ± 1.2) %	
Γ_3 $2\pi^+ 2\pi^-$	not seen	
Γ_4 $K^+ K^- \pi^+ \pi^-$	not seen	
Γ_5 $2K^+ 2K^-$	not seen	
Γ_6 $p\bar{p}$	not seen	
Γ_7 $\gamma\gamma$	$< 5 \times 10^{-4}$	90%

$\eta_c(2S)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	Γ_7
1.3±0.6	⁷ ASNER 04 CLEO $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
⁷ They measure $\Gamma(\eta_c(2S)\gamma\gamma)$ $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02)$ $\Gamma(\eta_c(1S)\gamma\gamma)$ $B(\eta_c(1S) \rightarrow K\bar{K}\pi)$. The value for $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$ is derived assuming that the branching fractions for $\eta_c(2S)$ and $\eta_c(1S)$ decays to $K_S K\pi$ are equal and using $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$ keV.	

$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(2\pi^+ 2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3 \Gamma_7 / \Gamma$
VALUE (eV)	CL\%
<6.5	90
UEHARA	08
BELL	

$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$			$\Gamma_4 \Gamma_7/\Gamma$	
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(2K^+ 2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$			$\Gamma_5 \Gamma_7/\Gamma$	
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+ K^-)$

$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$			$\Gamma_6/\Gamma \times \Gamma_7/\Gamma$	
VALUE (units 10^{-8})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.6	90	8,9,10	AMBROGIANI 01	E835 $\bar{p}p \rightarrow \gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 8.0	90	8,9,11	AMBROGIANI 01	E835 $\bar{p}p \rightarrow \gamma\gamma$
<12.0	90	9,11	AMBROGIANI 01	E835 $\bar{p}p \rightarrow \gamma\gamma$

⁸ Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

⁹ For a total width $\Gamma=5$ MeV.

¹⁰ For the resonance mass region $3589\text{--}3599$ MeV/ c^2 .

¹¹ For the resonance mass region $3575\text{--}3660$ MeV/ c^2 .

$\eta_c(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$			Γ_1/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	ABREU	980	DLPH	$e^+ e^- \rightarrow e^+ e^- + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	12 EDWARDS	82C	CBAL	$e^+ e^- \rightarrow \gamma X$

¹² For a mass value of 3594 ± 5 MeV

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$			Γ_2/Γ	
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.9±0.4±1.1	59 ± 12	13 AUBERT	08AB	BABR $B \rightarrow \eta_c(2S) K \rightarrow K\bar{K}\pi K$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	39 ± 11	14 CHOI	02	BELL $B \rightarrow K K_S K^- \pi^+$

¹³ Derived from a measurement of $[B(B^+ \rightarrow \eta_c(2S) K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$ and using $B(B^+ \rightarrow \eta_c(2S) K^+) = (3.4 \pm 1.8) \times 10^{-4}$, and $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$.

¹⁴ For a mass value of 3654 ± 6 MeV

$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$			Γ_3/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

$\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_4/Γ				
<u>VALUE</u>					
not seen					
<u>DOCUMENT ID</u>					
UEHARA	08				
<u>TECN</u>					
BELL					
<u>COMMENT</u>					
$\gamma\gamma \rightarrow \eta_c(2S)$					
$\Gamma(2K^+2K^-)/\Gamma_{\text{total}}$	Γ_5/Γ				
<u>VALUE</u>					
not seen					
<u>DOCUMENT ID</u>					
UEHARA	08				
<u>TECN</u>					
BELL					
<u>COMMENT</u>					
$\gamma\gamma \rightarrow \eta_c(2S)$					
<u>VALUE</u>					
not seen					
<u>DOCUMENT ID</u>					
AMBROGIANI	01				
<u>TECN</u>					
E835					
<u>COMMENT</u>					
$\bar{p}p \rightarrow \gamma\gamma$					
$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	Γ_7/Γ				
<u>VALUE</u>	<u>CL%</u>				
<0.0005	90				
15	WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.01	90	LEE	85	CBAL	$\psi' \rightarrow \text{photons}$
¹⁵ WICHT 08 reports $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S)K^+)] < 0.18 \times 10^{-6}$. We divide by our best value $B(B^+ \rightarrow \eta_c(2S)K^+) = 3.4 \times 10^{-4}$.					

$\eta_c(2S)$ REFERENCES

AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101R	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)

OTHER RELATED PAPERS

PENNINGTON	07A	PR D76 077502	M.R. Pennington, DJ. Wilson	
SWANSON	06	PRPL 429 243	E.S. Swanson	(PITT)
BADALIAN	03	PR D67 071901	A.M. Badalian, B.L.G. Bakker	
EICHEN	02	PRL 89 162002	E.J. Eichten, K. Lane, C. Quigg	
ACCIARRI	99T	PL B461 155	M. Acciari <i>et al.</i>	(L3 Collab.)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
PORTER	81	SLAC Summer Inst. 355	F.C. Porter <i>et al.</i>	(CIT, HARV, PRIN+)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)