

$\Sigma(2030) F_{17}$

$I(J^P) = 1(\frac{7}{2}^+)$ Status: ****

Discovered by COOL 66 and by WOHL 66. For most results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters **111B** 1 (1982).

This entry only includes results from partial-wave analyses. Parameters of peaks seen in cross sections and invariant-mass distributions around 2030 MeV may be found in our 1984 edition, Reviews of Modern Physics **56** S1 (1984).

$\Sigma(2030)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|--|
| 2025 to 2040 (\approx 2030) OUR ESTIMATE | | | |
| 2036 \pm 5 | GOPAL | 80 | DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| 2038 \pm 10 | CORDEN | 77B | $K^- N \rightarrow N\bar{K}^*$ |
| 2040 \pm 5 | GOPAL | 77 | DPWA $\bar{K}N$ multichannel |
| 2030 \pm 3 | ¹ CORDEN | 76 | DPWA $K^- n \rightarrow \Lambda\pi^-$ |
| 2035 \pm 15 | BAILLON | 75 | IPWA $\bar{K}N \rightarrow \Lambda\pi$ |
| 2038 \pm 10 | HEMINGWAY | 75 | DPWA $K^- p \rightarrow \bar{K}N$ |
| 2042 \pm 11 | VANHORN | 75 | DPWA $K^- p \rightarrow \Lambda\pi^0$ |
| 2020 \pm 6 | KANE | 74 | DPWA $K^- p \rightarrow \Sigma\pi$ |
| 2035 \pm 10 | LITCHFIELD | 74B | DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$ |
| 2020 \pm 30 | LITCHFIELD | 74C | DPWA $K^- p \rightarrow \Delta(1232)\bar{K}$ |
| 2025 \pm 10 | LITCHFIELD | 74D | DPWA $K^- p \rightarrow \Lambda(1820)\pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 2027 to 2057 | GOYAL | 77 | DPWA $K^- N \rightarrow \Sigma\pi$ |
| 2030 | DEBELLEFON | 76 | IPWA $K^- p \rightarrow \Lambda\pi^0$ |

$\Sigma(2030)$ WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|--|
| 150 to 200 (\approx 180) OUR ESTIMATE | | | |
| 172 \pm 10 | GOPAL | 80 | DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| 137 \pm 40 | CORDEN | 77B | $K^- N \rightarrow N\bar{K}^*$ |
| 190 \pm 10 | GOPAL | 77 | DPWA $\bar{K}N$ multichannel |
| 201 \pm 9 | ¹ CORDEN | 76 | DPWA $K^- n \rightarrow \Lambda\pi^-$ |
| 180 \pm 20 | BAILLON | 75 | IPWA $\bar{K}N \rightarrow \Lambda\pi$ |
| 172 \pm 15 | HEMINGWAY | 75 | DPWA $K^- p \rightarrow \bar{K}N$ |
| 178 \pm 13 | VANHORN | 75 | DPWA $K^- p \rightarrow \Lambda\pi^0$ |
| 111 \pm 5 | KANE | 74 | DPWA $K^- p \rightarrow \Sigma\pi$ |
| 160 \pm 20 | LITCHFIELD | 74B | DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$ |
| 200 \pm 30 | LITCHFIELD | 74C | DPWA $K^- p \rightarrow \Delta(1232)\bar{K}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------|------------|-----|------|---|
| 260 | DECLAIS | 77 | DPWA | $\bar{K}N \rightarrow \bar{K}N$ |
| 126 to 195 | GOYAL | 77 | DPWA | $K^- N \rightarrow \Sigma \pi$ |
| 160 | DEBELLEFON | 76 | IPWA | $K^- p \rightarrow \Lambda \pi^0$ |
| 70 to 125 | LITCHFIELD | 74D | DPWA | $K^- p \rightarrow \Lambda(1820) \pi^0$ |

$\Sigma(2030)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|--|--------------------------------|
| $\Gamma_1 N\bar{K}$ | 17–23 % |
| $\Gamma_2 \Lambda\pi$ | 17–23 % |
| $\Gamma_3 \Sigma\pi$ | 5–10 % |
| $\Gamma_4 \Xi K$ | <2 % |
| $\Gamma_5 \Sigma(1385)\pi$ | 5–15 % |
| $\Gamma_6 \Sigma(1385)\pi$, <i>F</i> -wave | |
| $\Gamma_7 \Lambda(1520)\pi$ | 10–20 % |
| $\Gamma_8 \Lambda(1520)\pi$, <i>D</i> -wave | |
| $\Gamma_9 \Lambda(1520)\pi$, <i>G</i> -wave | |
| $\Gamma_{10} \Delta(1232)\bar{K}$ | 10–20 % |
| $\Gamma_{11} \Delta(1232)\bar{K}$, <i>F</i> -wave | |
| $\Gamma_{12} \Delta(1232)\bar{K}$, <i>H</i> -wave | |
| $\Gamma_{13} N\bar{K}^*(892)$ | <5 % |
| $\Gamma_{14} N\bar{K}^*(892)$, $S=1/2$, <i>F</i> -wave | |
| $\Gamma_{15} N\bar{K}^*(892)$, $S=3/2$, <i>F</i> -wave | |
| $\Gamma_{16} \Lambda(1820)\pi$, <i>P</i> -wave | |

The above branching fractions are our estimates, not fits or averages.

$\Sigma(2030)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

| $\Gamma(N\bar{K})/\Gamma_{\text{total}}$ | Γ_1/Γ |
|---|---|
| <u>VALUE</u> | <u>DOCUMENT ID</u> |
| 0.17 to 0.23 OUR ESTIMATE | <u>TECN</u> |
| 0.19±0.03 | GOPAL 80 DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| 0.18±0.03 | HEMINGWAY 75 DPWA $K^- p \rightarrow \bar{K}N$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | |
| 0.15 | DECLAIS 77 DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| 0.24±0.02 | GOPAL 77 DPWA See GOPAL 80 |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda\pi$ $(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------|---------------------|------|---|
| +0.18 ± 0.02 | GOPAL | 77 | DPWA $\bar{K}N$ multichannel |
| +0.20 ± 0.01 | ¹ CORDEN | 76 | DPWA $K^- n \rightarrow \Lambda\pi^-$ |
| +0.18 ± 0.02 | BAILLON | 75 | IPWA $\bar{K}N \rightarrow \Lambda\pi$ |
| +0.20 ± 0.01 | VANHORN | 75 | DPWA $K^- p \rightarrow \Lambda\pi^0$ |
| +0.195 ± 0.053 | DEVENISH | 74B | Fixed- t dispersion rel. • • • We do not use the following data for averages, fits, limits, etc. • • • |
| 0.20 | DEBELLEFON | 76 | IPWA $K^- p \rightarrow \Lambda\pi^0$ |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Sigma\pi$ $(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|------------------------------------|
| -0.09 ± 0.01 | ² CORDEN | 77C | $K^- n \rightarrow \Sigma\pi$ |
| -0.06 ± 0.01 | ² CORDEN | 77C | $K^- n \rightarrow \Sigma\pi$ |
| -0.15 ± 0.03 | GOPAL | 77 | DPWA $\bar{K}N$ multichannel |
| -0.10 ± 0.01 | KANE | 74 | DPWA $K^- p \rightarrow \Sigma\pi$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| -0.085 ± 0.02 | ³ GOYAL | 77 | DPWA $K^- N \rightarrow \Sigma\pi$ |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Xi K$ $(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|--------------------------------|
| 0.023 | MULLER | 69B | DPWA $K^- p \rightarrow \Xi K$ |
| <0.05 | BURGUN | 68 | DPWA $K^- p \rightarrow \Xi K$ |
| <0.05 | TRIPP | 67 | RVUE $K^- p \rightarrow \Xi K$ |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda(1820)\pi$, *P-wave* $(\Gamma_1 \Gamma_{16})^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|---|
| 0.14 ± 0.02 | CORDEN | 75B | DBC $K^- n \rightarrow N\bar{K}\pi^-$ |
| 0.18 ± 0.04 | LITCHFIELD | 74D | DPWA $K^- p \rightarrow \Lambda(1820)\pi^0$ |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda(1520)\pi$, *D-wave* $(\Gamma_1 \Gamma_8)^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|------|---|
| +0.114 ± 0.010 | ⁴ CAMERON | 77 | DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$ |
| 0.14 ± 0.03 | LITCHFIELD | 74B | DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.10 ± 0.03 | ⁵ CORDEN | 75B | DBC $K^- n \rightarrow N\bar{K}\pi^-$ |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda(1520)\pi$, *G-wave* $(\Gamma_1 \Gamma_9)^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------|----------------------|------|---|
| +0.146 ± 0.010 | ⁴ CAMERON | 77 | DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$ |
| 0.02 ± 0.02 | LITCHFIELD | 74B | DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$ |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Delta(1232)\bar{K}$, *F-wave* $(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|---------------------|------|--|
| 0.16 ± 0.03 | LITCHFIELD | 74C | DPWA $K^- p \rightarrow \Delta(1232)\bar{K}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.17 ± 0.03 | ⁵ CORDEN | 75B | DBC $K^- n \rightarrow N\bar{K}\pi^-$ |

| $(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Delta(1232)\bar{K}$, H-wave | $(\Gamma_1 \Gamma_{12})^{1/2} / \Gamma$ | | |
|--|---|-------------|--|
| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.00 \pm 0.02 | LITCHFIELD | 74C | DPWA $K^- p \rightarrow \Delta(1232)\bar{K}$ |
| $(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Sigma(1385)\pi$ | $(\Gamma_1 \Gamma_5)^{1/2} / \Gamma$ | | |
| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| +0.153 \pm 0.026 | ⁴ CAMERON | 78 | DPWA $K^- p \rightarrow \Sigma(1385)\pi$ |
| $(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow N\bar{K}^*(892)$, S=1/2, F-wave | $(\Gamma_1 \Gamma_{14})^{1/2} / \Gamma$ | | |
| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| +0.06 \pm 0.03 | ⁴ CAMERON | 78B | DPWA $K^- p \rightarrow N\bar{K}^*$ |
| -0.02 \pm 0.01 | CORDEN | 77B | $K^- d \rightarrow NN\bar{K}^*$ |
| $(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow N\bar{K}^*(892)$, S=3/2, F-wave | $(\Gamma_1 \Gamma_{15})^{1/2} / \Gamma$ | | |
| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| +0.04 \pm 0.03 | ⁶ CAMERON | 78B | DPWA $K^- p \rightarrow N\bar{K}^*$ |
| -0.12 \pm 0.02 | CORDEN | 77B | $K^- d \rightarrow NN\bar{K}^*$ |

$\Sigma(2030)$ FOOTNOTES

¹ Preferred solution 3; see CORDEN 76 for other possibilities.

² The two entries for CORDEN 77C are from two different acceptable solutions.

³ This coupling is extracted from unnormalized data.

⁴ The published sign has been changed to be in accord with the baryon-first convention.

⁵ An upper limit.

⁶ The upper limit on the G_3 wave is 0.03.

$\Sigma(2030)$ REFERENCES

| | | | | |
|------------|-----|-------------------|---|-------------------------|
| PDG | 84 | RMP 56 S1 | C.G. Wohl <i>et al.</i> | (LBL, CIT, CERN) |
| PDG | 82 | PL 111B 1 | M. Roos <i>et al.</i> | (HELS, CIT, CERN) |
| GOPAL | 80 | Toronto Conf. 159 | G.P. Gopal | (RHEL) IJP |
| CAMERON | 78 | NP B143 189 | W. Cameron <i>et al.</i> | (RHEL, LOIC) IJP |
| CAMERON | 78B | NP B146 327 | W. Cameron <i>et al.</i> | (RHEL, LOIC) IJP |
| CAMERON | 77 | NP B131 399 | W. Cameron <i>et al.</i> | (RHEL, LOIC) IJP |
| CORDEN | 77B | NP B121 365 | M.J. Corden <i>et al.</i> | (BIRM) IJP |
| CORDEN | 77C | NP B125 61 | M.J. Corden <i>et al.</i> | (BIRM) IJP |
| DECLAIS | 77 | CERN 77-16 | Y. Declais <i>et al.</i> | (CAEN, CERN) IJP |
| GOPAL | 77 | NP B119 362 | G.P. Gopal <i>et al.</i> | (LOIC, RHEL) IJP |
| GOYAL | 77 | PR D16 2746 | D.P. Goyal, A.V. Sodhi | (DELH) IJP |
| CORDEN | 76 | NP B104 382 | M.J. Corden <i>et al.</i> | (BIRM) IJP |
| DEBELLEFON | 76 | NP B109 129 | A. de Bellefon, A. Berthon | (CDEF) IJP |
| BAILLON | 75 | NP B94 39 | P.H. Baillon, P.J. Litchfield | (CERN, RHEL) IJP |
| CORDEN | 75B | NP B92 365 | M.J. Corden <i>et al.</i> | (BIRM) IJP |
| HEMINGWAY | 75 | NP B91 12 | R.J. Hemingway <i>et al.</i> | (CERN, HEIDH, MPIM) IJP |
| VANHORN | 75 | NP B87 145 | A.J. van Horn | (LBL) IJP |
| Also | | NP B87 157 | A.J. van Horn | (LBL) IJP |
| DEVENISH | 74B | NP B81 330 | R.C.E. Devenish, C.D. Froggatt, B.R. Martin | (DESY+) (LBL) IJP |
| KANE | 74 | LBL-2452 | D.F. Kane | (LBL) IJP |

| | | | | |
|------------|-----|-------------------|---|--------------------|
| LITCHFIELD | 74B | NP B74 19 | P.J. Litchfield <i>et al.</i> | (CERN, HEIDH) IJP |
| LITCHFIELD | 74C | NP B74 39 | P.J. Litchfield <i>et al.</i> | (CERN, HEIDH) IJP |
| LITCHFIELD | 74D | NP B74 12 | P.J. Litchfield <i>et al.</i> | (CERN, HEIDH) IJP |
| MULLER | 69B | Thesis UCRL 19372 | R.A. Muller | (LRL) |
| BURGUN | 68 | NP B8 447 | G. Burgun <i>et al.</i> | (SACL, CDEF, RHEL) |
| TRIPP | 67 | NP B3 10 | R.D. Tripp <i>et al.</i> | (LRL, SLAC, CERN+) |
| COOL | 66 | PRL 16 1228 | R.L. Cool <i>et al.</i> | (BNL) |
| WOHL | 66 | PRL 17 107 | C.G. Wohl, F.T. Solmitz, M.L. Stevenson | (LRL) IJP |
