

**$\Sigma(3170)$  Bumps** $I(J^P) = 1(?^?)$  Status: \*

## OMITTED FROM SUMMARY TABLE

Seen by AMIRZADEH 79 as a narrow 6.5-standard-deviation enhancement in the reaction  $K^- p \rightarrow Y^{*+} \pi^-$  using data from independent high statistics bubble chamber experiments at 8.25 and 6.5 GeV/c. The dominant decay modes are multibody, multistrange final states and the production is via isospin-3/2 baryon exchange. Isospin 1 is favored.

Not seen in a  $K^- p$  experiment in LASS at 11 GeV/c (ASTON 85B).

 **$\Sigma(3170)$  MASS  
(PRODUCTION EXPERIMENTS)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>≈ 3170 OUR ESTIMATE</b>				
3170±5	35	AMIRZADEH 79	HBC	$K^- p \rightarrow Y^{*+} \pi^-$

 **$\Sigma(3170)$  WIDTH  
(PRODUCTION EXPERIMENTS)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<20	35	1 AMIRZADEH 79	HBC	$K^- p \rightarrow Y^{*+} \pi^-$

 **$\Sigma(3170)$  DECAY MODES  
(PRODUCTION EXPERIMENTS)**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \Lambda K \bar{K} \pi'$ 's	seen
$\Gamma_2 \Sigma K \bar{K} \pi'$ 's	seen
$\Gamma_3 \Xi K \pi'$ 's	seen

 **$\Sigma(3170)$  BRANCHING RATIOS  
(PRODUCTION EXPERIMENTS)**

$\Gamma(\Lambda K \bar{K} \pi')$ /Γ <sub>total</sub>	$\Gamma_1/\Gamma$
seen	AMIRZADEH 79 HBC $K^- p \rightarrow Y^{*+} \pi^-$

$\Gamma(\Sigma K \bar{K} \pi')$ /Γ <sub>total</sub>	$\Gamma_2/\Gamma$
seen	AMIRZADEH 79 HBC $K^- p \rightarrow Y^{*+} \pi^-$

$\Gamma(\Xi K\pi')$ / $\Gamma_{\text{total}}$	$\Gamma_3/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	AMIRZADEH 79	HBC	$K^- p \rightarrow Y^* + \pi^-$

## **$\Sigma(3170)$ FOOTNOTES (PRODUCTION EXPERIMENTS)**

<sup>1</sup> Observed width consistent with experimental resolution.

## **$\Sigma(3170)$ REFERENCES (PRODUCTION EXPERIMENTS)**

ASTON	85B	PR D32 2270	D. Aston <i>et al.</i>	(SLAC, CARL, CNRC, CINC)
AMIRZADEH	79	PL 89B 125	J. Amirzadeh <i>et al.</i>	(BIRM, CERN, GLAS+) I
Also		Toronto Conf. 263	J.B. Kinson <i>et al.</i>	(BIRM, CERN, GLAS+) I