

$\Upsilon(3S)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\Upsilon(3S)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
10.3552±0.0005	¹ ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10.3553±0.0005	^{2,3} BARU	86B REDE	$e^+ e^- \rightarrow$ hadrons
¹ Reanalysis of BARU 86B using new electron mass (COHEN 87).			
² Reanalysis of ARTAMONOV 84.			
³ Superseded by ARTAMONOV 00.			

$\Upsilon(3S)$ WIDTH

VALUE (keV)	DOCUMENT ID
20.32±1.85 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

$\Upsilon(3S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \Upsilon(2S)$ anything	(10.6 ± 0.8) %	
$\Gamma_2 \Upsilon(2S)\pi^+\pi^-$	(2.45 ± 0.23) %	S=1.1
$\Gamma_3 \Upsilon(2S)\pi^0\pi^0$	(1.85 ± 0.14) %	
$\Gamma_4 \Upsilon(2S)\gamma\gamma$	(5.0 ± 0.7) %	
$\Gamma_5 \Upsilon(2S)\pi^0$	< 5.1 × 10 ⁻⁴	CL=90%
$\Gamma_6 \Upsilon(1S)\pi^+\pi^-$	(4.40 ± 0.10) %	
$\Gamma_7 \Upsilon(1S)\pi^0\pi^0$	(2.20 ± 0.13) %	
$\Gamma_8 \Upsilon(1S)\eta$	< 1.8 × 10 ⁻⁴	CL=90%
$\Gamma_9 \Upsilon(1S)\pi^0$	< 7 × 10 ⁻⁵	CL=90%
$\Gamma_{10} \tau^+\tau^-$	(2.29 ± 0.30) %	
$\Gamma_{11} \mu^+\mu^-$	(2.18 ± 0.21) %	S=2.1
$\Gamma_{12} e^+e^-$	seen	

Radiative decays

$\Gamma_{13} \gamma\chi b_2(2P)$	(13.1 ± 1.6) %	S=3.4
$\Gamma_{14} \gamma\chi b_1(2P)$	(12.6 ± 1.2) %	S=2.4
$\Gamma_{15} \gamma\chi b_0(2P)$	(5.9 ± 0.6) %	S=1.4
$\Gamma_{16} \gamma\chi b_2(1P)$	< 1.9 %	CL=90%
$\Gamma_{17} \gamma\chi b_1(1P)$	< 1.7 × 10 ⁻³	CL=90%
$\Gamma_{18} \gamma\chi b_0(1P)$	(3.0 ± 1.1) × 10 ⁻³	
$\Gamma_{19} \gamma\eta_b(2S)$	< 6.2 × 10 ⁻⁴	CL=90%
$\Gamma_{20} \gamma\eta_b(1S)$	(4.8 ± 1.3) × 10 ⁻⁴	
$\Gamma_{21} \gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 2.2 × 10 ⁻⁴	CL=95%

Lepton Flavor (*LF*) violating decays

$$\Gamma_{22} \quad \mu^\pm \tau^\mp \quad LF \quad < 2.03 \quad \times 10^{-5} \quad CL=95\%$$

[a] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

$\Upsilon(3S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_0\Gamma_{12}/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

0.414±0.007 OUR AVERAGE

$0.413 \pm 0.004 \pm 0.006$	ROSNER 06	CLEO 10.4	$e^+e^- \rightarrow \text{hadrons}$
$0.45 \pm 0.03 \pm 0.03$	⁴ GILES 84B	CLEO	$e^+e^- \rightarrow \text{hadrons}$

⁴ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

$$\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_6\Gamma_{12}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

18.46±0.27±0.77

6.4K	⁵ AUBERT 08BP	BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
------	--------------------------	------	---

⁵ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

$\Upsilon(3S)$ PARTIAL WIDTHS

$$\Gamma(e^+e^-) \quad \Gamma_{12}$$

VALUE (keV)	DOCUMENT ID
-------------	-------------

0.443±0.008 OUR EVALUATION

$\Upsilon(3S)$ BRANCHING RATIOS

$$\Gamma(\Upsilon(2S)\text{anything})/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.106 ±0.008 OUR AVERAGE

0.1023 ± 0.0105	4625	6,7,8 BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+\ell^-X$
0.111 ± 0.012	4891	7,8,9 BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X, \pi^+\pi^-\ell^+\ell^-$

⁶ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.

⁷ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

⁸ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (18.5 \pm 0.8)\%$.

⁹ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_2/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.45±0.23 OUR AVERAGE				Error includes scale factor of 1.1.
2.40±0.10±0.26	800	10 AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-e^+e^-$
3.12±0.49	980	11,12 BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$
2.13±0.38	974	13 BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X, \pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.82±0.65±0.53	138	13 WU	93 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
3.1 ±2.0	5	MAGERAS	82 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

10 Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

11 From the exclusive mode.

12 Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.

13 Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

 $\Gamma(\Upsilon(2S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.85±0.14 OUR AVERAGE				

1.82±0.09±0.12	4391	14 BHARI	09 CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.16±0.39		15,16 BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

1.7 ± 0.5 ± 0.2	10	17 HEINTZ	92 CSB2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
-----------------	----	-----------	---------	---

14 Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.

15 $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$ and assuming $e\mu$ universality.

16 From the exclusive mode.

17 $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

 $\Gamma(\Upsilon(2S)\gamma\gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0502±0.0069	18 BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+\ell^-2\gamma$

18 From the exclusive mode.

 $\Gamma(\Upsilon(2S)\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.51	90	19 HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

19 Authors assume $B(\Upsilon(2S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.

 $\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

Abbreviation MM in the *COMMENT* field below stands for missing mass.

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.40±0.10 OUR AVERAGE				

4.46±0.01±0.13	190k	20 BHARI	09 CLEO	$e^+e^- \rightarrow \pi^+\pi^- \text{MM}$
----------------	------	----------	---------	---

$4.17 \pm 0.06 \pm 0.19$	6.4K	²¹ AUBERT	08BP BABR	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
4.52 ± 0.35	11830	²² BUTLER	94B CLE2	$e^+ e^- \rightarrow \pi^+\pi^- X, \pi^+\pi^-\ell^+\ell^-$
$4.46 \pm 0.34 \pm 0.50$	451	²² WU	93 CUSB	$\gamma(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46 ± 0.30	11221	²² BROCK	91 CLEO	$e^+ e^- \rightarrow \pi^+\pi^- X, \pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.9 ± 1.0	22	GREEN	82 CLEO	$\gamma(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
3.9 ± 1.3	26	MAGERAS	82 CUSB	$\gamma(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

²⁰ A weighted average of the inclusive and exclusive results.

²¹ Using $B(\gamma(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\gamma(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$, and $\Gamma_{ee}(\gamma(3S)) = 0.443 \pm 0.008$ keV.

²² Using $B(\gamma(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

$\Gamma(\gamma(2S)\pi^+\pi^-)/\Gamma(\gamma(1S)\pi^+\pi^-)$ Γ_2/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.577 $\pm 0.026 \pm 0.060$	800	²³ AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
-----------------------------	-----	----------------------	-----------	---

²³ Using $B(\gamma(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\gamma(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, $B(\gamma(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, and $B(\gamma(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$. Not independent of other values reported by AUBERT 08BP.

$\Gamma(\gamma(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.20 ± 0.13 OUR AVERAGE

2.24 $\pm 0.09 \pm 0.11$	6584	²⁴ BHARI	09 CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.99 ± 0.34	56	²⁵ BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.2 $\pm 0.4 \pm 0.3$	33	²⁶ HEINTZ	92 CSB2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

²⁴ Authors assume $B(\gamma(1S) \rightarrow e^+e^-) + B(\gamma(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

²⁵ Using $B(\gamma(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$ and assuming $e\mu$ universality.

²⁶ Using $B(\gamma(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\gamma(1S)\pi^0\pi^0)/\Gamma(\gamma(1S)\pi^+\pi^-)$ Γ_7/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.501 ± 0.043	²⁷ BHARI	09 CLEO	$e^+e^- \rightarrow \gamma(3S)$
-------------------	---------------------	---------	---------------------------------

²⁷ Not independent of other values reported by BHARI 09.

$\Gamma(\gamma(1S)\eta)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.18	90	²⁸ HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$
-------	----	------------------	----------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8	90	²⁹ AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$
<2.2	90	BROCK	91 CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$

²⁸ Authors assume $B(\gamma(1S) \rightarrow e^+e^-) + B(\gamma(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

²⁹ Using $B(\gamma(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\gamma(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, and $\Gamma_{ee}(\gamma(3S)) = 0.443 \pm 0.008$ keV.

$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_8/Γ_6

<u>VALUE</u> (units 10^{-2})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.9 90 ³⁰AUBERT 08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

³⁰ Not independent of other values reported by AUBERT 08BP.

 $\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	------------	--------------------	-------------	----------------

<0.07 90 ³¹HE 08A CLEO $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

³¹ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE</u> (units 10^{-2})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	-------------	--------------------	-------------	----------------

2.29±0.21±0.22 15k ³²BESSON 07 CLEO $e^+e^- \rightarrow \Upsilon(3S) \rightarrow \tau^+\tau^-$

³² BESSON 07 reports $[\Gamma(\Upsilon(3S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(3S) \rightarrow \mu^+\mu^-)] = 1.05 \pm 0.08 \pm 0.05$. We multiply by our best value $B(\Upsilon(3S) \rightarrow \mu^+\mu^-) = (2.18 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$ Γ_{10}/Γ_{11}

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------


1.05±0.08±0.05 15k BESSON 07 CLEO $e^+e^- \rightarrow \Upsilon(3S)$

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

0.0218±0.0021 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.


0.0239±0.0007±0.0010	81k	ADAMS	05	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
0.0202±0.0019±0.0033		CHEN	89B	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
0.0173±0.0015±0.0011		KAARSBERG	89	CSB2 $e^+e^- \rightarrow \mu^+\mu^-$
0.033 ± 0.013 ± 0.007	1096	ANDREWS	83	CLEO $e^+e^- \rightarrow \mu^+\mu^-$



$\Gamma(\gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
0.131 ±0.016 OUR AVERAGE		Error includes scale factor of 3.4. See the ideogram below.			
0.1579±0.0017±0.0073	568k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$	
0.111 ± 0.005 ± 0.004	10319	³³ HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X$	
0.135 ± 0.003 ± 0.017	30741	MORRISON	91	CLE2 $e^+ e^- \rightarrow \gamma X$	


³³ Supersedes NARAIN 91.



$\Gamma(\gamma\chi_{b1}(2P))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
0.126 ±0.012 OUR AVERAGE	537k	ARTUSO 05	CLEO	e ⁺ e ⁻ → γX	
0.1454±0.0018±0.0073	11147	HEINTZ 92	CSB2	e ⁺ e ⁻ → γX	
0.115 ±0.005 ±0.005	34	MORRISON 91	CLE2	e ⁺ e ⁻ → γX	
0.105 +0.003 -0.002 ±0.013	25759				


³⁴ Supersedes NARAIN 91.



$\Gamma(\gamma\chi_{b0}(2P))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
0.059 ±0.006 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.			
0.0677±0.0020±0.0065	225k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$	
0.060 ±0.004 ±0.006	4959	³⁵ HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X$	
0.049 ^{+0.003} _{-0.004} ±0.006	9903	MORRISON	91	CLE2 $e^+ e^- \rightarrow \gamma X$	

³⁵ Supersedes NARAIN 91.



$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<190	90	36 ASNER 08A	CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$

36 ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))]$ $< 27.1 \times 10^{-2}$. We multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<17	90	37 ASNER 08A	CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$

37 ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))]$ $< 2.5 \times 10^{-2}$. We multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = 6.9 \times 10^{-2}$.

$\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.30±0.04±0.10	8.7k	ARTUSO 05	CLEO	$e^+ e^- \rightarrow \gamma X$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8 90 38 ASNER 08A CLEO $\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$

38 ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))]$ $< 21.9 \times 10^{-2}$. We multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$.

$\Gamma(\gamma\eta_b(2S))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6.2	90	ARTUSO 05	CLEO	$e^+ e^- \rightarrow \gamma X$

$\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{20}/Γ
4.8±0.5±1.2		$19 \pm 3k$	AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.3	90	ARTUSO	05	CLEO	$e^+ e^- \rightarrow \gamma X$
------	----	--------	----	------	--------------------------------

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ (1.5 GeV < m_X < 5.0 GeV)

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{21}/Γ
<2.2	95	ROSNER	07A	CLEO	$e^+ e^- \rightarrow \gamma X$

 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{22}/Γ
<20.3	95	LOVE	08A	CLEO	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

 $\Upsilon(3S)$ REFERENCES

BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)
WU	93	PL B301 307	Q.W. Wu <i>et al.</i>	(CUSB Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
BROCK	91	PR D43 1448	I.C. Brock <i>et al.</i>	(CLEO Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE $e^+ e^-$ Physics 412	W. Buchmuller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
		Translated from YAF 41 733.		
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GREEN	82	PRL 49 617	J. Green <i>et al.</i>	(CLEO Collab.)
MAGERAS	82	PL 118B 453	G. Mageras <i>et al.</i>	(COLU, CORN, LSU+)

OTHER RELATED PAPERS

BRIERE	07	PR D76 012005	R.A. Briere <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN...	07	PR D76 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
GUO	05	NP A761 269	F.-K. Guo <i>et al.</i>	
ROSNER	03	PR D67 097504	J.L. Rosner	
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)

HAN	82	PRL 49 1612	K. Han <i>et al.</i>	(CUSB Collab.)
PETERSON	82	PL 114B 277	D. Peterson <i>et al.</i>	(CUSB Collab.)
KAPLAN	78	PRL 40 435	D.M. Kaplan <i>et al.</i>	(STON, FNAL, COLU)
YOH	78	PRL 41 684	J.K. Yoh <i>et al.</i>	(COLU, FNAL, STON)
COBB	77	PL 72B 273	J.H. Cobb <i>et al.</i>	(BNL, CERN, SYRA, YALE)
HERB	77	PRL 39 252	S.W. Herb <i>et al.</i>	(COLU, FNAL, STON)
INNES	77	PRL 39 1240	W.R. Innes <i>et al.</i>	(COLU, FNAL, STON)