

BOTTOM BARYONS ($B = -1$)

$$\Lambda_b^0 = udb, \Xi_b^0 = usb, \Xi_b^- = dsb, \Omega_b^- = ssb$$

Λ_b^0

$$I(J^P) = 0(\frac{1}{2}^+)$$

$I(J^P)$ not yet measured; $0(\frac{1}{2}^+)$ is the quark model prediction.

$$\text{Mass } m = 5620.2 \pm 1.6 \text{ MeV}$$

$$m_{\Lambda_b} - m_{B^0} = 339.2 \pm 1.4 \text{ MeV}$$

$$\text{Mean life } \tau = (1.383_{-0.048}^{+0.049}) \times 10^{-12} \text{ s}$$

$$c\tau = 415 \mu\text{m}$$

These branching fractions are actually an average over weakly decaying b -baryons weighted by their production rates in Z decay (or high-energy $p\bar{p}$), branching ratios, and detection efficiencies. They scale with the LEP b -baryon production fraction $B(b \rightarrow b\text{-baryon})$ and are evaluated for our value $B(b \rightarrow b\text{-baryon}) = (9.2 \pm 1.8)\%$.

The branching fractions $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{ anything})$ and $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})$ are not pure measurements because the underlying measured products of these with $B(b \rightarrow b\text{-baryon})$ were used to determine $B(b \rightarrow b\text{-baryon})$, as described in the note "Production and Decay of b -Flavored Hadrons."

For inclusive branching fractions, e.g., $B \rightarrow D^\pm \text{ anything}$, the values usually are multiplicities, not branching fractions. They can be greater than one.

Λ_b^0 DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	P (MeV/c)
$J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0)$	$(4.7 \pm 2.3) \times 10^{-5}$		1741
$\Lambda_c^+ \pi^-$	$(8.8 \pm 3.2) \times 10^{-3}$		2343
$\Lambda_c^+ a_1(1260)^-$	seen		2153
$\Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything}$	[a] $(10.6 \pm 3.1) \%$		—
$\Lambda_c^+ \ell^- \bar{\nu}_\ell$	$(5.0_{-1.4}^{+1.9}) \%$		2345
$\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell$	$(5.6 \pm 3.1) \%$		2335
$\Lambda_c(2595)^+ \ell^- \bar{\nu}_\ell$	$(6.3_{-3.1}^{+4.0}) \times 10^{-3}$		2211
$\Lambda_c(2625)^+ \ell^- \bar{\nu}_\ell$	$(1.1_{-0.4}^{+0.6}) \%$		2196

$p h^-$	$[b] < 2.3$	$\times 10^{-5}$	90%	2730
$p \pi^-$	< 5.0	$\times 10^{-5}$	90%	2730
$p K^-$	< 5.0	$\times 10^{-5}$	90%	2709
$\Lambda \gamma$	< 1.3	$\times 10^{-3}$	90%	2699

Σ_b

$$I(J^P) = 1(\frac{1}{2}^+)$$

I, J, P need confirmation.

Mass $m(\Sigma_b^+) = 5807.8 \pm 2.7$ MeV

Mass $m(\Sigma_b^-) = 5815.2 \pm 2.0$ MeV

Σ_b DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_b^0 \pi$	dominant	128

Σ_b^*

$$I(J^P) = 1(\frac{3}{2}^+)$$

I, J, P need confirmation.

Mass $m(\Sigma_b^{*+}) = 5829.0 \pm 3.4$ MeV

Mass $m(\Sigma_b^{*-}) = 5836.4 \pm 2.8$ MeV

$m_{\Sigma_b^*} - m_{\Sigma_b} = 21.2 \pm 2.0$ MeV

Σ_b^* DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda_b^0 \pi$	dominant	156

Ξ_b^0, Ξ_b^-

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

I, J, P need confirmation.

Mass $m = 5792.4 \pm 3.0$ MeV

Mean life $\tau = (1.42^{+0.28}_{-0.24}) \times 10^{-12}$ s

Ξ_b DECAY MODES	Fraction (Γ_i/Γ)	Scale factor	p (MeV/c)
$\Xi_b^- \rightarrow \Xi^- \ell^- \bar{\nu}_\ell X \times B(\bar{b} \rightarrow \Xi_b^-)$	$(3.9 \pm 1.2) \times 10^{-4}$	1.4	-
$\Xi_b^- \rightarrow J/\psi \Xi^- \times B(b \rightarrow \Xi_b^-)$	$(1.3 \pm 0.9) \times 10^{-5}$		-

Ω_b^-

$$I(J^P) = 0(\frac{1}{2}^+)$$

I, J, P need confirmation.

Mass $m = 6165 \pm 16$ MeV

Ω_b^- DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$J/\psi \Omega^- \times B(b \rightarrow \Omega_b)$	$(1.1 \pm 0.8) \times 10^{-5}$	1900

b -baryon ADMIXTURE ($\Lambda_b, \Xi_b, \Sigma_b, \Omega_b$)

$$\text{Mean life } \tau = (1.319_{-0.038}^{+0.039}) \times 10^{-12} \text{ s}$$

These branching fractions are actually an average over weakly decaying b -baryons weighted by their production rates in Z decay (or high-energy $p\bar{p}$), branching ratios, and detection efficiencies. They scale with the LEP b -baryon production fraction $B(b \rightarrow b\text{-baryon})$ and are evaluated for our value $B(b \rightarrow b\text{-baryon}) = (9.2 \pm 1.8)\%$.

The branching fractions $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{ anything})$ and $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})$ are not pure measurements because the underlying measured products of these with $B(b \rightarrow b\text{-baryon})$ were used to determine $B(b \rightarrow b\text{-baryon})$, as described in the note "Production and Decay of b -Flavored Hadrons."

For inclusive branching fractions, *e.g.*, $B \rightarrow D^\pm \text{ anything}$, the values usually are multiplicities, not branching fractions. They can be greater than one.

b -baryon ADMIXTURE DECAY MODES

$(\Lambda_b, \Xi_b, \Sigma_b, \Omega_b)$	Fraction (Γ_i/Γ)	p (MeV/c)
$p \mu^- \bar{\nu}$ anything	$(5.7_{-2.3}^{+2.6})\%$	—
$p \ell \bar{\nu}_\ell$ anything	$(5.5 \pm 1.6)\%$	—
p anything	$(69 \pm 26)\%$	—
$\Lambda \ell^- \bar{\nu}_\ell$ anything	$(3.7 \pm 0.9)\%$	—
$\Lambda/\bar{\Lambda}$ anything	$(38 \pm 10)\%$	—
$\Xi^- \ell^- \bar{\nu}_\ell$ anything	$(6.4 \pm 2.1) \times 10^{-3}$	—

NOTES

[a] Not a pure measurement. See note at head of Λ_b^0 Decay Modes.

[b] Here h^- means π^- or K^- .