

**X(3940)** $I^G(J^{PC}) = ??(???)$ 

## OMITTED FROM SUMMARY TABLE

Reported by ABE 07, observed in  $e^+ e^- \rightarrow J/\psi X$ .**X(3940) MASS**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3942^{+7}_{-6} \pm 6</math></b>	52	PAKHLOV	08	$e^+ e^- \rightarrow J/\psi X$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
3943 $\pm 6 \pm 6$	25	<sup>1</sup> ABE	07	$e^+ e^- \rightarrow J/\psi X$
3936 $\pm 14$	266	<sup>2</sup> ABE	07	$e^+ e^- \rightarrow J/\psi(c\bar{c})$

<sup>1</sup> From a fit to  $D^* D^-$  and  $D^* \bar{D}^0$  events.<sup>2</sup> From the inclusive fit. Not independent of the exclusive measurement by ABE 07.**X(3940) WIDTH**

<u>VALUE (MeV)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>37^{+26}_{-15} \pm 8</math></b>		52	PAKHLOV	08	$e^+ e^- \rightarrow J/\psi X$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>					
<52	90	25	ABE	07	$e^+ e^- \rightarrow J/\psi X$

**X(3940) DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 D \bar{D}^* + c.c.$	seen
$\Gamma_2 D \bar{D}$	not seen
$\Gamma_3 J/\psi \omega$	not seen

**X(3940) BRANCHING RATIOS**

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u><math>\Gamma_1/\Gamma</math></u>
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>						
>0.45	90	25	<sup>3,4</sup> ABE	07	$e^+ e^- \rightarrow J/\psi X$	
<sup>3</sup> For $X(3940)$ decaying to final states with more than two tracks.						
<sup>4</sup> PAKHLOV 08 finds that the inclusive peak near $3940 \text{ MeV}/c^2$ may consist of several states.						

### $\Gamma(D\bar{D})/\Gamma_{\text{total}}$

### $\Gamma_2/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.41                  90            5,<sup>6</sup> ABE                  07            BELL             $e^+ e^- \rightarrow J/\psi X$

<sup>5</sup> For  $X(3940)$  decaying to final states with more than two tracks.

<sup>6</sup> PAKHLOV 08 finds that the inclusive peak near  $3940 \text{ MeV}/c^2$  may consist of several states.

### $\Gamma(J/\psi\omega)/\Gamma_{\text{total}}$

### $\Gamma_3/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.26                  90            7,<sup>8</sup> ABE                  07            BELL             $e^+ e^- \rightarrow J/\psi X$

<sup>7</sup> For  $X(3940)$  decaying to final states with more than two tracks.

<sup>8</sup> PAKHLOV 08 finds that the inclusive peak near  $3940 \text{ MeV}/c^2$  may consist of several states.

## X(3940) REFERENCES

PAKHLOV        08        PRL 100 202001  
ABE              07        PRL 98 082001

P. Pakhlov *et al.*  
K. Abe *et al.*

(BELLE Collab.)  
(BELLE Collab.)