

$\chi_{b1}(2P)$
 $I^G(J^{PC}) = 0^+(1^{++})$
J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

 $\chi_{b1}(2P)$ MASS

<u>VALUE</u> (GeV)	<u>DOCUMENT ID</u>
$10.25546 \pm 0.00022 \pm 0.00050$ OUR EVALUATION	From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV

 $m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)}$

<u>VALUE</u> (MeV)	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$23.5 \pm 0.7 \pm 0.7$	¹ HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X, \ell^+ \ell^- \gamma \gamma$


¹ From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

 γ ENERGY IN $\Upsilon(3S)$ DECAY

<u>VALUE</u> (MeV)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
99.26 ± 0.22 OUR EVALUATION				Treating systematic errors as correlated
99.53 ± 0.23 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
$99.15 \pm 0.07 \pm 0.25$		ARTUSO	05	CLEO $\Upsilon(3S) \rightarrow \gamma X$
99 ± 1	169	CRAWFORD	92B	CLE2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
100.1 ± 0.4	11147	² HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X$
100.2 ± 0.5	223	³ HEINTZ	92	CSB2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
$99.5 \pm 0.1 \pm 0.5$	25759	MORRISON	91	CLE2 $e^+ e^- \rightarrow \gamma X$

² A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

³ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.



$\chi_{b1}(2P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor
$\Gamma_1 \omega \eta(1S)$	($1.63^{+0.40}_{-0.34}$) %	
$\Gamma_2 \gamma \eta(2S)$	(19.9 ± 1.9) %	
$\Gamma_3 \gamma \eta(1S)$	(9.2 ± 0.8) %	1.1
$\Gamma_4 \pi\pi\chi_{b1}(1P)$	(9.1 ± 1.3) $\times 10^{-3}$	
$\Gamma_5 D^0 X$	(8.8 ± 1.7) %	
$\Gamma_6 \pi^+\pi^- K^+ K^- \pi^0$	(3.1 ± 1.0) $\times 10^{-4}$	
$\Gamma_7 2\pi^+\pi^- K^- K_S^0$	(1.1 ± 0.5) $\times 10^{-4}$	
$\Gamma_8 2\pi^+\pi^- K^- K_S^0 2\pi^0$	(7.7 ± 3.2) $\times 10^{-4}$	
$\Gamma_9 2\pi^+ 2\pi^- 2\pi^0$	(5.9 ± 2.0) $\times 10^{-4}$	
$\Gamma_{10} 2\pi^+ 2\pi^- K^+ K^-$	(10 ± 4) $\times 10^{-5}$	
$\Gamma_{11} 2\pi^+ 2\pi^- K^+ K^- \pi^0$	(5.5 ± 1.8) $\times 10^{-4}$	
$\Gamma_{12} 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(10 ± 4) $\times 10^{-4}$	
$\Gamma_{13} 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	(6.7 ± 2.6) $\times 10^{-4}$	
$\Gamma_{14} 3\pi^+ 3\pi^-$	(1.2 ± 0.4) $\times 10^{-4}$	
$\Gamma_{15} 3\pi^+ 3\pi^- 2\pi^0$	(1.2 ± 0.4) $\times 10^{-3}$	
$\Gamma_{16} 3\pi^+ 3\pi^- K^+ K^-$	(2.0 ± 0.8) $\times 10^{-4}$	

Γ_{17}	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(6.1 \pm 2.2) \times 10^{-4}$
Γ_{18}	$4\pi^+ 4\pi^-$	$(1.7 \pm 0.6) \times 10^{-4}$
Γ_{19}	$4\pi^+ 4\pi^- 2\pi^0$	$(1.9 \pm 0.7) \times 10^{-3}$

$\chi_{b1}(2P)$ BRANCHING RATIOS

$\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$	Γ_1/Γ
$\text{VALUE (units } 10^{-2}\text{)}$	EVTS

1.63 $^{+0.35}_{-0.31}{}^{+0.16}_{-0.15}$ $32.6^{+6.9}_{-6.1}$ ⁴ CRONIN-HEN..04 CLE3 $\Upsilon(3S) \rightarrow \gamma \omega \Upsilon(1S)$

⁴ Using $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (11.3 \pm 0.6)\%$ and $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 2 (2.48 \pm 0.06)\%$.

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$	Γ_2/Γ
VALUE	EVTS

0.199 ± 0.019 OUR AVERAGE

$0.190 \pm 0.018 \pm 0.017$ 4.3k
 $0.356 \pm 0.042 \pm 0.092$
 $0.199 \pm 0.020 \pm 0.022$

⁵ LEES 11J reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (2.4 \pm 0.1 \pm 0.2) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (10.23 \pm 1.20 \pm 1.26) \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = 0.105^{+0.003}_{-0.002} \pm 0.013$.

⁷ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (11.5 \pm 0.5 \pm 0.5)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$	Γ_3/Γ
VALUE	EVTS

0.092 ± 0.008 OUR AVERAGE

$0.098 \pm 0.005 \pm 0.009$ 15k
 $0.120 \pm 0.021 \pm 0.021$
 $0.080 \pm 0.009 \pm 0.007$

⁸ LEES 11J reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (12.4 \pm 0.3 \pm 0.6) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (6.47 \pm 1.12 \pm 0.82) \times 10^{-4}$ and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = 0.105^{+0.003}_{-0.002} \pm 0.013$.

¹⁰ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (11.5 \pm 0.5 \pm 0.5)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\pi\pi\chi_{b1}(1P))/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.1±1.3 OUR AVERAGE				
$9.2 \pm 1.1 \pm 0.8$	31k	¹¹ LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$
$8.6 \pm 2.3 \pm 2.1$		¹² CAWLFIELD	06 CLE3	$\gamma(3S) \rightarrow 2(\gamma\pi\ell)$
11 LEES 11C measures $B(\gamma(3S) \rightarrow \chi_{b1}(2P)X) \times B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) = (1.16 \pm 0.07 \pm 0.12) \times 10^{-3}$. We derive the value assuming $B(\gamma(3S) \rightarrow \chi_{b1}(2P)X) = B(\gamma(3S) \rightarrow \chi_{b1}(2P)\gamma) = (12.6 \pm 1.2) \times 10^{-2}$.				
12 CAWLFIELD 06 quote $\Gamma(\chi_b(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$ keV assuming l-spin conservation, no D-wave contribution, $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$ keV, and $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$ keV.				

 $\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.8±1.5±0.8	2243	¹³ BRIERE	08 CLEO	$\gamma(3S) \rightarrow \gamma D^0 X$
13 For $p_{D^0} > 2.5$ GeV/c.				

 $\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.1±1.0±0.3	30	¹⁴ ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$
14 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (39 \pm 8 \pm 9) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.1±0.5±0.1	10	¹⁵ ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$
15 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (14 \pm 5 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.7±3.1±0.7	15	¹⁶ ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$
16 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (97 \pm 30 \pm 26) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.9±2.0±0.5	36	¹⁷ ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$

¹⁷ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))]$ $= (74 \pm 16 \pm 19) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$

Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.0±0.4±0.1	12	18 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$

¹⁸ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$

Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.5±1.7±0.5	38	19 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$

¹⁹ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (69 \pm 13 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$

Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.6±3.5±0.9	27	20 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$

²⁰ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (121 \pm 29 \pm 33) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$

Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.7±2.5±0.6	17	21 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$

²¹ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (85 \pm 23 \pm 22) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$

Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2±0.4±0.1	18	22 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$

²² ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (15 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12±4±1	44	23 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$

²³ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(150 \pm 30 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))$ = $(12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.0±0.7±0.2	16	24 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$

²⁴ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(25 \pm 7 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))$ = $(12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.1±2.1±0.6	25	25 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$

²⁵ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(77 \pm 17 \pm 21) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))$ = $(12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.7±0.6±0.2	16	26 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$

²⁶ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(22 \pm 6 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))$ = $(12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
19±7±2	41	27 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$

²⁷ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(241 \pm 47 \pm 72) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))$ = $(12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\chi_{b1}(2P)$ Cross-Particle Branching Ratios $\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Gamma(1S))/\Gamma_{\text{total}} \times \Gamma(\Gamma(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_{20}^{T(3S)}/\Gamma^{T(3S)}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
12.4±0.3±0.6	15k	LEES	11J BABR	$\Gamma(3S) \rightarrow X\gamma$

|

$$\frac{\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}}}{\Gamma_2/\Gamma \times \Gamma_{20}^{\Upsilon(3S)}/\Gamma^{\Upsilon(3S)}}$$

<u>VALUE</u> (units 10^{-2})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.4±0.1±0.2	4.3k	LEES	11J	BABR $\Upsilon(3S) \rightarrow X \gamma$

$$\mathbf{B}(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) \times \mathbf{B}(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X)$$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.16±0.07±0.12	31k	LEES	11C	BABR $e^+e^- \rightarrow \pi^+\pi^-X$

$$\mathbf{B}(\chi_{b2}(2P) \rightarrow pX + \bar{p}X)/\mathbf{B}(\chi_{b1}(2P) \rightarrow pX + \bar{p}X)$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.109±0.007±0.040	BRIERE	07	CLEO $\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)$

$$\mathbf{B}(\chi_{b0}(2P) \rightarrow pX + \bar{p}X)/\mathbf{B}(\chi_{b1}(2P) \rightarrow pX + \bar{p}X)$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.082±0.025±0.060	BRIERE	07	CLEO $\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)$

$\chi_{b1}(2P)$ REFERENCES

LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
BRIERE	07	PR D76 012005	R.A. Briere <i>et al.</i>	(CLEO Collab.)
CAWLFIELD	06	PR D73 012003	C. Cawlfield <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN... 04	04	PRL 92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92B	PL B294 139	G. Crawford, R. Fulton	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)