

$f_0(1370)$

$I^G(J^{PC}) = 0^+(0^{++})$

See also the mini-reviews on scalar mesons under $f_0(500)$ (see the index for the page number) and on non- $q\bar{q}$ candidates in PDG 06, Journal of Physics, G **33** 1 (2006).

$f_0(1370)$ T-MATRIX POLE POSITION

Note that $\Gamma \approx 2 \operatorname{Im}(\sqrt{s_{\text{pole}}})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1200–1500)–i(150–250) OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$(1290 \pm 50) - i(170^{+20}_{-40})$	¹ ANISOVICH 09 RVUE $0.0 \bar{p}p, \pi N$		
$(1373 \pm 15) - i(137 \pm 10)$	² BARGIOTTI 03 OBLX $\bar{p}p$		
$(1302 \pm 17) - i(166 \pm 18)$	³ BARBERIS 00C $450 pp \rightarrow p_f 4\pi p_s$		
$(1312 \pm 25 \pm 10) - i(109 \pm 22 \pm 15)$	BARBERIS 99D OMEG $450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$		
$(1406 \pm 19) - i(80 \pm 6)$	⁴ KAMINSKI 99 RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$		
$(1300 \pm 20) - i(120 \pm 20)$	ANISOVICH 98B RVUE Compilation		
$(1290 \pm 15) - i(145 \pm 15)$	BARBERIS 97B OMEG $450 pp \rightarrow pp 2(\pi^+ \pi^-)$		
$(1548 \pm 40) - i(560 \pm 40)$	BERTIN 97C OBLX $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$		
$(1380 \pm 40) - i(180 \pm 25)$	ABELE 96B CBAR $0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$		
$(1300 \pm 15) - i(115 \pm 8)$	BUGG 96 RVUE		
$(1330 \pm 50) - i(150 \pm 40)$	⁵ AMSLER 95B CBAR $\bar{p}p \rightarrow 3\pi^0$		
$(1360 \pm 35) - i(150–300)$	⁵ AMSLER 95C CBAR $\bar{p}p \rightarrow \pi^0 \eta\eta$		
$(1390 \pm 30) - i(190 \pm 40)$	⁶ AMSLER 95D CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$		
$1346 - i249$	^{7,8} JANSEN 95 RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$		
$1214 - i168$	^{8,9} TORNQVIST 95 RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$		
$1364 - i139$	AMSLER 94D CBAR $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$		
$(1365^{+20}_{-55}) - i(134 \pm 35)$	ANISOVICH 94 CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$		
$(1340 \pm 40) - i(127^{+30}_{-20})$	¹⁰ BUGG 94 RVUE $\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0 \pi^0$		
$(1430 \pm 5) - i(73 \pm 13)$	¹¹ KAMINSKI 94 RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$		
$1420 - i220$	¹² AU 87 RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$		

¹ Another pole is found at $(1510 \pm 130) - i(800^{+100}_{-150})$ MeV.

² Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

³ Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$.

⁴ T-matrix pole on sheet ——.

⁵ Supersedes ANISOVICH 94.

⁶ Coupled-channel analysis of $\bar{p}p \rightarrow 3\pi^0$, $\pi^0 \eta\eta$, and $\pi^0 \pi^0 \eta$ on sheet IV. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

⁷ Analysis of data from FALVARD 88.

⁸ The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

- ⁹ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.
¹⁰ Reanalysis of ANISOVICH 94 data.
¹¹ T-matrix pole on sheet III.
¹² Analysis of data from OCHS 73, GRAYER 74, BECKER 79, and CASON 83.

$f_0(1370)$ BREIT-WIGNER MASS OR K-MATRIX POLE PARAMETER

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>
1200 to 1500 OUR ESTIMATE	

$\pi\pi$ MODE

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1400 ± 40		13 AUBERT	09L BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
1470 $^{+6}_{-7}$ $^{+72}_{-255}$		14 UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1259 ± 55	2.6k	BONVICINI	07 CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
1309 ± 1 ± 15		15 BUGG	07A RVUE	$0.0 p\bar{p} \rightarrow 3\pi^0$
1449 ± 13	4286	16 GARMASH	06 BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$
1350 ± 50		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
1265 ± 30 $^{+20}_{-35}$		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
1434 ± 18 ± 9	848	AITALA	01A E791	$D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
1308 ± 10		BARBERIS	99B OMEG	$450 pp \rightarrow p_s p_f \pi^+ \pi^-$
1315 ± 50		BELLAZZINI	99 GAM4	$450 pp \rightarrow pp \pi^0 \pi^0$
1315 ± 30		ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$
1280 ± 55		BERTIN	98 OBLX	$0.05-0.405 \bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
1186	17,18	TORNQVIST	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1472 ± 12		ARMSTRONG	91 OMEG	$300 pp \rightarrow pp\pi\pi, ppK\bar{K}$
1275 ± 20		BREAKSTONE	90 SFM	$62 pp \rightarrow pp\pi^+ \pi^-$
1420 ± 20		AKESSON	86 SPEC	$63 pp \rightarrow pp\pi^+ \pi^-$
1256		FROGGATT	77 RVUE	$\pi^+ \pi^-$ channel

¹³ Breit-Wigner mass.

¹⁴ Breit-Wigner mass. May also be the $f_0(1500)$.

¹⁵ Reanalysis of ABELE 96C data.

¹⁶ Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays. Supersedes GARMASH 05.

¹⁷ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

¹⁸ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$ decays

$K\bar{K}$ MODE

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1440 ± 6	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1391 ± 10	TIKHOMIROV 03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1440 ± 50	BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1463 ± 9	ETKIN 82B	MPS	$23 \pi^- p \rightarrow n2K_S^0$
1425 ± 15	WICKLUND 80	SPEC	$6 \pi N \rightarrow K^+ K^- N$
~ 1300	POLYCHRO... 79	STRC	$7 \pi^- p \rightarrow n2K_S^0$

4π MODE $2(\pi\pi)_S + \rho\rho$

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1395 \pm 40		ABELE	01	CBAR $0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
1374 \pm 38		AMSLER	94	CBAR $0.0 \bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
1345 \pm 12		ADAMO	93	OBIX $\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
1386 \pm 30		GASPERO	93	DBC $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
~ 1410	5751	¹⁹ BETTINI	66	DBC $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
$19 \rho\rho$ dominant.				

 $\eta\eta$ MODE

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1262^{+51+82}_{-78-103}$	20 UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
1430	AMSLER	92 CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta\eta$
1220 \pm 40	ALDE	86D GAM4	$100 \pi^- p \rightarrow n 2\eta$
20 Breit-Wigner mass. May also be the $f_0(1500)$.			

COUPLED CHANNEL MODE

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •		
1306 \pm 20	21 ANISOVICH	03 RVUE
21 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.		

 $f_0(1370)$ BREIT-WIGNER WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>
200 to 500 OUR ESTIMATE	

 $\pi\pi$ MODE

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
300 \pm 80		22 AUBERT	09L BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
90^{+2+50}_{-1-22}		23 UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
298 \pm 21	2.6k	BONVICINI	07 CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
126 \pm 25	4286	²⁴ GARMASH	06 BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$
265 \pm 40		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
$350 \pm 100^{+105}_{-60}$		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
173 \pm 32 \pm 6	848	AITALA	01A E791	$D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
222 \pm 20		BARBERIS	99B OMEG	$450 pp \rightarrow p_s p_f \pi^+ \pi^-$

255 ± 60	BELLAZZINI	99	GAM4	$450 \text{ } pp \rightarrow pp\pi^0\pi^0$
190 ± 50	ALDE	98	GAM4	$100 \text{ } \pi^- p \rightarrow \pi^0\pi^0 n$
323 ± 13	BERTIN	98	OBLX	$0.05\text{--}0.405 \bar{n}p \rightarrow \pi^+\pi^+\pi^-$
350	25,26 TORNQVIST	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
195 ± 33	ARMSTRONG	91	OMEG	$300 \text{ } pp \rightarrow pp\pi\pi, ppK\bar{K}$
285 ± 60	BREAKSTONE	90	SFM	$62 \text{ } pp \rightarrow pp\pi^+\pi^-$
460 ± 50	AKESSON	86	SPEC	$63 \text{ } pp \rightarrow pp\pi^+\pi^-$
~ 400	27 FROGGATT	77	RVUE	$\pi^+\pi^- \text{ channel}$

²² The systematic errors are not reported.²³ Breit-Wigner width. May also be the $f_0(1500)$.²⁴ Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0\pi^+\pi^-$ decays. Supersedes GARMASH 05.²⁵ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.²⁶ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decays²⁷ Width defined as distance between 45 and 135° phase shift.

$K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
121 ± 15	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
55 ± 26	TIKHOLOMOV 03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
250 ± 80	BOLONKIN	88	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
118^{+138}_{-16}	ETKIN	82B	MPS $23 \pi^- p \rightarrow n2K_S^0$
160 ± 30	WICKLUND	80	SPEC $6 \pi N \rightarrow K^+ K^- N$
~ 150	POLYCHRO...	79	STRC $7 \pi^- p \rightarrow n2K_S^0$

4π MODE $2(\pi\pi)_S + \rho\rho$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
275 ± 55		ABELE	01	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
375 ± 61		AMSLER	94	$0.0 \bar{p}p \rightarrow \pi^+\pi^- 3\pi^0$
398 ± 26		ADAMO	93	$\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
310 ± 50		GASPERO	93	$0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
~ 90	5751	28 BETTINI	66	$0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$

²⁸ $\rho\rho$ dominant.

$\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$484^{+246+246}_{-170-263}$	29 UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
250		AMSLER	$92 \text{ CBAR } 0.0 \bar{p}p \rightarrow \pi^0\eta\eta$
320 ± 40		ALDE	$86D \text{ GAM4 } 100 \pi^- p \rightarrow n2\eta$

²⁹ Breit-Wigner width. May also be the $f_0(1500)$.

COUPLED CHANNEL MODE

<i>VALUE</i> (MeV)	<i>DOCUMENT ID</i>	<i>TECN</i>
--------------------	--------------------	-------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

147^{+30}_{-50}	30 ANISOVICH	03 RVUE
-------------------	--------------	---------

30 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

 $f_0(1370)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi \pi$	seen
$\Gamma_2 4\pi$	seen
$\Gamma_3 4\pi^0$	seen
$\Gamma_4 2\pi^+ 2\pi^-$	seen
$\Gamma_5 \pi^+ \pi^- 2\pi^0$	seen
$\Gamma_6 \rho \rho$	dominant
$\Gamma_7 2(\pi\pi)_S$ -wave	seen
$\Gamma_8 \pi(1300)\pi$	seen
$\Gamma_9 a_1(1260)\pi$	seen
$\Gamma_{10} \eta \eta$	seen
$\Gamma_{11} K\bar{K}$	seen
$\Gamma_{12} K\bar{K}n\pi$	not seen
$\Gamma_{13} 6\pi$	not seen
$\Gamma_{14} \omega \omega$	not seen
$\Gamma_{15} \gamma \gamma$	seen
$\Gamma_{16} e^+ e^-$	not seen

 $f_0(1370)$ PARTIAL WIDTHS **$\Gamma(\gamma\gamma)$** See $\gamma\gamma$ widths under $f_0(500)$ and MORGAN 90. **Γ_{15}** **$\Gamma(e^+ e^-)$** **Γ_{16}**

<i>VALUE</i> (eV)	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<20	90	VOROBYEV	88	$e^+ e^- \rightarrow \pi^0 \pi^0$

 $f_0(1370) \Gamma(i) \Gamma(\gamma\gamma)/\Gamma(\text{total})$ **$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$** **$\Gamma_{10}\Gamma_{15}/\Gamma$**

<i>VALUE</i> (eV)	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
-------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$121^{+133+169}_{-53-106}$	31 UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
----------------------------	-----------	----------	---

³¹ Including interference with the $f_2'(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.

$f_0(1370)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_1/Γ
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.26 \pm 0.09	BUGG 96	RVUE		
<0.15	³² AMSLER 94	CBAR	$\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$	
<0.06	GASPERO 93	DBC	0.0 $\bar{p}n \rightarrow$ hadrons	

³² Using AMSLER 95B ($3\pi^0$).

$\Gamma(4\pi)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_2/\Gamma = (\Gamma_3 + \Gamma_4 + \Gamma_5)/\Gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
>0.72	GASPERO 93	DBC	0.0 $\bar{p}n \rightarrow$ hadrons	

$\Gamma(4\pi^0)/\Gamma(4\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ_2
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
seen	ABELE 96	CBAR	0.0 $\bar{p}p \rightarrow 5\pi^0$	
0.068 \pm 0.005	³³ GASPERO 93	DBC	0.0 $\bar{p}n \rightarrow$ hadrons	

³³ Model-dependent evaluation.

$\Gamma(2\pi^+ 2\pi^-)/\Gamma(4\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_4/\Gamma_2 = \Gamma_4/(\Gamma_3 + \Gamma_4 + \Gamma_5)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.420 \pm 0.014	³⁴ GASPERO 93	DBC	0.0 $\bar{p}n \rightarrow 2\pi^+ 3\pi^-$	

³⁴ Model-dependent evaluation.

$\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma(4\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_5/\Gamma_2 = \Gamma_5/(\Gamma_3 + \Gamma_4 + \Gamma_5)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.512 \pm 0.019	³⁵ GASPERO 93	DBC	0.0 $\bar{p}n \rightarrow$ hadrons	

³⁵ Model-dependent evaluation.

$\Gamma(\rho\rho)/\Gamma(4\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_6/Γ_2
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.26 \pm 0.07	ABELE 01B	CBAR	0.0 $\bar{p}d \rightarrow 5\pi p$	

$\Gamma(2(\pi\pi)_S\text{-wave})/\Gamma(\pi\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_7/Γ_1
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
5.6 \pm 2.6	³⁶ ABELE 01	CBAR	0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$	

³⁶ From the combined data of ABELE 96 and ABELE 96C.

$\Gamma(2(\pi\pi)s\text{-wave})/\Gamma(4\pi)$ Γ_7/Γ_2

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.51 \pm 0.09	ABELE	01B CBAR	$0.0 \bar{p}d \rightarrow 5\pi p$

$\Gamma(\rho\rho)/\Gamma(2(\pi\pi)s\text{-wave})$ Γ_6/Γ_7

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
large	BARBERIS	00C	$450 pp \rightarrow p_f 4\pi p_s$
1.6 ± 0.2	AMSLER	94	$\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
~ 0.65	GASPERO	93	$0.0 \bar{p}n \rightarrow \text{hadrons}$

$\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ Γ_8/Γ_2

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.17 \pm 0.06	ABELE	01B CBAR	$0.0 \bar{p}d \rightarrow 5\pi p$

$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$ Γ_9/Γ_2

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.06 \pm 0.02	ABELE	01B CBAR	$0.0 \bar{p}d \rightarrow 5\pi p$

$\Gamma(\eta\eta)/\Gamma(4\pi)$ $\Gamma_{10}/\Gamma_2 = \Gamma_{10}/(\Gamma_3 + \Gamma_4 + \Gamma_5)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$(28 \pm 11) \times 10^{-3}$	³⁷ ANISOVICH	02D SPEC	Combined fit
$(4.7 \pm 2.0) \times 10^{-3}$	BARBERIS	00E	$450 pp \rightarrow p_f \eta\eta p_s$
³⁷ From a combined K-matrix analysis of Crystal Barrel ($0. p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n$, $\eta\eta n$, $\eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K}n$) data.			

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.35 \pm 0.13	BUGG	96 RVUE	

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$ Γ_{11}/Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.08 \pm 0.08	ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$, $\phi K^+ K^-$
0.91 \pm 0.20	³⁸ BARGIOTTI	03 OBLX	$\bar{p}p$
0.12 \pm 0.06	³⁹ ANISOVICH	02D SPEC	Combined fit
$0.46 \pm 0.15 \pm 0.11$	BARBERIS	99D OMEG	$450 pp \rightarrow K^+ K^-$, $\pi^+ \pi^-$
³⁸ Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.			
³⁹ From a combined K-matrix analysis of Crystal Barrel ($0. p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n$, $\eta\eta n$, $\eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K}n$) data.			

$\Gamma(K\bar{K}n\pi)/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.03	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons

 $\Gamma(6\pi)/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.22	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.13	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons

 $f_0(1370)$ REFERENCES

UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
AUBERT	09L	PR D79 072006	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BUGG	07A	JPG 34 151	D.V. Bugg <i>et al.</i>	
GARMASH	07	PR D75 012006	A. Garmash <i>et al.</i>	(BELLE Collab.)
GARMASH	06	PRL 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)
PDG	06	JPG 33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirska <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>	
		Translated from YAF 65 1583.		
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>	
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 62 446.		
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	
		Translated from UFN 168 481.		
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96B	PL B385 425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BUGG	96	NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)

AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)
TORNQVIST	95	ZPHY C68 647	N.A. Tornqvist	(HELS)
AMSLER	94	PL B322 431	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) JPC
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.) JPC
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.) JPC
GASPERO	93	NP A562 407	M. Gaspero	(ROMAI) JPC
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
VOROBIEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)
Translated from YAF 48 436.				
AU	87	PR D35 1633	K.L. Au, D. Morgan, M.R. Pennington	(DURH, RAL)
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
WICKLUND	80	PRL 45 1469	A.B. Wicklund <i>et al.</i>	(ANL)
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)
BETTINI	66	NC 42A 695	A. Bettini <i>et al.</i>	(PADO, PISA)