

# Neutrino Mixing

A REVIEW GOES HERE – Check our WWW List of Reviews

## (A) Neutrino fluxes and event ratios

### Events (observed/expected) from accelerator $\nu_\mu$ experiments.

Some neutrino oscillation experiments compare the flux in two or more detectors. This is usually quoted as the ratio of the event rate in the far detector to the expected rate based on an extrapolation from the near detector in the absence of oscillations.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.71 \pm 0.08$	<sup>1</sup> AHN	06A K2K	K2K to Super-K
$0.64 \pm 0.05$	<sup>2</sup> MICHAEL	06 MINS	All charged current events
$0.71^{+0.08}_{-0.09}$	<sup>3</sup> ALIU	05 K2K	KEK to Super-K
$0.70^{+0.10}_{-0.11}$	<sup>4</sup> AHN	03 K2K	KEK to Super-K

<sup>1</sup> Based on the observation of 112 events when  $158.1^{+9.2}_{-8.6}$  were expected without oscillations. Including not only the number of events but also the shape of the energy distribution, the evidence for oscillation is at the level of about  $4.3 \sigma$ . Supersedes ALIU 05.

<sup>2</sup> This ratio is based on the observation of 215 events compared to an expectation of  $336 \pm 14$  without oscillations. See also ADAMSON 08.

<sup>3</sup> This ratio is based on the observation of 107 events at the far detector 250 km away from KEK, and an expectation of  $151^{+12}_{-10}$ .

<sup>4</sup> This ratio is based on the observation of 56 events with an expectation of  $80.1^{+6.2}_{-5.4}$ .

### Events (observed/expected) from reactor $\bar{\nu}_e$ experiments.

The quoted values are the ratios of the measured reactor  $\bar{\nu}_e$  event rate at the quoted distances, and the rate expected without oscillations. The expected rate is based on the experimental data for the most significant reactor fuels ( $^{235}\text{U}$ ,  $^{239}\text{Pu}$ ,  $^{241}\text{Pu}$ ) and on calculations for  $^{238}\text{U}$ .

A recent re-evaluation of the spectral conversion of electron to  $\bar{\nu}_e$  in MUELLER 11 results in an upward shift of the reactor  $\bar{\nu}_e$  spectrum by 3% and, thus, might require revisions to the ratios listed in this table.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.944 \pm 0.016 \pm 0.040$	<sup>1</sup> ABE	12 DCHZ	Chooz reactors
$0.920 \pm 0.009 \pm 0.014$	<sup>2</sup> AHN	12 RENO	Yonggwang reactors
$0.940 \pm 0.011 \pm 0.004$	<sup>3</sup> AN	12 DAYA	Daya Bay, Ling Ao, Ling Ao-II reactors
$1.08 \pm 0.21 \pm 0.16$	<sup>4</sup> DENIZ	10 TEXO	Kuo-Sheng reactor, 28 m
$0.658 \pm 0.044 \pm 0.047$	<sup>5</sup> ARAKI	05 KLND	Japanese react. $\sim 180$ km
$0.611 \pm 0.085 \pm 0.041$	<sup>6</sup> EGUCHI	03 KLND	Japanese react. $\sim 180$ km
$1.01 \pm 0.024 \pm 0.053$	<sup>7</sup> BOEHM	01	Palo Verde react. 0.75–0.89 km
$1.01 \pm 0.028 \pm 0.027$	<sup>8</sup> APOLLONIO	99 CHOZ	Chooz reactors 1 km

$0.987 \pm 0.006 \pm 0.037$	<sup>9</sup> GREENWOOD 96		Savannah River, 18.2 m
$0.988 \pm 0.004 \pm 0.05$	ACHKAR 95	CNTR	Bugey reactor, 15 m
$0.994 \pm 0.010 \pm 0.05$	ACHKAR 95	CNTR	Bugey reactor, 40 m
$0.915 \pm 0.132 \pm 0.05$	ACHKAR 95	CNTR	Bugey reactor, 95 m
$0.987 \pm 0.014 \pm 0.027$	<sup>10</sup> DECLAIS 94	CNTR	Bugey reactor, 15 m
$0.985 \pm 0.018 \pm 0.034$	KUVSHINN... 91	CNTR	Rovno reactor
$1.05 \pm 0.02 \pm 0.05$	VUILLEUMIER 82		Gösgen reactor
$0.955 \pm 0.035 \pm 0.110$	<sup>11</sup> KWON 81		$\bar{\nu}_e p \rightarrow e^+ n$
$0.89 \pm 0.15$	<sup>11</sup> BOEHM 80		$\bar{\nu}_e p \rightarrow e^+ n$

- <sup>1</sup> ABE 12 determine the  $\bar{\nu}_e$  interaction rate in a single detector, located 1050 m from the cores of two reactors. The rate normalization is fixed by the results of the Bugey4 reactor experiment, thus avoiding any dependence on possible very short baseline oscillations.
- <sup>2</sup> AHN 12 use two identical detectors, placed at flux weighted distances of 408.56 m and 1433.99m from six reactor cores, to determine the  $\bar{\nu}_e$  interaction rate ratio.
- <sup>3</sup> AN 12 use six identical detectors with three placed near the reactor cores (flux-weighted baselines of 470 m and 576 m) and the remaining three at the far hall (at the flux averaged distance of 1648 m from all six reactor cores) to determine the  $\bar{\nu}_e$  interaction rate ratios.
- <sup>4</sup> DENIZ 10 observe reactor  $\bar{\nu}_e e$  scattering with recoil kinetic energies 3–8 MeV using CsI(Tl) detectors. The observed rate is consistent with the Standard Model prediction, leading to a constraint on  $\sin^2 \theta_{\nu W} = 0.251 \pm 0.031(\text{stat}) \pm 0.024(\text{sys})$ .
- <sup>5</sup> Updated result of KamLAND, including the data used in EGUCHI 03. Note that the survival probabilities for different periods are not directly comparable because the effective baseline varies with power output of the reactor sources involved, and there were large variations in the reactor power production in Japan in 2003.
- <sup>6</sup> EGUCHI 03 observe reactor neutrino disappearance at  $\sim 180$  km baseline to various Japanese nuclear power reactors.
- <sup>7</sup> BOEHM 01 search for neutrino oscillations at 0.75 and 0.89 km distance from the Palo Verde reactors.
- <sup>8</sup> APOLLONIO 99, APOLLONIO 98 search for neutrino oscillations at 1.1 km fixed distance from Chooz reactors. They use  $\bar{\nu}_e p \rightarrow e^+ n$  in Gd-loaded scintillator target. APOLLONIO 99 supersedes APOLLONIO 98. See also APOLLONIO 03 for detailed description.
- <sup>9</sup> GREENWOOD 96 search for neutrino oscillations at 18 m and 24 m from the reactor at Savannah River.
- <sup>10</sup> DECLAIS 94 result based on integral measurement of neutrons only. Result is ratio of measured cross section to that expected in standard V-A theory. Replaced by ACHKAR 95.
- <sup>11</sup> KWON 81 represents an analysis of a larger set of data from the same experiment as BOEHM 80.

### ———— Atmospheric neutrinos ————

Neutrinos and antineutrinos produced in the atmosphere induce  $\mu$ -like and  $e$ -like events in underground detectors. The ratio of the numbers of the two kinds of events is defined as  $\mu/e$ . It has the advantage that systematic effects, such as flux uncertainty, tend to cancel, for both experimental and theoretical values of the ratio. The “ratio of the ratios” of experimental to theoretical  $\mu/e$ ,  $R(\mu/e)$ , or that of experimental to theoretical  $\mu/\text{total}$ ,  $R(\mu/\text{total})$  with  $\text{total} = \mu + e$ , is reported below. If the actual value is not unity, the value obtained in a given experiment may depend on the experimental conditions. In addition, the measured “up-down asymmetry” for  $\mu$  ( $N_{up}(\mu)/N_{down}(\mu)$ ) or  $e$  ( $N_{up}(e)/N_{down}(e)$ ) is reported. The expected “up-down asymmetry” is nearly unity if there is no neutrino oscillation.

**$R(\mu/e) = (\text{Measured Ratio } \mu/e) / (\text{Expected Ratio } \mu/e)$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.658 \pm 0.016 \pm 0.035$	<sup>1</sup> ASHIE	05	SKAM sub-GeV
$0.702^{+0.032}_{-0.030} \pm 0.101$	<sup>2</sup> ASHIE	05	SKAM multi-GeV
$0.69 \pm 0.10 \pm 0.06$	<sup>3</sup> SANCHEZ	03	SOU2 Calorimeter raw data
	<sup>4</sup> FUKUDA	96B	KAMI Water Cherenkov
$1.00 \pm 0.15 \pm 0.08$	<sup>5</sup> DAUM	95	FREJ Calorimeter
$0.60^{+0.06}_{-0.05} \pm 0.05$	<sup>6</sup> FUKUDA	94	KAMI sub-GeV
$0.57^{+0.08}_{-0.07} \pm 0.07$	<sup>7</sup> FUKUDA	94	KAMI multi-GeV
	<sup>8</sup> BECKER-SZ...	92B	IMB Water Cherenkov

<sup>1</sup> ASHIE 05 results are based on an exposure of 92 kton yr during the complete Super-Kamiokande I running period. The analyzed data sample consists of fully-contained single-ring  $e$ -like events with  $0.1 \text{ GeV}/c < p_e$  and  $\mu$ -like events  $0.2 \text{ GeV}/c < p_\mu$ , both having a visible energy  $< 1.33 \text{ GeV}$ . These criteria match the definition used by FUKUDA 94.

<sup>2</sup> ASHIE 05 results are based on an exposure of 92 kton yr during the complete Super-Kamiokande I running period. The analyzed data sample consists of fully-contained single-ring events with visible energy  $> 1.33 \text{ GeV}$  and partially-contained events. All partially-contained events are classified as  $\mu$ -like.

<sup>3</sup> SANCHEZ 03 result is based on an exposure of 5.9 kton yr, and updates ALLISON 99 result. The analyzed data sample consists of fully-contained  $e$ -flavor and  $\mu$ -flavor events having lepton momentum  $> 0.3 \text{ GeV}/c$ .

<sup>4</sup> FUKUDA 96B studied neutron background in the atmospheric neutrino sample observed in the Kamiokande detector. No evidence for the background contamination was found.

<sup>5</sup> DAUM 95 results are based on an exposure of 2.0 kton yr which includes the data used by BERGER 90B. This ratio is for the contained and semicontained events. DAUM 95 also report  $R(\mu/e) = 0.99 \pm 0.13 \pm 0.08$  for the total neutrino induced data sample which includes upward going stopping muons and horizontal muons in addition to the contained and semicontained events.

<sup>6</sup> FUKUDA 94 result is based on an exposure of 7.7 kton yr and updates the HIRATA 92 result. The analyzed data sample consists of fully-contained  $e$ -like events with  $0.1 < p_e < 1.33 \text{ GeV}/c$  and fully-contained  $\mu$ -like events with  $0.2 < p_\mu < 1.5 \text{ GeV}/c$ .

<sup>7</sup> FUKUDA 94 analyzed the data sample consisting of fully contained events with visible energy  $> 1.33 \text{ GeV}$  and partially contained  $\mu$ -like events.

<sup>8</sup> BECKER-SZENDY 92B reports the fraction of nonshowering events (mostly muons from atmospheric neutrinos) as  $0.36 \pm 0.02 \pm 0.02$ , as compared with expected fraction  $0.51 \pm 0.01 \pm 0.05$ . After cutting the energy range to the Kamiokande limits, BEIER 92 finds  $R(\mu/e)$  very close to the Kamiokande value.

 **$R(\nu_\mu) = (\text{Measured Flux of } \nu_\mu) / (\text{Expected Flux of } \nu_\mu)$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.84 \pm 0.12$	<sup>1</sup> ADAMSON	06	MINS MINOS atmospheric
$0.72 \pm 0.026 \pm 0.13$	<sup>2</sup> AMBROSIO	01	MCRO upward through-going
$0.57 \pm 0.05 \pm 0.15$	<sup>3</sup> AMBROSIO	00	MCRO upgoing partially contained
$0.71 \pm 0.05 \pm 0.19$	<sup>4</sup> AMBROSIO	00	MCRO downgoing partially contained + upgoing stopping
$0.74 \pm 0.036 \pm 0.046$	<sup>5</sup> AMBROSIO	98	MCRO Streamer tubes

	<sup>6</sup> CASPER	91	IMB	Water Cherenkov
	<sup>7</sup> AGLIETTA	89	NUSX	
0.95 ± 0.22	<sup>8</sup> BOLIEV	81		Baksan
0.62 ± 0.17	CROUCH	78		Case Western/UCI

<sup>1</sup> ADAMSON 06 uses a measurement of 107 total neutrinos compared to an expected rate of  $127 \pm 13$  without oscillations.

<sup>2</sup> AMBROSIO 01 result is based on the upward through-going muon tracks with  $E_\mu > 1$  GeV. The data came from three different detector configurations, but the statistics is largely dominated by the full detector run, from May 1994 to December 2000. The total live time, normalized to the full detector configuration, is 6.17 years. The first error is the statistical error, the second is the systematic error, dominated by the theoretical error in the predicted flux.

<sup>3</sup> AMBROSIO 00 result is based on the upgoing partially contained event sample. It came from 4.1 live years of data taking with the full detector, from April 1994 to February 1999. The average energy of atmospheric muon neutrinos corresponding to this sample is 4 GeV. The first error is statistical, the second is the systematic error, dominated by the 25% theoretical error in the rate (20% in the flux and 15% in the cross section, added in quadrature). Within statistics, the observed deficit is uniform over the zenith angle.

<sup>4</sup> AMBROSIO 00 result is based on the combined samples of downgoing partially contained events and upgoing stopping events. These two subsamples could not be distinguished due to the lack of timing information. The result came from 4.1 live years of data taking with the full detector, from April 1994 to February 1999. The average energy of atmospheric muon neutrinos corresponding to this sample is 4 GeV. The first error is statistical, the second is the systematic error, dominated by the 25% theoretical error in the rate (20% in the flux and 15% in the cross section, added in quadrature). Within statistics, the observed deficit is uniform over the zenith angle.

<sup>5</sup> AMBROSIO 98 result is for all nadir angles and updates AHLEN 95 result. The lower cutoff on the muon energy is 1 GeV. In addition to the statistical and systematic errors, there is a Monte Carlo flux error (theoretical error) of  $\pm 0.13$ . With a neutrino oscillation hypothesis, the fit either to the flux or zenith distribution independently yields  $\sin^2 2\theta = 1.0$  and  $\Delta(m^2) \sim$  a few times  $10^{-3} \text{ eV}^2$ . However, the fit to the observed zenith distribution gives a maximum probability for  $\chi^2$  of only 5% for the best oscillation hypothesis.

<sup>6</sup> CASPER 91 correlates showering/nonshowering signature of single-ring events with parent atmospheric-neutrino flavor. They find nonshowering ( $\approx \nu_\mu$  induced) fraction is  $0.41 \pm 0.03 \pm 0.02$ , as compared with expected  $0.51 \pm 0.05$  (syst).

<sup>7</sup> AGLIETTA 89 finds no evidence for any anomaly in the neutrino flux. They define  $\rho = (\text{measured number of } \nu_e \text{'s}) / (\text{measured number of } \nu_\mu \text{'s})$ . They report  $\rho(\text{measured}) = \rho(\text{expected}) = 0.96^{+0.32}_{-0.28}$ .

<sup>8</sup> From this data BOLIEV 81 obtain the limit  $\Delta(m^2) \leq 6 \times 10^{-3} \text{ eV}^2$  for maximal mixing,  $\nu_\mu \leftrightarrow \nu_\mu$  type oscillation.

### **R( $\mu$ /total) = (Measured Ratio $\mu$ /total) / (Expected Ratio $\mu$ /total)**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.1 <sup>+0.07</sup> <sub>-0.12</sub> ± 0.11	<sup>1</sup> CLARK	97	IMB	multi-GeV
----------------------------------------------	--------------------	----	-----	-----------

<sup>1</sup> CLARK 97 obtained this result by an analysis of fully contained and partially contained events in the IMB water-Cherenkov detector with visible energy  $> 0.95$  GeV.

**$N_{\text{up}}(\mu)/N_{\text{down}}(\mu)$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.551^{+0.035}_{-0.033} \pm 0.004$	<sup>1</sup> ASHIE	05	SKAM multi-GeV
-------------------------------------	--------------------	----	----------------

<sup>1</sup> ASHIE 05 results are based on an exposure of 92 kton yr during the complete Super-Kamiokande I running period. The analyzed data sample consists of fully-contained single-ring  $\mu$ -like events with visible energy  $> 1.33$  GeV and partially-contained events. All partially-contained events are classified as  $\mu$ -like. Upward-going events are those with  $-1 < \cos(\text{zenith angle}) < -0.2$  and downward-going events are those with  $0.2 < \cos(\text{zenith angle}) < 1$ . The  $\mu$ -like up-down ratio for the multi-GeV data deviates from 1 (the expectation for no atmospheric  $\nu_{\mu}$  oscillations) by more than 12 standard deviations.

 **$N_{\text{up}}(e)/N_{\text{down}}(e)$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.961^{+0.086}_{-0.079} \pm 0.016$	<sup>1</sup> ASHIE	05	SKAM multi-GeV
-------------------------------------	--------------------	----	----------------

<sup>1</sup> ASHIE 05 results are based on an exposure of 92 kton yr during the complete Super-Kamiokande I running period. The analyzed data sample consists of fully-contained single-ring  $e$ -like events with visible energy  $> 1.33$  GeV. Upward-going events are those with  $-1 < \cos(\text{zenith angle}) < -0.2$  and downward-going events are those with  $0.2 < \cos(\text{zenith angle}) < 1$ . The  $e$ -like up-down ratio for the multi-GeV data is consistent with 1 (the expectation for no atmospheric  $\nu_e$  oscillations).

 **$R(\text{up/down}; \mu) = (\text{Measured up/down}; \mu) / (\text{Expected up/down}; \mu)$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.62^{+0.19}_{-0.14} \pm 0.02$	<sup>1</sup> ADAMSON	06	MINS atmospheric $\nu$ with far detector
---------------------------------	----------------------	----	------------------------------------------

<sup>1</sup> ADAMSON 06 result is obtained with the MINOS far detector with an exposure of 4.54 kton yr. The expected ratio is calculated with no neutrino oscillation.

 **$R(\mu^+/\mu^-) = (\text{Measured } N(\mu^+)/N(\mu^-)) / (\text{Expected } N(\mu^+)/N(\mu^-))$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.39^{+0.35}_{-0.46} \pm 0.08$	<sup>1</sup> ADAMSON	07	MINS Upward and horizontal $\mu$ with far detector
$0.96^{+0.38}_{-0.27} \pm 0.15$	<sup>2</sup> ADAMSON	06	MINS atmospheric $\nu$ with far detector

<sup>1</sup> ADAMSON 07 result is obtained with the MINOS far detector in 854.24 live days, based on neutrino-induced upward-going and horizontal muons. This result is consistent with *CPT* conservation.

<sup>2</sup> ADAMSON 06 result is obtained with the MINOS far detector with an exposure of 4.54 kton yr, based on contained events. The expected ratio is calculated by assuming the same oscillation parameters for neutrinos and antineutrinos.

## Solar neutrinos

Solar neutrinos are produced by thermonuclear fusion reactions in the Sun. Radiochemical experiments measure particular combinations of fluxes from various neutrino-producing reactions, whereas water-Cherenkov experiments mainly measure a flux of neutrinos from decay of  $^8\text{B}$ . Solar neutrino fluxes are composed of all active neutrino species,  $\nu_e$ ,  $\nu_\mu$ , and  $\nu_\tau$ . In addition, some other mechanisms may cause antineutrino components in solar neutrino fluxes. Each measurement method is sensitive to a particular component or a combination of components of solar neutrino fluxes. For details, see Section 13.4 of Reviews, Tables, and Plots.

### $\nu_e$ Capture Rates from Radiochemical Experiments

1 SNU (Solar Neutrino Unit) =  $10^{-36}$  captures per atom per second.

<u>VALUE (SNU)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
73.4 $\begin{smallmatrix} +6.1 & +3.7 \\ -6.0 & -4.1 \end{smallmatrix}$	<sup>1</sup> KAETHER	10	GALX reanalysis
67.6 $\pm 4.0 \pm 3.2$	<sup>2</sup> KAETHER	10	GNO+GALX reanalysis combined
65.4 $\begin{smallmatrix} +3.1 & +2.6 \\ -3.0 & -2.8 \end{smallmatrix}$	<sup>3</sup> ABDURASHI...	09 SAGE	$^{71}\text{Ga} \rightarrow ^{71}\text{Ge}$
62.9 $\begin{smallmatrix} +5.5 & \pm 2.5 \\ -5.3 & \end{smallmatrix}$	<sup>4</sup> ALTMANN	05 GNO	$^{71}\text{Ga} \rightarrow ^{71}\text{Ge}$
69.3 $\pm 4.1 \pm 3.6$	<sup>5</sup> ALTMANN	05 GNO	GNO + GALX combined
77.5 $\pm 6.2 \begin{smallmatrix} +4.3 \\ -4.7 \end{smallmatrix}$	<sup>6</sup> HAMPEL	99 GALX	$^{71}\text{Ga} \rightarrow ^{71}\text{Ge}$
2.56 $\pm 0.16 \pm 0.16$	<sup>7</sup> CLEVELAND	98 HOME	$^{37}\text{Cl} \rightarrow ^{37}\text{Ar}$

<sup>1</sup> KAETHER 10 reports the reanalysis results of a complete GALLEX data (GALLEX I+II+III+IV, reported in HAMPEL 99) based on the event selection with a new pulse shape analysis, which provides a better background reduction than the rise time analysis adopted in HAMPEL 99.

<sup>2</sup> Combined result of GALLEX I+II+III+IV reanalysis and GNO I+II+III (ALTMANN 05).

<sup>3</sup> ABDURASHITOV 09 reports a combined analysis of 168 extractions of the SAGE solar neutrino experiment during the period January 1990 through December 2007, and updates the ABDURASHITOV 02 result. The data are consistent with the assumption that the solar neutrino production rate is constant in time. Note that a  $\sim 15\%$  systematic uncertainty in the overall normalization may be added to the ABDURASHITOV 09 result, because calibration experiments for gallium solar neutrino measurements using intense  $^{51}\text{Cr}$  (twice by GALLEX and once by SAGE) and  $^{37}\text{Ar}$  (by SAGE) result in an average ratio of  $0.87 \pm 0.05$  of the observed to calculated rates.

<sup>4</sup> ALTMANN 05 reports the complete result from the GNO solar neutrino experiment (GNO I+II+III), which is the successor project of GALLEX. Experimental technique of GNO is essentially the same as that of GALLEX. The run data cover the period 20 May 1998 through 9 April 2003.

<sup>5</sup> Combined result of GALLEX I+II+III+IV (HAMPEL 99) and GNO I+II+III.

<sup>6</sup> HAMPEL 99 report the combined result for GALLEX I+II+III+IV (65 runs in total), which update the HAMPEL 96 result. The GALLEX IV result (12 runs) is  $118.4 \pm 17.8 \pm 6.6$  SNU. (HAMPEL 99 discuss the consistency of partial results with the mean.) The GALLEX experimental program has been completed with these runs. The total run data cover the period 14 May 1991 through 23 January 1997. A total of 300  $^{71}\text{Ge}$  events were observed. Note that a  $\sim 15\%$  systematic uncertainty in the overall normalization may be added to the HAMPEL 99 result, because calibration experiments for gallium solar neutrino measurements using intense  $^{51}\text{Cr}$  (twice by GALLEX and once by SAGE)

and  $^{37}\text{Ar}$  (by SAGE) result in an average ratio of  $0.87 \pm 0.05$  of the observed to calculated rates.

<sup>7</sup> CLEVELAND 98 is a detailed report of the  $^{37}\text{Cl}$  experiment at the Homestake Mine. The average solar neutrino-induced  $^{37}\text{Ar}$  production rate from 108 runs between 1970 and 1994 updates the DAVIS 89 result.

### $\phi_{ES} (^8\text{B})$

$^8\text{B}$  solar-neutrino flux measured via  $\nu e$  elastic scattering. This process is sensitive to all active neutrino flavors, but with reduced sensitivity to  $\nu_\mu, \nu_\tau$  due to the cross-section difference,  $\sigma(\nu_{\mu,\tau} e) \sim 0.16\sigma(\nu_e e)$ . If the  $^8\text{B}$  solar-neutrino flux involves nonelectron flavor active neutrinos, their contribution to the flux is  $\sim 0.16$  times of  $\nu_e$ .

<u>VALUE (<math>10^6 \text{ cm}^{-2}\text{s}^{-1}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$2.32 \pm 0.04 \pm 0.05$	<sup>1</sup> ABE	11	SKAM SK-III average flux
$2.41 \pm 0.05^{+0.16}_{-0.15}$	<sup>2</sup> ABE	11	SKAM SK-II average flux
$2.38 \pm 0.02 \pm 0.08$	<sup>3</sup> ABE	11	SKAM SK-I average flux
$2.77 \pm 0.26 \pm 0.32$	<sup>4</sup> ABE	11B	KLND average flux
$2.4 \pm 0.4 \pm 0.1$	<sup>5</sup> BELLINI	10A	BORX average flux
$1.77^{+0.24+0.09}_{-0.21-0.10}$	<sup>6</sup> AHARMIM	08	SNO Phase III
$2.38 \pm 0.05^{+0.16}_{-0.15}$	<sup>7</sup> CRAVENS	08	SKAM average flux
$2.35 \pm 0.02 \pm 0.08$	<sup>8</sup> HOSAKA	06	SKAM average flux
$2.35 \pm 0.22 \pm 0.15$	<sup>9</sup> AHARMIM	05A	SNO Salty $\text{D}_2\text{O}$ ; $^8\text{B}$ shape not constrained
$2.34 \pm 0.23^{+0.15}_{-0.14}$	<sup>9</sup> AHARMIM	05A	SNO Salty $\text{D}_2\text{O}$ ; $^8\text{B}$ shape constrained
$2.39^{+0.24}_{-0.23} \pm 0.12$	<sup>10</sup> AHMAD	02	SNO average flux
$2.39 \pm 0.34^{+0.16}_{-0.14}$	<sup>11</sup> AHMAD	01	SNO average flux
$2.80 \pm 0.19 \pm 0.33$	<sup>12</sup> FUKUDA	96	KAMI average flux
$2.70 \pm 0.27$	<sup>12</sup> FUKUDA	96	KAMI day flux
$2.87^{+0.27}_{-0.26}$	<sup>12</sup> FUKUDA	96	KAMI night flux

<sup>1</sup> ABE 11 reports the Super-Kamiokande-III results for 548 live days from August 4, 2006 to August 18, 2008. The analysis threshold is 5.0 MeV, but the event sample in the 5.0–6.5 MeV total electron range has a total live time of 298 days.

<sup>2</sup> ABE 11 recalculated the Super-Kamiokande-II results using  $^8\text{B}$  spectrum of WINTER 06A.

<sup>3</sup> ABE 11 recalculated the Super-Kamiokande-I results using  $^8\text{B}$  spectrum of WINTER 06A.

<sup>4</sup> ABE 11B use a 123 kton-day exposure of the KamLAND liquid scintillation detector to measure the  $^8\text{B}$  solar neutrino flux. They utilize  $\nu - e$  elastic scattering above a reconstructed-energy threshold of 5.5 MeV, corresponding to 5 MeV electron recoil energy. 299 electron recoil candidate events are reported, of which  $157 \pm 23.6$  are assigned to background.

<sup>5</sup> BELLINI 10A reports the Borexino result with 3 MeV energy threshold for scattered electrons. The data correspond to 345.3 live days with a target mass of 100 t, between July 15, 2007 and August 23, 2009.

<sup>6</sup> AHARMIM 08 reports the results from SNO Phase III measurement using an array of  $^3\text{He}$  proportional counters to measure the rate of NC interactions in heavy water, over the period between November 27, 2004 and November 28, 2006, corresponding to 385.17

live days. A simultaneous fit was made for the number of NC events detected by the proportional counters and the numbers of NC, CC, and ES events detected by the PMTs, where the spectral distributions of the ES and CC events were not constrained to the  $^8\text{B}$  shape.

- 7 CRAVENS 08 reports the Super-Kamiokande-II results for 791 live days from December 2002 to October 2005. The photocathode coverage of the detector is 19% (reduced from 40% of that of Super-Kamiokande-I due to an accident in 2001). The analysis threshold for the average flux is 7 MeV.
- 8 HOSAKA 06 reports the final results for 1496 live days with Super-Kamiokande-I between May 31, 1996 and July 15, 2001, and replace FUKUDA 02 results. The analysis threshold is 5 MeV except for the first 280 live days (6.5 MeV).
- 9 AHARMIM 05A measurements were made with dissolved NaCl (0.195% by weight) in heavy water over the period between July 26, 2001 and August 28, 2003, corresponding to 391.4 live days, and update AHMED 04A. The CC, ES, and NC events were statistically separated. In one method, the  $^8\text{B}$  energy spectrum was not constrained. In the other method, the constraint of an undistorted  $^8\text{B}$  energy spectrum was added for comparison with AHMAD 02 results.
- 10 AHMAD 02 reports the  $^8\text{B}$  solar-neutrino flux measured via  $\nu e$  elastic scattering above the kinetic energy threshold of 5 MeV. The data correspond to 306.4 live days with SNO between November 2, 1999 and May 28, 2001, and updates AHMAD 01 results.
- 11 AHMAD 01 reports the  $^8\text{B}$  solar-neutrino flux measured via  $\nu e$  elastic scattering above the kinetic energy threshold of 6.75 MeV. The data correspond to 241 live days with SNO between November 2, 1999 and January 15, 2001.
- 12 FUKUDA 96 results are for a total of 2079 live days with Kamiokande II and III from January 1987 through February 1995, covering the entire solar cycle 22, with threshold  $E_e > 9.3$  MeV (first 449 days),  $> 7.5$  MeV (middle 794 days), and  $> 7.0$  MeV (last 836 days). These results update the HIRATA 90 result for the average  $^8\text{B}$  solar-neutrino flux and HIRATA 91 result for the day-night variation in the  $^8\text{B}$  solar-neutrino flux. The total data sample was also analyzed for short-term variations: within experimental errors, no strong correlation of the solar-neutrino flux with the sunspot numbers was found.

### $\phi_{\text{CC}} (^8\text{B})$

$^8\text{B}$  solar-neutrino flux measured with charged-current reaction which is sensitive exclusively to  $\nu_e$ .

<u>VALUE (<math>10^6 \text{ cm}^{-2}\text{s}^{-1}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$1.67^{+0.05+0.07}_{-0.04-0.08}$	1 AHARMIM	08 SNO	Phase III
$1.68 \pm 0.06^{+0.08}_{-0.09}$	2 AHARMIM	05A SNO	Salty $\text{D}_2\text{O}$ ; $^8\text{B}$ shape not const.
$1.72 \pm 0.05 \pm 0.11$	2 AHARMIM	05A SNO	Salty $\text{D}_2\text{O}$ ; $^8\text{B}$ shape constrained
$1.76^{+0.06}_{-0.05} \pm 0.09$	3 AHMAD	02 SNO	average flux
$1.75 \pm 0.07^{+0.12}_{-0.11} \pm 0.05$	4 AHMAD	01 SNO	average flux

- 1 AHARMIM 08 reports the results from SNO Phase III measurement using an array of  $^3\text{He}$  proportional counters to measure the rate of NC interactions in heavy water, over the period between November 27, 2004 and November 28, 2006, corresponding to 385.17 live days. A simultaneous fit was made for the number of NC events detected by the proportional counters and the numbers of NC, CC, and ES events detected by the PMTs, where the spectral distributions of the ES and CC events were not constrained to the  $^8\text{B}$  shape.

- <sup>2</sup> AHARMIM 05A measurements were made with dissolved NaCl (0.195% by weight) in heavy water over the period between July 26, 2001 and August 28, 2003, corresponding to 391.4 live days, and update AHMED 04A. The *CC*, *ES*, and *NC* events were statistically separated. In one method, the <sup>8</sup>B energy spectrum was not constrained. In the other method, the constraint of an undistorted <sup>8</sup>B energy spectrum was added for comparison with AHMAD 02 results.
- <sup>3</sup> AHMAD 02 reports the SNO result of the <sup>8</sup>B solar-neutrino flux measured with charged-current reaction on deuterium,  $\nu_e d \rightarrow ppe^-$ , above the kinetic energy threshold of 5 MeV. The data correspond to 306.4 live days with SNO between November 2, 1999 and May 28, 2001, and updates AHMAD 01 results. The complete description of the SNO Phase I data set is given in AHARMIM 07.
- <sup>4</sup> AHMAD 01 reports the first SNO result of the <sup>8</sup>B solar-neutrino flux measured with the charged-current reaction on deuterium,  $\nu_e d \rightarrow ppe^-$ , above the kinetic energy threshold of 6.75 MeV. The data correspond to 241 live days with SNO between November 2, 1999 and January 15, 2001.

### $\phi_{NC} (^8B)$

<sup>8</sup>B solar neutrino flux measured with neutral-current reaction, which is equally sensitive to  $\nu_e$ ,  $\nu_\mu$ , and  $\nu_\tau$ .

VALUE ( $10^6 \text{ cm}^{-2}\text{s}^{-1}$ )	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
5.140 <sup>+0.160+0.132</sup> <sub>-0.158-0.117</sub>	1 AHARMIM	10 SNO	Phase I+II, low threshold
5.54 <sup>+0.33+0.36</sup> <sub>-0.31-0.34</sub>	2 AHARMIM	08 SNO	Phase III, prop. counter + PMT
4.94 <sup>±0.21+0.38</sup> <sub>-0.34</sub>	3 AHARMIM	05A SNO	Salty D <sub>2</sub> O; <sup>8</sup> B shape not const.
4.81 <sup>±0.19+0.28</sup> <sub>-0.27</sub>	3 AHARMIM	05A SNO	Salty D <sub>2</sub> O; <sup>8</sup> B shape constrained
5.09 <sup>+0.44+0.46</sup> <sub>-0.43-0.43</sub>	4 AHMAD	02 SNO	average flux; <sup>8</sup> B shape const.
6.42 <sup>±1.57+0.55</sup> <sub>-0.58</sub>	4 AHMAD	02 SNO	average flux; <sup>8</sup> B shape not const.

- <sup>1</sup> AHARMIM 10 reports this result from a joint analysis of SNO Phase I+II data with the "effective electron kinetic energy" threshold of 3.5 MeV. This result is obtained with a "binned-histogram unconstrained fit" where binned probability distribution functions of the neutrino signal observables were used without any model constraints on the shape of the neutrino spectrum.
- <sup>2</sup> AHARMIM 08 reports the results from SNO Phase III measurement using an array of <sup>3</sup>He proportional counters to measure the rate of NC interactions in heavy water, over the period between November 27, 2004 and November 28, 2006, corresponding to 385.17 live days. A simultaneous fit was made for the number of NC events detected by the proportional counters and the numbers of NC, CC, and ES events detected by the PMTs, where the spectral distributions of the ES and CC events were not constrained to the <sup>8</sup>B shape.
- <sup>3</sup> AHARMIM 05A measurements were made with dissolved NaCl (0.195% by weight) in heavy water over the period between July 26, 2001 and August 28, 2003, corresponding to 391.4 live days, and update AHMED 04A. The *CC*, *ES*, and *NC* events were statistically separated. In one method, the <sup>8</sup>B energy spectrum was not constrained. In the other method, the constraint of an undistorted <sup>8</sup>B energy spectrum was added for comparison with AHMAD 02 results.
- <sup>4</sup> AHMAD 02 reports the first SNO result of the <sup>8</sup>B solar-neutrino flux measured with the neutral-current reaction on deuterium,  $\nu_\ell d \rightarrow n p \nu_\ell$ , above the neutral-current reaction threshold of 2.2 MeV. The data correspond to 306.4 live days with SNO between November 2, 1999 and May 28, 2001. The complete description of the SNO Phase I data set is given in AHARMIM 07.

## $\phi_{\nu_{\mu}+\nu_{\tau}}$ ( $^8\text{B}$ )

Nonelectron-flavor active neutrino component ( $\nu_{\mu}$  and  $\nu_{\tau}$ ) in the  $^8\text{B}$  solar-neutrino flux.

<u>VALUE (<math>10^6 \text{ cm}^{-2}\text{s}^{-1}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$3.26 \pm 0.25^{+0.40}_{-0.35}$	<sup>1</sup> AHARMIM	05A SNO	From $\phi_{NC}$ , $\phi_{CC}$ , and $\phi_{ES}$ ; $^8\text{B}$ shape not const.
$3.09 \pm 0.22^{+0.30}_{-0.27}$	<sup>1</sup> AHARMIM	05A SNO	From $\phi_{NC}$ , $\phi_{CC}$ , and $\phi_{ES}$ ; $^8\text{B}$ shape constrained
$3.41 \pm 0.45^{+0.48}_{-0.45}$	<sup>2</sup> AHMAD	02 SNO	From $\phi_{NC}$ , $\phi_{CC}$ , and $\phi_{ES}$
$3.69 \pm 1.13$	<sup>3</sup> AHMAD	01	Derived from SNO+SuperKam, water Cherenkov

<sup>1</sup> AHARMIM 05A measurements were made with dissolved NaCl (0.195% by weight) in heavy water over the period between July 26, 2001 and August 28, 2003, corresponding to 391.4 live days, and update AHMED 04A. The *CC*, *ES*, and *NC* events were statistically separated. In one method, the  $^8\text{B}$  energy spectrum was not constrained. In the other method, the constraint of an undistorted  $^8\text{B}$  energy spectrum was added for comparison with AHMAD 02 results.

<sup>2</sup> AHMAD 02 deduced the nonelectron-flavor active neutrino component ( $\nu_{\mu}$  and  $\nu_{\tau}$ ) in the  $^8\text{B}$  solar-neutrino flux, by combining the charged-current result, the  $\nu e$  elastic-scattering result and the neutral-current result. The complete description of the SNO Phase I data set is given in AHARMIM 07.

<sup>3</sup> AHMAD 01 deduced the nonelectron-flavor active neutrino component ( $\nu_{\mu}$  and  $\nu_{\tau}$ ) in the  $^8\text{B}$  solar-neutrino flux, by combining the SNO charged-current result (AHMAD 01) and the Super-Kamiokande  $\nu e$  elastic-scattering result (FUKUDA 01).

## Total Flux of Active $^8\text{B}$ Solar Neutrinos

Total flux of active neutrinos ( $\nu_e$ ,  $\nu_{\mu}$ , and  $\nu_{\tau}$ ).

<u>VALUE (<math>10^6 \text{ cm}^{-2}\text{s}^{-1}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$5.046^{+0.159+0.107}_{-0.152-0.123}$	<sup>1</sup> AHARMIM	10 SNO	From $\phi_{NC}$ in Phase III
$5.54^{+0.33+0.36}_{-0.31-0.34}$	<sup>2</sup> AHARMIM	08 SNO	$\phi_{NC}$ in Phase III
$4.94 \pm 0.21^{+0.38}_{-0.34}$	<sup>3</sup> AHARMIM	05A SNO	From $\phi_{NC}$ ; $^8\text{B}$ shape not const.
$4.81 \pm 0.19^{+0.28}_{-0.27}$	<sup>3</sup> AHARMIM	05A SNO	From $\phi_{NC}$ ; $^8\text{B}$ shape constrained
$5.09^{+0.44+0.46}_{-0.43-0.43}$	<sup>4</sup> AHMAD	02 SNO	Direct measurement from $\phi_{NC}$
$5.44 \pm 0.99$	<sup>5</sup> AHMAD	01	Derived from SNO+SuperKam, water Cherenkov

<sup>1</sup> AHARMIM 10 reports this result from a joint analysis of SNO Phase I+II data with the "effective electron kinetic energy" threshold of 3.5 MeV. This result is obtained with the assumption of unitarity, which relates the NC, CC, and ES rates. The data were fit with the free parameters directly describing the total  $^8\text{B}$  neutrino flux and the energy-dependent  $\nu_e$  survival probability.

<sup>2</sup> AHARMIM 08 reports the results from SNO Phase III measurement using an array of  $^3\text{He}$  proportional counters to measure the rate of NC interactions in heavy water, over

the period between November 27, 2004 and November 28, 2006, corresponding to 385.17 live days. A simultaneous fit was made for the number of NC events detected by the proportional counters and the numbers of NC, CC, and ES events detected by the PMTs, where the spectral distributions of the ES and CC events were not constrained to the  $^8\text{B}$  shape.

<sup>3</sup> AHARMIM 05A measurements were made with dissolved NaCl (0.195% by weight) in heavy water over the period between July 26, 2001 and August 28, 2003, corresponding to 391.4 live days, and update AHMED 04A. The CC, ES, and NC events were statistically separated. In one method, the  $^8\text{B}$  energy spectrum was not constrained. In the other method, the constraint of an undistorted  $^8\text{B}$  energy spectrum was added for comparison with AHMAD 02 results.

<sup>4</sup> AHMAD 02 determined the total flux of active  $^8\text{B}$  solar neutrinos by directly measuring the neutral-current reaction,  $\nu_\ell d \rightarrow n p \nu_\ell$ , which is equally sensitive to  $\nu_e$ ,  $\nu_\mu$ , and  $\nu_\tau$ . The complete description of the SNO Phase I data set is given in AHARMIM 07.

<sup>5</sup> AHMAD 01 deduced the total flux of active  $^8\text{B}$  solar neutrinos by combining the SNO charged-current result (AHMAD 01) and the Super-Kamiokande  $\nu e$  elastic-scattering result (FUKUDA 01).

### Day-Night Asymmetry ( $^8\text{B}$ )

$$A = (\phi_{\text{night}} - \phi_{\text{day}}) / \phi_{\text{average}}$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.063 \pm 0.042 \pm 0.037$	<sup>1</sup> CRAVENS	08 SKAM	Based on $\phi_{ES}$
$0.021 \pm 0.020 \begin{smallmatrix} +0.012 \\ -0.013 \end{smallmatrix}$	<sup>2</sup> HOSAKA	06 SKAM	Based on $\phi_{ES}$
$0.017 \pm 0.016 \begin{smallmatrix} +0.012 \\ -0.013 \end{smallmatrix}$	<sup>3</sup> HOSAKA	06 SKAM	Fitted in the LMA region
$-0.056 \pm 0.074 \pm 0.053$	<sup>4</sup> AHARMIM	05A SNO	From salty SNO $\phi_{CC}$
$-0.037 \pm 0.063 \pm 0.032$	<sup>4</sup> AHARMIM	05A SNO	From salty SNO $\phi_{CC}$ ; const. of no $\phi_{NC}$ asymmetry
$0.14 \pm 0.063 \begin{smallmatrix} +0.015 \\ -0.014 \end{smallmatrix}$	<sup>5</sup> AHMAD	02B SNO	Derived from SNO $\phi_{CC}$
$0.07 \pm 0.049 \begin{smallmatrix} +0.013 \\ -0.012 \end{smallmatrix}$	<sup>6</sup> AHMAD	02B SNO	Const. of no $\phi_{NC}$ asymmetry

<sup>1</sup> CRAVENS 08 reports the Super-Kamiokande-II results for 791 live days from December 2002 to October 2005. The photocathode coverage of the detector is 19% (reduced from 40% of that of Super-Kamiokande-I due to an accident in 2001). The analysis threshold for the day and night fluxes is 7.5 MeV.

<sup>2</sup> HOSAKA 06 reports the final results for 1496 live days with Super-Kamiokande-I between May 31, 1996 and July 15, 2001, and replace FUKUDA 02 results. The analysis threshold is 5 MeV except for the first 280 live days (6.5 MeV).

<sup>3</sup> This result with reduced statistical uncertainty is obtained by assuming two-neutrino oscillations within the LMA (large mixing angle) region and by fitting the time variation of the solar neutrino flux measured via  $\nu_e$  elastic scattering to the variations expected from neutrino oscillations. For details, see SMY 04. There is an additional small systematic error of  $\pm 0.0004$  coming from uncertainty of oscillation parameters.

<sup>4</sup> AHARMIM 05A measurements were made with dissolved NaCl (0.195% by weight) in heavy water over the period between July 26, 2001 and August 28, 2003, with 176.5 days of the live time recorded during the day and 214.9 days during the night. This result is obtained with the spectral distribution of the CC events not constrained to the  $^8\text{B}$  shape.

<sup>5</sup> AHMAD 02B results are based on the charged-current interactions recorded between November 2, 1999 and May 28, 2001, with the day and night live times of 128.5 and 177.9 days, respectively. The complete description of the SNO Phase I data set is given in AHARMIM 07.

<sup>6</sup> AHMAD 02B results are derived from the charged-current interactions, neutral-current interactions, and  $\nu e$  elastic scattering, with the total flux of active neutrinos constrained to have no asymmetry. The data were recorded between November 2, 1999 and May 28, 2001, with the day and night live times of 128.5 and 177.9 days, respectively. The complete description of the SNO Phase I data set is given in AHARMIM 07.

### $\phi_{ES} (^7\text{Be})$

<sup>7</sup>Be solar-neutrino flux measured via  $\nu_e$  elastic scattering. This process is sensitive to all active neutrino flavors, but with reduced sensitivity to  $\nu_\mu, \nu_\tau$  due to the cross-section difference,  $\sigma(\nu_{\mu,\tau} e) \sim 0.2 \sigma(\nu_e e)$ . If the <sup>7</sup>Be solar-neutrino flux involves nonelectron flavor active neutrinos, their contribution to the flux is  $\sim 0.2$  times that of  $\nu_e$ .

<u>VALUE (<math>10^9 \text{ cm}^{-2} \text{ s}^{-1}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-----------------------------------------------------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.10 \pm 0.15$	<sup>1</sup> BELLINI	11A BORX	average flux
-----------------	----------------------	----------	--------------

<sup>1</sup> BELLINI 11A reports the <sup>7</sup>Be solar neutrino flux measured via  $\nu - e$  elastic scattering. The data correspond to 740.7 live days between May 16, 2007 and May 8, 2010, and also correspond to 153.6 ton-year fiducial exposure. BELLINI 11A measured the 862 keV <sup>7</sup>Be solar neutrino flux, which is an 89.6% branch of the <sup>7</sup>Be solar neutrino flux, to be  $(2.78 \pm 0.13) \times 10^9 \text{ cm}^{-2} \text{ s}^{-1}$ . Supercedes ARPESELLA 08A.

### $\phi_{CC}(pp)$

$pp$  solar-neutrino flux measured with charged-current reaction which is sensitive exclusively to  $\nu_e$ .

<u>VALUE (<math>10^{10} \text{ cm}^{-2} \text{ s}^{-1}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------------------------------------------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.38 \pm 0.47$	<sup>1</sup> ABDURASHI...	09 FIT	Fit existing solar- $\nu$ data
-----------------	---------------------------	--------	--------------------------------

<sup>1</sup> ABDURASHITOV 09 reports the  $pp$  solar-neutrino flux derived from the Ga solar neutrino capture rate by subtracting contributions from <sup>8</sup>B, <sup>7</sup>Be,  $p e p$  and CNO solar neutrino fluxes determined by other solar neutrino experiments as well as neutrino oscillation parameters determined from available world neutrino oscillation data.

### $\phi_{ES} (\text{hep})$

hep solar-neutrino flux measured via  $\nu e$  elastic scattering. This process is sensitive to all active neutrino flavors, but with reduced sensitivity to  $\nu_\mu, \nu_\tau$  due to the cross-section difference,  $\sigma(\nu_{\mu,\tau} e) \sim 0.16 \sigma(\nu_e e)$ . If the hep solar-neutrino flux involves nonelectron flavor active neutrinos, their contribution to the flux is  $\sim 0.16$  times of  $\nu_e$ .

<u>VALUE (<math>10^3 \text{ cm}^{-2} \text{ s}^{-1}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
-----------------------------------------------------------------	------------	--------------------	-------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 73$	90	<sup>1</sup> HOSAKA	06 SKAM
--------	----	---------------------	---------

<sup>1</sup> HOSAKA 06 result is obtained from the recoil electron energy window of 18–21 MeV, and updates FUKUDA 01 result.

### $\phi_{\bar{\nu}_e} (^8\text{B})$

Searches are made for electron antineutrino flux from the Sun. Flux limits listed here are derived relative to the BS05(OP) Standard Solar Model  $^8\text{B}$  solar neutrino flux ( $5.69 \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$ ), with an assumption that solar  $\bar{\nu}_e$ s follow an unoscillated  $^8\text{B}$  neutrino spectrum.

<u>VALUE (%)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.013	90	BELLINI	11	BORX $E_{\bar{\nu}_e} > 1.8 \text{ MeV}$
<1.9	90	<sup>1</sup> BALATA	06	CNTR $1.8 < E_{\bar{\nu}_e} < 20.0 \text{ MeV}$
<0.72	90	AHARMIM	04	SNO $4.0 < E_{\bar{\nu}_e} < 14.8 \text{ MeV}$
<0.022	90	EGUCHI	04	KLND $8.3 < E_{\bar{\nu}_e} < 14.8 \text{ MeV}$
<0.7	90	GANDO	03	SKAM $8.0 < E_{\bar{\nu}_e} < 20.0 \text{ MeV}$
<1.7	90	AGLIETTA	96	LSD $7 < E_{\bar{\nu}_e} < 17 \text{ MeV}$

<sup>1</sup>BALATA 06 obtained this result from the search for  $\bar{\nu}_e$  interactions with Counting Test Facility (the prototype of the Borexino detector).

### (B) Three-neutrino mixing parameters

A REVIEW GOES HERE – Check our WWW List of Reviews

### $\sin^2(2\theta_{12})$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.857^{+0.023}_{-0.025}</math></b>	<sup>1</sup> GANDO	11	FIT KamLAND + solar: $3\nu$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.85 \pm 0.02$	<sup>2</sup> ABE	11	FIT KamLAND + global solar: $2\nu$
$0.84^{+0.03}_{-0.02}$	<sup>3</sup> ABE	11	FIT global solar: $2\nu$
$0.85^{+0.04}_{-0.03}$	<sup>4</sup> ABE	11	FIT KamLAND + global solar: $3\nu$
$0.85^{+0.04}_{-0.05}$	<sup>5</sup> ABE	11	FIT global solar: $3\nu$
$0.861^{+0.022}_{-0.018}$	<sup>6</sup> BELLINI	11A	FIT KamLAND + global solar: $2\nu$
$0.869^{+0.024}_{-0.022}$	<sup>7</sup> BELLINI	11A	FIT global solar: $2\nu$
$0.846^{+0.064}_{-0.073}$	<sup>8</sup> GANDO	11	FIT KamLAND: $3\nu$
$0.861^{+0.026}_{-0.022}$	<sup>9,10</sup> AHARMIM	10	FIT KamLAND + global solar: $2\nu$
$0.861^{+0.024}_{-0.031}$	<sup>9,11</sup> AHARMIM	10	FIT global solar: $2\nu$
$0.869^{+0.026}_{-0.024}$	<sup>9,12</sup> AHARMIM	10	FIT KamLAND + global solar: $3\nu$
$0.869^{+0.031}_{-0.037}$	<sup>9,13</sup> AHARMIM	10	FIT global solar: $3\nu$
$0.92 \pm 0.05$	<sup>14</sup> ABE	08A	FIT KamLAND
$0.87 \pm 0.04$	<sup>15</sup> ABE	08A	FIT KamLAND + global fit
$0.87 \pm 0.03$	<sup>16</sup> AHARMIM	08	FIT KamLAND + global solar

0.85 $\begin{smallmatrix} +0.04 \\ -0.06 \end{smallmatrix}$	17 HOSAKA	06	FIT	KamLAND + global solar
0.85 $\begin{smallmatrix} +0.06 \\ -0.05 \end{smallmatrix}$	18 HOSAKA	06	FIT	SKAM+SNO+KamLAND
0.86 $\begin{smallmatrix} +0.05 \\ -0.07 \end{smallmatrix}$	19 HOSAKA	06	FIT	SKAM+SNO
0.86 $\begin{smallmatrix} +0.03 \\ -0.04 \end{smallmatrix}$	20 AHARMIM	05A	FIT	KamLAND + global solar
0.75–0.95	21 AHARMIM	05A	FIT	global solar
0.82 $\pm 0.05$	22 ARAKI	05	FIT	KamLAND + global solar
0.82 $\pm 0.04$	23 AHMED	04A	FIT	KamLAND + global solar
0.71–0.93	24 AHMED	04A	FIT	global solar
0.85 $\begin{smallmatrix} +0.05 \\ -0.07 \end{smallmatrix}$	25 SMY	04	FIT	KamLAND + global solar
0.83 $\begin{smallmatrix} +0.06 \\ -0.08 \end{smallmatrix}$	26 SMY	04	FIT	global solar
0.87 $\begin{smallmatrix} +0.07 \\ -0.08 \end{smallmatrix}$	27 SMY	04	FIT	SKAM + SNO
0.62–0.88	28 AHMAD	02B	FIT	global solar
0.62–0.95	29 FUKUDA	02	FIT	global solar

<sup>1</sup> GANDO 11 obtain this result with three-neutrino fit using the KamLAND + solar data. Supersedes ABE 08A.

<sup>2</sup> ABE 11 obtained this result by a two-neutrino oscillation analysis using solar neutrino data including Super-Kamiokande, SNO, Borexino (ARPESELLA 08A), Homestake, GALLEX/GNO, SAGE, and KamLAND data. CPT invariance is assumed.

<sup>3</sup> ABE 11 obtained this result by a two-neutrino oscillation analysis using solar neutrino data including Super-Kamiokande, SNO, Borexino (ARPESELLA 08A), Homestake, GALLEX/GNO, and SAGE data.

<sup>4</sup> ABE 11 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{32}^2$  fixed to  $2.4 \times 10^{-3} \text{ eV}^2$ , using solar neutrino data including Super-Kamiokande, SNO, Borexino (ARPESELLA 08A), Homestake, GALLEX/GNO, SAGE, and KamLAND data. The normal neutrino mass hierarchy and CPT invariance are assumed.

<sup>5</sup> ABE 11 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{32}^2$  fixed to  $2.4 \times 10^{-3} \text{ eV}^2$ , using solar neutrino data including Super-Kamiokande, SNO, Borexino (ARPESELLA 08A), Homestake, and GALLEX/GNO data. The normal neutrino mass hierarchy is assumed.

<sup>6</sup> BELLINI 11A obtained this result by a two-neutrino oscillation analysis using KamLAND, Homestake, SAGE, Gallex, GNO, Kamiokande, Super-Kamiokande, SNO, and Borexino (BELLINI 11A) data and the SSM flux prediction in SERENELLI 11 (Astrophysical Journal **743** 24 (2011)) with the exception that the  $^8\text{B}$  flux was left free. CPT invariance is assumed.

<sup>7</sup> BELLINI 11A obtained this result by a two-neutrino oscillation analysis using Homestake, SAGE, Gallex, GNO, Kamiokande, Super-Kamiokande, SNO, and Borexino (BELLINI 11A) data and the SSM flux prediction in SERENELLI 11 (Astrophysical Journal **743** 24 (2011)) with the exception that the  $^8\text{B}$  flux was left free.

<sup>8</sup> GANDO 11 obtain this result with three-neutrino fit using the KamLAND data only. Supersedes ABE 08A.

<sup>9</sup> AHARMIM 10 global solar neutrino data include SNO's low-energy-threshold analysis survival probability day/night curves, SNO Phase III integral rates (AHARMIM 08), CI (CLEVELAND 98), SAGE (ABDURASHITOV 09), Gallex/GNO (HAMPEL 99, ALTMANN 05), Borexino (ARPESELLA 08A), SK-I zenith (HOSAKA 06), and SK-II day/night spectra (CRAVENS 08).

<sup>10</sup> AHARMIM 10 obtained this result by a two-neutrino oscillation analysis using global solar neutrino data and KamLAND data (ABE 08A). CPT invariance is assumed.

- 11 AHARMIM 10 obtained this result by a two-neutrino oscillation analysis using global solar neutrino data.
- 12 AHARMIM 10 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{31}^2$  fixed to  $2.3 \times 10^{-3} \text{ eV}^2$ , using global solar neutrino data and KamLAND data (ABE 08A). *CPT* invariance is assumed.
- 13 AHARMIM 10 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{31}^2$  fixed to  $2.3 \times 10^{-3} \text{ eV}^2$ , using global solar neutrino data.
- 14 ABE 08A obtained this result by a rate + shape + time combined geoneutrino and reactor two-neutrino fit for  $\Delta m_{21}^2$  and  $\tan^2 \theta_{12}$ , using KamLAND data only.
- 15 ABE 08A obtained this result by means of a two-neutrino fit using KamLAND, Homestake, SAGE, GALLEX, GNO, SK (zenith angle and E-spectrum), the SNO  $\chi^2$ -map, and solar flux data. *CPT* invariance is assumed.
- 16 The result given by AHARMIM 08 is  $\theta = (34.4_{-1.2}^{+1.3})^\circ$ . This result is obtained by a two-neutrino oscillation analysis using solar neutrino data including those of Borexino (ARPESELLA 08A) and Super-Kamiokande-I (HOSAKA 06), and KamLAND data (ABE 08A). *CPT* invariance is assumed.
- 17 HOSAKA 06 obtained this result by a two-neutrino oscillation analysis using SK  $\nu_e$  data, CC data from other solar neutrino experiments, and KamLAND data (ARAKI 05). *CPT* invariance is assumed.
- 18 HOSAKA 06 obtained this result by a two-neutrino oscillation analysis using the data from Super-Kamiokande, SNO (AHMAD 02 and AHMAD 02B), and KamLAND (ARAKI 05) experiments. *CPT* invariance is assumed.
- 19 HOSAKA 06 obtained this result by a two-neutrino oscillation analysis using the Super-Kamiokande and SNO (AHMAD 02 and AHMAD 02B) solar neutrino data.
- 20 The result given by AHARMIM 05A is  $\theta = (33.9 \pm 1.6)^\circ$ . This result is obtained by a two-neutrino oscillation analysis using SNO pure deuteron and salt phase data, SK  $\nu_e$  data, Cl and Ga CC data, and KamLAND data (ARAKI 05). *CPT* invariance is assumed. AHARMIM 05A also quotes  $\theta = (33.9_{-2.2}^{+2.4})^\circ$  as the error enveloping the 68% CL two-dimensional region. This translates into  $\sin^2 2\theta = 0.86_{-0.06}^{+0.05}$ .
- 21 AHARMIM 05A obtained this result by a two-neutrino oscillation analysis using the data from all solar neutrino experiments. The listed range of the parameter envelops the 95% CL two-dimensional region shown in figure 35a of AHARMIM 05A. AHARMIM 05A also quotes  $\tan^2 \theta = 0.45_{-0.08}^{+0.09}$  as the error enveloping the 68% CL two-dimensional region. This translates into  $\sin^2 2\theta = 0.86_{-0.07}^{+0.05}$ .
- 22 ARAKI 05 obtained this result by a two-neutrino oscillation analysis using KamLAND and solar neutrino data. *CPT* invariance is assumed. The  $1\sigma$  error shown here is translated from the number provided by the KamLAND collaboration,  $\tan^2 \theta = 0.40_{-0.05}^{+0.07}$ . The corresponding number quoted in ARAKI 05 is  $\tan^2 \theta = 0.40_{-0.07}^{+0.10}$  ( $\sin^2 2\theta = 0.82 \pm 0.07$ ), which envelops the 68% CL two-dimensional region.
- 23 The result given by AHMED 04A is  $\theta = (32.5_{-1.6}^{+1.7})^\circ$ . This result is obtained by a two-neutrino oscillation analysis using solar neutrino and KamLAND data (EGUCHI 03). *CPT* invariance is assumed. AHMED 04A also quotes  $\theta = (32.5_{-2.3}^{+2.4})^\circ$  as the error enveloping the 68% CL two-dimensional region. This translates into  $\sin^2 2\theta = 0.82 \pm 0.06$ .
- 24 AHMED 04A obtained this result by a two-neutrino oscillation analysis using the data from all solar neutrino experiments. The listed range of the parameter envelops the 95% CL two-dimensional region shown in Fig. 5(a) of AHMED 04A. The best-fit point is  $\Delta(m^2) = 6.5 \times 10^{-5} \text{ eV}^2$ ,  $\tan^2 \theta = 0.40$  ( $\sin^2 2\theta = 0.82$ ).
- 25 The result given by SMY 04 is  $\tan^2 \theta = 0.44 \pm 0.08$ . This result is obtained by a two-neutrino oscillation analysis using solar neutrino and KamLAND data (IANNI 03). *CPT* invariance is assumed.

- <sup>26</sup> SMY 04 obtained this result by a two-neutrino oscillation analysis using the data from all solar neutrino experiments. The  $1\sigma$  errors are read from Fig. 6(a) of SMY 04.
- <sup>27</sup> SMY 04 obtained this result by a two-neutrino oscillation analysis using the Super-Kamiokande and SNO (AHMAD 02 and AHMAD 02B) solar neutrino data. The  $1\sigma$  errors are read from Fig. 6(a) of SMY 04.
- <sup>28</sup> AHMAD 02B obtained this result by a two-neutrino oscillation analysis using the data from all solar neutrino experiments. The listed range of the parameter envelops the 95% CL two-dimensional region shown in Fig. 4(b) of AHMAD 02B. The best fit point is  $\Delta(m^2) = 5.0 \times 10^{-5} \text{ eV}^2$  and  $\tan\theta = 0.34$  ( $\sin^2 2\theta = 0.76$ ).
- <sup>29</sup> FUKUDA 02 obtained this result by a two-neutrino oscillation analysis using the data from all solar neutrino experiments. The listed range of the parameter envelops the 95% CL two-dimensional region shown in Fig. 4 of FUKUDA 02. The best fit point is  $\Delta(m^2) = 6.9 \times 10^{-5} \text{ eV}^2$  and  $\tan^2\theta = 0.38$  ( $\sin^2 2\theta = 0.80$ ).

## $\Delta m_{21}^2$

VALUE ( $10^{-5} \text{ eV}^2$ )	DOCUMENT ID	TECN	COMMENT
<b>7.50<sup>+0.19</sup><sub>-0.20</sub></b>	1 GANDO	11 FIT	KamLAND + solar: $3\nu$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
7.6 $\pm$ 0.2	2 ABE	11 FIT	KamLAND + global solar: $2\nu$
6.2 <sup>+1.1</sup> <sub>-1.9</sub>	3 ABE	11 FIT	global solar: $2\nu$
7.7 $\pm$ 0.3	4 ABE	11 FIT	KamLAND + global solar: $3\nu$
6.0 <sup>+2.2</sup> <sub>-2.5</sub>	5 ABE	11 FIT	global solar: $3\nu$
7.50 <sup>+0.16</sup> <sub>-0.24</sub>	6 BELLINI	11A FIT	KamLAND + global solar: $2\nu$
5.2 <sup>+1.5</sup> <sub>-0.9</sub>	7 BELLINI	11A FIT	global solar: $2\nu$
7.49 $\pm$ 0.20	8 GANDO	11 FIT	KamLAND: $3\nu$
7.59 <sup>+0.20</sup> <sub>-0.21</sub>	9,10 AHARMIM	10 FIT	KamLAND + global solar: $2\nu$
5.89 <sup>+2.13</sup> <sub>-2.16</sub>	9,11 AHARMIM	10 FIT	global solar: $2\nu$
7.59 $\pm$ 0.21	9,12 AHARMIM	10 FIT	KamLAND + global solar: $3\nu$
6.31 <sup>+2.49</sup> <sub>-2.58</sub>	9,13 AHARMIM	10 FIT	global solar: $3\nu$
7.58 <sup>+0.14</sup> <sub>-0.13</sub> $\pm$ 0.15	14 ABE	08A FIT	KamLAND
7.59 $\pm$ 0.21	15 ABE	08A FIT	KamLAND + global solar
7.59 <sup>+0.19</sup> <sub>-0.21</sub>	16 AHARMIM	08 FIT	KamLAND + global solar
8.0 $\pm$ 0.3	17 HOSAKA	06 FIT	KamLAND + global solar
8.0 $\pm$ 0.3	18 HOSAKA	06 FIT	SKAM+SNO+KamLAND
6.3 <sup>+3.7</sup> <sub>-1.5</sub>	19 HOSAKA	06 FIT	SKAM+SNO
5–12	20 HOSAKA	06 FIT	SKAM day/night in the LMA region
8.0 <sup>+0.4</sup> <sub>-0.3</sub>	21 AHARMIM	05A FIT	KamLAND + global solar LMA
3.3–14.4	22 AHARMIM	05A FIT	global solar
7.9 <sup>+0.4</sup> <sub>-0.3</sub>	23 ARAKI	05 FIT	KamLAND + global solar
7.1 <sup>+1.0</sup> <sub>-0.3</sub>	24 AHMED	04A FIT	KamLAND + global solar

3.2–13.7	25 AHMED	04A	FIT	global solar
7.1 $+0.6$ $-0.5$	26 SMY	04	FIT	KamLAND + global solar
6.0 $+1.7$ $-1.6$	27 SMY	04	FIT	global solar
6.0 $+2.5$ $-1.6$	28 SMY	04	FIT	SKAM + SNO
2.8–12.0	29 AHMAD	02B	FIT	global solar
3.2–19.1	30 FUKUDA	02	FIT	global solar

<sup>1</sup> GANDO 11 obtain this result with three-neutrino fit using the KamLAND + solar data. Supersedes ABE 08A.

<sup>2</sup> ABE 11 obtained this result by a two-neutrino oscillation analysis using solar neutrino data including Super-Kamiokande, SNO, Borexino (ARPESELLA 08A), Homestake, GALLEX/GNO, SAGE, and KamLAND data. CPT invariance is assumed.

<sup>3</sup> ABE 11 obtained this result by a two-neutrino oscillation analysis using solar neutrino data including Super-Kamiokande, SNO, Borexino (ARPESELLA 08A), Homestake, GALLEX/GNO, and SAGE data.

<sup>4</sup> ABE 11 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{32}^2$  fixed to  $2.4 \times 10^{-3} \text{ eV}^2$ , using solar neutrino data including Super-Kamiokande, SNO, Borexino (ARPESELLA 08A), Homestake, GALLEX/GNO, SAGE, and KamLAND data. The normal neutrino mass hierarchy and CPT invariance are assumed.

<sup>5</sup> ABE 11 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{32}^2$  fixed to  $2.4 \times 10^{-3} \text{ eV}^2$ , using solar neutrino data including Super-Kamiokande, SNO, Borexino (ARPESELLA 08A), Homestake, and GALLEX/GNO data. The normal neutrino mass hierarchy is assumed.

<sup>6</sup> BELLINI 11A obtained this result by a two-neutrino oscillation analysis using KamLAND, Homestake, SAGE, Gallex, GNO, Kamiokande, Super-Kamiokande, SNO, and Borexino (BELLINI 11A) data and the SSM flux prediction in SERENELLI 11 (Astrophysical Journal **743** 24 (2011)) with the exception that the  $^8\text{B}$  flux was left free. CPT invariance is assumed.

<sup>7</sup> BELLINI 11A obtained this result by a two-neutrino oscillation analysis using Homestake, SAGE, Gallex, GNO, Kamiokande, Super-Kamiokande, SNO, and Borexino (BELLINI 11A) data and the SSM flux prediction in SERENELLI 11 (Astrophysical Journal **743** 24 (2011)) with the exception that the  $^8\text{B}$  flux was left free.

<sup>8</sup> GANDO 11 obtain this result with three-neutrino fit using the KamLAND data only. Supersedes ABE 08A.

<sup>9</sup> AHARMIM 10 global solar neutrino data include SNO's low-energy-threshold analysis survival probability day/night curves, SNO Phase III integral rates (AHARMIM 08), CI (CLEVELAND 98), SAGE (ABDURASHITOV 09), Gallex/GNO (HAMPEL 99, ALTMANN 05), Borexino (ARPESELLA 08A), SK-I zenith (HOSAKA 06), and SK-II day/night spectra (CRAVENS 08).

<sup>10</sup> AHARMIM 10 obtained this result by a two-neutrino oscillation analysis using global solar neutrino data and KamLAND data (ABE 08A). CPT invariance is assumed.

<sup>11</sup> AHARMIM 10 obtained this result by a two-neutrino oscillation analysis using global solar neutrino data.

<sup>12</sup> AHARMIM 10 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{31}^2$  fixed to  $2.3 \times 10^{-3} \text{ eV}^2$ , using global solar neutrino data and KamLAND data (ABE 08A). CPT invariance is assumed.

<sup>13</sup> AHARMIM 10 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{31}^2$  fixed to  $2.3 \times 10^{-3} \text{ eV}^2$ , using global solar neutrino data.

<sup>14</sup> ABE 08A obtained this result by a rate + shape + time combined geoneutrino and reactor two-neutrino fit for  $\Delta m_{21}^2$  and  $\tan^2\theta_{12}$ , using KamLAND data only.

- <sup>15</sup> ABE 08A obtained this result by means of a two-neutrino fit using KamLAND, Homestake, SAGE, GALLEX, GNO, SK (zenith angle and E-spectrum), the SNO  $\chi^2$ -map, and solar flux data. *CPT* invariance is assumed.
- <sup>16</sup> AHARMIM 08 obtained this result by a two-neutrino oscillation analysis using all solar neutrino data including those of Borexino (ARPESELLA 08A) and Super-Kamiokande-I (HOSAKA 06), and KamLAND data (ABE 08A). *CPT* invariance is assumed.
- <sup>17</sup> HOSAKA 06 obtained this result by a two-neutrino oscillation analysis using solar neutrino and KamLAND data (ARAKI 05). *CPT* invariance is assumed.
- <sup>18</sup> HOSAKA 06 obtained this result by a two-neutrino oscillation analysis using the data from Super-Kamiokande, SNO (AHMAD 02 and AHMAD 02B), and KamLAND (ARAKI 05) experiments. *CPT* invariance is assumed.
- <sup>19</sup> HOSAKA 06 obtained this result by a two-neutrino oscillation analysis using the Super-Kamiokande and SNO (AHMAD 02 and AHMAD 02B) solar neutrino data.
- <sup>20</sup> HOSAKA 06 obtained this result from the consistency between the observed and expected day-night flux asymmetry amplitude. The listed 68% CL range is derived from the  $1\sigma$  boundary of the amplitude fit to the data. Oscillation parameters are constrained to be in the LMA region. The mixing angle is fixed at  $\tan^2\theta = 0.44$  because the fit depends only very weakly on it.
- <sup>21</sup> AHARMIM 05A obtained this result by a two-neutrino oscillation analysis using solar neutrino and KamLAND data (ARAKI 05). *CPT* invariance is assumed. AHARMIM 05A also quotes  $\Delta(m^2) = (8.0^{+0.6}_{-0.4}) \times 10^{-5} \text{ eV}^2$  as the error enveloping the 68% CL two-dimensional region.
- <sup>22</sup> AHARMIM 05A obtained this result by a two-neutrino oscillation analysis using the data from all solar neutrino experiments. The listed range of the parameter envelops the 95% CL two-dimensional region shown in figure 35a of AHARMIM 05A. AHARMIM 05A also quotes  $\Delta(m^2) = (6.5^{+4.4}_{-2.3}) \times 10^{-5} \text{ eV}^2$  as the error enveloping the 68% CL two-dimensional region.
- <sup>23</sup> ARAKI 05 obtained this result by a two-neutrino oscillation analysis using KamLAND and solar neutrino data. *CPT* invariance is assumed. The  $1\sigma$  error shown here is provided by the KamLAND collaboration. The error quoted in ARAKI 05,  $\Delta(m^2) = (7.9^{+0.6}_{-0.5}) \times 10^{-5}$ , envelops the 68% CL two-dimensional region.
- <sup>24</sup> AHMED 04A obtained this result by a two-neutrino oscillation analysis using solar neutrino and KamLAND data (EGUCHI 03). *CPT* invariance is assumed. AHMED 04A also quotes  $\Delta(m^2) = (7.1^{+1.2}_{-0.6}) \times 10^{-5} \text{ eV}^2$  as the error enveloping the 68% CL two-dimensional region.
- <sup>25</sup> AHMED 04A obtained this result by a two-neutrino oscillation analysis using the data from all solar neutrino experiments. The listed range of the parameter envelops the 95% CL two-dimensional region shown in Fig. 5(a) of AHMED 04A. The best-fit point is  $\Delta(m^2) = 6.5 \times 10^{-5} \text{ eV}^2$ ,  $\tan^2\theta = 0.40$  ( $\sin^2 2\theta = 0.82$ ).
- <sup>26</sup> SMY 04 obtained this result by a two-neutrino oscillation analysis using solar neutrino and KamLAND data (IANNI 03). *CPT* invariance is assumed.
- <sup>27</sup> SMY 04 obtained this result by a two-neutrino oscillation analysis using the data from all solar neutrino experiments. The  $1\sigma$  errors are read from Fig. 6(a) of SMY 04.
- <sup>28</sup> SMY 04 obtained this result by a two-neutrino oscillation analysis using the Super-Kamiokande and SNO (AHMAD 02 and AHMAD 02B) solar neutrino data. The  $1\sigma$  errors are read from Fig. 6(a) of SMY 04.
- <sup>29</sup> AHMAD 02B obtained this result by a two-neutrino oscillation analysis using the data from all solar neutrino experiments. The listed range of the parameter envelops the 95% CL two-dimensional region shown in Fig. 4(b) of AHMAD 02B. The best fit point is  $\Delta(m^2) = 5.0 \times 10^{-5} \text{ eV}^2$  and  $\tan\theta = 0.34$  ( $\sin^2 2\theta = 0.76$ ).
- <sup>30</sup> FUKUDA 02 obtained this result by a two-neutrino oscillation analysis using the data from all solar neutrino experiments. The listed range of the parameter envelops the 95% CL two-dimensional region shown in Fig. 4 of FUKUDA 02. The best fit point is  $\Delta(m^2) = 6.9 \times 10^{-5} \text{ eV}^2$  and  $\tan^2\theta = 0.38$  ( $\sin^2 2\theta = 0.80$ ).

**$\sin^2(2\theta_{23})$** 

The ranges below correspond to the projection onto the  $\sin^2(2\theta_{23})$  axis of the 90% CL contours in the  $\sin^2(2\theta_{23}) - \Delta m_{32}^2$  plane presented by the authors. The values are reported with one standard deviation uncertainty.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>&gt;0.95</b>	<sup>1</sup> ABE	11C	SKAM Super-Kamiokande
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
>0.90	ADAMSON	11	MINS $2\nu$ oscillation; maximal mixing
0.86 $^{+0.11}_{-0.12}$	<sup>2</sup> ADAMSON	11B	MINS $\bar{\nu}$ beam
>0.965	<sup>3</sup> WENDELL	10	SKAM $3\nu$ oscillation with solar terms; $\theta_{13}=0$
>0.95	<sup>4</sup> WENDELL	10	SKAM $3\nu$ oscillation; normal mass hierarchy
>0.93	<sup>5</sup> WENDELL	10	SKAM $3\nu$ oscillation; inverted mass hierarchy
>0.85	ADAMSON	08A	MINS MINOS
>0.2	<sup>6</sup> ADAMSON	06	MINS atmospheric $\nu$ with far detector
>0.59	<sup>7</sup> AHN	06A	K2K KEK to Super-K
>0.7	<sup>8</sup> MICHAEL	06	MINS MINOS
>0.58	<sup>9</sup> ALIU	05	K2K KEK to Super-K
>0.6	<sup>10</sup> ALLISON	05	SOU2
>0.92	<sup>11</sup> ASHIE	05	SKAM Super-Kamiokande
>0.80	<sup>12</sup> AMBROSIO	04	MCRO MACRO
>0.90	<sup>13</sup> ASHIE	04	SKAM L/E distribution
>0.30	<sup>14</sup> AHN	03	K2K KEK to Super-K
>0.45	<sup>15</sup> AMBROSIO	03	MCRO MACRO
>0.77	<sup>16</sup> AMBROSIO	03	MCRO MACRO
>0.50	<sup>17</sup> SANCHEZ	03	SOU2 Soudan-2 Atmospheric
>0.80	<sup>18</sup> AMBROSIO	01	MCRO upward $\mu$
>0.82	<sup>19</sup> AMBROSIO	01	MCRO upward $\mu$
>0.45	<sup>20</sup> FUKUDA	99C	SKAM upward $\mu$
>0.70	<sup>21</sup> FUKUDA	99D	SKAM upward $\mu$
>0.30	<sup>22</sup> FUKUDA	99D	SKAM stop $\mu$ / through
>0.82	<sup>23</sup> FUKUDA	98C	SKAM Super-Kamiokande
>0.30	<sup>24</sup> HATAKEYAMA98	KAMI	Kamiokande
>0.73	<sup>25</sup> HATAKEYAMA98	KAMI	Kamiokande
>0.65	<sup>26</sup> FUKUDA	94	KAMI Kamiokande

<sup>1</sup> ABE 11C obtained this result by a two-neutrino oscillation analysis using the Super-Kamiokande-I+II+III atmospheric neutrino data. ABE 11C also reported results under a two-neutrino disappearance model with separate mixing parameters between  $\nu$  and  $\bar{\nu}$ , and obtained  $\sin^2 2\theta > 0.93$  for  $\nu$  and  $\sin^2 2\theta > 0.83$  for  $\bar{\nu}$  at 90% C.L.

<sup>2</sup> ADAMSON 11B obtained this result by a two-neutrino oscillation analysis of antineutrinos in an antineutrino enhanced beam with  $1.71 \times 10^{20}$  protons on target. This result is consistent with the neutrino measurements of ADAMSON 11 at 2% C.L.

<sup>3</sup> WENDELL 10 obtained this result ( $\sin^2 \theta_{23} = 0.407\text{--}0.583$ ) by a three-neutrino oscillation analysis using the Super-Kamiokande-I+II+III atmospheric neutrino data, assuming  $\theta_{13} = 0$  but including the solar oscillation parameters  $\Delta m_{21}^2$  and  $\sin^2 \theta_{12}$  in the fit.

<sup>4</sup> WENDELL 10 obtained this result ( $\sin^2 \theta_{23} = 0.43\text{--}0.61$ ) by a three-neutrino oscillation analysis with one mass scale dominance ( $\Delta m_{21}^2 = 0$ ) using the Super-Kamiokande-I+II+III atmospheric neutrino data, and updates the HOSAKA 06A result.

- 5 WENDELL 10 obtained this result ( $\sin^2\theta_{23} = 0.44\text{--}0.63$ ) by a three-neutrino oscillation analysis with one mass scale dominance ( $\Delta m_{21}^2 = 0$ ) using the Super-Kamiokande-I+II+III atmospheric neutrino data, and updates the HOSAKA 06A result.
- 6 ADAMSON 06 obtained this result by a two-neutrino oscillation analysis of the L/E distribution using 4.54 kton yr atmospheric neutrino data with the MINOS far detector.
- 7 Supercedes ALIU 05.
- 8 MICHAEL 06 best fit is for maximal mixing. See also ADAMSON 08.
- 9 The best fit is for maximal mixing.
- 10 ALLISON 05 result is based upon atmospheric neutrino interactions including upward-stopping muons, with an exposure of 5.9 kton yr. From a two-flavor oscillation analysis the best-fit point is  $\Delta m^2 = 0.0017 \text{ eV}^2$  and  $\sin^2(2\theta) = 0.97$ .
- 11 ASHIE 05 obtained this result by a two-neutrino oscillation analysis using 92 kton yr atmospheric neutrino data from the complete Super-Kamiokande I running period.
- 12 AMBROSIO 04 obtained this result, without using the absolute normalization of the neutrino flux, by combining the angular distribution of upward through-going muon tracks with  $E_\mu > 1 \text{ GeV}$ ,  $N_{low}$  and  $N_{high}$ , and the numbers of InDown + UpStop and InUp events. Here,  $N_{low}$  and  $N_{high}$  are the number of events with reconstructed neutrino energies  $< 30 \text{ GeV}$  and  $> 130 \text{ GeV}$ , respectively. InDown and InUp represent events with downward and upward-going tracks starting inside the detector due to neutrino interactions, while UpStop represents entering upward-going tracks which stop in the detector. The best fit is for maximal mixing.
- 13 ASHIE 04 obtained this result from the L(flight length)/E(estimated neutrino energy) distribution of  $\nu_\mu$  disappearance probability, using the Super-Kamiokande-I 1489 live-day atmospheric neutrino data.
- 14 There are several islands of allowed region from this K2K analysis, extending to high values of  $\Delta m^2$ . We only include the one that overlaps atmospheric neutrino analyses. The best fit is for maximal mixing.
- 15 AMBROSIO 03 obtained this result on the basis of the ratio  $R = N_{low}/N_{high}$ , where  $N_{low}$  and  $N_{high}$  are the number of upward through-going muon events with reconstructed neutrino energy  $< 30 \text{ GeV}$  and  $> 130 \text{ GeV}$ , respectively. The data came from the full detector run started in 1994. The method of FELDMAN 98 is used to obtain the limits.
- 16 AMBROSIO 03 obtained this result by using the ratio  $R$  and the angular distribution of the upward through-going muons.  $R$  is given in the previous note and the angular distribution is reported in AMBROSIO 01. The method of FELDMAN 98 is used to obtain the limits. The best fit is to maximal mixing.
- 17 SANCHEZ 03 is based on an exposure of 5.9 kton yr. The result is obtained using a likelihood analysis of the neutrino L/E distribution for a selection  $\mu$  flavor sample while the  $e$ -flavor sample provides flux normalization. The method of FELDMAN 98 is used to obtain the allowed region. The best fit is  $\sin^2(2\theta) = 0.97$ .
- 18 AMBROSIO 01 result is based on the angular distribution of upward through-going muon tracks with  $E_\mu > 1 \text{ GeV}$ . The data came from three different detector configurations, but the statistics is largely dominated by the full detector run, from May 1994 to December 2000. The total live time, normalized to the full detector configuration is 6.17 years. The best fit is obtained outside the physical region. The method of FELDMAN 98 is used to obtain the limits. The best fit is for maximal mixing.
- 19 AMBROSIO 01 result is based on the angular distribution and normalization of upward through-going muon tracks with  $E_\mu > 1 \text{ GeV}$ . See the previous footnote.
- 20 FUKUDA 99C obtained this result from a total of 537 live days of upward through-going muon data in Super-Kamiokande between April 1996 to January 1998. With a threshold of  $E_\mu > 1.6 \text{ GeV}$ , the observed flux is  $(1.74 \pm 0.07 \pm 0.02) \times 10^{-13} \text{ cm}^{-2}\text{s}^{-1}\text{sr}^{-1}$ . The best fit is  $\sin^2(2\theta) = 0.95$ .
- 21 FUKUDA 99D obtained this result from a simultaneous fitting to zenith angle distributions of upward-stopping and through-going muons. The flux of upward-stopping muons of

- minimum energy of 1.6 GeV measured between April 1996 and January 1998 is  $(0.39 \pm 0.04 \pm 0.02) \times 10^{-13} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ . This is compared to the expected flux of  $(0.73 \pm 0.16 \text{ (theoretical error)}) \times 10^{-13} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ . The best fit is to maximal mixing.
- <sup>22</sup> FUKUDA 99D obtained this result from the zenith dependence of the upward-stopping/through-going flux ratio. The best fit is to maximal mixing.
- <sup>23</sup> FUKUDA 98C obtained this result by an analysis of 33.0 kton yr atmospheric neutrino data. The best fit is for maximal mixing.
- <sup>24</sup> HATAKEYAMA 98 obtained this result from a total of 2456 live days of upward-going muon data in Kamiokande between December 1985 and May 1995. With a threshold of  $E_\mu > 1.6 \text{ GeV}$ , the observed flux of upward through-going muons is  $(1.94 \pm 0.10^{+0.07}_{-0.06}) \times 10^{-13} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ . This is compared to the expected flux of  $(2.46 \pm 0.54 \text{ (theoretical error)}) \times 10^{-13} \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ . The best fit is for maximal mixing.
- <sup>25</sup> HATAKEYAMA 98 obtained this result from a combined analysis of Kamiokande contained events (FUKUDA 94) and upward going muon events. The best fit is  $\sin^2(2\theta) = 0.95$ .
- <sup>26</sup> FUKUDA 94 obtained the result by a combined analysis of sub- and multi-GeV atmospheric neutrino events in Kamiokande. The best fit is for maximal mixing.

## $\Delta m_{32}^2$

The sign of  $\Delta m_{32}^2$  is not known at this time. Only the absolute value is quoted below. Unless otherwise specified, the ranges below correspond to the projection onto the  $\Delta m_{32}^2$  axis of the 90% CL contours in the  $\sin^2(2\theta_{23}) - \Delta m_{32}^2$  plane presented by the authors. The values are reported with one standard deviation uncertainty.

VALUE ( $10^{-3} \text{ eV}^2$ )	DOCUMENT ID	TECN	COMMENT
<b><math>2.32^{+0.12}_{-0.08}</math></b>	ADAMSON	11	MINS $2\nu$ oscillation; maximal mixing
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.3 - 4.0	1 ABE	11C	SKAM atmospheric $\bar{\nu}$
$3.36^{+0.46}_{-0.40}$	2 ADAMSON	11B	MINS $\bar{\nu}$ beam
<3.37	3 ADAMSON	11C	MINS MINOS
1.9 - 2.6	4 WENDELL	10	SKAM $3\nu$ osc.; normal mass hierarchy
1.7 - 2.7	4 WENDELL	10	SKAM $3\nu$ osc.; inverted mass hierarchy
$2.43 \pm 0.13$	ADAMSON	08A	MINS MINOS
0.07-50	5 ADAMSON	06	MINS atmospheric $\nu$ with far detector
1.9-4.0	6,7 AHN	06A	K2K KEK to Super-K
2.2-3.8	8 MICHAEL	06	MINS MINOS
1.9-3.6	6 ALIU	05	K2K KEK to Super-K
0.3-12	9 ALLISON	05	SOU2
1.5-3.4	10 ASHIE	05	SKAM atmospheric neutrino
0.6-8.0	11 AMBROSIO	04	MCRO MACRO
1.9 to 3.0	12 ASHIE	04	SKAM L/E distribution
1.5-3.9	13 AHN	03	K2K KEK to Super-K
0.25-9.0	14 AMBROSIO	03	MCRO MACRO
0.6-7.0	15 AMBROSIO	03	MCRO MACRO
0.15-15	16 SANCHEZ	03	SOU2 Soudan-2 Atmospheric
0.6-15	17 AMBROSIO	01	MCRO upward $\mu$
1.0-6.0	18 AMBROSIO	01	MCRO upward $\mu$
1.0-50	19 FUKUDA	99C	SKAM upward $\mu$

1.5–15.0	20	FUKUDA	99D	SKAM	upward $\mu$
0.7–18	21	FUKUDA	99D	SKAM	stop $\mu$ / through
0.5–6.0	22	FUKUDA	98C	SKAM	Super-Kamiokande
0.55–50	23	HATAKEYAMA98	KAMI	Kamiokande	
4–23	24	HATAKEYAMA98	KAMI	Kamiokande	
5–25	25	FUKUDA	94	KAMI	Kamiokande

- <sup>1</sup> ABE 11C obtained this result by a two-neutrino oscillation analysis with separate mixing parameters between neutrinos and antineutrinos, using the Super-Kamiokande-I+II+III atmospheric neutrino data. The corresponding 90% CL neutrino oscillation parameter range obtained from this analysis is  $\Delta m^2 = 1.7\text{--}3.0 \times 10^{-3} \text{ eV}^2$ .
- <sup>2</sup> ADAMSON 11B obtained this result by a two-neutrino oscillation analysis of antineutrinos in an antineutrino enhanced beam with  $1.71 \times 10^{20}$  protons on target. This result is consistent with the neutrino measurements of ADAMSON 11 at 2% C.L.
- <sup>3</sup> ADAMSON 11C obtains this result based on a study of antineutrinos in a neutrino beam and assumes maximal mixing in the two-flavor approximation.
- <sup>4</sup> WENDELL 10 obtained this result by a three-neutrino oscillation analysis with one mass scale dominance ( $\Delta m_{21}^2 = 0$ ) using the Super-Kamiokande-I+II+III atmospheric neutrino data, and updates the HOSAKA 06A result.
- <sup>5</sup> ADAMSON 06 obtained this result by a two-neutrino oscillation analysis of the L/E distribution using 4.54 kton yr atmospheric neutrino data with the MINOS far detector.
- <sup>6</sup> The best fit in the physical region is for  $\Delta m^2 = 2.8 \times 10^{-3} \text{ eV}^2$ .
- <sup>7</sup> Supercedes ALIU 05.
- <sup>8</sup> MICHAEL 06 best fit is  $2.74 \times 10^{-3} \text{ eV}^2$ . See also ADAMSON 08.
- <sup>9</sup> ALLISON 05 result is based on an atmospheric neutrino observation with an exposure of 5.9 kton yr. From a two-flavor oscillation analysis the best-fit point is  $\Delta m^2 = 0.0017 \text{ eV}^2$  and  $\sin^2 2\theta = 0.97$ .
- <sup>10</sup> ASHIE 05 obtained this result by a two-neutrino oscillation analysis using 92 kton yr atmospheric neutrino data from the complete Super-Kamiokande I running period. The best fit is for  $\Delta m^2 = 2.1 \times 10^{-3} \text{ eV}^2$ .
- <sup>11</sup> AMBROSIO 04 obtained this result, without using the absolute normalization of the neutrino flux, by combining the angular distribution of upward through-going muon tracks with  $E_\mu > 1 \text{ GeV}$ ,  $N_{low}$  and  $N_{high}$ , and the numbers of InDown + UpStop and InUp events. Here,  $N_{low}$  and  $N_{high}$  are the number of events with reconstructed neutrino energies  $< 30 \text{ GeV}$  and  $> 130 \text{ GeV}$ , respectively. InDown and InUp represent events with downward and upward-going tracks starting inside the detector due to neutrino interactions, while UpStop represents entering upward-going tracks which stop in the detector. The best fit is for  $\Delta m^2 = 2.3 \times 10^{-3} \text{ eV}^2$ .
- <sup>12</sup> ASHIE 04 obtained this result from the L(flight length)/E(estimated neutrino energy) distribution of  $\nu_\mu$  disappearance probability, using the Super-Kamiokande-I 1489 live-day atmospheric neutrino data. The best fit is for  $\Delta m^2 = 2.4 \times 10^{-3} \text{ eV}^2$ .
- <sup>13</sup> There are several islands of allowed region from this K2K analysis, extending to high values of  $\Delta m^2$ . We only include the one that overlaps atmospheric neutrino analyses. The best fit is for  $\Delta m^2 = 2.8 \times 10^{-3} \text{ eV}^2$ .
- <sup>14</sup> AMBROSIO 03 obtained this result on the basis of the ratio  $R = N_{low}/N_{high}$ , where  $N_{low}$  and  $N_{high}$  are the number of upward through-going muon events with reconstructed neutrino energy  $< 30 \text{ GeV}$  and  $> 130 \text{ GeV}$ , respectively. The data came from the full detector run started in 1994. The method of FELDMAN 98 is used to obtain the limits. The best fit is for  $\Delta m^2 = 2.5 \times 10^{-3} \text{ eV}^2$ .
- <sup>15</sup> AMBROSIO 03 obtained this result by using the ratio R and the angular distribution of the upward through-going muons. R is given in the previous note and the angular distribution is reported in AMBROSIO 01. The method of FELDMAN 98 is used to obtain the limits. The best fit is for  $\Delta m^2 = 2.5 \times 10^{-3} \text{ eV}^2$ .
- <sup>16</sup> SANCHEZ 03 is based on an exposure of 5.9 kton yr. The result is obtained using a likelihood analysis of the neutrino L/E distribution for a selection  $\mu$  flavor sample while

the e-flavor sample provides flux normalization. The method of FELDMAN 98 is used to obtain the allowed region. The best fit is for  $\Delta m^2 = 5.2 \times 10^{-3} \text{ eV}^2$ .

- 17 AMBROSIO 01 result is based on the angular distribution of upward through-going muon tracks with  $E_\mu > 1 \text{ GeV}$ . The data came from three different detector configurations, but the statistics is largely dominated by the full detector run, from May 1994 to December 2000. The total live time, normalized to the full detector configuration is 6.17 years. The best fit is obtained outside the physical region. The method of FELDMAN 98 is used to obtain the limits.
- 18 AMBROSIO 01 result is based on the angular distribution and normalization of upward through-going muon tracks with  $E_\mu > 1 \text{ GeV}$ . See the previous footnote.
- 19 FUKUDA 99C obtained this result from a total of 537 live days of upward through-going muon data in Super-Kamiokande between April 1996 to January 1998. With a threshold of  $E_\mu > 1.6 \text{ GeV}$ , the observed flux is  $(1.74 \pm 0.07 \pm 0.02) \times 10^{-13} \text{ cm}^{-2}\text{s}^{-1}\text{sr}^{-1}$ . The best fit is for  $\Delta m^2 = 5.9 \times 10^{-3} \text{ eV}^2$ .
- 20 FUKUDA 99D obtained this result from a simultaneous fitting to zenith angle distributions of upward-stopping and through-going muons. The flux of upward-stopping muons of minimum energy of 1.6 GeV measured between April 1996 and January 1998 is  $(0.39 \pm 0.04 \pm 0.02) \times 10^{-13} \text{ cm}^{-2}\text{s}^{-1}\text{sr}^{-1}$ . This is compared to the expected flux of  $(0.73 \pm 0.16 \text{ (theoretical error)}) \times 10^{-13} \text{ cm}^{-2}\text{s}^{-1}\text{sr}^{-1}$ . The best fit is for  $\Delta m^2 = 3.9 \times 10^{-3} \text{ eV}^2$ .
- 21 FUKUDA 99D obtained this result from the zenith dependence of the upward-stopping/through-going flux ratio. The best fit is for  $\Delta m^2 = 3.1 \times 10^{-3} \text{ eV}^2$ .
- 22 FUKUDA 98C obtained this result by an analysis of 33.0 kton yr atmospheric neutrino data. The best fit is for  $\Delta m^2 = 2.2 \times 10^{-3} \text{ eV}^2$ .
- 23 HATAKEYAMA 98 obtained this result from a total of 2456 live days of upward-going muon data in Kamiokande between December 1985 and May 1995. With a threshold of  $E_\mu > 1.6 \text{ GeV}$ , the observed flux of upward through-going muons is  $(1.94 \pm 0.10^{+0.07}_{-0.06}) \times 10^{-13} \text{ cm}^{-2}\text{s}^{-1}\text{sr}^{-1}$ . This is compared to the expected flux of  $(2.46 \pm 0.54 \text{ (theoretical error)}) \times 10^{-13} \text{ cm}^{-2}\text{s}^{-1}\text{sr}^{-1}$ . The best fit is for  $\Delta m^2 = 2.2 \times 10^{-3} \text{ eV}^2$ .
- 24 HATAKEYAMA 98 obtained this result from a combined analysis of Kamiokande contained events (FUKUDA 94) and upward going muon events. The best fit is for  $\Delta m^2 = 13 \times 10^{-3} \text{ eV}^2$ .
- 25 FUKUDA 94 obtained the result by a combined analysis of sub- and multi-GeV atmospheric neutrino events in Kamiokande. The best fit is for  $\Delta m^2 = 16 \times 10^{-3} \text{ eV}^2$ .

### $\sin^2(2\theta_{13})$

At present time direct measurements of  $\sin^2(2\theta_{13})$  are derived from the reactor  $\bar{\nu}_e$  disappearance at distances corresponding to the  $\Delta m_{32}^2$  value, i.e.  $L \sim 1\text{km}$ . Alternatively, limits can also be obtained from the analysis of the solar neutrino data and accelerator-based  $\nu_\mu \rightarrow \nu_e$  experiments.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>0.098 ± 0.013 OUR AVERAGE</b>				
0.086 ± 0.041 ± 0.030	1	ABE	12	DCHZ Chooz reactors
0.113 ± 0.013 ± 0.019	2	AHN	12	RENO Yonggwang reactors
0.092 ± 0.016 ± 0.005	3	AN	12	DAYA Daya Bay, Ling Ao, Ling Ao-II reactors

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.098^{+0.067}_{-0.062}$	68	4 ABE	11	FIT	KamLAND + global solar
$< 0.23$	95	5 ABE	11	FIT	Global solar
$0.05 - 0.21$	68	6 ABE	11A	T2K	Normal mass hierarchy
$0.06 - 0.25$	68	7 ABE	11A	T2K	Inverted mass hierarchy
$0.01 - 0.09$	68	8 ADAMSON	11D	MINS	Normal mass hierarchy
$0.03 - 0.15$	68	9 ADAMSON	11D	MINS	Inverted mass hierarchy
$0.08 \pm 0.03$	68	10 FOGLI	11	FIT	Global neutrino data
$0.078 \pm 0.062$	68	11 GANDO	11	FIT	KamLAND + solar: $3\nu$
$0.124 \pm 0.133$	68	12 GANDO	11	FIT	KamLAND: $3\nu$
$0.03^{+0.09}_{-0.07}$	90	13 ADAMSON	10A	MINS	Normal mass hierarchy
$0.06^{+0.14}_{-0.06}$	90	14 ADAMSON	10A	MINS	Inverted mass hierarchy
$0.08^{+0.08}_{-0.07}$		15,16 AHARMIM	10	FIT	KamLAND + global solar: $3\nu$
$< 0.30$	95	15,17 AHARMIM	10	FIT	global solar: $3\nu$
$< 0.15$	90	18 WENDELL	10	SKAM	$3\nu$ osc.; normal $m$ hierarchy
$< 0.33$	90	18 WENDELL	10	SKAM	$3\nu$ osc.; inverted $m$ hierarchy
$0.11^{+0.11}_{-0.08}$		19 ADAMSON	09	MINS	Normal mass hierarchy
$0.18^{+0.15}_{-0.11}$		20 ADAMSON	09	MINS	Inverted mass hierarchy
$0.06 \pm 0.04$		21 FOGLI	08	FIT	Global neutrino data
$0.08 \pm 0.07$		22 FOGLI	08	FIT	Solar + KamLAND data
$0.05 \pm 0.05$		23 FOGLI	08	FIT	Atmospheric+LBL+CHOOZ
$< 0.36$	90	24 YAMAMOTO	06	K2K	Accelerator experiment
$< 0.48$	90	25 AHN	04	K2K	Accelerator experiment
$< 0.36$	90	26 BOEHM	01		Palo Verde react.
$< 0.45$	90	27 BOEHM	00		Palo Verde react.
$< 0.15$	90	28 APOLLONIO	99	CHOZ	Reactor Experiment

<sup>1</sup> ABE 12 determine the  $\bar{\nu}_e$  interaction rate in a single detector, located 1050 m from the cores of two reactors. The rate normalization is fixed by the results of the Bugey4 reactor experiment, thus avoiding any dependence on possible very short baseline oscillations. The value of  $\Delta m_{31}^2 = 2.4 \times 10^{-3} \text{ eV}^2$  is used in the analysis.

<sup>2</sup> AHN 12 use two identical detectors, placed at flux weighted distances of 408.56 m and 1433.99 m from six reactor cores, to determine the mixing angle  $\theta_{13}$ . This rate-only analysis excludes the no-oscillation hypothesis at 4.9 standard deviations. The value of  $\Delta m_{31}^2 = (2.32^{+0.12}_{-0.08}) \times 10^{-3} \text{ eV}^2$  was assumed in the analysis.

<sup>3</sup> AN 12 use six identical detectors with three placed near the reactor cores (flux-weighted baselines of 470 m and 576 m) and the remaining three at the far hall (at the flux averaged distance of 1648 m from all six reactor cores) to determine the mixing angle  $\theta_{13}$  using the  $\bar{\nu}_e$  observed interaction rate ratios. This rate-only analysis excludes the no-oscillation hypothesis at 5.2 standard deviations. The value of  $\Delta m_{31}^2 = (2.32^{+0.12}_{-0.08}) \times 10^{-3} \text{ eV}^2$  was assumed in the analysis.

<sup>4</sup> ABE 11 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{32}^2$  fixed to  $2.4 \times 10^{-3} \text{ eV}^2$ , using solar neutrino data including Super-Kamiokande, SNO, Borexino (ARPESELLA 08A), Homestake, GALLEX/GNO, SAGE, and KamLAND data. This result implies an upper bound of  $\sin^2\theta_{13} < 0.059$  (95% CL) or  $\sin^2 2\theta_{13} < 0.22$  (95% CL). The normal neutrino mass hierarchy and CPT invariance are assumed.

- 5 ABE 11 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{32}^2$  fixed to  $2.4 \times 10^{-3} \text{ eV}^2$ , using solar neutrino data including Super-Kamiokande, SNO, Borexino (ARPESELLA 08A), Homestake, and GALLEX/GNO data. The normal neutrino mass hierarchy is assumed.
- 6 The quoted limit is for  $\Delta m_{32}^2 = 2.4 \times 10^{-3} \text{ eV}^2$ ,  $\theta_{23} = \pi/2$ ,  $\delta = 0$ , and the normal mass hierarchy. For other values of  $\delta$ , the 68% region spans from 0.03 to 0.25, and the 90% region from 0.02 to 0.32.
- 7 The quoted limit is for  $\Delta m_{32}^2 = 2.4 \times 10^{-3} \text{ eV}^2$ ,  $\theta_{23} = \pi/2$ ,  $\delta = 0$ , and the inverted mass hierarchy. For other values of  $\delta$ , the 68% region spans from 0.04 to 0.30, and the 90% region from 0.02 to 0.39.
- 8 The quoted limit is for  $\Delta m_{32}^2 = 2.32 \times 10^{-3} \text{ eV}^2$ ,  $\theta_{23} = \pi/2$ ,  $\delta = 0$ , and the normal mass hierarchy. For other values of  $\delta$ , the 68% region spans from 0.02 to 0.12, and the 90% region from 0 to 0.16.
- 9 The quoted limit is for  $\Delta m_{32}^2 = 2.32 \times 10^{-3} \text{ eV}^2$ ,  $\theta_{23} = \pi/2$ ,  $\delta = 0$ , and the inverted mass hierarchy. For other values of  $\delta$ , the 68% region spans from 0.02 to 0.16, and the 90% region from 0 to 0.21.
- 10 FOGLI 11 obtained this result from an analysis using the atmospheric, accelerator long baseline, CHOOZ, solar, and KamLAND data. Recently, MUELLER 11 suggested an average increase of about 3.5% in normalization of the reactor  $\bar{\nu}_e$  fluxes, and using these fluxes, the fitted result becomes  $0.10 \pm 0.03$ .
- 11 GANDO 11 report  $\sin^2\theta_{13} = 0.020 \pm 0.016$ . This result was obtained with three-neutrino fit using the KamLAND + solar data.
- 12 GANDO 11 report  $\sin^2\theta_{13} = 0.032 \pm 0.037$ . This result was obtained with three-neutrino fit using the KamLAND data only.
- 13 This result corresponds to the limit of  $<0.12$  at 90% CL for  $\Delta m_{32}^2 = 2.43 \times 10^{-3} \text{ eV}^2$ ,  $\theta_{23} = \pi/2$ , and  $\delta = 0$ . For other values of  $\delta$ , the 90% CL region spans from 0 to 0.16.
- 14 This result corresponds to the limit of  $<0.20$  at 90% CL for  $\Delta m_{32}^2 = 2.43 \times 10^{-3} \text{ eV}^2$ ,  $\theta_{23} = \pi/2$ , and  $\delta = 0$ . For other values of  $\delta$ , the 90% CL region spans from 0 to 0.21.
- 15 AHARMIM 10 global solar neutrino data include SNO's low-energy-threshold analysis survival probability day/night curves, SNO Phase III integral rates (AHARMIM 08), CI (CLEVELAND 98), SAGE (ABDURASHITOV 09), Gallex/GNO (HAMPEL 99, ALTMANN 05), Borexino (ARPESELLA 08A), SK-I zenith (HOSAKA 06), and SK-II day/night spectra (CRAVENS 08).
- 16 AHARMIM 10 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{31}^2$  fixed to  $2.3 \times 10^{-3} \text{ eV}^2$ , using global solar neutrino data and KamLAND data (ABE 08A). *CPT* invariance is assumed. This result implies an upper bound of  $\sin^2\theta_{13} < 0.057$  (95% CL) or  $\sin^2 2\theta_{13} < 0.22$  (95% CL).
- 17 AHARMIM 10 obtained this result by a three-neutrino oscillation analysis with the value of  $\Delta m_{31}^2$  fixed to  $2.3 \times 10^{-3} \text{ eV}^2$ , using global solar neutrino data.
- 18 WENDELL 10 obtained this result by a three-neutrino oscillation analysis with one mass scale dominance ( $\Delta m_{21}^2 = 0$ ) using the Super-Kamiokande-I+II+III atmospheric neutrino data, and updates the HOSAKA 06A result.
- 19 The quoted limit is for  $\Delta m_{32}^2 = 2.43 \times 10^{-3} \text{ eV}^2$ ,  $\theta_{23} = \pi/2$ , and  $\delta = 0$ . For other values of  $\delta$ , the 68% CL region spans from 0.02 to 0.26.
- 20 The quoted limit is for  $\Delta m_{32}^2 = 2.43 \times 10^{-3} \text{ eV}^2$ ,  $\theta_{23} = \pi/2$ , and  $\delta = 0$ . For other values of  $\delta$ , the 68% CL region spans from 0.04 to 0.34.
- 21 FOGLI 08 obtained this result from a global analysis of all neutrino oscillation data, that is, solar + KamLAND + atmospheric + accelerator long baseline + CHOOZ.
- 22 FOGLI 08 obtained this result from an analysis using the solar and KamLAND neutrino oscillation data.
- 23 FOGLI 08 obtained this result from an analysis using the atmospheric, accelerator long baseline, and CHOOZ neutrino oscillation data.

- <sup>24</sup> YAMAMOTO 06 searched for  $\nu_\mu \rightarrow \nu_e$  appearance. Assumes  $2 \sin^2(2\theta_{\mu e}) = \sin^2(2\theta_{13})$ . The quoted limit is for  $\Delta m_{32}^2 = 1.9 \times 10^{-3} \text{ eV}^2$ . That value of  $\Delta m_{32}^2$  is the one- $\sigma$  low value for AHN 06A. For the AHN 06A best fit value of  $2.8 \times 10^{-3} \text{ eV}^2$ , the  $\sin^2(2\theta_{13})$  limit is  $< 0.26$ . Supersedes AHN 04.
- <sup>25</sup> AHN 04 searched for  $\nu_\mu \rightarrow \nu_e$  appearance. Assuming  $2 \sin^2(2\theta_{\mu e}) = \sin^2(2\theta_{13})$ , a limit on  $\sin^2(2\theta_{\mu e})$  is converted to a limit on  $\sin^2(2\theta_{13})$ . The quoted limit is for  $\Delta m_{32}^2 = 1.9 \times 10^{-3} \text{ eV}^2$ . That value of  $\Delta m_{32}^2$  is the one- $\sigma$  low value for ALIU 05. For the ALIU 05 best fit value of  $2.8 \times 10^{-3} \text{ eV}^2$ , the  $\sin^2(2\theta_{13})$  limit is  $< 0.30$ .
- <sup>26</sup> The quoted limit is for  $\Delta m_{32}^2 = 1.9 \times 10^{-3} \text{ eV}^2$ . That value of  $\Delta m_{32}^2$  is the 1- $\sigma$  low value for ALIU 05. For the ALIU 05 best fit value of  $2.8 \times 10^{-3} \text{ eV}^2$ , the  $\sin^2 2\theta_{13}$  limit is  $< 0.19$ . In this range, the  $\theta_{13}$  limit is larger for lower values of  $\Delta m_{32}^2$ , and smaller for higher values of  $\Delta m_{32}^2$ .
- <sup>27</sup> The quoted limit is for  $\Delta m_{32}^2 = 1.9 \times 10^{-3} \text{ eV}^2$ . That value of  $\Delta m_{32}^2$  is the 1- $\sigma$  low value for ALIU 05. For the ALIU 05 best fit value of  $2.8 \times 10^{-3} \text{ eV}^2$ , the  $\sin^2 2\theta_{13}$  limit is  $< 0.23$ .
- <sup>28</sup> The quoted limit is for  $\Delta m_{32}^2 = 2.43 \times 10^{-3} \text{ eV}^2$ . That value of  $\Delta m_{32}^2$  is the central value for ADAMSON 08. For the ADAMSON 08 1- $\sigma$  low value of  $2.30 \times 10^{-3} \text{ eV}^2$ , the  $\sin^2 2\theta_{13}$  limit is  $< 0.16$ . See also APOLLONIO 03 for a detailed description of the experiment.

### (C) Other neutrino mixing results

The LSND collaboration reported in AGUILAR 01 a signal which is consistent with  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$  oscillations. In a three neutrino framework, this would be a measurement of  $\theta_{12}$  and  $\Delta m_{21}^2$ . This does not appear to be consistent with the interpretation of other neutrino data. The MiniBooNE experiment, reported in AGUILAR-AREVALO 07, does a two-neutrino analysis which, assuming *CPT* conservation, rules out AGUILAR 01. The following listings include results which might be relevant towards understanding these observations. They include searches for  $\nu_\mu \rightarrow \nu_e$ ,  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ , sterile neutrino oscillations, and *CPT* violation.

#### $\Delta(m^2)$ for $\sin^2(2\theta) = 1$ ( $\nu_\mu \rightarrow \nu_e$ )

VALUE (eV <sup>2</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.034	90	AGUILAR-AR...07	MBOO	MiniBooNE
<0.0008	90	AHN 04	K2K	Water Cherenkov
<0.4	90	ASTIER 03	NOMD	CERN SPS
<2.4	90	AVVAKUMOV 02	NTEV	NUTEV FNAL
		<sup>1</sup> AGUILAR 01	LSND	$\nu_\mu \rightarrow \nu_e$ osc.prob.
0.03 to 0.3	95	<sup>2</sup> ATHANASSO...98	LSND	$\nu_\mu \rightarrow \nu_e$
<2.3	90	<sup>3</sup> LOVERRE 96		CHARM/CDHS
<0.9	90	VILAIN 94C	CHM2	CERN SPS
<0.09	90	ANGELINI 86	HLBC	BEBC CERN PS

- <sup>1</sup> AGUILAR 01 is the final analysis of the LSND full data set. Search is made for the  $\nu_\mu \rightarrow \nu_e$  oscillations using  $\nu_\mu$  from  $\pi^+$  decay in flight by observing beam-on electron events from  $\nu_e C \rightarrow e^- X$ . Present analysis results in  $8.1 \pm 12.2 \pm 1.7$  excess events in the  $60 < E_e < 200$  MeV energy range, corresponding to oscillation probability of  $0.10 \pm 0.16 \pm 0.04\%$ . This is consistent, though less significant, with the previous result of ATHANASSOPOULOS 98, which it supersedes. The present analysis uses selection criteria developed for the decay at rest region, and is less effective in removing the background above 60 MeV than ATHANASSOPOULOS 98.
- <sup>2</sup> ATHANASSOPOULOS 98 is a search for the  $\nu_\mu \rightarrow \nu_e$  oscillations using  $\nu_\mu$  from  $\pi^+$  decay in flight. The 40 observed beam-on electron events are consistent with  $\nu_e C \rightarrow e^- X$ ; the expected background is  $21.9 \pm 2.1$ . Authors interpret this excess as evidence for an oscillation signal corresponding to oscillations with probability  $(0.26 \pm 0.10 \pm 0.05)\%$ . Although the significance is only  $2.3 \sigma$ , this measurement is an important and consistent cross check of ATHANASSOPOULOS 96 who reported evidence for  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$  oscillations from  $\mu^+$  decay at rest. See also ATHANASSOPOULOS 98B.
- <sup>3</sup> LOVERRE 96 uses the charged-current to neutral-current ratio from the combined CHARM (ALLABY 86) and CDHS (ABRAMOWICZ 86) data from 1986.

### $\sin^2(2\theta)$ for "Large" $\Delta(m^2)$ ( $\nu_\mu \rightarrow \nu_e$ )

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 1.8	90	<sup>1</sup> AGUILAR-AR...07	MBOO	MiniBooNE
<110	90	<sup>2</sup> AHN 04	K2K	Water Cherenkov
< 1.4	90	ASTIER 03	NOMD	CERN SPS
< 1.6	90	AVVAKUMOV 02	NTEV	NUTEV FNAL
0.5 to 30	95	<sup>3</sup> AGUILAR 01	LSND	$\nu_\mu \rightarrow \nu_e$ osc.prob.
< 3.0	90	<sup>4</sup> ATHANASSO...98	LSND	$\nu_\mu \rightarrow \nu_e$
< 9.4	90	<sup>5</sup> LOVERRE 96		CHARM/CDHS
< 5.6	90	<sup>6</sup> VILAIN 94C	CHM2	CERN SPS

- <sup>1</sup> The limit is  $\sin^2 2\theta < 0.9 \times 10^{-3}$  at  $\Delta m^2 = 2 \text{ eV}^2$ . That value of  $\Delta m^2$  corresponds to the smallest mixing angle consistent with the reported signal from LSND in AGUILAR 01.
- <sup>2</sup> The limit becomes  $\sin^2 2\theta < 0.15$  at  $\Delta m^2 = 2.8 \times 10^{-3} \text{ eV}^2$ , the best-fit value of the  $\nu_\mu$  disappearance analysis in K2K.
- <sup>3</sup> AGUILAR 01 is the final analysis of the LSND full data set of the search for the  $\nu_\mu \rightarrow \nu_e$  oscillations. See footnote in preceding table for further details.
- <sup>4</sup> ATHANASSOPOULOS 98 report  $(0.26 \pm 0.10 \pm 0.05)\%$  for the oscillation probability; the value of  $\sin^2 2\theta$  for large  $\Delta m^2$  is deduced from this probability. See footnote in preceding table for further details, and see the paper for a plot showing allowed regions. If effect is due to oscillation, it is most likely to be intermediate  $\sin^2 2\theta$  and  $\Delta m^2$ . See also ATHANASSOPOULOS 98B.
- <sup>5</sup> LOVERRE 96 uses the charged-current to neutral-current ratio from the combined CHARM (ALLABY 86) and CDHS (ABRAMOWICZ 86) data from 1986.
- <sup>6</sup> VILAIN 94C limit derived by combining the  $\nu_\mu$  and  $\bar{\nu}_\mu$  data assuming  $CP$  conservation.

$\Delta(m^2)$  for  $\sin^2(2\theta) = 1$  ( $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ )

VALUE (eV <sup>2</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.03–0.09	90	<sup>1</sup> AGUILAR-AR...10	MBOO	$E_\nu > 475$ MeV
0.03–0.07	90	<sup>2</sup> AGUILAR-AR...10	MBOO	$E_\nu > 200$ MeV
<0.06	90	AGUILAR-AR...09B	MBOO	MiniBooNE
<0.055	90	<sup>3</sup> ARMBRUSTER02	KAR2	Liquid Sci. calor.
<2.6	90	AVVAKUMOV 02	NTEV	NUTEV FNAL
0.03–0.05		<sup>4</sup> AGUILAR 01	LSND	LAMPF
0.05–0.08	90	<sup>5</sup> ATHANASSO...96	LSND	LAMPF
0.048–0.090	80	<sup>6</sup> ATHANASSO...95		
<0.07	90	<sup>7</sup> HILL 95		
<0.9	90	VILAIN 94C	CHM2	CERN SPS
<0.14	90	<sup>8</sup> FREEDMAN 93	CNTR	LAMPF

<sup>1</sup> This value is for a two neutrino oscillation analysis for excess antineutrino events with  $E_\nu > 475$  MeV. The best fit is at 0.07. The allowed region is consistent with LSND reported by AGUILAR 01. Supersedes AGUILAR-AREVALO 09B.

<sup>2</sup> This value is for a two neutrino oscillation analysis for excess antineutrino events with  $E_\nu > 200$  MeV with subtraction of the expected 12 events low energy excess seen in the neutrino component of the beam. The best fit value is 0.007 for  $\Delta(m^2) = 4.4$  eV<sup>2</sup>.

<sup>3</sup> ARMBRUSTER 02 is the final analysis of the KARMEN 2 data for 17.7 m distance from the ISIS stopped pion and muon neutrino source. It is a search for  $\bar{\nu}_e$ , detected by the inverse  $\beta$ -decay reaction on protons and <sup>12</sup>C. 15 candidate events are observed, and  $15.8 \pm 0.5$  background events are expected, hence no oscillation signal is detected. The results exclude large regions of the parameter area favored by the LSND experiment.

<sup>4</sup> AGUILAR 01 is the final analysis of the LSND full data set. It is a search for  $\bar{\nu}_e$  30 m from LAMPF beam stop. Neutrinos originate mainly for  $\pi^+$  decay at rest.  $\bar{\nu}_e$  are detected through  $\bar{\nu}_e p \rightarrow e^+ n$  ( $20 < E_{e^+} < 60$  MeV) in delayed coincidence with  $np \rightarrow d\gamma$ . Authors observe  $87.9 \pm 22.4 \pm 6.0$  total excess events. The observation is attributed to  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$  oscillations with the oscillation probability of  $0.264 \pm 0.067 \pm 0.045\%$ , consistent with the previously published result. Taking into account all constraints, the most favored allowed region of oscillation parameters is a band of  $\Delta(m^2)$  from 0.2–2.0 eV<sup>2</sup>. Supersedes ATHANASSOPOULOS 95, ATHANASSOPOULOS 96, and ATHANASSOPOULOS 98.

<sup>5</sup> ATHANASSOPOULOS 96 is a search for  $\bar{\nu}_e$  30 m from LAMPF beam stop. Neutrinos originate mainly from  $\pi^+$  decay at rest.  $\bar{\nu}_e$  could come from either  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$  or  $\nu_e \rightarrow \bar{\nu}_e$ ; our entry assumes the first interpretation. They are detected through  $\bar{\nu}_e p \rightarrow e^+ n$  ( $20 \text{ MeV} < E_{e^+} < 60 \text{ MeV}$ ) in delayed coincidence with  $np \rightarrow d\gamma$ . Authors observe  $51 \pm 20 \pm 8$  total excess events over an estimated background  $12.5 \pm 2.9$ . ATHANASSOPOULOS 96B is a shorter version of this paper.

<sup>6</sup> ATHANASSOPOULOS 95 error corresponds to the  $1.6\sigma$  band in the plot. The expected background is  $2.7 \pm 0.4$  events. Corresponds to an oscillation probability of  $(0.34^{+0.20}_{-0.18} \pm 0.07)\%$ . For a different interpretation, see HILL 95. Replaced by ATHANASSOPOULOS 96.

<sup>7</sup> HILL 95 is a report by one member of the LSND Collaboration, reporting a different conclusion from the analysis of the data of this experiment (see ATHANASSOPOULOS 95). Contrary to the rest of the LSND Collaboration, Hill finds no evidence for the neutrino oscillation  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$  and obtains only upper limits.

<sup>8</sup> FREEDMAN 93 is a search at LAMPF for  $\bar{\nu}_e$  generated from any of the three neutrino types  $\nu_\mu$ ,  $\bar{\nu}_\mu$ , and  $\nu_e$  which come from the beam stop. The  $\bar{\nu}_e$ 's would be detected by the reaction  $\bar{\nu}_e p \rightarrow e^+ n$ . FREEDMAN 93 replaces DURKIN 88.

**$\sin^2(2\theta)$  for "Large"  $\Delta(m^2)$  ( $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ )**

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.4–9.0	99	<sup>1</sup> AGUILAR-AR...10	MBOO	$E_\nu > 475$ MeV
0.4–9.0	99	<sup>2</sup> AGUILAR-AR...10	MBOO	$E_\nu > 200$ MeV
<3.3	90	<sup>3</sup> AGUILAR-AR...09B	MBOO	MiniBooNE
<1.7	90	<sup>4</sup> ARMBRUSTER02	KAR2	Liquid Sci. calor.
<1.1	90	AVVAKUMOV 02	NTEV	NUTEV FNAL
$5.3 \pm 1.3 \pm 9.0$		<sup>5</sup> AGUILAR 01	LSND	LAMPF
$6.2 \pm 2.4 \pm 1.0$		<sup>6</sup> ATHANASSO...96	LSND	LAMPF
3–12	80	<sup>7</sup> ATHANASSO...95		
<6	90	<sup>8</sup> HILL 95		

<sup>1</sup> This value is for a two neutrino oscillation analysis for excess antineutrino events with  $E_\nu > 475$  MeV. At 90% CL there is no solution at high  $\Delta(m^2)$ . The best fit is at maximal mixing. The allowed region is consistent with LSND reported by AGUILAR 01. Supersedes AGUILAR-AREVALO 09B.

<sup>2</sup> This value is for a two neutrino oscillation analysis for excess antineutrino events with  $E_\nu > 200$  MeV with subtraction of the expected 12 events low energy excess seen in the neutrino component of the beam. At 90% CL there is no solution at high  $\Delta(m^2)$ . The best fit value is 0.007 for  $\Delta(m^2) = 4.4$  eV<sup>2</sup>.

<sup>3</sup> This result is inconclusive with respect to small amplitude mixing suggested by LSND.

<sup>4</sup> ARMBRUSTER 02 is the final analysis of the KARMEN 2 data. See footnote in the preceding table for further details, and the paper for the exclusion plot.

<sup>5</sup> AGUILAR 01 is the final analysis of the LSND full data set. The deduced oscillation probability is  $0.264 \pm 0.067 \pm 0.045\%$ ; the value of  $\sin^2 2\theta$  for large  $\Delta(m^2)$  is twice this probability (although these values are excluded by other constraints). See footnote in preceding table for further details, and the paper for a plot showing allowed regions. Supersedes ATHANASSOPOULOS 95, ATHANASSOPOULOS 96, and ATHANASSOPOULOS 98.

<sup>6</sup> ATHANASSOPOULOS 96 reports  $(0.31 \pm 0.12 \pm 0.05)\%$  for the oscillation probability; the value of  $\sin^2 2\theta$  for large  $\Delta(m^2)$  should be twice this probability. See footnote in preceding table for further details, and see the paper for a plot showing allowed regions.

<sup>7</sup> ATHANASSOPOULOS 95 error corresponds to the  $1.6\sigma$  band in the plot. The expected background is  $2.7 \pm 0.4$  events. Corresponds to an oscillation probability of  $(0.34^{+0.20}_{-0.18} \pm 0.07)\%$ . For a different interpretation, see HILL 95. Replaced by ATHANASSOPOULOS 96.

<sup>8</sup> HILL 95 is a report by one member of the LSND Collaboration, reporting a different conclusion from the analysis of the data of this experiment (see ATHANASSOPOULOS 95). Contrary to the rest of the LSND Collaboration, Hill finds no evidence for the neutrino oscillation  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$  and obtains only upper limits.

 **$\Delta(m^2)$  for  $\sin^2(2\theta) = 1$  ( $\nu_\mu(\bar{\nu}_\mu) \rightarrow \nu_e(\bar{\nu}_e)$ )**

VALUE (eV <sup>2</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.075</b>	90	BORODOV... 92	CNTR	BNL E776

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6	90	<sup>1</sup> ROMOSAN 97	CCFR	FNAL
------	----	-------------------------	------	------

<sup>1</sup> ROMOSAN 97 uses wideband beam with a 0.5 km decay region.

### $\sin^2(2\theta)$ for "Large" $\Delta(m^2)$ ( $\nu_\mu(\bar{\nu}_\mu) \rightarrow \nu_e(\bar{\nu}_e)$ )

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	<sup>1</sup> ROMOSAN	97	CCFR FNAL
<3.8	90	<sup>2</sup> MCFARLAND	95	CCFR FNAL
<3	90	BORODOV...	92	CNTR BNL E776

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup> ROMOSAN 97 uses wideband beam with a 0.5 km decay region.
- <sup>2</sup> MCFARLAND 95 state that "This result is the most stringent to date for  $250 < \Delta(m^2) < 450$  eV<sup>2</sup> and also excludes at 90%CL much of the high  $\Delta(m^2)$  region favored by the recent LSND observation." See ATHANASSOPOULOS 95 and ATHANASSOPOULOS 96.

### $\Delta(m^2)$ for $\sin^2(2\theta) = 1$ ( $\bar{\nu}_e \not\leftrightarrow \bar{\nu}_e$ )

VALUE (eV <sup>2</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	90	<sup>1</sup> ACHKAR	95	CNTR Bugey reactor

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup> ACHKAR 95 bound is for  $L=15, 40,$  and  $95$  m.

### $\sin^2(2\theta)$ for "Large" $\Delta(m^2)$ ( $\bar{\nu}_e \not\leftrightarrow \bar{\nu}_e$ )

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.02	90	<sup>1</sup> ACHKAR	95	CNTR For $\Delta(m^2) = 0.6$ eV <sup>2</sup>

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup> ACHKAR 95 bound is from data for  $L=15, 40,$  and  $95$  m distance from the Bugey reactor.

## ———— Sterile neutrino limits from atmospheric neutrino studies ————

### $\Delta(m^2)$ for $\sin^2(2\theta) = 1$ ( $\nu_\mu \rightarrow \nu_s$ )

$\nu_s$  means  $\nu_\tau$  or any sterile (noninteracting)  $\nu$ .

VALUE ( $10^{-5}$ eV <sup>2</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<3000 (or <550)	90	<sup>1</sup> OYAMA	89	KAMI Water Cherenkov
< 4.2 or > 54.	90	BIONTA	88	IMB Flux has $\nu_\mu, \bar{\nu}_\mu, \nu_e,$ and $\bar{\nu}_e$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup> OYAMA 89 gives a range of limits, depending on assumptions in their analysis. They argue that the region  $\Delta(m^2) = (100-1000) \times 10^{-5}$  eV<sup>2</sup> is not ruled out by any data for large mixing.

### Search for $\nu_\mu \rightarrow \nu_s$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
	<sup>1</sup> AMBROSIO 01	MCRO	matter effects
	<sup>2</sup> FUKUDA 00	SKAM	neutral currents + matter effects

<sup>1</sup> AMBROSIO 01 tested the pure 2-flavor  $\nu_\mu \rightarrow \nu_s$  hypothesis using matter effects which change the shape of the zenith-angle distribution of upward through-going muons. With maximum mixing and  $\Delta(m^2)$  around  $0.0024 \text{ eV}^2$ , the  $\nu_\mu \rightarrow \nu_s$  oscillation is disfavored with 99% confidence level with respect to the  $\nu_\mu \rightarrow \nu_\tau$  hypothesis.

<sup>2</sup> FUKUDA 00 tested the pure 2-flavor  $\nu_\mu \rightarrow \nu_s$  hypothesis using three complementary atmospheric-neutrino data samples. With this hypothesis, zenith-angle distributions are expected to show characteristic behavior due to neutral currents and matter effects. In the  $\Delta(m^2)$  and  $\sin^2 2\theta$  region preferred by the Super-Kamiokande data, the  $\nu_\mu \rightarrow \nu_s$  hypothesis is rejected at the 99% confidence level, while the  $\nu_\mu \rightarrow \nu_\tau$  hypothesis consistently fits all of the data sample.

————— **CPT tests** —————

**$\langle \Delta m_{21}^2 - \Delta \bar{m}_{21}^2 \rangle$**

VALUE ( $10^{-4} \text{ eV}^2$ )	CL%	DOCUMENT ID	TECN	COMMENT
----------------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.1	99.7	<sup>1</sup> DEGOUVEA 05	FIT	solar vs. reactor
------	------	--------------------------	-----	-------------------

<sup>1</sup> DEGOUVEA 05 obtained this bound at the  $3\sigma$  CL from the KamLAND (ARAKI 05) and solar neutrino data.

**REFERENCES FOR Neutrino Mixing**

ABE	12	PRL 108 131801	Y. Abe <i>et al.</i>	(Double Chooz Collab.)
AHN	12	PRL 108 191802	J.K. Ahn <i>et al.</i>	(RENO Collab.)
AN	12	PRL 108 171803	F.P. An <i>et al.</i>	(Daya Bay Collab.)
ABE	11	PR D83 052010	K. Abe <i>et al.</i>	(Super-Kamiokande Collab.)
ABE	11A	PRL 107 041801	K. Abe <i>et al.</i>	(T2K Collab.)
ABE	11B	PR C84 035804	S. Abe <i>et al.</i>	(KamLAND Collab.)
ABE	11C	PRL 107 241801	K. Abe <i>et al.</i>	(Super-Kamiokande Collab.)
ADAMSON	11	PRL 106 181801	P. Adamson <i>et al.</i>	(MINOS Collab.)
ADAMSON	11B	PRL 107 021801	P. Adamson <i>et al.</i>	(MINOS Collab.)
ADAMSON	11C	PR D84 071103	P. Adamson <i>et al.</i>	(MINOS Collab.)
ADAMSON	11D	PRL 107 181802	P. Adamson <i>et al.</i>	(MINOS Collab.)
BELLINI	11	PL B696 191	G. Bellini <i>et al.</i>	(Borexino Collab.)
BELLINI	11A	PRL 107 141302	G. Bellini <i>et al.</i>	(Borexino Collab.)
FOGLI	11	PR D84 053007	G.L. Fogli <i>et al.</i>	
GANDO	11	PR D83 052002	A. Gando <i>et al.</i>	(KamLAND Collab.)
MUELLER	11	PR C83 054615	Th.A Mueller <i>et al.</i>	
SERENELLI	11	APJ 743 24	A.M. Serenelli, W.C. Haxton, C. Pena-Garay	
ADAMSON	10A	PR D82 051102R	P. Adamson <i>et al.</i>	(MINOS Collab.)
AGUILAR-AR...	10	PRL 105 181801	A.A. Aguilar-Arevalo <i>et al.</i>	(MiniBooNE Collab.)
AHARMIM	10	PR C81 055504	B. Aharmim <i>et al.</i>	(SNO Collab.)
BELLINI	10A	PR D82 033006	G. Bellini <i>et al.</i>	(Borexino Collab.)
DENIZ	10	PR D81 072001	M. Deniz <i>et al.</i>	(TEXONO Collab.)
KAETHER	10	PL B685 47	F. Kaether <i>et al.</i>	
WENDELL	10	PR D81 092004	R. Wendell <i>et al.</i>	(Super-Kamiokande Collab.)
ABDURASHI...	09	PR C80 015807	J.N. Abdurashitov <i>et al.</i>	(SAGE Collab.)
ADAMSON	09	PRL 103 261802	P. Adamson <i>et al.</i>	(MINOS Collab.)
AGUILAR-AR...	09B	PRL 103 111801	A.A. Aguilar-arevalo <i>et al.</i>	(MiniBooNE Collab.)
ABE	08A	PRL 100 221803	S. Abe <i>et al.</i>	(KamLAND Collab.)
Also		PRL 101 119904E	S. Abe <i>et al.</i>	(KamLAND Collab.)
ADAMSON	08	PR D77 072002	P. Adamson <i>et al.</i>	(MINOS Collab.)
ADAMSON	08A	PRL 101 131802	P. Adamson <i>et al.</i>	(MINOS Collab.)
AHARMIM	08	PRL 101 111301	B. Aharmim <i>et al.</i>	(SNO Collab.)
ARPESELLA	08A	PRL 101 091302	C. Arpesella <i>et al.</i>	(Borexino Collab.)
CRAVENS	08	PR D78 032002	J.P. Cravens <i>et al.</i>	(Super-Kamiokande Collab.)
FOGLI	08	PRL 101 141801	G.L. Fogli, <i>et al.</i>	
ADAMSON	07	PR D75 092003	P. Adamson <i>et al.</i>	(MINOS Collab.)
AGUILAR-AR...	07	PRL 98 231801	A.A. Aguilar-Arevalo <i>et al.</i>	(MiniBooNE Collab.)
AHARMIM	07	PR C75 045502	B. Aharmim <i>et al.</i>	(SNO Collab.)
ADAMSON	06	PR D73 072002	P. Adamson <i>et al.</i>	(MINOS Collab.)

AHN	06A	PR D74 072003	M.H. Ahn <i>et al.</i>	(K2K Collab.)
BALATA	06	EPJ C47 21	M. Balata <i>et al.</i>	(Borexino Collab.)
HOSAKA	06	PR D73 112001	J. Hosaka <i>et al.</i>	(Super-Kamiokande Collab.)
HOSAKA	06A	PR D74 032002	J. Hosaka <i>et al.</i>	(Super-Kamiokande Collab.)
MICHAEL	06	PRL 97 191801	D. Michael <i>et al.</i>	(MINOS Collab.)
WINTER	06A	PR C73 025503	W.T. Winter <i>et al.</i>	
YAMAMOTO	06	PRL 96 181801	S. Yamamoto <i>et al.</i>	(K2K Collab.)
AHARMIM	05A	PR C72 055502	B. Aharmim <i>et al.</i>	(SNO Collab.)
ALIU	05	PRL 94 081802	E. Aliu <i>et al.</i>	(K2K Collab.)
ALLISON	05	PR D72 052005	W.W.M. Allison <i>et al.</i>	(SOUDAN-2 Collab.)
ALTMANN	05	PL B616 174	M. Altmann <i>et al.</i>	(GNO Collab.)
ARAKI	05	PRL 94 081801	T. Araki <i>et al.</i>	(KamLAND Collab.)
ASHIE	05	PR D71 112005	Y. Ashie <i>et al.</i>	(Super-Kamiokande Collab.)
DEGOUVEA	05	PR D71 093002	A. de Gouvea, C. Pena-Garay	
AHARMIM	04	PR D70 093014	B. Aharmim <i>et al.</i>	(SNO Collab.)
AHMED	04A	PRL 92 181301	S.N. Ahmed <i>et al.</i>	(SNO Collab.)
AHN	04	PRL 93 051801	M.H. Ahn <i>et al.</i>	(K2K Collab.)
AMBROSIO	04	EPJ C36 323	M. Ambrosio <i>et al.</i>	(MACRO Collab.)
ASHIE	04	PRL 93 101801	Y. Ashie <i>et al.</i>	(Super-Kamiokande Collab.)
EGUCHI	04	PRL 92 071301	K. Eguchi <i>et al.</i>	(KamLAND Collab.)
SMY	04	PR D69 011104R	M.B. Smy <i>et al.</i>	(Super-Kamiokande Collab.)
AHN	03	PRL 90 041801	M.H. Ahn <i>et al.</i>	(K2K Collab.)
AMBROSIO	03	PL B566 35	M. Ambrosio <i>et al.</i>	(MACRO Collab.)
APOLLONIO	03	EPJ C27 331	M. Apollonio <i>et al.</i>	(CHOOZ Collab.)
ASTIER	03	PL B570 19	P. Astier <i>et al.</i>	(NOMAD Collab.)
EGUCHI	03	PRL 90 021802	K. Eguchi <i>et al.</i>	(KamLAND Collab.)
GANDO	03	PRL 90 171302	Y. Gando <i>et al.</i>	(Super-Kamiokande Collab.)
IANNI	03	JPG 29 2107	A. Ianni	(INFN Gran Sasso)
SANCHEZ	03	PR D68 113004	M. Sanchez <i>et al.</i>	(Soudan 2 Collab.)
ABDURASHI...	02	JETP 95 181	J.N. Abdurashitov <i>et al.</i>	(SAGE Collab.)
		Translated from ZETF 122 211.		
AHMAD	02	PRL 89 011301	Q.R. Ahmad <i>et al.</i>	(SNO Collab.)
AHMAD	02B	PRL 89 011302	Q.R. Ahmad <i>et al.</i>	(SNO Collab.)
ARMBRUSTER	02	PR D65 112001	B. Armbruster <i>et al.</i>	(KARMEN 2 Collab.)
AVVAKUMOV	02	PRL 89 011804	S. Avvakumov <i>et al.</i>	(NuTeV Collab.)
FUKUDA	02	PL B539 179	S. Fukuda <i>et al.</i>	(Super-Kamiokande Collab.)
AGUILAR	01	PR D64 112007	A. Aguilar <i>et al.</i>	(LSND Collab.)
AHMAD	01	PRL 87 071301	Q.R. Ahmad <i>et al.</i>	(SNO Collab.)
AMBROSIO	01	PL B517 59	M. Ambrosio <i>et al.</i>	(MACRO Collab.)
BOEHM	01	PR D64 112001	F. Boehm <i>et al.</i>	
FUKUDA	01	PRL 86 5651	S. Fukuda <i>et al.</i>	(Super-Kamiokande Collab.)
AMBROSIO	00	PL B478 5	M. Ambrosio <i>et al.</i>	(MACRO Collab.)
BOEHM	00	PRL 84 3764	F. Boehm <i>et al.</i>	
FUKUDA	00	PRL 85 3999	S. Fukuda <i>et al.</i>	(Super-Kamiokande Collab.)
ALLISON	99	PL B449 137	W.W.M. Allison <i>et al.</i>	(Soudan 2 Collab.)
APOLLONIO	99	PL B466 415	M. Apollonio <i>et al.</i>	(CHOOZ Collab.)
Also		PL B472 434 (erratum)	M. Apollonio <i>et al.</i>	(CHOOZ Collab.)
FUKUDA	99C	PRL 82 2644	Y. Fukuda <i>et al.</i>	(Super-Kamiokande Collab.)
FUKUDA	99D	PL B467 185	Y. Fukuda <i>et al.</i>	(Super-Kamiokande Collab.)
HAMPEL	99	PL B447 127	W. Hampel <i>et al.</i>	(GALLEX Collab.)
AMBROSIO	98	PL B434 451	M. Ambrosio <i>et al.</i>	(MACRO Collab.)
APOLLONIO	98	PL B420 397	M. Apollonio <i>et al.</i>	(CHOOZ Collab.)
ATHANASSO...	98	PRL 81 1774	C. Athanassopoulos <i>et al.</i>	(LSND Collab.)
ATHANASSO...	98B	PR C58 2489	C. Athanassopoulos <i>et al.</i>	(LSND Collab.)
CLEVELAND	98	APJ 496 505	B.T. Cleveland <i>et al.</i>	(Homestake Collab.)
FELDMAN	98	PR D57 3873	G.J. Feldman, R.D. Cousins	
FUKUDA	98C	PRL 81 1562	Y. Fukuda <i>et al.</i>	(Super-Kamiokande Collab.)
HATAKEYAMA	98	PRL 81 2016	S. Hatakeyama <i>et al.</i>	(Kamiokande Collab.)
CLARK	97	PRL 79 345	R. Clark <i>et al.</i>	(IMB Collab.)
ROMOSAN	97	PRL 78 2912	A. Romosan <i>et al.</i>	(CCFR Collab.)
AGLIETTA	96	JETPL 63 791	M. Aglietta <i>et al.</i>	(LSD Collab.)
		Translated from ZETFP 63 753.		
ATHANASSO...	96	PR C54 2685	C. Athanassopoulos <i>et al.</i>	(LSND Collab.)
ATHANASSO...	96B	PRL 77 3082	C. Athanassopoulos <i>et al.</i>	(LSND Collab.)
FUKUDA	96	PRL 77 1683	Y. Fukuda <i>et al.</i>	(Kamiokande Collab.)
FUKUDA	96B	PL B388 397	Y. Fukuda <i>et al.</i>	(Kamiokande Collab.)
GREENWOOD	96	PR D53 6054	Z.D. Greenwood <i>et al.</i>	(UCI, SVR, SCUC)
HAMPEL	96	PL B388 384	W. Hampel <i>et al.</i>	(GALLEX Collab.)
LOVERRE	96	PL B370 156	P.F. Loverre	
ACHKAR	95	NP B434 503	B. Achkar <i>et al.</i>	(SING, SACL D, CPPM, CDEF+)
AHLEN	95	PL B357 481	S.P. Ahlen <i>et al.</i>	(MACRO Collab.)

ATHANASSO...	95	PRL 75 2650	C. Athanassopoulos <i>et al.</i>	(LSND Collab.)
DAUM	95	ZPHY C66 417	K. Daum <i>et al.</i>	(FREJUS Collab.)
HILL	95	PRL 75 2654	J.E. Hill	(PENN)
MCFARLAND	95	PRL 75 3993	K.S. McFarland <i>et al.</i>	(CCFR Collab.)
DECLAIS	94	PL B338 383	Y. Declais <i>et al.</i>	
FUKUDA	94	PL B335 237	Y. Fukuda <i>et al.</i>	(Kamiokande Collab.)
VILAIN	94C	ZPHY C64 539	P. Vilain <i>et al.</i>	(CHARM II Collab.)
FREEDMAN	93	PR D47 811	S.J. Freedman <i>et al.</i>	(LAMPF E645 Collab.)
BECKER-SZ...	92B	PR D46 3720	R.A. Becker-Szendy <i>et al.</i>	(IMB Collab.)
BEIER	92	PL B283 446	E.W. Beier <i>et al.</i>	(KAM2 Collab.)
Also		PTRSL A346 63	E.W. Beier, E.D. Frank	(PENN)
BORODOV...	92	PRL 68 274	L. Borodovsky <i>et al.</i>	(COLU, JHU, ILL)
HIRATA	92	PL B280 146	K.S. Hirata <i>et al.</i>	(Kamiokande II Collab.)
CASPER	91	PRL 66 2561	D. Casper <i>et al.</i>	(IMB Collab.)
HIRATA	91	PRL 66 9	K.S. Hirata <i>et al.</i>	(Kamiokande II Collab.)
KUVSHINN...	91	JETPL 54 253	A.A. Kuvshinnikov <i>et al.</i>	(KIAE)
BERGER	90B	PL B245 305	C. Berger <i>et al.</i>	(FREJUS Collab.)
HIRATA	90	PRL 65 1297	K.S. Hirata <i>et al.</i>	(Kamiokande II Collab.)
AGLIETTA	89	EPL 8 611	M. Aglietta <i>et al.</i>	(FREJUS Collab.)
DAVIS	89	ARNPS 39 467	R. Davis, A.K. Mann, L. Wolfenstein	(BNL, PENN+)
OYAMA	89	PR D39 1481	Y. Oyama <i>et al.</i>	(Kamiokande II Collab.)
BIONTA	88	PR D38 768	R.M. Bionta <i>et al.</i>	(IMB Collab.)
DURKIN	88	PRL 61 1811	L.S. Durkin <i>et al.</i>	(OSU, ANL, CIT+)
ABRAMOWICZ	86	PRL 57 298	H. Abramowicz <i>et al.</i>	(CDHS Collab.)
ALLABY	86	PL B177 446	J.V. Allaby <i>et al.</i>	(CHARM Collab.)
ANGELINI	86	PL B179 307	C. Angelini <i>et al.</i>	(PISA, ATHU, PADO+)
VUILLEUMIER	82	PL 114B 298	J.L. Vuilleumier <i>et al.</i>	(CIT, SIN, MUNI)
BOLIEV	81	SJNP 34 787	M.M. Boliev <i>et al.</i>	(INRM)
		Translated from YAF 34 1418.		
KWON	81	PR D24 1097	H. Kwon <i>et al.</i>	(CIT, ISNG, MUNI)
BOEHM	80	PL 97B 310	F. Boehm <i>et al.</i>	(ILLG, CIT, ISNG, MUNI)
CROUCH	78	PR D18 2239	M.F. Crouch <i>et al.</i>	(CASE, UCI, WITW)