

- 12 From the fit to $e^+e^- \rightarrow \pi^+\pi^-$ data from the compilations of HEYN 81 and BARKOV 85, including the GOUNARIS 68 parametrization of the pion form factor.
 13 A fit of BARKOV 85 data assuming the direct $\omega\pi\pi$ coupling.
 14 Applying the S-matrix formalism to the BARKOV 85 data.
 15 Includes BARKOV 85 data. Model-dependent width definition.
 16 $|F_\pi(0)|^2$ fixed to 1.
 17 From the GOUNARIS 68 parametrization of the pion form factor.
 18 The error combines statistical and systematic uncertainties. Supersedes BARATE 97M.
 19 $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.
 20 From the GOUNARIS 68 parametrization of the pion form factor. The second error is a model error taking into account different parametrizations of the pion form factor.
 21 Using the data of BARATE 97M and the effective chiral Lagrangian.
 22 From a fit of the model-independent parameterization of the pion form factor to the data of BARATE 97M.
 23 Assuming the equality of ρ^+ and ρ^- masses and widths.
 24 Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.
 25 Phase shift analysis. Systematic errors added corresponding to spread of different fits.
 26 From fit of 3-parameter relativistic P -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.
 27 From the parametrization according to SOEDING 66.
 28 From the parametrization according to ROSS 66.
 29 HEYN 81 includes all spacelike and timelike F_π values until 1978.
 30 From pole extrapolation.
 31 From phase shift analysis of GRAYER 74 data.
 32 Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, GOLDHABER 64, ABOLINS 63.
 33 Breit-Wigner mass from a phase-shift analysis of HYAMS 73 and PROTOPOPESCU 73 data.
 34 Using relativistic Breit-Wigner and taking into account $\rho\omega$ interference.
 35 Systematic errors not evaluated.
 36 Systematic effects not studied.
 37 From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P -wave intensity. CHABAUD 83 includes data of GRAYER 74.
-

$m_{\rho(770)^0} - m_{\rho(770)^\pm}$


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-0.7 ± 0.8 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.				
-2.4 \pm 0.8		38 SCHAEL	05C ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	
0.4 \pm 0.7 \pm 0.6	1.98M	39 ALOISIO	03 KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	
1.3 \pm 1.1 \pm 2.0	500k	39 ACHASOV	02 SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	
1.6 \pm 0.6 \pm 1.7	600k	ABELE	99E CBAR	$0 \pm 0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$	
-4 \pm 4	3000	40 REYNOLDS	69 HBC	-0 2.26 $\pi^- p$	
-5 \pm 5	3600	40 FOSTER	68 HBC	$\pm 0 0.0 \bar{p}p$	
2.4 \pm 2.1	22950	41 PISUT	68 RVUE	$\pi N \rightarrow \rho N$	

³⁸ From the combined fit of the τ^- data from ANDERSON 00A and SCHael 05C and $e^+ e^-$ data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. Supersedes BARATE 97M.

³⁹ Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

⁴⁰ From quoted masses of charged and neutral modes.

⁴¹ Includes MALAMUD 69, ARMENISE 68, BATON 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65, CARMONY 64, GOLDHABER 64, ABOLINS 63.



$m_{\rho(770)^+} - m_{\rho(770)^-}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5±0.8±0.7 1.98M ⁴² ALOISIO 03 KLOE 1.02 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

⁴² Without limitations on masses and widths.

Γ_{12}	$e^+ e^-$	[a]	(4.72 ± 0.05) $\times 10^{-5}$
Γ_{13}	$\pi^+ \pi^- \pi^0$		($1.01^{+0.54}_{-0.36} \pm 0.34$) $\times 10^{-4}$
Γ_{14}	$\pi^+ \pi^- \pi^+ \pi^-$		(1.8 ± 0.9) $\times 10^{-5}$
Γ_{15}	$\pi^+ \pi^- \pi^0 \pi^0$		(1.6 ± 0.8) $\times 10^{-5}$
Γ_{16}	$\pi^0 e^+ e^-$	< 1.2	$\times 10^{-5}$ CL=90%
Γ_{17}	$\eta e^+ e^-$		

[a] The $\omega \rho$ interference is then due to $\omega \rho$ mixing only, and is expected to be small. If $e\mu$ universality holds, $\Gamma(\rho^0 \rightarrow \mu^+ \mu^-) = \Gamma(\rho^0 \rightarrow e^+ e^-) \times 0.99785$.

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 10 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 10.7$ for 8 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|cc} x_3 & -100 & \\ \hline \Gamma & 15 & -15 \\ \hline x_2 & x_3 \end{array}$$

	Mode	Rate (MeV)	Scale factor
Γ_2	$\pi^\pm \pi^0$	150.2 ± 2.4	
Γ_3	$\pi^\pm \gamma$	0.068 ± 0.007	2.3

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 7 branching ratios uses 21 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 6.0$ for 13 degrees of freedom.


The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_7	-100							
x_8	-5	0						
x_9	-1	0	1					
x_{10}	-1	0	0	0				
x_{11}	2	-3	0	0	0			
x_{12}	0	0	-8	-9	0	0		
x_{14}	-1	0	0	0	0	0	0	
Γ	0	0	4	5	0	0	-54	
	x_6	x_7	x_8	x_9	x_{10}	x_{11}	x_{12}	x_{14}

	Mode	Rate (MeV)	
Γ_6	$\pi^+ \pi^-$	147.5	± 0.9
Γ_7	$\pi^+ \pi^- \gamma$	1.48	± 0.24
Γ_8	$\pi^0 \gamma$	0.089	± 0.012
Γ_9	$\eta \gamma$	0.0447	± 0.0031
Γ_{10}	$\pi^0 \pi^0 \gamma$	0.0066	± 0.0012
Γ_{11}	$\mu^+ \mu^-$	[a]	0.0068 ± 0.0004
Γ_{12}	$e^+ e^-$	[a]	0.00704 ± 0.00006
Γ_{14}	$\pi^+ \pi^- \pi^+ \pi^-$	0.0027	± 0.0014

$\rho(770)$ PARTIAL WIDTHS

$\Gamma(\pi^\pm \gamma)$	Γ_3
VALUE (keV)	DOCUMENT ID TECN CHG COMMENT
68 ± 7 OUR FIT	Error includes scale factor of 2.3.
68 ± 7 OUR AVERAGE	Error includes scale factor of 2.2. See the ideogram below.
81 ± 4 ± 4	CAPRARO 87 SPEC - 200 $\pi^- A \rightarrow \pi^- \pi^0 A$
59.8 ± 4.0	HUSTON 86 SPEC + 202 $\pi^+ A \rightarrow \pi^+ \pi^0 A$
71 ± 7	JENSEN 83 SPEC - 156–260 $\pi^- A \rightarrow \pi^- \pi^0 A$



$\Gamma(e^+ e^-)$

Γ_{12}

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
7.04 \pm0.06 OUR FIT				
7.04 \pm0.06 OUR AVERAGE				
7.048 \pm 0.057 \pm 0.050	900k	80 AKHMETSHIN 07	e ⁺ e ⁻ \rightarrow $\pi^+ \pi^-$	
7.06 \pm 0.11 \pm 0.05	114k	81,82 AKHMETSHIN 04	CMD2 e ⁺ e ⁻ \rightarrow $\pi^+ \pi^-$	
6.77 \pm 0.10 \pm 0.30		BARKOV 85	OLYA e ⁺ e ⁻ \rightarrow $\pi^+ \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.12 \pm 0.02 \pm 0.11	800k	83 ACHASOV 06	SND e ⁺ e ⁻ \rightarrow $\pi^+ \pi^-$	
6.3 \pm 0.1		84 BENAYOUN 98	RVUE e ⁺ e ⁻ \rightarrow $\pi^+ \pi^-$, $\mu^+ \mu^-$	

$\Gamma(\pi^0 \gamma)$

Γ_8

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
77 \pm 17 \pm 11	36500	85 ACHASOV 03	SND $0.60\text{--}0.97$ e ⁺ e ⁻ \rightarrow $\pi^0 \gamma$	
121 \pm 31		DOLINSKY 89	ND e ⁺ e ⁻ \rightarrow $\pi^0 \gamma$	

$\Gamma(\eta \gamma)$

Γ_9

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
62 \pm 17	86 DOLINSKY 89	ND e ⁺ e ⁻ \rightarrow $\eta \gamma$	

